



LUND UNIVERSITY

Prostaglandin D(2) induces contractions through activation of TP receptors in peripheral lung tissue from the guinea pig.

Larsson Callerfelt, Anna-Karin; Hagfjärd, Annika; Dahlén, Sven-Erik; Adner, Mikael

Published in:
European Journal of Pharmacology

DOI:
[10.1016/j.ejphar.2011.07.046](https://doi.org/10.1016/j.ejphar.2011.07.046)

2011

[Link to publication](#)

Citation for published version (APA):
Larsson Callerfelt, A-K., Hagfjärd, A., Dahlén, S-E., & Adner, M. (2011). Prostaglandin D(2) induces contractions through activation of TP receptors in peripheral lung tissue from the guinea pig. *European Journal of Pharmacology*, 669(1-3), 136-142. <https://doi.org/10.1016/j.ejphar.2011.07.046>

Total number of authors:
4

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Prostaglandin D₂ induces contractions through activation of TP receptors in peripheral lung tissue from the guinea pig

Anna-Karin Larsson^{a, b*}, Annika Hagfjård^a, Sven-Erik Dahlén^a and Mikael Adner^a

^a Experimental Asthma and Allergy Research, The Institute of Environmental Medicine, Karolinska Institutet, 171 77 Stockholm, ^b Lung Biology, Department of Experimental Medical Science, Lund University, 221 84 Lund, Sweden

E-mail addresses:

Anna-Karin.Larsson@med.lu.se

Annika.Hagfjard@gmail.com

Sven-Erik.Dahlen@ki.se

Mikael.Adner@ki.se

** Corresponding author:*

Anna-Karin Larsson, Lung Biology, Department of Experimental Medical Science, BMC D12, Lund University, 221 84 Lund, Sweden. Tel: +46 462229441

E-mail: Anna-Karin_L.Larsson@med.lu.se

1 **Abstract**

2 Prostaglandin D₂ (PGD₂), released through mast cell activation, is used as a non-invasive
3 biomarker in patients with asthma. Since PGD₂ can elicit opposing effects on airway tone via
4 activation of the PGD₂ receptors DP₁ and DP₂ as well as the thromboxane receptor TP, the
5 aim of this study was to characterize the receptors that are activated by PGD₂ in the guinea
6 pig lung parenchyma. PGD₂ and the thromboxane analogue U46619 induced concentration-
7 dependent contractions. U46619 was more potent and caused stronger effect than PGD₂. The
8 specific TP receptor antagonist SQ-29548 and the combined TP and DP₂ receptor antagonist
9 BAYu3405 concentration-dependently shifted the curves for both agonists to the right. The
10 DP₁ receptor agonist BW245 induced a weak relaxation at high concentrations, whereas the
11 DP₁ receptor antagonist BWA868C did not affect the PGD₂ induced contractions. The
12 specific DP₂ receptor agonist 13,14-dihydro-15-keto -PGD₂ showed neither contractile nor
13 relaxant effect in the parenchyma. Furthermore, studies in precision-cut lung slices specified
14 that airways as well as pulmonary arteries and veins contracted to both PGD₂ and U46619.
15 When the lung parenchyma from ovalbumin sensitized guinea pigs were exposed to
16 ovalbumin, both thromboxane B₂ and PGD₂ were released. Ovalbumin also induced maximal
17 contractions at similar level as PGD₂ in the parenchyma, which was partly reduced by SQ-
18 29548. These data show that PGD₂ should be recognized as a TP receptor agonist in the
19 peripheral lung inducing contraction on airways, arteries and veins. Therefore, a TP receptor
20 antagonist can be useful in combination treatment of allergic responses in asthma.

21

22

23 **Keywords:** Guinea pig; lung parenchyma; ovalbumin; precision cut lung slices; prostaglandin
24 D₂; thromboxane

25 **1. Introduction**

26 Prostaglandin D₂ (PGD₂), secreted during mast cell activation (Dahlen and Kumlin, 2004),
27 and its metabolite 9 α -11 β PGF_{2 α} are used as a non-invasive biomarkers in patients with
28 asthma (O'Sullivan et al., 1996). PGD₂ is part of the acute asthmatic airway response; levels
29 of this mediator can be found within minutes in BAL fluid and at 150-fold higher biologically
30 active levels than before the exacerbation (Liu et al., 1991). In addition to the acute reaction,
31 PGD₂ has through recruitment of inflammatory cells been suggested to contribute to the
32 formation of the chronic asthmatic inflammation and subsequent airway remodelling (Balzar
33 et al., 2011).

34

35 PGD₂, generated from arachidonic acid, is converted to PG via cyclooxygenase (COX) (Vane,
36 1971) and PGD synthase (Urade and Eguchi, 2002). There are two distinct types of PGDS;
37 hematopoietic (H-PGDS) and lipocalin-type (L-PGDS). H-PGDS is highly expressed in mast
38 cells, eosinophils, macrophages, and lymphocytes as well as structural cells such as epithelial
39 cells and fibroblasts, whereas L-PGDS is mainly expressed in the central nervous system and
40 heart (Okano et al., 2006). PGD₂ exits the cell via a carrier-mediated process and activates
41 specific G-protein coupled receptors on target cells. PGD₂ is classified to mediate its effect via
42 the DP₁ (Coleman et al., 1994) and the DP₂ (CRTH₂) receptors (Abe et al., 1999), but also
43 known to act via the receptor for tromboxane A₂ (TXA₂), the TP receptor (Hamid-Bloomfield
44 et al., 1990). The DP₁ receptor is widely distributed in airway and vascular smooth muscle,
45 blood platelets, airway epithelium and nervous tissue (Coleman et al., 1994; Matsuoka et al.,
46 2000; Norel et al., 1999). PGD₂ has also an important chemotactic role via activation of the
47 DP₂ receptor (Abe et al., 1999), which is mainly expressed on Th2 cells and eosinophils (Abe
48 et al., 1999) but also on human airway smooth muscle (Abe et al., 1999; Parameswaran et al.,
49 2007). The TP receptors are expressed on bronchial and vascular smooth muscle cells, blood

50 platelets and myofibroblasts (Capra et al., 2003; Coleman et al., 1994) and are known to
51 mediate a strong and long-lasting contraction in these tissues (Held et al., 1999; Ressmeyer et
52 al., 2006). PGD₂ may thus have broad actions since activation of multiple receptors can elicit
53 theoretically opposing effects on airway tone.

54

55 Although the lung parenchyma is a complex tissue, the action in the peripheral lung is of
56 importance to study since asthma is suggested to be a disease of the small airways (van den
57 Berge et al., 2011). Especially the action of PGD₂ is of interest since it has been shown that
58 mast cells are located peripherally around small bronchi, vessels and further out to the alveoli
59 (Andersson et al., 2009) and thus may not only affect airways. The aim of this study was
60 therefore to characterize the receptors that are activated by PGD₂ in the peripheral lung and
61 subsequently investigate the significance of this effect in allergen-induced contractions. The
62 guinea pig parenchyma is particularly suitable as it has been shown to respond to many
63 agonists similar to human (Canning and Chou, 2008; Ressmeyer et al., 2006).

64 **2. Methods**

65 *2.1. Animals and ovalbumin-sensitization*

66 Male Dunkin Hartley guinea pigs (300–350 g b.w.) were used. In one part of the experiments
67 the guinea pigs were sensitized to ovalbumin at least four weeks prior to experiments as
68 previously described (Larsson et al., 2005). The study was approved by the regional
69 committee of animal experimentation ethics (N127/04, N63/07).

70

71 *2.2. Lung parenchymal strips and organ bath experiments*

72 The animals were sacrificed by an overdose of inhaled CO₂ and the heart-lung-package was
73 quickly removed and placed in ice-cold Tyrode's solution (prepared each day, containing
74 NaCl 149.2 mM, KCl 2.7 mM, NaHCO₃ 11.9 mM, glucose 5.5 mM, CaCl₂ 1.8 mM, MgCl₂
75 0.5 mM, NaH₂PO₄ 0.4 mM). The lung parenchyma was cut parallel to the peripheral
76 margins, yielding four to eight strips, each having a size of 2×2×20 mm and a weight of
77 approximately 60 mg. The parenchymal strips were set up at a resting tension of 4.0 mN in 5
78 ml organ baths filled with Tyrode's solution, bubbled with carbogen gas (6.5% CO₂ in O₂) to
79 keep a pH of 7.4 at 37°C. Changes in smooth muscle tension, contractions and relaxations,
80 were recorded via isometric force-displacement transducers connected to a Grass polygraph.
81 After an equilibration period of 90 min and washes each 15 min, histamine was added as a
82 control of the parenchymal strip reactivity. Preparations displaying contraction responses less
83 than 1.0 mN to 30 μM of histamine were excluded from further experiments. Another wash
84 and equilibration period between histamine and treatment period was performed. All
85 antagonists were given 15 min before the challenges. For study the effect of different
86 agonists, the parenchymal strip was exposed to cumulative concentrations. To study the
87 relaxation the parenchyma was pre-contracted with 10 nM of LTD₄ generating a 50%
88 contraction. To study the allergic early phase reaction, ovalbumin was added as cumulative

89 challenge of increasing concentrations every 10 min without changing bath fluid. Maximum
90 contractions of the preparation were determined with histamine (1 mM), acetylcholine (1
91 mM) and potassium chloride (KCl; 50 mM) at the end of each experiment, and other
92 responses were expressed as percent of maximum contractions.

93

94 *2.3. Measurements of released mediators with enzyme immunoassays*

95 A 1 mL aliquot of organ bath fluid was collected from each organ bath and immediately
96 frozen at -20°C. The samples were taken at the end of the equilibration period to obtain basal
97 mediator release from the tissue and at the obtained contractile plateau after challenge with
98 ovalbumin 1000 ng/ml. Enzyme immunoassay (EIA) analyses of the prostanoids TXA₂ and
99 PGD₂ were performed according to the manufacturer's instructions. TXA₂ was measured as
100 the stable metabolite TXB₂. PGD₂ was measured as PGD₂-mox. The assay detection limits in
101 the bath fluid levels were 7.8 pg/ml. The EIA specificity for the different mediators to
102 interfere with each other was less than 0.01%, with the exception of the EIA kit for TXB₂ that
103 cross reacted with PGD₂ (0.53%).

104

105 *2.4. Precision-cut lung slices*

106 Guinea pig precision-cut lung slices were prepared as previously described (Ressmeyer et al.,
107 2006). Briefly, the lung was filled through the trachea with a low melting-point agarose
108 solution (0.75%) containing salbutamol (1 mM). Lung tissue cores were prepared and cut into
109 220-mm-thick slices with a Krumdieck tissue slicer (Alabama Research and Development,
110 Munford, AL, USA). Tissue slices were incubated at 37°C in a humid atmosphere in minimal
111 essential medium supplemented with sodium pyruvate, amino acids, vitamins and glutamine.
112 The medium was changed on a regular basis during four hours in order to remove the agarose
113 and cell debris from the tissue. Salbutamol was added to the medium during the first three

114 hours. The slices were then imaged using an analogue (JAI 2040; JAI Pulnix, Alzenau,
115 Germany) or digital camera (IRB640; Visitron Systems, Munich, Germany). For
116 measurements, slices with comparable airway and vessel size were selected.
117 Bronchoconstriction or vasoconstriction was expressed as airway/vessel area as the
118 percentage of the initial area. A control image was taken before cumulative addition of
119 U46619 or PGD₂ and frames were recorded every 30 sec for 20 min.

120

121 *2.5. Data analysis and statistical procedures*

122 All data are presented as mean \pm standard error of the mean (S.E.M.). Statistical analyses
123 were made for paired and unpaired observations by Student's t-test or analyses of variances
124 (ANOVA) followed by the post hoc tests Bonferroni's t-test. A P-value of less than 0.05 was
125 considered significant. To provide estimates of maximal effect (E_{\max}), midpoint location
126 (pEC_{50}) and Hill slope (n_H), agonist concentration-effect curve data from individual tissues
127 were fitted to the Hill equation using an iterative, least square method (GraphPad Prism, San
128 Diego, USA). For the Schild plot analysis, the concentration-response curves with antagonists
129 were set at a global shared maximum assuming competitive antagonism. If the value of the
130 slope was found not to be significantly different from unity a second fit was performed for the
131 calculation of pK_B values with the slope constrained to unity.

132

133 *2.6. Drugs and chemical reagents*

134 NaCl, KCl, CaCl₂, MgSO₄, NaHCO₃, KH₂PO₄ and glucose were obtained from VWR
135 International (West Chester, Pennsylvania, USA). Histamine dihydrochloride, acetylcholine,
136 ovalbumin (chicken egg albumin, grade II), agarose, salbutamol, dimethylsulfoxid (DMSO)
137 were purchased from Sigma-Aldrich (St. Louis, Missouri, USA). 3R-[[4-
138 fluorophenyl)sulfonyl]amino]-1,2,3,4-tetrahydro-9H-carbazole-9-propanoic acid (BAYu3405,

139 Ramatroban) was purchased from Bayer AG (Wuppertal, Germany). PGD₂, (4S)-(3-[(3R,S)-
140 3-cyclohexyl-3-hydroxypropyl]-2,5-dioxo)-4-imidazolidineheptanoic acid (BW245C), LTD₄,
141 9,11-dideoxy-9 α ,11 α -methanoepoxy PGF_{2 α} (U46619), PGF_{2 α} , [1S-[1 α ,2 α (Z),3 α ,4 α]]-7-[3-[[2-
142 [(phenylamino)carbonyl]hydrazino]methyl]-7-oxabicyclo[2.2.1]hept-2-yl]-5-heptenoic acid
143 (SQ-29548), 3-[(2-cyclohexyl-2-hydroxyethyl)amino]-2,5-dioxo-1-(phenylmethyl)-4-
144 imidazolidineheptanoic acid (BWA868C), 9 α ,11 β -PGF₂ and 13,14-dihydro-15-keto (DK)-
145 PGD₂ were bought from Cayman Chemical (Ann Arbor, MI, USA). LTD₄ was from Cascade
146 Biochemicals Ltd. (Reading, UK). The EIA kits for TXB₂ and PGD₂-mox were obtained from
147 Cayman Chemicals (Ann Arbor, Michigan, USA). Stock solutions of 1 mM LTD₄ and
148 prostanoids were dissolved in 50% ethanol-water and then diluted in 20% ethanol-water.
149 Ovalbumin was dissolved in 0.9% NaCl. The other drugs were dissolved and diluted in
150 Tyrode's solution or millipure water. Dilutions of drugs were freshly made from the stocks for
151 each experiment. The drugs were present in the organ bath fluid during the remaining
152 experiment. 0.1% DMSO was added as a control and did not influence the baseline or
153 cumulative contractions to ovalbumin.

154 **3. Results**

155 *3.1. The TP receptor is the main contractile receptor for PGD₂, TXA₂ and PGF_{2α}*

156 PGD₂ induced concentration-dependent contractions in the lung parenchyma (Fig. 1A and B).
157 The contraction ($79.8 \pm 8.5\%$), obtained at the highest concentration used (100 μM), was
158 according to the non-linear regression analysis not the maximal effect. The inability of PGD₂
159 to reach the maximum capacity of the tissue contrasted to U46619 (Fig. 1C and D), which
160 induced a contraction reaching similar or higher maximum effect as the high concentrations
161 of histamine, acetylcholine and potassium chloride used as the reference at the end of each
162 experiment ($108 \pm 4.6\%$) and with a 30-fold greater potency than PGD₂ (pEC_{50} : 6.68 ± 0.24
163 and 5.14 ± 0.22 , respectively). However, the concentration-response curves for both PGD₂ and
164 U44619 were markedly shallow with Hill slopes significantly below 1 (0.58 ± 0.04 and $0.53 \pm$
165 0.05).

166
167 When preparations were treated with the competitive TP receptor antagonist SQ-29548, the
168 concentration-response curve to PGD₂ was shifted to the right (Fig. 1A). Pretreatment with
169 0.1 or 1 μM of SQ-29548 gave rise to significantly different pEC_{50} values (4.7 ± 0.1 and $4.2 \pm$
170 0.1 , respectively) compared to control (5.4 ± 0.3 ; table 1). At these concentrations of SQ-
171 29548, the Hill slope was significantly higher than for the control. Further experiments with
172 the combined TP and DP₂ receptor antagonist BAYu3405 (Fig. 1B) also produced
173 concentration-dependent rightward shifts of the PGD₂ induced concentration-response curve.
174 Preparations treated with 0.1 or 1 μM BAYu3405 displayed significantly different pEC_{50}
175 values (4.3 ± 0.1 and 4.0 ± 0.2 , respectively) compared to controls (5.2 ± 0.2 ; table 1).

176
177 In preparations pre-treated with 0.1 or 1 μM SQ-29548, U46619 displayed a concentration-
178 dependent shift of the concentration-response curve and the pEC_{50} values (6.0 ± 0.2 and $5.3 \pm$

179 0.2, respectively) were significantly lower than control preparations (6.8 ± 0.2 ; Fig. 1C).
180 When BAYu3405 was used as TP receptor antagonist, the same pattern was shown again
181 with significantly lower pEC_{50} values for 0.1 or 1 μ M BAYu3405 (5.4 ± 0.1 and 5.0 ± 0.1 ,
182 respectively), compared to control (7.2 ± 0.4 ; Fig. 1D). To quantify the antagonistic capacity
183 for SQ-29548 and BAYu3405, Schild plot analysis was performed, assuming that the agonist
184 curves reached the similar maximum and ignoring the absence of parallel shifts. Although the
185 antagonists showed a linear regression statistically not deviating from unity it was a tendency
186 for lower Schild slopes for both SQ-29548 (0.89 ± 0.08 for PGD_2 ; $P = 0.250$; and 0.85 ± 0.18
187 for U46619; $P = 0.521$) and BAYu3405 (0.71 ± 0.25 for PGD_2 ; $P = 0.375$; and 0.84 ± 0.13 for
188 U46619; $P = 0.364$). The pK_B values for the experiments with SQ-2954 rendered a 10-fold
189 differences between PGD_2 and U46619 (7.14 ± 0.08 and 8.17 ± 0.18 , respectively) whereas
190 no significant difference was seen in the experiments with BAYu3405 (7.82 ± 0.15 and $7.60 \pm$
191 0.16 for PGD_2 and U46619, respectively).

192

193 To test if the difference between PGD_2 and U46619 could relate to metabolism of PGD_2 , into
194 a compound activating other receptors, the early PGD_2 metabolite $9\alpha,11\beta$ - PGF_2 was studied.
195 $9\alpha,11\beta$ - PGF_2 yielded a weaker response than both U46619 and PGD_2 , reaching only about 30
196 % of the maximum contraction (Fig. 2A). The low efficacy of $9\alpha,11\beta$ - PGF_2 did not make it
197 meaningful to calculate pEC_{50} values. Since there is structural similarity between $9\alpha,11\beta$ -
198 PGF_2 and the FP receptor agonist $PGF_{2\alpha}$ (Komoto et al., 2004), $PGF_{2\alpha}$ may also be involved
199 in contractions mediated by PGD_2 and its metabolites. The effect of $PGF_{2\alpha}$ was almost
200 identical to the effect of its stereoisomer $9\alpha,11\beta$ - PGF_2 (Fig. 2B). Pretreatment with the TP
201 receptor antagonist SQ-29548 significantly abolished the contractions induced by $9\alpha,11\beta$ -
202 PGF_2 ($P < 0.001$) (Fig. 2A) and $PGF_{2\alpha}$ ($P < 0.001$) (Fig. 2B), indicating that $PGF_{2\alpha}$ acted as a
203 TP receptor agonist in the guinea pig peripheral lung preparation.

204

205 *3.2. The DP₁ receptor induce a weak relaxation of peripheral lung tissue*

206 The DP₁ receptor has been described to mediate relaxation of vascular and bronchial SMCs
207 (Norel, 2007). Therefore, it was investigated if the DP₁ receptor may mediate relaxation of
208 peripheral lung tissue. Lung strips were treated with cumulative concentrations of the DP₁
209 receptor agonist BW245C after pre-contraction with 10 nM of LTD₄. The pre-contracted
210 strips relaxed at the highest concentration of BW245C (10 μM) (Fig 3A). Based on these
211 results, experiments with the DP₁ receptor antagonist BWA868C (0.1 and 1 μM) were
212 performed with the hypothesis that inhibition of the DP₁ receptor should enhance the PGD₂
213 induced contractions. However, the BWA868C treated preparations yielded pEC₅₀ values for
214 PGD₂ that were not significant different from controls (Fig. 3B and table 2).

215

216 *3.3. The DP₂ receptor induce neither contraction nor relaxation*

217 Since BAYu3405 is both a TP and DP₂ receptor antagonist, the DP₂ receptor mediated
218 response in the parenchyma was investigated. Cumulative concentrations of the DP₂ receptor
219 agonist DK-PGD₂ were added to the lung preparations both before and after pre-contraction
220 with LTD₄. DK-PGD₂ up to 10 μM generated neither significant contraction nor relaxation of
221 the peripheral lung tissue (n=5, data not shown).

222

223 *3.4. The effects of PGD₂ and U46619 in airways and pulmonary vessels in precision-cut lung* 224 *slices*

225 Since the lung parenchyma preparation consists of both airways and vessels, precision cut
226 lung slices were examined to study the contractile effect of PGD₂ and U46619 in peripheral
227 airways and pulmonary arteries and veins. Indeed, PGD₂ induced contractile effects in
228 airways (pEC₅₀: 6.8 ± 0.1) as well as both pulmonary veins (pEC₅₀: 7.2 ± 0.2) and arteries

229 (pEC₅₀: 6.0 ± 0.2; Fig. 4A). U46619 induced similar strong contractile responses but was
230 significantly more potent in airways (pEC₅₀: 8.9 ± 0.2), veins (pEC₅₀: 9.2 ± 0.2) and arteries
231 (pEC₅₀: 8.1 ± 0.2) in the lung parenchymal preparation (Fig. 4B).

232

233 *3.5. Role of prostanoids on antigen-induced contractions*

234 Since the contractile effect of both PGD₂ and U46619 was mediated through the TP receptor,
235 experiments with SQ-29548 were performed to study the significance of this receptor in the
236 early allergic reaction. Thus, cumulative challenge with ovalbumin on parenchymal strips
237 from sensitized guinea pigs caused a concentration-dependent contraction that reached about
238 60-70% of the maximum contraction. Pre-treatment with SQ-29548 partly decreased the
239 ovalbumin-induced contraction (P < 0.01). Analysis of the bath fluid after challenge with
240 ovalbumin (1 µg/mL) showed that TXB₂ (1021 ± 325 fmol/g; n=6) was released in 20-fold
241 higher concentration than PGD₂ (53 ± 6 fmol/g; n=6). Synthesis before ovalbumin stimulation
242 was 61 ± 51 fmol/g for TXB₂ and not detectable for PGD₂.

243

244

245

246

247

248

249

250

251

252

253

254 **4. Discussion**

255 It was found that the predominant effect of PGD₂ in the peripheral lung is a contractile effect
256 which is mediated through activation of TP receptors situated on airways as well as arteries
257 and veins in the parenchymal lung tissue. A minor relaxant effect was found to be mediated
258 through the DP₁ receptor and no effect was found to be mediated through the DP₂ in the
259 present study. When inducing ovalbumin activation of sensitized parenchymal strips, both
260 TXA₂ and PGD₂ were released. Ovalbumin also induced contractions that were partly
261 mediated through activation of the TP receptor. Thus, a TP receptor antagonist can be useful
262 to block the contractile action of TXA₂ and PGD₂ in allergic reactions in the airways (Beasley
263 et al., 1989).

264

265 PGD₂ and the TP receptor agonist U46619 induced both concentration-response curves with
266 Hill slopes clearly lower than 1, indicating a complexity in the contractile action. One
267 possibility that could explain these shallow curves is if the effects were due to action through
268 more than one receptor. However, since the selective TP (SQ-29548 (Abramovitz et al.,
269 2000); and the dual TP/DP₂ (BAY u3405 (Sugimoto et al., 2005)) receptor antagonists caused
270 right-ward shifts of the concentration-response curve to PGD₂, this indicated, as in line with
271 previous observations (Hamid-Bloomfield et al., 1990; McKenniff et al., 1991), that the TP
272 receptor mediates the main part of the PGD₂-induced contraction. As U46619 induced both
273 stronger contraction and was more potent than PGD₂, the data indicated, in accordance with
274 the much lower affinity for PGD₂ to the TP receptor (Abramovitz et al., 2000), that the
275 responses were mediated through the TP receptor. On the other hand, the curves for the
276 antagonist were not shifted in parallel. Although fully concentration-response curves of PGD₂
277 were not able to obtain due to the high concentration needed, a clear trend to increased Hill
278 slope with increasing concentration of antagonists was found for both the agonists, mutually

279 with SQ-29548 and BAY u3405. It is possible that the reason for this increase is due to that
280 both PGD₂ and U46619 have shown the capacity to bind to almost all prostanoid receptors
281 (Abramovitz et al., 2000). Thus, at the higher concentrations of the agonists used for the
282 antagonist experiments, the activation of the TP receptor simultaneously with one or more
283 prostanoid receptors cause an additional or a synergistic effect.

284

285 Although the Schild plot slopes did not significantly deviate from unity there was a trend to
286 lower slope values, which may be due to actions of more than one receptor, especially at the
287 higher concentrations of the agonists. Also the discrepancy of the pK_B values for SQ-29548
288 and not BAY u3405 between the agonists can indicate that another receptor than the TP
289 receptor is activated. The pK_B values for BAY u3405 (7.82 and 7.60 for PGD₂ and U46619,
290 respectively) are in line with presented earlier in guinea pig lung strips (7.7 with U46619 as
291 agonist; (Norman et al., 1992)) whereas the pK_B values for SQ-29548 was 10 fold lower for
292 BAY u3405 (7.14) than for U46619 (8.17) as agonists (range from 7.7 to 8.7 in guinea pig;
293 (Dube et al., 1992; Norman et al., 1992). Since it has been shown that U46619 do not activate
294 the DP₂ receptor (Monneret et al., 2001), the lower pK_B value for SQ-29548 from the PGD₂
295 experiments can be due to activation of DP₂ receptors. However, as described in this study, a
296 negligible effect is shown when this receptor is selectively activated. Thus, from these
297 experiments the reason for the difference of the pK_B values for SQ-29548 cannot be
298 completely concluded. Taking all these complexities of the actions of the agonists both in
299 absence and presence of antagonism in consideration, the clear antagonism with both these
300 known TP receptor antagonists indicates that the main action of PGD₂ goes through the TP
301 receptor.

302

303 The major PGD₂ metabolite, 9 α ,11 β -PGF₂, induced a weak contraction of the peripheral lung
304 that was blocked by SQ-29548 indicating that the breakdown of PGD₂ in this assay do not
305 cause activation of any further receptor then PGD₂ by itself. PGF_{2 α} is a stereoisomer to
306 9 α ,11 β -PGF₂ and closely structurally related to PGD₂ (Sandig et al., 2006). Thus, one
307 possibility was that PGD₂ also acted through FP receptors (Kiriyaama et al., 1997). Since there
308 are no specific receptor antagonists available for the FP receptor we chose to investigate how
309 the response to PGF_{2 α} could be blocked by SQ-29548. However, as the effect of PGF_{2 α} was
310 abolished by SQ-29548 it is unlikely that PGD₂ mediate any major effect through the FP
311 receptor. Altogether, these data implicate that PGF_{2 α} , which along with PGD₂ has been shown
312 to be released after antigen challenge in guinea pig lung (Dawson et al., 1976), also is part of
313 the TP receptor mediated constriction of the peripheral lung.

314

315 The DP₁ receptor has been shown to mediate relaxation of the bronchioles in response to
316 PGD₂ and may thus serve as protection in a situation of bronchoconstriction (Norel et al.,
317 1999). Studies have also shown that rabbit jugular vein preparations treated with the DP₁
318 receptor agonist BW245C relaxed concentration-dependently (Lydford et al., 1996).
319 Nevertheless, the DP₁ receptor agonist only weakly relaxed the parenchymal lung tissue in the
320 present study. Furthermore, the DP₁ receptor antagonist BWA868C did not affect the PGD₂
321 induced contractions. Since the DP₂ receptor agonist did not induce contractions or relaxations
322 in this lung preparation, this point towards that the main action for PGD₂ on airway
323 inflammation is the known induction of chemotaxis of eosinophils, basophils and Th2 cells in
324 guinea pigs (Liu et al., 2005) rather than direct responses on the airway smooth muscle.

325

326 During the control situation, both agonists induced shallow concentration-response curves in
327 the parenchymal strips. Except for activation on more than one receptor, these shallow curves

328 can be due to the action of several smooth muscle components, such as small airways and
329 vessels (Evans and Adler, 1981), that not react with similar potencies and maximal effects. In
330 the precision-cut lung slice experiments, the potency difference between PGD₂ and U44619
331 of the three studied components was similar as in the parenchymal strips (approximately 100-
332 fold) indicating similar contractile actions for the agonists during the control situation.
333 However, a clear difference was obtained between the components, especially the pulmonary
334 arteries compared to the airways and pulmonary veins with a both lower potency and maximal
335 effect in the arteries, suggesting that this is the reason for the shallow control curves for both
336 PGD₂ and U46619. Actually, it is indicated that not only airways and vessels but also pleural
337 cells, alveolar ducts and interstitial cells are activated by TP receptor agonists (Wong et al.,
338 1992). Thus, it is possible that these more peripheral located contractile units, which not
339 easily can be measured in precision-cut lung slices, together with the airways and vessels are
340 activated in a serial manner causing the described shallow concentration-response curves for
341 the two agonists.

342

343 Allergen-challenge with ovalbumin showed that both PGD₂ and TXA₂, the latter measured as
344 TXB₂, are released from the distal lung tissue and may be important mediators of the allergen-
345 induced bronchoconstriction in the peripheral lung. Previous studies in this model have shown
346 that also histamine, leukotrienes and PGE₂ are generated after ovalbumin stimulation (Larsson
347 et al., 2009). The 20-fold higher level of TXB₂ in this study is similar to the levels which have
348 been seen in effluent from ovalbumin exposed isolated perfused and ventilated guinea pig
349 lung (Selg et al., 2008). That TXA₂ also is released from mast cells has been shown in studies
350 of the human mast cell line HMC-1 (Macchia et al., 1995). However, human purified sinus
351 mast cells showed a 10-fold higher level of PGD₂ than TXB₂ after IgE stimulation (Mita et
352 al., 1999) suggesting that clear species or anatomical differences exist.

353

354 Previously it has been shown that anti-histamine or 5-LO inhibitor alone has no inhibitory
355 effect on the antigen-induced contraction in guinea pig lung parenchyma (Jonsson and
356 Dahlen, 1994; Larsson et al., 2007). Furthermore, it has been shown that the allergen induced
357 contraction in the peripheral lung of the guinea pig needs to be antagonised or inhibited via
358 several mediator pathways in order to significantly attenuate the contraction (Larsson et al.,
359 2007; Ressmeyer et al., 2006). However, in the present study we found that the TP receptor
360 antagonist significantly inhibited part of the ovalbumin-induced contractions. Thus, the
361 experimental results suggest that TP receptors mediate a significant component of the
362 allergen-induced contractions in this model of the peripheral lung. The explanation, indicated
363 by our findings, may be that several COX-products released by the antigen-challenge (PGD₂,
364 TXA₂ and PGF_{2α}; (Dawson et al., 1976) all act on TP receptors.

365

366 The present study highlights that the parenchymal constriction induced by PGD₂ should be
367 attributed to its properties as a TP receptor agonist. Even though PGD₂ may have minor role
368 as an agonist in the allergen-induced contraction in guinea pigs, since it is both released in
369 lower amount and have lesser effect than TXA₂, the role in human may be of great importance
370 due to the high amount of mast cell release (Mita et al., 1999). Accordingly, the contractile
371 effect of PGD₂ in human airways has also been shown to be antagonized by BAY u3405
372 (Magnussen et al., 1992), which can be important to bear in mind when considering the
373 treatment of early asthmatic responses. Furthermore, the results here are consistent with other
374 studies showing that the allergen response needs to be antagonised or inhibited via several
375 mechanisms to attenuate the contraction (Jonsson and Dahlen, 1994; Roquet et al., 1997; Selg
376 et al., 2009). In this concept, therapy with TP receptor antagonists should be considered as
377 one important component to reduce early asthmatic responses in patients.

378 **Acknowledgement**

379 We would like to express our gratitude to Margareta Andersson for skillful technical
380 assistance and we would like to thank Swedish Research Council in Medicine and Health,
381 Swedish Heart and Lung foundation, Vinnova Chronic inflammation - diagnostic and therapy
382 (CIDaT), the Crafoord Foundation, the Stockholm County Council Research Funds (ALF),
383 Karolinska Institutet , the Tore Nilsson Foundation and the Swedish Society of Medicine for
384 financial support.

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403 **References**

- 404 Abe, H., Takeshita, T., Nagata, K., Arita, T., Endo, Y., Fujita, T., Takayama, H., Kubo, M.,
405 Sugamura, K., 1999. Molecular cloning, chromosome mapping and characterization of
406 the mouse CRTH2 gene, a putative member of the leukocyte chemoattractant receptor
407 family. *Gene* 227, 71-77.
- 408 Abramovitz, M., Adam, M., Boie, Y., Carriere, M., Denis, D., Godbout, C., Lamontagne, S.,
409 Rochette, C., Sawyer, N., Tremblay, N.M., Belley, M., Gallant, M., Dufresne, C., Gareau,
410 Y., Ruel, R., Juteau, H., Labelle, M., Ouimet, N., Metters, K.M., 2000. The utilization of
411 recombinant prostanoid receptors to determine the affinities and selectivities of
412 prostaglandins and related analogs. *Biochim. Biophys. Acta* 1483, 285-293.
- 413 Andersson, C.K., Mori, M., Bjermer, L., Lofdahl, C.G., Erjefalt, J.S., 2009. Novel site-
414 specific mast cell subpopulations in the human lung. *Thorax* 64, 297-305.
- 415 Balzar, S., Fajt, M.L., Comhair, S.A., Erzurum, S.C., Bleecker, E., Busse, W.W., Castro, M.,
416 Gaston, B., Israel, E., Schwartz, L.B., Curran-Everett, D., Moore, C.G., Wenzel, S.E.,
417 Mast cell phenotype, location, and activation in severe asthma: data from the severe
418 asthma research program. *Am. J. Respir. Crit. Care Med.* 183, 299-309.
- 419 Beasley, R.C., Featherstone, R.L., Church, M.K., Rafferty, P., Varley, J.G., Harris, A.,
420 Robinson, C., Holgate, S.T., 1989. Effect of a thromboxane receptor antagonist on
421 PGD₂- and allergen-induced bronchoconstriction. *J. Appl. Physiol.* 66, 1685-1693.
- 422 Canning, B.J., Chou, Y., 2008. Using guinea pigs in studies relevant to asthma and COPD.
423 *Pulm. Pharmacol. Ther.* 21, 702-720.
- 424 Capra, V., Habib, A., Accomazzo, M.R., Ravasi, S., Citro, S., Levy-Toledano, S., Nicosia, S.,
425 Rovati, G.E., 2003. Thromboxane prostanoid receptor in human airway smooth muscle
426 cells: a relevant role in proliferation. *Eur. J. Pharmacol.* 474, 149-159.

427 Coleman, R.A., Smith, W.L., Narumiya, S., 1994. International Union of Pharmacology
428 classification of prostanoid receptors: properties, distribution, and structure of the
429 receptors and their subtypes. *Pharmacol. Rev.* 46, 205-229.

430 Dahlen, S.E., Kumlin, M., 2004. Monitoring mast cell activation by prostaglandin D2 in vivo.
431 *Thorax* 59, 453-455.

432 Dawson, W., Boot, J.R., Cockerill, A.F., Mallen, D.N., Osborne, D.J., 1976. Release of novel
433 prostaglandins and thromboxanes after immunological challenge of guinea pig lung.
434 *Nature* 262, 699-702.

435 Dube, G.P., Mais, D.E., Jakubowski, J.A., Brune, K.A., Utterback, B.G., True, T.A.,
436 Rinkema, L.E., Kurtz, W.L., 1992. In vitro characterization of a novel TXA2/PGH2
437 receptor ligand (S-145) in platelets and vascular and airway smooth muscle. *J.*
438 *Pharmacol. Exp. Ther.* 262, 784-791.

439 Evans, J.N., Adler, K.B., 1981. The lung strip: evaluation of a method to study contractility of
440 pulmonary parenchyma. *Exp. Lung. Res.* 2, 187-195.

441 Hamid-Bloomfield, S., Payne, A.N., Petrovic, A.A., Whittle, B.J., 1990. The role of
442 prostanoid TP- and DP-receptors in the bronchoconstrictor effect of inhaled PGD2 in
443 anaesthetized guinea-pigs: effect of the DP-antagonist BW A868C. *Br. J. Pharmacol.*
444 100, 761-766.

445 Held, H.D., Martin, C., Uhlig, S., 1999. Characterization of airway and vascular responses in
446 murine lungs. *Br. J. Pharmacol.* 126, 1191-1199.

447 Jonsson, E.W., Dahlen, S.E., 1994. Interactions between leukotrienes and histamine in the
448 anaphylactic contraction of guinea pig lung parenchyma. *J. Pharmacol. Exp. Ther.* 271, 615-
449 623.

450 Kiriyaama, M., Ushikubi, F., Kobayashi, T., Hirata, M., Sugimoto, Y., Narumiya, S., 1997.
451 Ligand binding specificities of the eight types and subtypes of the mouse prostanoid
452 receptors expressed in Chinese hamster ovary cells. *Br. J. Pharmacol.* 122, 217-224.

453 Komoto, J., Yamada, T., Watanabe, K., Takusagawa, F., 2004. Crystal structure of human
454 prostaglandin F synthase (AKR1C3). *Biochemistry* 43, 2188-2198.

455 Larsson, A.K., Back, M., Hjoberg, J., Dahlen, S.E., 2005. Inhibition of nitric-oxide synthase
456 enhances antigen-induced contractions and increases release of cysteinyl-leukotrienes in
457 guinea pig lung parenchyma: nitric oxide as a protective factor. *J. Pharmacol. Exp. Ther.*
458 315, 458-465.

459 Larsson, A.K., Back, M., Lundberg, J.O., Dahlen, S.E., 2009. Specific mediator inhibition by
460 the NO donors SNP and NCX 2057 in the peripheral lung: implications for allergen-
461 induced bronchoconstriction. *Respir. Res.* 10, 46.

462 Larsson, A.K., Fumagalli, F., DiGennaro, A., Andersson, M., Lundberg, J., Edenius, C.,
463 Govoni, M., Monopoli, A., Sala, A., Dahlen, S.E., Folco, G.C., 2007. A new class of
464 nitric oxide-releasing derivatives of cetirizine; pharmacological profile in vascular and
465 airway smooth muscle preparations. *Br. J. Pharmacol.* 151, 35-44.

466 Liu, F., Gonzalo, J.A., Manning, S., O'Connell, L.E., Fedyk, E.R., Burke, K.E., Elder, A.M.,
467 Pulido, J.C., Cao, W., Tayber, O., Qiu, Y., Ghosh, S., Ocain, T.D., Hodge, M.R., Suzuki-
468 Yagawa, Y., 2005. Pharmacological characterization of guinea pig chemoattractant
469 receptor-homologous molecule expressed on Th2 cells (CRTH2). *Prostaglandins Other*
470 *Lipid Mediat.* 76, 133-147.

471 Liu, M.C., Hubbard, W.C., Proud, D., Stealey, B.A., Galli, S.J., Kagey-Sobotka, A., Bleecker,
472 E.R., Lichtenstein, L.M., 1991. Immediate and late inflammatory responses to ragweed
473 antigen challenge of the peripheral airways in allergic asthmatics. Cellular, mediator, and
474 permeability changes. *Am. Rev. Respir. Dis.* 144, 51-58.

475 Lydford, S.J., McKechnie, K.C., Leff, P., 1996. Interaction of BW A868C, a prostanoid DP-
476 receptor antagonist, with two receptor subtypes in the rabbit isolated saphenous vein.
477 Prostaglandins 52, 125-139.

478 Macchia, L., Hamberg, M., Kumlin, M., Butterfield, J.H., Haeggstrom, J.Z., 1995.
479 Arachidonic acid metabolism in the human mast cell line HMC-1: 5-lipoxygenase gene
480 expression and biosynthesis of thromboxane. Biochim. Biophys. Acta 1257, 58-74.

481 Magnussen, H., Boerger, S., Templin, K., Baunack, A.R., 1992. Effects of a thromboxane-
482 receptor antagonist, BAY u 3405, on prostaglandin D2- and exercise-induced
483 bronchoconstriction. J. Allergy Clin. Immunol. 89, 1119-1126.

484 Matsuoka, T., Hirata, M., Tanaka, H., Takahashi, Y., Murata, T., Kabashima, K., Sugimoto,
485 Y., Kobayashi, T., Ushikubi, F., Aze, Y., Eguchi, N., Urade, Y., Yoshida, N., Kimura, K.,
486 Mizoguchi, A., Honda, Y., Nagai, H., Narumiya, S., 2000. Prostaglandin D2 as a
487 mediator of allergic asthma. Science 287, 2013-2017.

488 McKenniff, M.G., Norman, P., Cuthbert, N.J., Gardiner, P.J., 1991. BAY u3405, a potent and
489 selective thromboxane A2 receptor antagonist on airway smooth muscle in vitro. Br. J.
490 Pharmacol. 104, 585-590.

491 Mita, H., Ishii, T., Akiyama, K., 1999. Generation of thromboxane A2 from highly purified
492 human sinus mast cells after immunological stimulation. Prostaglandins Leukot. Essent.
493 Fatty Acids 60, 175-180.

494 Monneret, G., Gravel, S., Diamond, M., Rokach, J., Powell, W.S., 2001. Prostaglandin D2 is
495 a potent chemoattractant for human eosinophils that acts via a novel DP receptor. Blood
496 98, 1942-1948.

497 Norel, X., 2007. Prostanoid receptors in the human vascular wall. ScientificWorldJournal 7,
498 1359-1374.

499 Norel, X., Walch, L., Labat, C., Gascard, J.P., Dulmet, E., Brink, C., 1999. Prostanoid
500 receptors involved in the relaxation of human bronchial preparations. *Br. J. Pharmacol.*
501 126, 867-872.

502 Norman, P., Cuthbert, N.J., McKenniff, M.G., Gardiner, P.J., 1992. The thromboxane
503 receptors of rat and guinea-pig lung. *Eur. J. Pharmacol.* 229, 171-178.

504 O'Sullivan, S., Dahlen, B., Dahlen, S.E., Kumlin, M., 1996. Increased urinary excretion of the
505 prostaglandin D2 metabolite 9 alpha, 11 beta-prostaglandin F2 after aspirin challenge
506 supports mast cell activation in aspirin-induced airway obstruction. *J. Allergy Clin.*
507 *Immunol.* 98, 421-432.

508 Okano, M., Fujiwara, T., Sugata, Y., Gotoh, D., Masaoka, Y., Sogo, M., Tanimoto, W.,
509 Yamamoto, M., Matsumoto, R., Eguchi, N., Kiniwa, M., Isik, A.U., Urade, Y., Nishizaki,
510 K., 2006. Presence and characterization of prostaglandin D2-related molecules in nasal
511 mucosa of patients with allergic rhinitis. *Am. J. Rhinol.* 20, 342-348.

512 Parameswaran, K., Radford, K., Fanat, A., Stephen, J., Bonnans, C., Levy, B.D., Janssen, L.J.,
513 Cox, P.G., 2007. Modulation of human airway smooth muscle migration by lipid
514 mediators and Th-2 cytokines. *Am. J. Respir. Cell Mol. Biol.* 37, 240-247.

515 Ressmeyer, A.R., Larsson, A.K., Vollmer, E., Dahlen, S.E., Uhlig, S., Martin, C., 2006.
516 Characterisation of guinea pig precision-cut lung slices: comparison with human tissues.
517 *Eur. Respir. J.* 28, 603-611.

518 Roquet, A., Dahlen, B., Kumlin, M., Ihre, E., Anstren, G., Binks, S., Dahlen, S.E., 1997.
519 Combined antagonism of leukotrienes and histamine produces predominant inhibition of
520 allergen-induced early and late phase airway obstruction in asthmatics. *Am. J. Respir.*
521 *Crit. Care Med.* 155, 1856-1863.

522 Sandig, H., Andrew, D., Barnes, A.A., Sabroe, I., Pease, J., 2006. 9alpha,11beta-PGF2 and its
523 stereoisomer PGF2alpha are novel agonists of the chemoattractant receptor, CRTH2.
524 FEBS Lett. 580, 373-379.

525 Selg, E., Andersson, M., Lastbom, L., Ryrfeldt, A., Dahlen, S.E., 2009. Two different
526 mechanisms for modulation of bronchoconstriction in guinea-pigs by cyclooxygenase
527 metabolites. Prostaglandins Other Lipid Mediat. 88, 101-110.

528 Selg, E., Lastbom, L., Ryrfeldt, A., Kumlin, M., Dahlen, S.E., 2008. Effects of selective and
529 non-selective COX inhibitors on antigen-induced release of prostanoid mediators and
530 bronchoconstriction in the isolated perfused and ventilated guinea pig lung.
531 Prostaglandins Leukot. Essent. Fatty Acids 78, 89-97.

532 Sugimoto, H., Shichijo, M., Okano, M., Bacon, K.B., 2005. CRTH2-specific binding
533 characteristics of [3H]ramatroban and its effects on PGD2-, 15-deoxy-Delta12, 14-PGJ2-
534 and indomethacin-induced agonist responses. Eur. J. Pharmacol. 524, 30-37.

535 Urade, Y., Eguchi, N., 2002. Lipocalin-type and hematopoietic prostaglandin D synthases as a
536 novel example of functional convergence. Prostaglandins Other Lipid Mediat. 68-69,
537 375-382.

538 van den Berge, M., ten Hacken, N.H., Cohen, J., Douma, W.R., Postma, D.S., 2011. Small
539 airway disease in asthma and COPD: clinical implications. Chest 139, 412-423.

540 Vane, J.R., 1971. Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-
541 like drugs. Nat. New Biol. 231, 232-235.

542 Wong, W.S., Bloomquist, S.L., Bendele, A.M., Fleisch, J.H., 1992. Pharmacological and
543 histological examinations of regional differences of guinea-pig lung: a role of pleural
544 surface smooth muscle in lung strip contraction. Br. J. Pharmacol. 105, 620-626.

545

546

547 **Figure legends**

548 **Fig. 1.** Contractions induced by cumulative concentrations of (A and B) PGD₂ and (C and D)
549 TP receptor agonist U46619 in guinea pig lung parenchymal strips. The experiments were
550 performed in absence or presence of (A and C) the TP receptor antagonists SQ-29548 or (B
551 and D) BAY u3405 (n=5-6). Data are presented as mean ± S.E.M..

552

553 **Fig. 2.** Contractions induced by cumulative concentrations of (A) 9 α ,11 β - PGF₂ and (B)
554 PGF_{2 α} in guinea pig lung parenchymal strips. The experiments were performed in absence or
555 presence of 1 μ M of the TP receptor antagonist SQ-29548 (n=5). Data are presented as mean ±
556 S.E.M..

557

558 **Fig. 3.** Studies of the DP₁ receptor in guinea pig lung parenchymal strips. (A) Effect of
559 cumulative concentrations of the DP₁ receptor agonist BW245C after precontraction with
560 LTD₄ 10 nM and (B) effect of the DP₁ receptor antagonist BWA868C on PGD₂ induced
561 contractions (n=5). Data are presented as mean ± S.E.M..

562

563 **Fig. 4.** Contractile responses to (A) PGD₂ and (B) the TP receptor agonist U46619 in airway
564 and, pulmonary artery and vein in guinea pig precision cut lung slices. Data presented as %
565 change of initial luminal area (n=4). Data are presented as mean ± S.E.M..

566

567 **Fig. 5.** Effect of the TP receptor antagonist SQ-29548 on cumulative doses of ovalbumin
568 (0.001-10 μ g/ml) in absence and presence of 1 μ M SQ-29548 (n=4) in guinea pig lung
569 parenchymal strips. Data are presented as mean ± S.E.M..

Table 1: Effects of the TP receptor antagonists on PGD₂ and U46619 induced contractions

Treatment	N	E _{max}	pEC ₅₀	Hill slope
PGD ₂ (Control)	6	65.2±4.8	5.4± 0.3	0.6 ± 0.0
PGD ₂ + SQ-29548 0.01µM	6	78.9±0.7	4.9±0.1	0.6±0.0
PGD ₂ + SQ-29548 0.1µM	6	72.5±5.2	4.7±0.1 ^a	0.9±0.1 ^a
PGD ₂ + SQ-29548 1µM	6	50.1±7.1	4.2±0.1 ^c	1.7±0.1 ^c
PGD ₂ (Control)	5	75.7±3.5	5.2±0.2	0.6±0.0
PGD ₂ + BAYu3405 0.1µM	5	64.7±8.2	4.3±0.1 ^b	1.4±0.2 ^b
PGD ₂ + BAYu3405 1µM	5	49.2±15.1	4.0±0.2 ^c	1.9±0.1 ^c
U46619 (Control)	6	99.1±0.9	6.8±0.2	0.6±0.1
U46619 + SQ-29548 0.01µM	5	99.1±1.0	6.6±0.2	0.7±0.1
U46619 + SQ-29548 0.1µM	5	100.0±0.0	6.0±0.2 ^a	0.8±0.1
U46619 + SQ-29548 1µM	5	100.0±0.0	5.3±0.2 ^c	1.6±0.3 ^c
U46619 (Control)	6	98.6±1.2	7.2±0.4	0.5±0.1
U46619 + BAYu3405 0.01µM	5	99.8±0.2	6.5±0.2	0.7±0.2
U46619 + BAYu3405 0.1µM	5	100.0±0.0	5.4±0.1 ^c	1.1±0.2
U46619 + BAYu3405 1µM	5	97.9±2.1	5.0±0.1 ^c	1.6±0.2 ^c

Calculations of E_{max}, pEC₅₀ and Hill slope presented as mean ± S.E.M.. Significant differences from the agonists (controls) are indicated as ^aP<0.05, ^bP<0.01 or ^cP<0.001.

Table 2. Effects of the DP₁ receptor antagonist on PGD₂ induced contractions

Treatment	N	E_{max}	pEC₅₀	Hill slope
PGD ₂ (Control)	5	83.8 ± 6.4	4.6 ± 0.3	0.4 ± 0.0
PGD ₂ + BWA868C 0.1µM	5	81.5 ± 5.9	4.9 ± 0.1	0.5 ± 0.1
PGD ₂ + BWA868C 1µM	5	76.9 ± 7.4	5.0 ± 0.3	0.5 ± 0.1

Calculations for E_{max}, pEC₅₀ and Hill slope are presented as mean ± S.E.M..

Figure 1




Figure 2





Figure 3





Figure 4





Figure 5

