



# LUND UNIVERSITY

## Multiple anthropometric measures in relation to incidence of diabetes: a Swedish population-based cohort study.

Borné, Yan; Nilsson, Peter; Melander, Olle; Hedblad, Bo; Engström, Gunnar

*Published in:*  
European Journal of Public Health

*DOI:*  
[10.1093/eurpub/ckv044](https://doi.org/10.1093/eurpub/ckv044)

2015

[Link to publication](#)

*Citation for published version (APA):*  
Borné, Y., Nilsson, P., Melander, O., Hedblad, B., & Engström, G. (2015). Multiple anthropometric measures in relation to incidence of diabetes: a Swedish population-based cohort study. *European Journal of Public Health*, 25(6), 1100-1105. <https://doi.org/10.1093/eurpub/ckv044>

*Total number of authors:*  
5

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

**Multiple anthropometric measures in relation to incidence of diabetes:  
a Swedish population-based cohort study**

Yan Borné<sup>1</sup>\*, Peter M Nilsson<sup>1</sup>, Olle Melander<sup>1</sup>, Bo Hedblad<sup>1</sup>, Gunnar Engström<sup>1</sup>

Department of Clinical Sciences, University Hospital, Lund University, Malmö, Sweden<sup>1</sup>

\* Corresponding author: Yan Borné: Department of Clinical Sciences, Cardiovascular  
Epidemiology, University Hospital, Lund University, CRC, 20502 Malmö, Sweden.  
Tel: +46 40391327, Fax: +46 40391322. Yan.Borne@med.lu.se

## **Abstract**

**Background:** Obesity is the major modifiable risk factor for diabetes. This study investigated the incidence of diabetes in relation to multiple anthropometric measures.

**Methods:** Body mass index (BMI), waist circumference (WC), waist-height ratio (WHtR), waist-hip ratio (WHR) and body fat percentage (BF %) were measured among 26604 subjects (aged 45-73 years) without history of diabetes from the Malmö Diet and Cancer cohort.

**Results:** During 14 years of follow-up, 2935 subjects (1519 men, 1416 women) were diagnosed with diabetes. In men, incidence of diabetes was 24.1 and 4.0 per 1000 person-years comparing the 4th vs 1st quartile of WHtR. The multivariate adjusted hazard ratios (HR; 4th vs 1st quartile) were 6.00 (95%CI: 5.02-7.16) for WHtR, 5.95 (CI: 4.96-7.14) for WC, 5.19 (CI: 4.38-6.15) for BMI, 4.71 (CI: 3.96-5.60) for WHR and 3.21 (CI: 2.75-3.76) for BF%. For women, incidence of diabetes was 15.1 and 1.4 per 1000 person-years for 4th vs 1st quartile of WHtR (HR: 10.19, CI: 8.10-12.82). HR was 9.16 (CI: 7.40-11.33) for WC, 6.42 (CI: 5.27-7.81) for BMI, 6.75 (CI: 5.52-8.25) for WHR and 5.39 (CI: 4.42-6.57) for BF%. Model discrimination was marginally increased when WC, WHtR or WHR was used in combination with BMI.

**Conclusion:** All measures of obesity were associated with substantially increased incidence of diabetes. Abdominal obesity was associated with higher incidence rates in men than in women, but in terms of relative risks the relationships were stronger in women. The combination of BMI and abdominal obesity measures had stronger association with diabetes than BMI alone.

**Keywords:** anthropometric measure, incidence of diabetes, risk factors, population-based cohort study.

## Introduction

Type 2 diabetes is an increasing health problem in most countries <sup>1</sup>. Age, sex and genetic factors play key roles for diabetes onset, however, diabetes is a preventable disease and obesity is the major modifiable risk factor in the population <sup>2-5</sup>. The relationships between obesity and blood glucose or diabetes have been demonstrated both in cross-sectional <sup>6-8</sup> and prospective studies <sup>9-12</sup>.

Body mass index (BMI) is the most practical and commonly used anthropometric measure for general fat <sup>6, 9, 10</sup>. However, the body fat distribution, especially abdominal obesity has been proposed to be a stronger risk factor for diabetes <sup>13, 14</sup>. A recent meta-analysis found that waist/height ratio (WHtR) is a better screening tool for cardiometabolic risk factors compared to waist circumference and BMI <sup>15</sup>. Other studies suggested that both general fat and abdominal fat are risk factor for diabetes <sup>7, 11</sup>. Prospective data on anthropometry and incidence of self-reported diabetes have been reported in a study from Germany. WC was a better predictor than other measures among men and women; WHtR improved the prediction most for men and WHtR and WC were similar predictors for women <sup>12</sup>. Most studies of obesity and incidence of diabetes have been performed in populations from Europe <sup>9, 11, 12</sup> or the US <sup>4, 6</sup>. However, cross-sectional analyses from the Obesity in Asia Collaboration, including more than 263 000 individuals from 21 studies, reported stronger relationships for waist circumference and type 2 diabetes than for BMI<sup>8</sup>.

The aim of this prospective study was to explore the relationship between different anthropometric measures, i.e., BMI, WC, WHtR, waist-hip ratio (WHR) or percent body fat (BF%), and incidence of diabetes in a large population-based cohort study.

## **Materials/subjects and Methods**

### **Study population**

Malmö Diet and Cancer (MDC) cohort, from the city of Malmö in southern Sweden, was used for the present study. All women born between 1923 and 1950 and men born between 1923 and 1945 living in Malmö city were invited to the MDC study during the period March 1991 to September 1996, participant rate was 41%<sup>16-18</sup>. A total of 28 449 subjects underwent measurement of blood pressure and anthropometric measures and filled out a self-administered questionnaire.

Subjects with history of diabetes (n= 958) at the baseline examination were excluded. In addition, 887 subjects were also excluded due to missing values of anthropometric measurements and other biological, life-style and socioeconomic variables. Thus, the final study population in the project consisted of 26 604 subjects (10 332 men (38.8%) and 16 272 women (61.2%)), aged 45-73 years.

The ethics committee at Lund University approved the study (LU 51/90) and all participants provided informed consent.

### **Baseline examinations**

The examinations were performed by trained nurses at the screening center. Standing height was measured with a fixed stadiometer calibrated in centimeters. Weight was measured to the nearest 0.1 kg using balance-beam scale with subjects wearing light clothing and no shoes. BMI was calculated as weight (kg) divided by the square of the height (m<sup>2</sup>). WC was measured as the circumference (cm) between the lowest rib margin and iliac crest and hip circumference (cm) as the largest circumference between waist and thighs. WHR was defined

as the ratio of circumference of waist to hip. WHtR was defined as the ratio of circumference of waist to height. Bioelectrical Impedance Analyzers (BIA) was used for estimating body composition and BF% was calculated using an algorithm, according to procedures provided by the manufacturer (BIA 103, JRL systems, single-frequency analyzer, Detroit, USA) <sup>19</sup>. BMI, WC, WHtR, WHR and BF % was categorized into quartiles Q1–4, separately for men and women.

Information on current use of lipid-lowering or anti-diabetic medications, smoking habits, alcohol consumption, leisure-time physical activity, education level, civil status and immigrant status were obtained from a self-administered questionnaire. Low level of leisure-time physical activity was defined as the lowest tertile of a score revealed through 18 questions covering a range of activities in the 4 seasons. The evaluation of the questionnaire has been previously reported <sup>20</sup>. Subjects were categorized into current smokers (i.e. those who smoked regularly or occasionally) or non-smokers (i.e. former smokers and never smokers). High alcohol consumption was defined as >40 g alcohol per day for men and >30 g per day for women. Education was divided into three groups: school year <9, 9-12 and > 12, respectively. Civil status was categorized into married or not. Immigrant status was grouped as Swedish-born and foreign-born.

### *Incidence of diabetes*

All subjects were followed from the baseline examination until first diagnosis of diabetes, death, emigration from Sweden or December 31<sup>st</sup>, 2009, whichever came first. A total of 5117 subjects were censored from the cohort due to death (n=4912) or emigration from Sweden (n=205). New-onset cases of diabetes in the MDC cohort were retrieved from several sources and have been described in detail previously <sup>21,22</sup>. In short, incident diabetes was identified in the Malmö HbA1c register (MHR) (56% of all cases), the Swedish National Diabetes Register

(NDR) (14%), the Swedish inpatient register (40%), the Swedish outpatient register (38%), the nationwide Swedish drug prescription register (65%), the regional Diabetes 2000 register of the Skåne region (22%). Of 113 cases (3.8%) that had diabetes as cause of death all except three were also found in other registers. In addition, 44% of the cases were identified at re-examinations of the cohort. At least two independent sources confirmed the diagnosis for 71.6% of the cases, and 53% of the cases were identified in three independent data sources. NDR and the Diabetes 2000 register required a physician diagnosis according to established diagnostic criteria (fasting plasma glucose concentration of  $\geq 7.0$  mmol/L, which corresponds to a fasting whole blood glucose of  $\geq 6.1$  mmol/L, measured on 2 different occasions). The MHR at the Department of Clinical Chemistry, Malmö University Hospital, analyzed and recorded all HbA1c samples taken in institutional and non-institutional care in the greater Malmö area from 1988 onwards. Individuals who had at least two HbA1c recordings  $\geq 6.0\%$  in the MHR with the Swedish Mono-S standardization system (corresponding to 7.0% according to the US National Glycohemoglobin Standardization Program) after the baseline examination were defined as incident diabetes cases.

### **Statistical analysis**

Cox proportional hazards regression was used to examine the association between anthropometric measures (in quartiles) and incidence of diabetes. Time axis was follow-up time until death, emigration, incident diabetes or end of follow-up. Hazard ratios (HR), with 95% confidence interval (CI) were calculated. Age was included as covariate in the basic model. Multivariable Cox proportional hazards models were adjusted for use of lipid-lowering medication, current smoking, high alcohol consumption, low leisure physical activity, low education, marital status and immigrant status. We calculated the Harrell's C-statistic to assess whether the anthropometric measures improved model discrimination<sup>23, 24</sup>.

The Harrell's C statistics<sup>25, 26</sup>, a generalization of the area under the receiver-operating characteristic (ROC) curve were calculated to assess the diabetes prediction efficiency of the models.

Interaction terms were added to the age- and sex-adjusted Cox model to explore the potential interactions between gender and quartiles of anthropometric measures. In addition we also performed analyses of incidence of diabetes in relation to anthropometric measures dichotomized according to previously suggested cut-off values<sup>15, 27</sup>. All analyses were performed using IBM SPSS Statistics (version 20; IBM Sweden AB, Stockholm, Sweden) and Stata software version 12.0 (StataCorp).

## **Results**

### **Study cohort**

The baseline characteristics of men and women in the MDC cohort (N=26 604) are presented in Table 1. As expected, men generally had higher levels of the anthropometric measures, except for BF%. Mean age was higher for men. Prevalence of smoking was similar in men and women.

### **Incidence of diabetes in relation to anthropometric measures**

During a mean follow-up of 14 years, a total of 2 935 subjects (1 519 men and 1 416 women) developed diabetes. Incidence (per 1000 person-years) was 10.8 for men and 6.1 for women. The overall analysis showed that BMI, WC, WHtR, WHR and BF % all were major risk factors for diabetes independently from several sociodemographic, lifestyle and biological factors, both among men and women. Among men, incidence of diabetes was 24 (95%CI: 22-26) vs 4.0 per 1000 person-years (CI:3.3-4.6) , comparing 4<sup>th</sup> vs 1<sup>st</sup> quartile of WHtR (Table 2). The adjusted HR of incident diabetes (4<sup>th</sup> vs 1<sup>st</sup> quartile) were 6.00 (95% CI: 5.02-7.16) for



WHtR, 5.95 (4.96-7.14) for WC, 5.19 (4.38-6.15) for BMI, 4.71 (3.96-5.60) for WHR, and 3.21 (2.75-3.76) for BF% after adjustment for several potential confounding factors, Table 2. For women, incidence of diabetes was 15.1 (95%CI: 14-16) and 1.4 per 1000 person-years (95%CI: 1.1-1.7), respectively, for 4<sup>th</sup> vs 1<sup>st</sup> quartile of WHtR. The adjusted HR was 10.19 (8.10-12.82) for WHtR, 9.16 (7.40-11.33) for WC, 6.75 (5.52-8.25) for WHR, 5.39 (4.42-6.57) for BF% and 6.42 (5.27-7.81) for BMI, Table 3. All anthropometric measures except BMI (p=0.058) showed statistically significant interactions with gender with respect to incidence of diabetes, indicating greater HRs in women.

We also performed analyses of incidence of diabetes in relation to anthropometric measures dichotomized according to previously suggested cut-off values (BMI >30 kg/m<sup>2</sup>, WC >102 cm for men and >88 cm for women, WHR >0.9 for men and >0.85 for women and WHtR >0.52 for men and > 0.53 for women) <sup>15, 27</sup>. In the final multivariate model, HR was 3.61(3.22-4.03) for men and 3.43(3.06-3.85) for women for BMI; HR was 3.41 (3.07-3.80) for men and 4.07(3.64-4.54) for women for WC; HR was 2.93(2.47-3.47) for men and 3.64(3.25-4.07) for women for WHR; HR was 3.27(2.89-3.70) for men and 4.03(3.62-4.50) for women for WHtR.

### **Anthropometric measures and incidence of diabetes discrimination**

All anthropometric measures significantly improved model discrimination of diabetes compared to model adjusted for age only, Table 4. The C-statistics were significantly increased when WC, WHtR or WHR was added on top of BMI and age, Table 4.

### **Discussion**

In the present population-based cohort study, we found that all measures of obesity were associated with a substantially increased incidence of diabetes, with somewhat higher point

estimates for abdominal obesity measures than for general obesity measures, i.e., BMI and BF%. The relationships were independent of measured potential confounders, including multiple biological, lifestyle, and socio-economic factors. WC, WHtR or WHR significantly improved the association with diabetes on top of BMI.

Type 2 diabetes is an increasing health problem in most countries and obesity is the major modifiable risk factor in the population. It is therefore important to identify and utilize the most appropriate measures to assess the future risk. In our study, the highest HR for diabetes was found for WHtR. This is in line with results from other studies<sup>12, 15</sup>. The suggested cut-off value for WHtR based on 34 analyses in 16 studies was 0.52 and 0.53 for diabetes, in men and women, respectively<sup>15</sup>. It has been suggested that WHtR might be a simple, more effective indicator of obesity; “keep your waist circumference to less than half your height”<sup>28</sup>.

Abdominal obesity was associated with higher HRs in women compared to men. In women, but not in men, the confidence intervals were non-overlapping between BMI and abdominal measures of obesity. There were also significant interaction terms for WC, WHR and WHtR, indicating greater HRs of abdominal obesity in women than in men. However, in a public health perspective, incidence rates and absolute numbers of diabetes are more important measures than relative risks. With this perspective, it is important to note that incidence for men in the 4th quartile of WHtR was 24 per 1000 person-years, and the corresponding rates for women 15 per 1000 person-years. Hence, even though HRs were higher in women, the absolute risk associated with abdominal obesity still is higher in men.

It is also noteworthy that abdominal measures added information on top of BMI, which suggests that both measures could be useful in some situations, in contrast to the general

obesity measures (e.g. BMI and BF%). Abdominal obesity, in terms of visceral fat tissue, which is metabolically more active than non-visceral fat, could influence the development of diabetes and increase the degree of insulin resistance<sup>29,30</sup>. Inflammation in fat tissue may link between obesity and diabetes<sup>31</sup>. It has been shown that tumor necrosis factor-alpha (TNF- $\alpha$ ) and interleukin-6 (IL-6) in obesity could alter the insulin signaling mechanism in adipose tissue and increase the susceptibility to develop insulin resistance<sup>32,33</sup>. This circumstance might explain our finding that the point estimates for abdominal anthropometric measures were higher than for BMI or BF%.

### **Strength and limitations**

The strength of the study was the large numbers of subjects and events during a long follow-up period. A main limitation of the present study is lack of information on type of diabetes. Since the subjects of the study were 45- 73 years old, it can be assumed that the vast majority of the incident cases developed type 2 diabetes since type 1 diabetes usually has early onset<sup>34</sup>. Another question is whether the study cohort was representative for the background population since the participation rate of MDC was approximately 41%. However, a previous study from the city of Malmö showed no substantial difference in basic characteristics, such as smoking and obesity, between participants in the MDC and a health survey within the Region of Skåne, Sweden, with 75% participation rate<sup>17</sup>. Another issue is to generalize our findings to other countries, especially non-European populations. Our study included 11.8% of foreign-born, of whom most were from European countries and less than 1% were born outside Europe. Since obesity is increasing in Sweden and in many other countries, while other cardiovascular risk factors, such as smoking and hypertension tend to decrease, the prevalence of obesity in this study is somewhat lower than in more recent studies<sup>35-37</sup>.

Weight and height were measured by trained nurses at baseline. These measures are easy to

measure and quite stable over time and it can be assumed that measurement errors were small. However, the impedance measurements in this study showed weaker associations with incidence of diabetes. We cannot rule out that lower precision of the bio-impedance measures, as compared to other anthropometric measures, could explain the lower HRs.

New cases of diabetes were identified from several national and local data sources<sup>38,39</sup>. The coverage of the registers is very good. Many cases were detected during re-examinations of the MDC cohort. However, diabetes can often go undetected for a long time before seeking medical care and cases that did not seek medical care might have been missed.

Change in exposure is an inherent problem in long-term cohort studies. We lack information on anthropometric measures during the follow-up. It is possible that body fat distribution in some cases changed during the 14 years follow-up. However, this is usually slow process<sup>40</sup> and, if anything weight changes should bias the estimates towards null.

## **Conclusions**

In conclusion, all anthropometric measures of obesity were associated with a substantially increased incidence of diabetes. Abdominal obesity was associated with higher incidence rates of diabetes in men, but in terms of relative risks the relationships were stronger in women. The combination of BMI and abdominal obesity measures had stronger association with diabetes than BMI alone.

## **Acknowledgments**

This work was supported by grants from the Swedish Research Council (Dnr 2011-3891; 2014-2265), the Swedish Heart and Lung Foundation (20100244; 20130249), funds from the

Skåne University Hospital and the Lundströms Foundation.

### **Conflict of interest**

The authors declare no conflict of interest.

### **Key points**

- All anthropometric measures of obesity were associated with a substantially increased incidence of diabetes, with higher point estimates for WHtR or WC than for BMI and BF%.
- Abdominal obesity was associated with higher incidence rates of diabetes in men than in women, but in terms of relative risks the relationships were stronger in women.
- The combination of BMI and abdominal obesity measures had stronger association with diabetes than BMI alone.

## References

- 1 Danaei G, Finucane MM, Lu Y, et al. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet* 2011;378:31-40.
- 2 Hu FB, Manson JE, Stampfer MJ, et al. Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *The New England journal of medicine* 2001;345:790-797.
- 3 Carey VJ, Walters EE, Colditz GA, et al. Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses' Health Study. *American journal of epidemiology* 1997;145:614-619.
- 4 Chan JM, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes care* 1994;17:961-969.
- 5 Klein S, Sheard NF, Pi-Sunyer X, et al. Weight management through lifestyle modification for the prevention and management of type 2 diabetes: rationale and strategies: a statement of the American Diabetes Association, the North American Association for the Study of Obesity, and the American Society for Clinical Nutrition. *Diabetes care* 2004;27:2067-2073.
- 6 Narayan KM, Boyle JP, Thompson TJ, Gregg EW, Williamson DF. Effect of BMI on lifetime risk for diabetes in the U.S. *Diabetes care* 2007;30:1562-1566.
- 7 de Koning L, Gerstein HC, Bosch J, et al. Anthropometric measures and glucose levels in a large multi-ethnic cohort of individuals at risk of developing type 2 diabetes. *Diabetologia* 2010;53:1322-1330.
- 8 Huxley R, James WP, Barzi F, et al. Ethnic comparisons of the cross-sectional relationships between measures of body size with diabetes and hypertension. *Obesity*

- reviews : an official journal of the International Association for the Study of Obesity  
2008;9 Suppl 1:53-61.
- 9 Taylor AE, Ebrahim S, Ben-Shlomo Y, et al. Comparison of the associations of body mass index and measures of central adiposity and fat mass with coronary heart disease, diabetes, and all-cause mortality: a study using data from 4 UK cohorts. *Am J Clin Nutr* 2010;91:547-556.
  - 10 Arnlöv J, Sundström J, Ingelsson E, Lind L. Impact of BMI and the metabolic syndrome on the risk of diabetes in middle-aged men. *Diabetes care* 2011;34:61-65.
  - 11 Lundgren H, Bengtsson C, Blohme G, Lapidus L, Sjöström L. Adiposity and adipose tissue distribution in relation to incidence of diabetes in women: results from a prospective population study in Gothenburg, Sweden. *Int J Obes* 1989;13:413-423.
  - 12 Schulze MB, Heidemann C, Schienkiewitz A, Bergmann MM, Hoffmann K, Boeing H. Comparison of anthropometric characteristics in predicting the incidence of type 2 diabetes in the EPIC-Potsdam study. *Diabetes care* 2006;29:1921-1923.
  - 13 McTernan CL, McTernan PG, Harte AL, Levick PL, Barnett AH, Kumar S. Resistin, central obesity, and type 2 diabetes. *Lancet* 2002;359:46-47.
  - 14 Despres JP, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature* 2006;444:881-887.
  - 15 Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 2012;13:275-286.
  - 16 Berglund G, Elmståhl S, Janzon L, Larsson SA. The Malmo Diet and Cancer Study. Design and feasibility. *J Intern Med* 1993;233:45-51.

- 17 Manjer J, Carlsson S, Elmståhl S, et al. The Malmo Diet and Cancer Study: representativity, cancer incidence and mortality in participants and non-participants. *Eur J Cancer Prev* 2001;10:489-499.
- 18 Borné Y, Smith JG, Melander O, Hedblad B, Engström G. Red cell distribution width and risk for first hospitalization due to heart failure: a population-based cohort study. *Eur J Heart Fail* 2011;13:1355-1361.
- 19 Lahmann PH, Lissner L, Gullberg B, Berglund G. Differences in body fat and central adiposity between Swedes and European immigrants: The Malmo diet and cancer study. *Obes Res* 2000;8:620-631.
- 20 Li C, Aronsson CA, Hedblad B, Gullberg B, Wirfält E, Berglund G. Ability of physical activity measurements to assess health-related risks. *Eur J Clin Nutr* 2009;63:1448-1451.
- 21 Enhörning S, Wang TJ, Nilsson PM, et al. Plasma copeptin and the risk of diabetes mellitus. *Circulation* 2010;121:2102-2108.
- 22 Melander O, Maisel AS, Almgren P, et al. Plasma proneurotensin and incidence of diabetes, cardiovascular disease, breast cancer, and mortality. *JAMA : the journal of the American Medical Association* 2012;308:1469-1475.
- 23 Melander O, Newton-Cheh C, Almgren P, et al. Novel and conventional biomarkers for prediction of incident cardiovascular events in the community. *JAMA : the journal of the American Medical Association* 2009;302:49-57.
- 24 Wang TJ, Gona P, Larson MG, et al. Multiple biomarkers for the prediction of first major cardiovascular events and death. *The New England journal of medicine* 2006;355:2631-2639.



- 25 Harrell FE, Jr., Califf RM, Pryor DB, Lee KL, Rosati RA. Evaluating the yield of medical tests. *JAMA : the journal of the American Medical Association* 1982;247:2543-2546.
- 26 Harrell FE, Jr., Lee KL, Mark DB. Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors. *Stat Med* 1996;15:361-387.
- 27 World Health Organization. Waist Circumference and Waist–Hip Ratio: Report of a WHO Expert Consultation Geneva, 8–11 December 2008.
- 28 Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *International journal of food sciences and nutrition* 2005;56:303-307.
- 29 Neeland IJ, Turer AT, Ayers CR, et al. Dysfunctional adiposity and the risk of prediabetes and type 2 diabetes in obese adults. *JAMA : the journal of the American Medical Association* 2012;308:1150-1159.
- 30 McKeigue PM, Shah B, Marmot MG. Relation of central obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in South Asians. *Lancet* 1991;337:382-386.
- 31 Wentworth JM, Naselli G, Brown WA, et al. Pro-inflammatory CD11c+CD206+ adipose tissue macrophages are associated with insulin resistance in human obesity. *Diabetes* 2010;59:1648-1656.
- 32 Hotamisligil GS. Inflammation and metabolic disorders. *Nature* 2006;444:860-867.
- 33 Vozarova B, Weyer C, Hanson K, Tataranni PA, Bogardus C, Pratley RE. Circulating interleukin-6 in relation to adiposity, insulin action, and insulin secretion. *Obes Res* 2001;9:414-417.

- 34 Maahs DM, West NA, Lawrence JM, Mayer-Davis EJ. Epidemiology of type 1 diabetes. *Endocrinology and metabolism clinics of North America* 2010;39:481-497.
- 35 Danielsson M, Talbäck M. Public health: an overview: Health in Sweden: The National Public Health Report 2012. Chapter 1. *Scand J Public Health* 2012;40:6-22.
- 36 Norberg M, Danielsson M. Overweight, cardiovascular diseases and diabetes: Health in Sweden: The National Public Health Report 2012. Chapter 7. *Scand J Public Health* 2012;40:135-163.
- 37 Giskes K, Kunst AE, Benach J, et al. Trends in smoking behaviour between 1985 and 2000 in nine European countries by education. *J Epidemiol Community Health* 2005;59:395-401.
- 38 Engström G, Smith JG, Persson M, Nilsson PM, Melander O, Hedblad B. Red cell distribution width, haemoglobin A1c and incidence of diabetes mellitus. *J Intern Med* 2014.
- 39 Enhörning S, Wang TJ, Nilsson PM, et al. Plasma copeptin and the risk of diabetes mellitus. *Circulation*;121:2102-2108.
- 40 Katzmarzyk PT, Perusse L, Malina RM, Bouchard C. Seven-year stability of indicators of obesity and adipose tissue distribution in the Canadian population. *Am J Clin Nutr* 1999;69:1123-1129.

Table 1. Baseline characteristics of the Malmö Diet and Cancer cohort (N=26 604).

<b>MDC (N=26 604)</b>	<b>Men (n=10 332)</b>	<b>Women (n=16 272)</b>
Age at screening (years)	59.1±7.1	57.3±7.9
BMI, body mass index (BMI)	26.2±3.4	25.3±4.1
Waist circumference (cm)	93.4±9.9	77.5±10.2
Waist-height ratio (WHtR)	0.53±0.1	0.47±0.1
Waist-hip ratio (WHR)	0.94±0.1	0.79±0.1
Body fat percentage (BF %)	20.7±4.9	30.7±4.9
Lipid-lowering medication (%)	4.3	1.8
Current smoker (%)	28.8	28.1
High alcohol consumption (%)	7.4	2.4
Low physical activity (%)	23.9	24.1
Married (%)	72.7	60.6
Low education (%)	45.8	39.0
Immigrant status		
Other European countries (%)	11.1	11.3
Non-European countries (%)	0.9	0.4

Values are means ± SD, unless stated otherwise.

MDC, Malmö Diet and Cancer.

Table 2: Incident diabetes in relation to different anthropometric measures among men in the MDC cohort (N=10 332).

	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Q4</b>	<i>p</i> for trend
<b>BMI (n)</b>	2583	2587	2583	2577	
Range, kg/m <sup>2</sup>	13.9-23.9	23.9-25.9	25.9-28.1	>28.1	
Diabetes, n (per 1000 p-y)	170 (4.7)	228 (6.2)	376 (10.7)	745 (23.7)	
HR <sup>1</sup>	1.00	1.32(1.08-1.61)	2.30(1.92-2.75)	5.23(4.43-6.18)	<0.001
Adjusted HR <sup>2</sup>	1.00	1.32(1.09-1.62)	2.33(1.94-2.80)	5.19(4.38-6.15)	<0.001
<b>WC (n)</b>	2448	2631	2749	2504	
Range, cm	62-86	87-92	93-99	>99	
Diabetes, n (per 1000 p-y)	141 (4.0)	267(7.2)	396(10.5)	715(23.8)	
HR <sup>1</sup>	1.00	1.79(1.46-2.20)	2.64(2.18-3.20)	6.13(5.12-7.35)	<0.001
Adjusted HR <sup>2</sup>	1.00	1.79(1.46-2.20)	2.61(2.16-3.17)	5.95(4.96-7.14)	<0.001
<b>WHtR (n)</b>	2590	2585	2570	2585	
Range	0.36-0.49	0.49-0.53	0.53-0.56	>0.56	
Diabetes, n (per 1000 p-y)	151 (4.0)	238 (6.5)	387 (11.1)	743 (24.1)	
HR <sup>1</sup>	1.00	1.60(1.31-1.97)	2.78(2.30-3.36)	6.24(5.25-7.44)	<0.001
Adjusted HR <sup>2</sup>	1.00	1.59(1.30-1.95)	2.74(2.27-3.81)	6.00(5.02-7.16)	<0.001
<b>WHR (n)</b>	2589	2582	2588	2571	
Range	0.65-0.90	0.90-0.94	0.94-0.98	>0.98	
Diabetes, n (per 1000 p-y)	162 (4.4)	297 (8.3)	394 (11.2)	666 (20.9)	
HR <sup>1</sup>	1.00	1.91 (1.57-3.31)	2.59(2.16-3.12)	5.00(4.21-5.93)	<0.001
Adjusted HR <sup>2</sup>	1.00	1.87 (1.54-2.26)	2.51(2.09-3.01)	4.71(3.96-5.60)	<0.001
<b>BF % (n)</b>	2734	2639	2375	2582	
Range, %	6-17	18-20	21-23	>24	
Diabetes, n (per 1000 p-y)	215 (5.6)	323 (8.7)	385 (11.9)	596 (18.5)	
HR <sup>1</sup>	1.00	1.57(1.32-1.87)	2.15(1.82-2.54)	3.38(2.89-3.95)	<0.001

Adjusted HR <sup>2</sup>	1.00	1.54(1.30-1.84)	2.08(1.76-2.46)	3.21(2.75-3.76)	<0.001
--------------------------	------	-----------------	-----------------	-----------------	--------

HR<sup>1</sup> hazard ratio adjusted for age.

HR<sup>2</sup> adjusted for age, use of lipid-lowering medications, current smoking, high alcohol consumption, low leisure time physical activity, low education, marital status, and immigrant status.

CI, confidence interval. p-y, person-years; BMI, body mass index; WC, waist circumference; WHtR, waist-height ratio; WHR, waist-hip ratio; BF %, body fat percentage. MDC, Malmö Diet and Cancer.

Table 3: Incident diabetes in relation to different anthropometric measures among women in the MDC cohort (N=16 272).

	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Q4</b>	<i>p</i> for trend
<b>BMI (n)</b>	4058	4090	4067	4057	
Range, kg/m <sup>2</sup>	14.0-22.4	22.4-24.6	24.6-27.5	>27.5	
Diabetes, n (per 1000 p-y)	120 (2.0)	196 (3.3)	346 (5.9)	754 (13.8)	
HR <sup>1</sup>	1.00	1.59(1.27-2.00)	2.83(2.30-3.49)	6.67(5.50-8.10)	<0.001
Adjusted HR <sup>2</sup>	1.00	1.60(1.27-2.01)	2.81(2.28-3.47)	6.42(5.27-7.81)	<0.001
<b>WC (n)</b>	4344	3545	4531	3852	
Range, cm	50-70	71-75	76-83	>84	
Diabetes, n (per 1000 p-y)	98(1.5)	170(3.2)	373(5.7)	775(15.2)	
HR <sup>1</sup>	1.00	2.09(1.63-2.68)	3.67(2.94-4.59)	9.76(7.90-12.06)	<0.001
Adjusted HR <sup>2</sup>	1.00	2.06(1.60-2.64)	3.56(2.84-4.45)	9.16(7.40-11.33)	<0.001
<b>WHtR (n)</b>	4075	4060	4076	4061	
Range	0.30-0.43	0.43-0.46	0.46-0.51	>0.51	
Diabetes, n (per 1000 p-y)	84 (1.4)	181 (3.0)	340 (5.8)	811 (15.1)	
HR <sup>1</sup>	1.00	2.15(1.66-2.72)	4.12(3.24-5.24)	10.83(8.63-13.59)	<0.001
Adjusted HR <sup>2</sup>	1.00	2.11(1.63-2.73)	3.98(3.13-5.06)	10.19(8.10-12.82)	<0.001
<b>WHR (n)</b>	4073	4073	4068	4058	
Range	0.52-0.76	0.76-0.79	0.79-0.82	>0.82	
Diabetes, n (per 1000 p-y)	111 (1.8)	250 (4.2)	351 (5.9)	704 (13.0)	
HR <sup>1</sup>	1.00	2.37 (1.90-2.97)	3.37(2.72-4.17)	7.29(5.96-8.90)	<0.001
Adjusted HR <sup>2</sup>	1.00	2.32 (1.86-2.91)	3.25(2.62-4.02)	6.75(5.52-8.25)	<0.001
<b>BF % (n)</b>	4111	3615	4927	3619	
Range, %	8-27	28-30	31-34	35-49	
Diabetes, n (per 1000 p-y)	125 (2.1)	181 (3.4)	481 (6.8)	629 (12.7)	
HR <sup>1</sup>	1.00	1.58(1.26-1.99)	3.09(2.54-3.77)	5.75(4.73-6.98)	<0.001

Adjusted HR <sup>2</sup>	1.00	1.57(1.25-1.98)	2.98(2.44-3.64)	5.39(4.42-6.57)	<0.001
--------------------------	------	-----------------	-----------------	-----------------	--------

Hazard ratio (HR)<sup>1</sup> adjusted for age.

HR<sup>2</sup> age, use of lipid-lowering medications, current smoking, high alcohol consumption, low leisure time physical activity, low education, marital status, and immigrant status.

CI, confidence interval. p-y, person-years; BMI, body mass index; WC, waist circumference; WHtR, waist-height ratio; WHR, waist-hip ratio; BF %, body fat percentage. MDC, Malmö Diet and Cancer.

Table 4. Model discrimination (Harrell's C statistic) with anthropometric measure added to age in a Cox regression model.

	Men		Women	
	C-statistic (CI)	ΔC	C-statistic (CI)	ΔC
Age	0.5278 (0.5137-0.5419)		0.5823 (0.5675-0.5971)	
Age+ BMI	0.6828 (0.6705-0.6972)	0.1550*	0.7116 (0.6985-0.7247)	0.1293*
Age + WC	0.6817 (0.6686-0.6948)	0.1539*	0.7333 (0.7208-0.7458)	0.1510*
Age + WHtR	0.6874 (0.6740-0.7007)	0.1596*	0.7366 (0.7244-0.7488)	0.1543*
Age + WHR	0.6630 (0.6498-0.6762)	0.1352*	0.7079 (0.6949-0.7210)	0.1256*
Age + BF%	0.6314 (0.6175-0.6453)	0.1036*	0.6893 (0.6759-0.7026)	0.1070*
Age + BMI+WC	0.6948 (0.6816-0.7080)	0.1670†	0.7365 (0.7239-0.7492)	0.1542†
Age + BMI+ WHtR	0.6986 (0.6855-0.7116)	0.1708†	0.7383 (0.7259-0.7508)	0.1560†
Age + BMI+ WHR	0.6978 (0.6847-0.7110)	0.1700†	0.7427 (0.7298-0.7556)	0.1604†

CI: confidence interval. ΔC: increment in C-statistic. BMI, body mass index; WC, waist circumference;

WHtR, waist-height ratio; WHR, waist-hip ratio; BF %, body fat percentage.

The significance levels are indicated as follows: \*p<0.001 compare to age. † p<0.001 compared to age+ BMI.