



# LUND UNIVERSITY

## 13C – A useful tool of source characterization of the carbonaceous aerosol at rural measurement sites?

Martinsson, Johan; Stenström, Kristina; Swietlicki, Erik; Olsson, Pål Axel

2015

[Link to publication](#)

*Citation for published version (APA):*

Martinsson, J., Stenström, K., Swietlicki, E., & Olsson, P. A. (2015). *δ13C – A useful tool of source characterization of the carbonaceous aerosol at rural measurement sites?*. Abstract from LUCCI Annual Report 2014/2015.

*Total number of authors:*

4

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# $\delta^{13}\text{C}$ – A useful tool of source characterization of the carbonaceous aerosol at rural measurement sites?

Johan Martinsson (a), Kristina Eriksson Stenström (a) Erik Swietlicki (a) and Pål-Axel Olsson (b)

(a) Dept of Nuclear Physics, Lund University

(b) Dept of Biology, Lund University

## Introduction

The ambient air contains various amounts of particles, aerosols. These can be of anthropogenic or natural origin. The carbonaceous fraction of the atmospheric aerosol has considerable effects on climate and human health. This fraction can be divided into organic carbon (OC) and elemental carbon (EC). EC or black carbon (BC) is formed during incomplete combustion and is strongly light absorbing and thus warms the climate. OC is emitted from various sources (fossil fuel combustion, biomass combustion and biogenic emissions) and was formerly believed to scatter solar radiation and thereby to have a cooling effect on the climate. However, recent studies [1] have shown that fractions of OC can be strongly light. Due to the different climate effects of OC and EC, and of different types of OC, large uncertainties are associated with the net effect of carbonaceous aerosols on the climate. If the sources of the atmospheric aerosols are known, both qualitatively and quantitatively, the uncertainty of the climate effect of carbonaceous aerosols can be lowered.

There are many techniques that are used to characterize the sources of atmospheric carbonaceous aerosols. Two of the most common are radiocarbon ( $^{14}\text{C}$ ) and levoglucosan [5]. These are tracers for biomass combustion and fossil fuel combustion, respectively. Stable carbon isotopes have widely been used to characterize the sources of atmospheric aerosol since the 1980's [2,3,4]. In principle, depending on the sources of particles they have specific isotopic signatures.

Several studies that have been carried out where  $\delta^{13}\text{C}$  has been measured in various environments [6,7,8,9]. When measuring  $\delta^{13}\text{C}$  next to the source the measured  $\delta^{13}\text{C}$  value reflects the  $\delta^{13}\text{C}$  value of the source. For example, if  $\delta^{13}\text{C}$  are measured in aerosols on a highly trafficated highway, the  $\delta^{13}\text{C}$  value will most probably reflect the  $\delta^{13}\text{C}$  signature of gasoline or diesel. However, many aerosol measurement stations are rural and thus influenced by a large number of various anthropogenic and biogenic activities. A problem is that many aerosol sources overlap in  $\delta^{13}\text{C}$  signatures (Figure 1). The case of overlapping values of  $\delta^{13}\text{C}$  for different sources can be difficult to handle when applying source characterization of the aerosol. One way to treat these difficulties is to measure  $\delta^{13}\text{C}$  on specific fractions of the carbonaceous aerosol. For instance, separate the OC from the EC and measure  $\delta^{13}\text{C}$  on both fractions, separately. However, at present, there are no robust standardized methods for separating OC from EC. Due to the lack of a robust separation method it is still desirable to measure  $\delta^{13}\text{C}$  on the total carbon (TC). The applicability of  $\delta^{13}\text{C}$  for source apportionment studies can thus be questioned at these rural sites.

Filters containing atmospheric aerosols were analyzed for levoglucosan and radiocarbon. Remaining filter material (TC) was analyzed for  $\delta^{13}\text{C}$  for comparison with the above mentioned tracers to see if the  $\delta^{13}\text{C}$  value is susceptible for influences of biomass and fossil fuel combustion and to evaluate if  $\delta^{13}\text{C}$  contributes to an improved source characterization at a rural sampling site.

## Methods

Sampling was conducted on the EUSAAR and EMEP background station Vavihill in southern Sweden (56°01' N, 13°09' E, 172 m.a.s.l.). The station is located on a pasture which is occasionally visited by grazing cattle. The surrounding land is mainly deciduous forest. The closest large pollution sources are the cities of Copenhagen, Malmö and Helsingborg which are located west and southwest of the station with distances of 50, 45 and 25 km, respectively.

Particles were collected weekly (168 h) from May 2008 to April 2009 on 47 mm quartz-fiber filters (n=25). After sampling, filters were placed in petri dishes, wrapped in aluminum foil and stored in a refrigerator at +5°C until analysis.

OC/EC analysis was conducted on a 0.5 cm<sup>2</sup> punch of the original filter using a DRI Model 2001 OC/EC Carbon Analyzer (Atmoslytic, Calabasas, CA, USA).

Stable isotopes of carbon were measured on a filter punch using an elemental analyzer connected to isotope ratio mass spectrometry (EA-IRMS) at the newly installed IRMS facility at the Department of Biology, Lund University.

## Results and Discussion




Figure 1. Data for particulate  $\delta^{13}\text{C}$  from various sources from the literature. <sup>1</sup>[6]; <sup>2</sup>[7]; <sup>3</sup>[8]; <sup>4</sup>[9]; <sup>5</sup>[10]; <sup>6</sup>[11]; <sup>7</sup>[12]; <sup>8</sup>[13]. The yellow area illustrate the measured  $\delta^{13}\text{C}$  range in this study.

Figure 1 and 2 shows that the  $\delta^{13}\text{C}$  varied between -26.73 ‰ and -25.54 ‰ with a mean of -26.16 ‰ during the sampling period. These values overlap with several sources, such as combustion of fossil fuels like gasoline, diesel, natural gas and fuel oil (figure 1). Within this interval we also find charcoal, C<sub>3</sub> plants and pollen. It is thus difficult to conclude any sources from such a narrow interval that overlap with the values for several potential sources.

Table 1 shows that the  $\delta^{13}\text{C}$  correlates poorly ( $0.012 < R^2 < 0.133$ ) with all the other measured parameters. The largest  $R^2$ -value (0.133) for  $\delta^{13}\text{C}$  is found when compared to  $f^{14}\text{C}$ . This is a negative relationship showing that  $\delta^{13}\text{C}$  tends to increase when  $f^{14}\text{C}$  decrease. A low  $f^{14}\text{C}$  indicates influence of fossil fuel combustion. From figure 2 we conclude that higher  $\delta^{13}\text{C}$  values also tend to correspond with fossil fuel combustion. Thus, in this data set,  $\delta^{13}\text{C}$  might act as a weak tracer for fossil fuel combustion.

The relationship between  $\delta^{13}\text{C}$  and the other parameters (levoglucosan, OC and EC) show very low  $R^2$ -values ( $0.012 < R^2 < 0.063$ ). Thus,  $\delta^{13}\text{C}$  does not contribute with any information regarding biomass combustion (levoglucosan), general combustion (EC) or organic aerosol (OC).




Figure 2.  $\delta^{13}\text{C}$  values for aerosols when the measurement periods were divided into seasons.

Table 1.  $R^2$ -values for measured parameters.

	$\delta^{13}\text{C}$	$f^{14}\text{C}$	Levoglucosan	OC	EC
$\delta^{13}\text{C}$					
$f^{14}\text{C}$	0.133				
Levoglucosan	0.047	0.210			
OC	0.012	0.079	0.344		
EC	0.063	0.193	0.518	0.413	

The overlap between sources with respect to  $\delta^{13}\text{C}$  signatures together with the poor correlation to other strong source tracers ( $^{14}\text{C}$  and levoglucosan) makes source identification and characterization of the rural aerosol TC fraction difficult using this method. Future work will aim to measure  $\delta^{13}\text{C}$  on the separated fractions of TC (OC and EC) for improved source characterization of the carbonaceous aerosol.

## References

- [1]Kirchstetter et al. (2004). Journal of Geophysical Research. 109, D21208.
- [2]Cachier, H. (1989). Aerosol Science and Technology, 10:2, 379-385.
- [3]Cachier et al. (1985). Journal of Atmospheric Chemistry. 3, 469-489.
- [4]Chesselet et al. (1981). Geophysical research letters. 8, 345-348.
- [5]Genberg et al. (2011). Atmos. Chem. Phys. 11, 11387-11400.
- [6]Widory et al. (2004). Atmospheric Environment, 38, 953-961.
- [7]Widory, D. (2006). Combustion Theory and Modelling, 10, 831-841.
- [8]Kawashima et al. (2012). Atmospheric Environment. 46, 568-579.
- [9]Agnihotri et al. (2011). Atmospheric Environment. 45, 2828-2835.
- [10]Cao et al. (2011). Atmospheric Environment. 45, 1359-1363.
- [11]Das et al. (2010). Organic Geochemistry, 41, 263-269.
- [12]Turekian et al. (1998). Chemical Geology, 152, 181-192.
- [13]Jung et al. (2011). Atmospheric Chemistry and Physics, 11, 10911-10928.

