



# LUND UNIVERSITY

## Lifetime and predissociation yield of N-14(2) b (1)Pi(u)(v=1) revisited: Effects of rotation

Lewis, BR; Gibson, ST; Sprengers, JP; Ubachs, W; Johansson, Ann; Wahlström, Claes-Göran

*Published in:*  
Journal of Chemical Physics

*DOI:*  
[10.1063/1.2137722](https://doi.org/10.1063/1.2137722)

2005

[Link to publication](#)

*Citation for published version (APA):*

Lewis, BR., Gibson, ST., Sprengers, JP., Ubachs, W., Johansson, A., & Wahlström, C-G. (2005). Lifetime and predissociation yield of N-14(2) b (1)Pi(u)(v=1) revisited: Effects of rotation. *Journal of Chemical Physics*, 123(23). <https://doi.org/10.1063/1.2137722>

*Total number of authors:*  
6

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

## Lifetime and predissociation yield of $b(v = 1)$ revisited: Effects of rotation

B. R. Lewis, S. T. Gibson, J. P. Sprengers, W. Ubachs, A. Johansson et al.

Citation: *J. Chem. Phys.* **123**, 236101 (2005); doi: 10.1063/1.2137722

View online: <http://dx.doi.org/10.1063/1.2137722>

View Table of Contents: <http://jcp.aip.org/resource/1/JCPSA6/v123/i23>

Published by the [American Institute of Physics](#).

---

### Additional information on *J. Chem. Phys.*

Journal Homepage: <http://jcp.aip.org/>

Journal Information: [http://jcp.aip.org/about/about\\_the\\_journal](http://jcp.aip.org/about/about_the_journal)

Top downloads: [http://jcp.aip.org/features/most\\_downloaded](http://jcp.aip.org/features/most_downloaded)

Information for Authors: <http://jcp.aip.org/authors>

### ADVERTISEMENT

**AIP**Advances

*Submit Now*

**Explore AIP's new  
open-access journal**

- **Article-level metrics  
now available**
- **Join the conversation!  
Rate & comment on articles**

## Lifetime and predissociation yield of $^{14}\text{N}_2 b^1\Pi_u(v=1)$ revisited: Effects of rotation

B. R. Lewis<sup>a)</sup> and S. T. Gibson

Research School of Physical Sciences and Engineering, The Australian National University, Canberra, Australian Capital Territory 0200, Australia

J. P. Sprengers and W. Ubachs

Laser Centre, Department of Physics and Astronomy, Vrije Universiteit, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands

A. Johansson and C. -G. Wahlström

Department of Physics, Lund Institute of Technology, P.O. Box 118, S-22100 Lund, Sweden

(Received 26 September 2005; accepted 24 October 2005; published online 21 December 2005)

[DOI: 10.1063/1.2137722]

The  $b^1\Pi_u(v=1)$  state leads to the second-strongest feature in the fluorescence-excitation spectrum of  $\text{N}_2$  excited by synchrotron radiation in the extreme ultraviolet (EUV).<sup>1</sup> The  $b^1\Pi_u-X^1\Sigma_g^+(1, v'')$  progression is also a prominent feature in the electron-impact-induced fluorescence spectrum of  $\text{N}_2$ ,<sup>2</sup> and is important in aeronomical processes in the terrestrial and planetary atmospheres, bands having been observed in the Earth's dayglow and aurora, and in the atmosphere of Titan.<sup>3</sup>

The particular importance of  $b(v=1)$  arises because it is the only  $b$ -state level which decays principally radiatively, rather than by predissociation. Recently, Sprengers *et al.*<sup>4</sup> determined experimentally a lifetime of 2.61(10) ns for this level, using an EUV-laser-based pump-probe scheme, and deduced a corresponding predissociation yield of  $\sim 28\%$  following a semiempirical calculation of 3.61 ns for the radiative lifetime. The lifetime of Ref. 4 differs significantly from the value 1.75(26) ns measured by Oertel *et al.*,<sup>1</sup> a difference for which there is currently no explanation.

The first quantitative model for the predissociation of the lowest  $b$ -state levels of  $\text{N}_2$ , based on a coupled-channel Schrödinger-equation (CSE) treatment of the interacting  $b, c, o^1\Pi_u$ , and  $C, C'^3\Pi_u$  states, was developed by Lewis *et al.*<sup>5</sup> Recently, Haverd *et al.*<sup>6</sup> have extended this model to consider the effects of molecular rotation on the predissociation linewidths of the  $^1\Pi_u$  levels and the band oscillator strengths for the  $^1\Pi_u-X^1\Sigma_g^+(v, 0)$  transitions, finding excellent agreement with the experimental data of Stark *et al.*<sup>7</sup>

Here, building on these recent developments, we perform CSE calculations of the  $J$  dependences of the radiative, predissociative, and total lifetimes of the  $b(v=1)$  level of  $\text{N}_2$ , and, by comparison with experiment, show that the apparent discrepancy between the results of Sprengers *et al.*<sup>4</sup> and Oertel *et al.*<sup>1</sup> is an artifact, caused by a significant  $J$  dependence of the lifetime, together with differing experimental conditions. This comparison is facilitated by the analysis of further EUV pump-probe data, not included in Ref. 4. In addition, the  $b(v=1)$  predissociation yield is shown to have an extremely strong  $J$  dependence, leading to significant conse-

quences for the interpretation of  $\text{N}_2$  fluorescence spectra and aeronomical data.

The CSE model employed here has been described in detail elsewhere.<sup>5,6</sup> Briefly, the model comprises the  $b^1\Pi_u$  valence state, the  $c, o^1\Pi_u$  Rydberg states, and the  $C, C'^3\Pi_u$  valence states, coupled by mutual electrostatic interactions within the singlet and triplet manifolds and spin-orbit interactions between the manifolds. The radial Schrödinger equation for the coupled  $^1, ^3\Pi_u$  states is solved numerically, yielding the coupled-channel wave functions for the excited states, which are then combined with the ground-state radial wave function and diabatic  $b, c, o-X$  electronic transition moments, in order to form the total photodissociation cross section. The rotational dependence of the cross section is evaluated by including centrifugal terms in the ground- and excited-state Hamiltonians. Oscillator strengths and predissociation linewidths are deduced from the computed CSE cross sections by fitting Fano profiles.

Using this model with the electronic transition moments of Ref. 8, but reduced by 13.4% in order to optimize agreement with experimental oscillator strengths,<sup>4,9</sup> we have computed the  $b(v=1, J)-X(v''=0, J)$  oscillator strengths, thereby extending the  $b(v=1)$  radiative lifetime calculation of Ref. 4 to nonzero  $J$ .<sup>10</sup> The results, shown in Fig. 1(a) (long-dashed curve), show only a small  $J$  dependence; the lifetime  $\tau_{\text{rad}}$  increasing from 3.6 ns at  $J=1$  to 4.0 ns at  $J=25$ . On the other hand, predissociation linewidths  $\Gamma_{\text{pre}}$  calculated for the  $b(v=1, J)$  levels<sup>11</sup> show an extremely strong rotational dependence, increasing from  $\sim 0.0003 \text{ cm}^{-1}$  full width at half maximum (FWHM) at  $J=1$  to  $0.10 \text{ cm}^{-1}$  FWHM at  $J=25$ . Predissociative lifetimes  $\tau_{\text{pre}}$  determined from these linewidths using the relation  $\tau_{\text{pre}}(\text{ps}) = 5.309 / \Gamma_{\text{pre}}(\text{cm}^{-1} \text{ FWHM})$  are also shown in Fig. 1(a) (dashed curve). "Total" effective  $b(v=1)$  lifetimes  $\tau$ , where  $1/\tau = 1/\tau_{\text{rad}} + 1/\tau_{\text{pre}}$ , also shown in Fig. 1(a) (solid curve), vary from 2.9 ns at  $J=1$  to 53 ps at  $J=25$ . This rapid lifetime shortening with increasing  $J$  translates into a rapidly increasing predissociation yield  $\eta^{\text{pre}}$ , where  $\tau = \tau_{\text{rad}}(1 - \eta^{\text{pre}})$ , shown in Fig. 1(b), with the yield increasing from  $< 20\%$  at  $J=1$  to  $\sim 100\%$  at  $J=25$ .<sup>12</sup>




FIG. 1. (a) Rotational dependence of CSE lifetimes for the  $b^1\Pi_u(v=1)$  level of  $^{14}\text{N}_2$ . Long-dashed curve: radiative lifetime, dashed curve: predissociative lifetime, and solid curve: "total" lifetime. (b) Rotational dependence of CSE predissociation yield.

Such a large rotational effect will obviously have to be taken into account when interpreting  $\text{N}_2$  fluorescence spectra, in particular, when attempting to attribute "rotational" temperatures, and will also have consequences in terrestrial and planetary aeronomy. A similar effect has recently been deduced indirectly for the other aeronomically important state of  $\text{N}_2$ ,  $c'^1\Sigma_u^+(v=0)$ , by Liu *et al.*,<sup>13</sup> from an analysis of rotationally resolved electron-impact emission spectra. Since the  $^1\Sigma_u^+$  states of  $\text{N}_2$  rotationally couple to the  $^1\Pi_u$  manifold, which spin-orbit couples to the  $^3\Pi_u$  manifold, which, in turn, provides the predissociation route,<sup>5</sup> a strong rotational dependence of the predissociation yield is not surprising for  $c'(v=0)$ . On the other hand, in the case of  $b(v=1)$ , the strong rotational dependence must be attributed to the effects of Rydberg-valence interactions, together with the indirect mechanism of predissociation by the  $C, C'^3\Pi_u$  states, effects which are already known to produce strong dependences of  $^1\Pi_u$  predissociation on vibrational quantum number and  $\text{N}_2$  isotopomer.<sup>5</sup> Such rotational effects would be expected to be strongest, in a relative sense, for the longest-lived levels, such as  $b(v=1)$ .

Finally, it is necessary to consider the experimental  $b(v=1)$  lifetimes. In view of the above, the concept of a single lifetime becomes invalid, and the crucial issue is the effective value of  $J$  associated with a particular measurement. The fluorescence measurements of Ref. 1 appear to have been performed on a gaseous  $\text{N}_2$  sample at room temperature. If room-temperature rotational line intensities<sup>14</sup> are weighted with the CSE fluorescence yield  $1 - \eta^{\text{pre}}$ , we obtain a weighted mean of  $J=5.8$  applicable to the measurements of Oertel *et al.*<sup>1</sup> Sprengers *et al.*<sup>4</sup> reported an effective rotational temperature of 80 K for their 1 EUV + 1 UV ionization spectrum, which implicitly includes the effects of  $\eta^{\text{pre}}$ , and reported only pump-probe decay traces taken at 98.57 nm, probing the (1,0) bandhead region. Using 80 K rotational line intensities for the lines in this region, we find a weighted mean of  $J=3.3$ . Here, we report the analysis of two additional traces taken in the 98.61 nm region, yielding a mean

lifetime of 1.97(10) ns and probing levels with a weighted mean of  $J=6.5$ . Total decay linewidths associated with all of these measurements, shown in Fig. 2 (circles), are in excellent agreement with our computed total linewidth (solid curve), thus removing the discrepancies that appear to exist if  $J$  dependence is not considered. Of course, direct measurements of the predissociation yield are highly desirable, in order to confirm in detail our prediction of linewidths and predissociation yields which increase rapidly as  $J$  increases (and lifetimes which decrease rapidly).

This work was supported by Australian Research Council Discovery Program Grant No. DP0558962 and the European Community, through the Access to Research Infrastructures action of the Improving Human Potential Program, Contract No. HPRI-CT-1999-00041.




FIG. 2. Comparison between experimental (circles) and CSE (curves) rotational dependences of linewidths for the  $b^1\Pi_u(v=1)$  level of  $^{14}\text{N}_2$ . Solid circles: widths from experimental lifetimes of Ref. 4 and this work, open circle: width from experimental lifetime of Ref. 1, long-dashed curve: radiative width, dashed curve: predissociative width, and solid curve: total width.

<sup>a</sup>)Electronic mail: brenton.lewis@anu.edu.au

<sup>1</sup>H. Oertel, M. Kratzat, J. Imschweiler, and T. Noll, Chem. Phys. Lett. **82**, 552 (1981).

<sup>2</sup>G. K. James, J. M. Ajello, B. Franklin, and D. E. Shemansky, J. Phys. B **23**, 2055 (1990).

<sup>3</sup>R. R. Meier, Space Sci. Rev. **58**, 1 (1991); P. D. Feldman, D. J. Sahnou, J. W. Kruk, E. M. Murphy, and H. W. Moos, J. Geophys. Res. **106**, 8119 (2001).

<sup>4</sup>J. P. Sprengers, W. Ubachs, A. Johansson, A. L'Huillier, C. -G. Wahlström, R. Lang, B. R. Lewis, and S. T. Gibson, J. Chem. Phys. **120**, 8973 (2004).

<sup>5</sup>B. R. Lewis, S. T. Gibson, W. Zhang, H. Lefebvre-Brion, and J. -M. Robbe, J. Chem. Phys. **122**, 144302 (2005).

<sup>6</sup>V. E. Haverd, B. R. Lewis, S. T. Gibson, and G. Stark, J. Chem. Phys. **123**, 214304 (2005).

<sup>7</sup>G. Stark, K. P. Huber, K. Yoshino, P. L. Smith, and K. Ito, J. Chem. Phys.

**123**, 214303 (2005).

<sup>8</sup>D. Spelsberg and W. Meyer, J. Chem. Phys. **115**, 6438 (2001).

<sup>9</sup>G. Stark, P. L. Smith, K. P. Huber, K. Yoshino, M. Stevens, and K. Ito, J. Chem. Phys. **197**, 4809 (1992).

<sup>10</sup>Lifetimes were calculated using Eq. (4) of Ref. 4, assuming the same  $\sim 7\%$  branching to the  $a^1\Pi_g$  state.

<sup>11</sup>These CSE predissociation linewidths were also summarized graphically in Ref. 6.

<sup>12</sup>Due to inevitable uncertainty in the calculation of very narrow predissociation linewidths, the relative uncertainty in the yield increases significantly as  $J$  decreases, reaching possibly  $\sim 50\%$  at  $J=1$ .

<sup>13</sup>X. Liu, D. E. Shemansky, M. Ciocca, I. Kanik, and J. M. Ajello, Astrophys. J. **623**, 579 (2005).

<sup>14</sup>The intensities used are the products of  $^1\Pi_u - ^1\Sigma_g^+$  Hönl-London factors and 295 K Boltzmann factors.