



LUND UNIVERSITY

The European budget of isoprene and monoterpene emissions and its change since the Last Glacial Maximum

Schurgers, Guy; Hickler, Thomas; Miller, Paul; Arneth, Almut

Published in:
Geophysical Research Abstracts

2009

[Link to publication](#)

Citation for published version (APA):
Schurgers, G., Hickler, T., Miller, P., & Arneth, A. (2009). The European budget of isoprene and monoterpene emissions and its change since the Last Glacial Maximum. *Geophysical Research Abstracts*, 11.

Total number of authors:
4

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



The European budget of isoprene and monoterpene emissions and its change since the Last Glacial Maximum

G. Schurgers, T. Hickler, P.A. Miller, and A. Arneth

Lund University, Department of Physical Geography and Ecosystems Analysis, Lund, Sweden (guy.schurgers@nateko.lu.se)

Volatile organic compounds emitted from the terrestrial biosphere (BVOC) play an important role in atmospheric processes. BVOC species are oxidized in the atmosphere and influence levels of ozone. BVOCs that are less volatile, as well as their oxidation products, can be important for the formation of biogenic aerosol. Both effects influence the earth's radiation balance and thus climate, but in polluted atmospheres they can affect human health as well.

In this study we present a novel simulation of the changes in BVOC emissions (isoprene, monoterpenes) from the Last Glacial Maximum (LGM) to the present as derived from the dynamic changes in species distribution of the main European tree species. These simulations were performed with a dynamic vegetation model, driven with climate anomalies from model simulations and reconstructed CO₂ concentrations.

We can quantify three main driving factors for the changes in emissions of isoprene and monoterpenes since the LGM: (1) The changes in climate, with temperature changes as most important factor affecting plant physiology and terpenoid production, (2) a change in species distribution related to the changes in climate, causing local shifts in emission characteristics of the vegetation, and (3) a change in CO₂ concentration, causing opposing effects on the availability of different substrates for terpenoid production. Particularly the effect of atmospheric CO₂ concentration is uncertain, and sensitivity simulations showed a strong dependence of the emissions on the application of a CO₂ inhibition.