

This is an author produced version of a paper published in Diabetologia.
This paper has been peer-reviewed but does not include the final
publisher proof-corrections or journal pagination.

Citation for the published paper:

Shaat, Nael.

"Association of the E23K polymorphism in the KCNJ11 gene with
gestational diabetes mellitus"

Diabetologia, 2005 Dec;48(12):2544-51.

<http://dx.doi.org/10.1007/s00125-005-0035-0>

Access to the published version may require journal subscription.

Published with permission from: Springer

Association of the E23K Polymorphism in the *KCNJ11* Gene with Gestational Diabetes Mellitus

Short running title: N. Shaat et al.: *KCNJ11* and risk of GDM

N. Shaat¹, M. Ekelund¹, Å. Lernmark^{1, 2}, S. Ivarsson³, P. Almgren¹, K. Berntorp¹ and L. Groop¹.

¹Department of Clinical Sciences/Diabetes & Endocrinology, Malmö University Hospital, Lund University, Malmö, Sweden.

²Robert H. Williams Laboratory, University of Washington, Seattle, WA, USA.

³Department of Pediatrics, Malmö University Hospital, Lund University, Malmö, Sweden.

Corresponding author:

Nael Shaat

Department of Clinical Sciences/Diabetes & Endocrinology

Malmö University Hospital

Lund University

Malmö, Sweden

Tel: +46-40-336022

Fax: +46-40-337042

E-mail: nael.shaat@med.lu.se

Words in Abstract: 246

Words in Text: 3245 plus references, tables and figure legend.

ABSTRACT

Aims/hypothesis. Gestational diabetes mellitus (GDM) and type 2 diabetes share common pathophysiological background such as beta cell dysfunction and insulin resistance. In addition, women with GDM are at increased risk of developing type 2 diabetes later in life. Our aim was to investigate whether GDM has a similar genetic predisposition as type 2 diabetes by studying 5 common polymorphisms in 4 candidate genes that have been previously associated with type 2 diabetes.

Material. We studied 1777 unrelated Scandinavian women (588 GDM and 1189 pregnant non-diabetic controls) for the potassium inwardly-rectifying channel, subfamily J, member 11 (*KCNJ11* E23K), insulin receptor substrate 1 (*IRS1* G972R), uncoupling protein 2 (*UCP2* -866G>A) and Calpain 10 (*CAPN10* SNP43 & SNP44) polymorphisms.

Results. The EE, EK and KK genotype frequencies of the *KCNJ11* E23K polymorphism differed significantly between GDM and control women (31.5%, 52.7%, 15.8% vs. 37.3%, 48.8%, 13.9% respectively, $p=0.050$). In addition, the K-allele was increased in GDM women (odds ratio [OR] 1.17, 95% CI 1.02–1.35, $p=0.027$) and the effect was greater under a dominant model [KK/EK vs. EE] (OR 1.3, 95% CI 1.05–1.60, $p=0.016$). Analysis of the *IRS1* G972R polymorphism showed that RR homozygosity was found exclusively in GDM women (91.0%, 8.3%, 0.7% vs. 90.7%, 9.3%, 0.0% for GG, GR and RR genotypes respectively, $p=0.014$). The genotype and allele frequencies of the other studied polymorphisms did not statistically differ between GDM and control women.

Conclusion. The E23K polymorphism of the *KCNJ11* gene seems to predispose to GDM in Scandinavian women.

Keywords. Association, *CAPN10*, E23K, gene, gestational diabetes mellitus, GDM, *IRS1*, *KCNJ11*, polymorphism, Scandinavian, type 2 diabetes and *UCP2*

Abbreviations. *CAPN10*: gene encoding calpain 10. DBS: dried blood spots. ESM: electronic supplementary material. GDM: gestational diabetes mellitus. *IRS1*: gene encoding insulin receptor substrate 1. *KCNJ11*: gene encoding potassium inwardly-rectifying channel, subfamily J, member 11. OR: odds ratio. SNP: single nucleotide polymorphism. *UCP2*: gene encoding uncoupling protein 2

INTRODUCTION

Gestational diabetes mellitus (GDM) is defined as glucose intolerance that is first diagnosed during pregnancy [1]. It complicates about 2% of pregnancies in Sweden [2]. However, the prevalence varies between populations [3]. GDM results when pancreatic beta cells fail to compensate for the increased insulin resistance during pregnancy [4, 5]. However, the degree of beta cell dysfunction seems to be the predominant determinant of who will develop GDM [6]. In support of this, several studies have demonstrated that insulin secretion was substantially reduced in women with NGT and a history of GDM compared with controls after pregnancy [7, 8]. In addition, impaired insulin secretions as well as a history of GDM have been shown to predict future type 2 diabetes [9, 10]. Epidemiological studies have shown that beta cell dysfunction and insulin resistance are the main determinants of type 2 diabetes [11,12]. Furthermore, both type 2 diabetes and GDM may share other risk factors and the same genetic susceptibility [13]. Also, women with a family history of type 2 diabetes are at increased risk of developing GDM [14]. Type 2 diabetes is considered a paradigm for a multifactorial polygenic disease where common variants in several genes interact with environmental factors to cause the disease [11, 15]. We have originally shown that the Ala allele of the peroxisome proliferator activated receptor gamma (*PPARG* Pro12Ala) polymorphism has been associated with reduced risk of type 2 diabetes [16]. In addition, we and others have reported association between the E23K polymorphism of the potassium inwardly rectifying channel subfamily J, member 11 (*KCNJ11*) gene and increased risk of type 2 diabetes [15, 17]. Although the G972R polymorphism of the insulin receptor substrate 1 (*IRS1*) gene has been associated with type 2 diabetes in several studies [15], no association was found in a recent large study [18]. Variations in the calpain 10 (*CAPN10*) gene have also been

associated with type 2 diabetes [15]. A promoter polymorphism (–866G>A) in the uncoupling protein 2 (*UCP2*) gene was originally associated with reduced risk of obesity [19] as well as with reduced [20, 21] or increased [22] risk of type 2 diabetes. Genetic predisposition to GDM has been reported for variations in the insulin receptor (*INSR*), insulin-like growth factor 2 (*IGF2*), β 3-adrenergic receptor (*ADRB3*), sulphonylurea receptor 1 (*ABCC8*), *CAPN10* and mannose-binding lectin (*MBL2*) genes [23–27], whereas no associations were found for the *PPARG* Pro12Ala polymorphism or insulin gene variable number of tandem repeats (*INS VNTR*) [28]. Also, an association with the *ADRB3* W64R variant could not be replicated in subsequent studies [29, 30]. However, this might be due to lack of power, given the small effect size of most common variants, or due to ethnic heterogeneity between different populations. There are few data on the role of the *KCNJ11* E23K, *IRS1* G972R, *UCP2* –866G>A and *CAPN10* (SNP43 and SNP44) variants in the risk of GDM. Therefore, in the present study we investigated whether GDM has a genetic predisposition similar to that of type 2 diabetes by genotyping these variants in a case–control study of 1,777 pregnant Scandinavian women, 33.1% of whom had GDM.

MATERIALS AND METHODS

Study population

In southern Sweden (Skåne), all pregnant women are routinely offered a 75-g OGTT at 27-28 weeks of pregnancy. Women at high risk (previous GDM or a family history of diabetes) are also offered a 75-g OGTT at 12-13 weeks. The tests are performed in the local antenatal care clinics, using HemoCue devices (HemoCue, Ängelholm, Sweden) for capillary whole blood analysis. GDM is defined as a 2-hour capillary glucose concentration (double test) of at least 9 mmol/l according to the proposal by the European Diabetic Pregnancy Study Group [31].

We studied 1777 unrelated Scandinavian women (588 GDM women and 1189 non-diabetic pregnant controls). Women were recruited from two different resources. Two hundred and twenty seven GDM women were recruited from women referred to Malmö or Lund University Hospitals during the period from March 1996 until December 2003. The other part of GDM women (n=361) and all non-diabetic pregnant controls (n=1189) were ascertained among women participating in the Diabetes Prediction in Skåne (DiPiS) study, which is a prospective, longitudinal study for the prediction of type 1 diabetes in all newborns in southern Sweden [32]. At delivery - for DiPiS subjects - and after oral consent, a blood sample was drawn and information obtained about possible GDM or diabetes status. When the child was 2 months old and has been entered into the population registry, the parents were invited by letter to participate with their child in the DiPiS study. If the parents agreed to do so, they gave their written consent and filled out psychosocial and hereditary questionnaire including information about diabetes status in the family and their country of birth. Ethnicity was also determined using both surname and given name. Since the DiPiS study was not restricted to Swedish subjects but included immigrants

as well, we chose only women with Scandinavian background for the present study. Most of the Scandinavian women were of Swedish origin and only few women of Danish, Norwegian or Finnish origin. An informed oral and/or written voluntary consent was obtained from all study subjects. The study was approved by the ethical committee of Lund University.

Genetic analyses

DNA extraction

Total DNA was isolated from peripheral blood lymphocytes or blood samples were collected as dried blood spots (DBS) on Schleicher & Schuell, Grade 2992 filters (Schleicher and Schuell; Dassel, Germany) and punch-outs in 96-well plates were soaked in PCR amplification buffer.

Genotyping using DNA

When peripheral blood DNA was available from the subjects, genotyping of all SNPs was carried out using TaqMan allelic discrimination assay. The assay was carried out on the ABI PRISM 7900 Sequence Detection System (Applied Biosystems, Foster City, CA, USA) in a 5 µl reaction according to the manufacturer's instructions. Primers and probes were designed using Assays-by-Design (Applied Biosystems, Foster City, CA, USA). The primers and probes used are listed in electronic supplementary material (ESM) Table S1.

Genotyping using DBS

When DBS were available from the subjects, SNP genotyping was carried out using PCR-RFLP, SNaPshot or TaqMan allelic discrimination assay.

The polymorphisms *KCNJ11* E23K (rs5219), *UCP2* -866G>A (rs659366) and *IRS1* G972R (rs1801278) were genotyped using PCR-RFLP or TaqMan allelic discrimination assay (see below). The primers used for template PCR amplification

are listed in ESM Table S2. The template PCR was performed with initial 2 cycles at (4°C for 30 s followed by 98°C for 3 min), followed by holding at 80°C while the PCR mix was added. Then the PCR program was continued with an initial denaturation (94°C for 5 min), followed by 45 cycles of denaturation (94°C for 30 s), annealing (30 s) and extension (72°C for 30 s), followed by final extension (72°C for 10 min). PCR amplification was carried out with a 3x3 mm of DBS in a total volume of 40 µl containing 1 x Pharmacia Amersham buffer (Amersham Pharmacia Biotech, Uppsala, Sweden), 4 nmol each dNTP (MBI Fermentas, St Leon-Rot, Germany), 20 pmol of each primer, 20 µmol Betaine (Sigma-Aldrich, Stockholm, Sweden) and 1.5 U Taq polymerase (New England Biolabs, Beverly, MA, USA). The following restriction enzymes were used to digest the PCR products, with name, origin, incubation conditions and agarose gel concentrations in parentheses: *KCNJ11* E23K (BanII; New England Biolabs; 37°C for 4 h; 3.5% agarose gel), *UCP2* -866G>A (MluI; MBI Fermentas; 37°C for 4 h; 3% agarose gel) and *IRS1* G972R (BstNI; New England Biolabs; 60°C for 2 h; 4.5% agarose gel). PCR products were separated on agarose gel (SeaKem, Rockland, ME, USA) and stained with ethidium bromide to visualise the fragments.

CAPN10 SNP43 (rs3792267) & SNP44 (rs2975760) were genotyped using SNaPshot assay on the ABI PRISM 3100 Sequence Detection System (Applied Biosystems, Foster City, CA, USA) according to the manufacturer's instructions or using TaqMan allelic discrimination assay (see below). The template PCR was carried out as described above (see description of PCR-RFLP above) and followed by primer extension (PE). The template PCR primers are listed in ESM Table S2. The primers used for PE were: *CAPN10* SNP43 5'-GGCTTAGCCTCACCTTCAA & SNP44 5'-GACTGCAGGGCGCTCACGCTTGCTG.

The majority of the samples (n=949) were genotyped using TaqMan allelic discrimination assay for all the SNPs. Initially, a template PCR was carried out as described above (see description of PCR-RFLP above) using primers listed in ESM Table S2. The template PCR was followed by TaqMan allelic discrimination assay, which was carried out with 2 µl of the PCR product according to the manufacturer's instructions.

Genotyping and quality control

The genotyping success rate was 99.2% for cases (*KCNJ11* E23K; 100%, *UCP2* –866G>A; 98.3%, *IRS1* G972R; 99.8%, *CAPN10* SNP43; 98.1% and SNP44; 99.6%) and 99.3% for controls (*KCNJ11* E23K; 99.2%, *UCP2* –866G>A; 98.7%, *IRS1* G972R; 100%, *CAPN10* SNP43; 99.3% and SNP44; 99.3%). Genotyping accuracy, as determined from re-genotyping a random 1124 (12.6%) duplicates for all SNPs [*KCNJ11* E23K; 170 (9.6%), *UCP2* –866G>A; 212 (11.9%), *IRS1* G972R; 176 (9.9%), *CAPN10* SNP43; 297 (16.7%) and SNP44; 269 (15.1%)], was 99.82%. In addition, 38 (6.5%) of GDM women had both peripheral blood DNA and DBS and their genotype results were compared to assess the concordance between the different genotyping methods and found no discrepancies. For all SNPs, both GDM and control groups were in Hardy-Weinberg equilibrium (HWE) ($\chi^2 P > 0.05$) apart from the control group for *UCP2* –866G>A polymorphism which showed mild deviation from equilibrium (p=0.029). Our quality control measures suggest that the deviation is due to chance variation rather than genotyping error.

Statistical analyses

Significance of difference in age (mean \pm SEM) between GDM and control groups was tested by analysis of variance (ANOVA) using the Number Cruncher Statistical Systems (NCSS, Kaysville, UT, USA). Chi-square or Fisher's exact tests were used

for comparison between group frequencies. Odds ratios (ORs) and 95% confidence interval (95%CI) were obtained from logistic regression analysis. The significance of difference in allele frequencies of the *KCNJ11* E23K polymorphism between GDM and controls was also tested by 1,000 permutations. Two-sided *p*-values equal to or less than 0.05 were considered statistically significant.

Power calculation

Power calculations were performed using the Genetic Power Calculator (available at <http://ibgwww.colorado.edu/~pshaun/gpc/> [33]). Our power estimates have shown that under a multiplicative model, the present study with a sample size of 588 cases and 1189 controls have 80% power to detect an effect size of 1.23 (as measured in terms of genotypic relative risk [GRR]) when the frequency of the predisposing allele equals to 30%, with a 5% type 1 error rate.

RESULTS

Table 1 shows some phenotypic characteristics of study subjects. GDM women were slightly older than non-diabetic control women (32.2 ± 0.2 vs. 30.5 ± 0.1 , $p<0.0001$) and gained more weight (5–10kg) during pregnancy (31 vs. 19.1%, $p<0.0001$). The genotype and allele frequency distributions of all polymorphisms are presented in Table 2.

KCNJ11 E23K. The EE, EK and KK genotype frequencies of the *KCNJ11* E23K polymorphism differed significantly between GDM and control women (31.5%, 52.7%, 15.8% vs. 37.3%, 48.8%, 13.9% respectively, $p=0.050$). In addition, the K-allele was increased in GDM women (OR 1.17, 95% CI 1.02–1.35, $p=0.027$) and the effect was greater under a dominant model [KK/EK vs. EE] (OR 1.3, 95% CI 1.05–1.60, $p=0.016$) [Figure 1]. The association became slightly stronger when women who were positive for GAD65Ab, IA-2Ab or both ($n=21$, data was not available for all subjects) or when GDM women who had low fasting C-peptide levels (< 0.3 nmol/l) ($n=15$, data was not available for all subjects) were excluded (Table 3). To verify the results obtained by a chi-square test (in an exact manner), we further analysed differences in allele frequency between cases and controls using 1000 permutations and the empirical two-tailed p-value was the same as what was found by chi-square test.

IRS1 G972R. RR homozygosity of the *IRS1* G972R polymorphism was found exclusively in GDM women (91.0%, 8.3%, 0.7% vs. 90.7%, 9.3%, 0.0% for GG, GR and RR genotypes respectively, $p=0.014$) and this was statistically significant under a recessive model [RR vs. GR/GG] (0.7% vs. 0.0%, $p=0.011$). However, the R972-allele frequency was similar in both groups (OR 1.04, 95% CI 0.75–1.44, $p=0.80$).

UCP2 –866G>A. There was no significant difference in genotype frequencies of the AA, GA and GG genotypes of the *UCP2* –866G>A polymorphism between GDM and control women (15.0%, 46.4%, 38.6% vs. 13.9%, 51.7%, 34.4% respectively, $p=0.11$). Also, the allele frequencies were similar in both groups (OR 1.07, 95% CI 0.92–1.23, $p=0.38$).

CAPN10 SNP43 & SNP44. To test for linkage disequilibrium (LD) between SNP43 and SNP44, we have calculated both pairwise LD measures (D' and r^2). The D' was 1.0 with high LOD score values in cases [$D'=1.0$; CI 0.9-1.0 (LOD=17.4)] and controls [$D'=1.0$; CI 0.94-1.0 (LOD=33.3)], while the r^2 was 0.09 in both groups. Both SNPs were in HWE ($p>0.4$) for GDM and controls. There was no significant difference in the frequencies of GG, GA and AA genotypes of SNP43 between GDM and controls (52.9%, 38.1%, 9.0% vs. 52.5%, 40.3%, 7.2% respectively, $p=0.34$) or in the allele frequencies of this SNP (OR 0.96, 95% CI 0.82–1.13, $p=0.65$). Neither was there any significant difference in the CC, TC and TT genotypes of SNP44 between GDM and control women (2.9%, 30.2%, 66.9% vs. 3.6%, 29.7%, 66.7% respectively, $p=0.71$) or in the allele frequencies (OR 0.97, 95% CI 0.81–1.16, $p=0.71$).

DISCUSSION

To our knowledge, this is the largest study evaluating the role of common variants in genes predisposing for type 2 diabetes for their putative role in GDM.

KCNJ11 E23K. The key finding of the present study is the modest association between the K-allele of the E23K polymorphism in the *KCNJ11* gene and GDM. This is in line with the dominating role of beta cell dysfunction in GDM [5-9]. *In vitro* the E23K variant leads to a modestly over-active pancreatic beta cell ATP-sensitive K⁺ (K_{ATP}) channel subunit (Kir6.2) with a decreased sensitivity toward ATP resulting in decreased insulin release [34]. We have previously shown that the E23K variant in the *KCNJ11* gene was associated with decreased insulin secretion in glucose-tolerant subjects [17]. Some caution is still warranted in the interpretation of the data. We did not correct for multiple comparisons, as we primarily tested the hypothesis whether a polymorphism increasing susceptibility to type 2 diabetes also would increase susceptibility to GDM.

IRS1 G972R. IRS1 is a major substrate for the insulin receptor and is present in insulin-sensitive tissues [35]. The G972R polymorphism of the *IRS1* gene, which is located between two potential tyrosine phosphorylation sites involved in binding of the p85 subunit of PI-3 kinase has previously been associated with type 2 diabetes [15], although we could not replicate this finding in our recent large study of 9000 individuals [18], which is a common problem in genetic association studies [36]. The G972R polymorphism has also been associated with impaired beta cell function in NGT subjects as well as with reduced insulin content and impaired insulin secretion in isolated human islets [37, 38]. Our finding that homozygosity for the G972R polymorphism was found only in GDM women might indicate an increased risk for GDM in Scandinavian women. This is consistent with a report on a healthy man

homozygous for the R-allele who showed 22% reduction of fasting insulin and 48% reduction of C-peptide values as well as ~25% reduction in acute responses of insulin and C-peptide to intravenous glucose compared to carriers of the wild-type [39]. Of note, IRS1 protein level is reduced in adipose tissue of obese women with GDM [40].

***UCP2* –866G>A.** *UCP2* is a member of the mitochondrial inner membrane carrier family that is expressed in a number of tissues and cell types including the pancreatic islets [41]. Increased expression of *UCP2* in pancreatic islets is associated with increased uncoupling, decreased formation of ATP and reduced insulin secretion [42]. The A-allele of the common (–866G>A) polymorphism in the promoter of the *UCP2* gene has originally been associated with reduced risk of obesity [19]. Subsequently, a study by Wang et al. has shown association of the G-allele with increased risk of type 2 diabetes (OR= 1.43) in individuals of Northern European ancestry [20]. This has been supported in the same study by the finding that the G-allele was associated with decreased insulin secretion adjusted for the degree of insulin resistance (i.e. disposition index) in non-diabetic individuals [20]. Another study has also shown association of the A-allele with decreased risk of type 2 diabetes in Caucasians from Italy [21]. On the contrary, the AA-genotype conferred an increased risk of type 2 diabetes (OR=1.84) in Italian women [22]. In line with that study, Sesti et al. found that the A-allele was associated with decreased glucose-stimulated insulin secretion in NGT subjects as well as in human islets [43]. Here, we could not find any association between the –866G>A polymorphism and GDM in Scandinavian women despite the fact that our study had 99% power to detect the OR reported for the AA-genotype in Italian women with type 2 diabetes [22], or for the G-allele reported by Wang et al. [20] as well as for the AA-genotype reported in Caucasians [21].

CAPN10 SNP43 and SNP44. In keeping with previous results from our laboratory, SNP43 and SNP44 were in LD [44]. Whereas D' reflects recombination events between two SNPs, r^2 reflects the absolute redundancy between them. The difference we observed between D' and r^2 is mainly because SNP44 arose on the same haplotype rarer than SNP43. *CAPN10* is a cysteine protease with the gene located on chromosome 2q37 [45]. It is widely expressed in different tissues including the pancreatic islets [45, 46]. Calpain inhibitors have been shown to increase insulin secretion by accelerating exocytosis of insulin granules in mouse pancreatic islets [47]. In addition, an isoform of *CAPN10* that is a Ca^{2+} -sensor has recently been shown to trigger exocytosis in pancreatic beta cells [46]. The GG-genotype of the SNP43 has been associated with reduced *CAPN10* mRNA expression in skeletal muscle and subcutaneous adipose tissue [48, 49]. Moreover, it has been associated with increased insulin secretion [50], insulin resistance [44] as well as decreased rate of glucose oxidation [48]. Consistent with the findings in the small study by Leipold et al. for SNP43, we did not observe any significant differences in allele or genotype frequencies between GDM and controls [26]. However, these authors reported association with SNP63 as well as a haplotype combination of SNP43, 19 and 63 (121/221) [26], but no data was available on the degree of LD between these SNPs. Of note, SNP63 has been shown to be in tight LD with SNP43 and SNP44 in Scandinavians [44].

Given the fact that both GDM and type 2 diabetes have beta cell dysfunction in common, we tested the hypothesis that common variants in candidate genes that have been associated with type 2 diabetes particularly with beta cell dysfunction might also be operative in GDM. We conclude that the K allele of the E23K polymorphism in *KCNJ11* gene seems to predispose to GDM in Scandinavian women. This is

compatible with its effect on insulin secretion and the crucial role of impaired beta cell function in the pathogenesis of GDM.

ACKNOWLEDGEMENTS

Acknowledgements This work was supported by grants from the JDF-Wallenberg Foundation, Swedish Medical Research Council, the European Commission, Lundberg Foundation, Novo Nordisk Foundation and grants to the Diabetes Prediction in Skåne (DiPiS) study. We thank all the subjects for their participation, and the DiPiS research group. We are indebted to M. Svensson, A. Berglund and A. Nilsson for excellent technical assistance, and to K. Lynch for helping with data analysis.

REFERENCES

1. Metzger BE, Coustan DR, the Organizing Committee (1998) Summary and recommendations of the Fourth International Workshop-Conference on Gestational Diabetes Mellitus. *Diabetes Care* 21 [Suppl 2]:B161-B167
2. Aberg A, Rydhstroem H, Frid A (2001) Impaired glucose tolerance associated with adverse pregnancy outcome: a population-based study in southern Sweden. *Am J Obstet Gynecol* 184:77-83
3. King H (1998) Epidemiology of glucose intolerance and gestational diabetes in women of childbearing age. *Diabetes Care* 21 [Suppl 2]:B9-B13
4. Catalano PM, Huston L, Amini SB, Kalhan SC (1999) Longitudinal changes in glucose metabolism during pregnancy in obese women with normal glucose tolerance and gestational diabetes mellitus. *Am J Obstet Gynecol* 180:903-916
5. Xiang AH, Peters RK, Trigo E, Kjos SL, Lee WP, Buchanan TA (1999) Multiple metabolic defects during late pregnancy in women at high risk for type 2 diabetes. *Diabetes* 48:848-854
6. Buchanan TA, Metzger BE, Freinkel N, Bergman RN (1990) Insulin sensitivity and B-cell responsiveness to glucose during late pregnancy in lean and moderately obese women with normal glucose tolerance or mild gestational diabetes. *Am J Obstet Gynecol* 162:1008-1014
7. Ryan EA, Imes S, Liu D et al (1995) Defects in insulin secretion and action in women with a history of gestational diabetes. *Diabetes* 44:506-512
8. Osei K, Gaillard TR, Schuster DP (1998) History of gestational diabetes leads to distinct metabolic alterations in nondiabetic African-American women with a parental history of type 2 diabetes. *Diabetes Care* 21:1250-1257
9. Buchanan TA, Xiang AH, Kjos SL, Trigo E, Lee WP, Peters RK (1999) Antepartum predictors of the development of type 2 diabetes in Latino women 11-26 months after pregnancies complicated by gestational diabetes. *Diabetes* 48:2430-2436
10. Kim C, Newton KM, Knopp RH (2002) Gestational diabetes and the incidence of type 2 diabetes: a systematic review. *Diabetes Care* 25:1862-1868
11. Groop L (2000) Pathogenesis of type 2 diabetes: the relative contribution of insulin resistance and impaired insulin secretion. *Int J Clin Pract* 113 [Supp 1]:3-13
12. Lyssenko V, Almgren P, Anevski D et al (2005) Predictors of and longitudinal changes in insulin sensitivity and secretion preceding onset of type 2 diabetes. *Diabetes* 54:166-174
13. Ben-Haroush A, Yogeve Y, Hod M (2004) Epidemiology of gestational diabetes mellitus and its association with Type 2 diabetes. *Diabet Med* 21:103-113
14. Williams MA, Qiu C, Dempsey JC, Luthy DA (2003) Familial aggregation of type 2 diabetes and chronic hypertension in women with gestational diabetes mellitus. *J Reprod Med* 48:955-962
15. Parikh H, Groop L (2004) Candidate genes for type 2 diabetes. *Rev Endocr Metab Disord* 5:151-176
16. Altshuler D, Hirschhorn JN, Klannemark M et al (2000) The common PPARgamma Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nat Genet* 26:76-80

17. Florez JC, Burt N, de Bakker PI et al (2004) Haplotype structure and genotype-phenotype correlations of the sulfonylurea receptor and the islet ATP-sensitive potassium channel gene region. *Diabetes* 53:1360-1368
18. Florez JC, Sjogren M, Burt N et al (2004) Association testing in 9,000 people fails to confirm the association of the insulin receptor substrate-1 G972R polymorphism with type 2 diabetes. *Diabetes* 53:3313-3318
19. Esterbauer H, Schneitler C, Oberkofler H et al (2001) A common polymorphism in the promoter of UCP2 is associated with decreased risk of obesity in middle-aged humans. *Nat Genet* 28:178-183
20. Wang H, Chu WS, Lu T, Hasstedt SJ, Kern PA, Elbein SC (2004) Uncoupling protein-2 polymorphisms in type 2 diabetes, obesity, and insulin secretion. *Am J Physiol Endocrinol Metab* 286:E1-E7
21. Bulotta A, Ludovico O, Coco A et al (2005) The common -866G/A polymorphism in the promoter region of the UCP-2 gene is associated with reduced risk of type 2 diabetes in Caucasians from Italy. *J Clin Endocrinol Metab* 90:1176-1180
22. D'Adamo M, Perego L, Cardellini M et al (2004) The -866A/A genotype in the promoter of the human uncoupling protein 2 gene is associated with insulin resistance and increased risk of type 2 diabetes. *Diabetes* 53:1905-1910
23. Ober C, Xiang KS, Thisted RA, Indovina KA, Wason CJ, Dooley S (1989) Increased risk for gestational diabetes mellitus associated with insulin receptor and insulin-like growth factor II restriction fragment length polymorphisms. *Genet Epidemiol* 6:559-569
24. Festa A, Krugluger W, Shnawa N, Hopmeier P, Haffner SM, Scherthaner G (1999) Trp64Arg polymorphism of the beta3-adrenergic receptor gene in pregnancy: association with mild gestational diabetes mellitus. *J Clin Endocrinol Metab* 84:1695-1699
25. Rissanen J, Markkanen A, Karkkainen P et al (2000) Sulfonylurea receptor 1 gene variants are associated with gestational diabetes and type 2 diabetes but not with altered secretion of insulin. *Diabetes Care* 23:70-73
26. Leipold H, Knofler M, Gruber C, Haslinger P, Bancher-Todesca D, Worda C (2004) Calpain-10 haplotype combination and association with gestational diabetes mellitus. *Obstet Gynecol* 103:1235-1240
27. Megia A, Gallart L, Fernandez-Real JM et al (2004) Mannose-binding lectin gene polymorphisms are associated with gestational diabetes mellitus. *J Clin Endocrinol Metab* 89:5081-5087
28. Shaat N, Ekelund M, Lernmark A et al (2004) Genotypic and phenotypic differences between Arabian and Scandinavian women with gestational diabetes mellitus. *Diabetologia* 47:878-884
29. Alevizaki M, Thalassinou L, Grigorakis SI et al (2000) Study of the Trp64Arg polymorphism of the beta3-adrenergic receptor in Greek women with gestational diabetes. *Diabetes Care* 23:1079-1083
30. Tsai PJ, Ho SC, Tsai LP et al (2004) Lack of relationship between beta3-adrenergic receptor gene polymorphism and gestational diabetes mellitus in a Taiwanese population. *Metabolism* 53:1136-1139
31. Lind T, Phillips PR (1991) Influence of pregnancy on the 75-g OGTT. A prospective multicenter study. The Diabetic Pregnancy Study Group of the European Association for the Study of Diabetes. *Diabetes* 40 (Suppl 2):8-13

32. Lernmark B, Elding-Larsson H, Hansson G, Lindberg B, Lynch K, Sjoblad S (2004) Parent responses to participation in genetic screening for diabetes risk. *Pediatr Diabetes* 5:174-181
33. Purcell S, Cherny SS, Sham PC (2003) Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics* 19:149-150. Available from <http://ibgwww.colorado.edu/~pshaun/gpc/> last accessed 10 August 2005
34. Schwanstecher C, Meyer U, Schwanstecher M (2002) K(IR)6.2 polymorphism predisposes to type 2 diabetes by inducing overactivity of pancreatic beta-cell ATP-sensitive K(+) channels. *Diabetes* 51:875-879
35. Hirayama I, Tamemoto H, Yokota H et al (1999) Insulin receptor-related receptor is expressed in pancreatic beta-cells and stimulates tyrosine phosphorylation of insulin receptor substrate-1 and -2. *Diabetes* 48:1237-1244
36. Hirschhorn JN, Lohmueller K, Byrne E, Hirschhorn K. (2002) A comprehensive review of genetic association studies. *Genet Med* 4:45-61
37. Stumvoll M, Fritsche A, Volk A et al (2001) The Gly972Arg polymorphism in the insulin receptor substrate-1 gene contributes to the variation in insulin secretion in normal glucose-tolerant humans. *Diabetes* 50:882-885
38. Marchetti P, Lupi R, Federici M et al (2002) Insulin secretory function is impaired in isolated human islets carrying the Gly(972)-->Arg IRS-1 polymorphism. *Diabetes* 51:1419-1424
39. Clausen JO, Hansen T, Bjorbaek C et al (1995) Insulin resistance: interactions between obesity and a common variant of insulin receptor substrate-1. *Lancet* 346:397-402
40. Catalano PM, Nizielski SE, Shao J, Preston L, Qiao L, Friedman JE (2002) Downregulated IRS-1 and PPARgamma in obese women with gestational diabetes: relationship to FFA during pregnancy. *Am J Physiol Endocrinol Metab* 282:E522-E533
41. Zhang CY, Baffy G, Perret P et al (2001) Uncoupling protein-2 negatively regulates insulin secretion and is a major link between obesity, beta cell dysfunction, and type 2 diabetes. *Cell* 105:745-755
42. Chan CB, De Leo D, Joseph JW et al (2001) Increased uncoupling protein-2 levels in beta-cells are associated with impaired glucose-stimulated insulin secretion: mechanism of action. *Diabetes* 50:1302-1310
43. Sesti G, Cardellini M, Marini MA et al (2003) A common polymorphism in the promoter of UCP2 contributes to the variation in insulin secretion in glucose-tolerant subjects. *Diabetes* 52:1280-1283
44. Orho-Melander M, Klannemark M, Svensson MK, Ridderstrale M, Lindgren CM, Groop L (2002) Variants in the calpain-10 gene predispose to insulin resistance and elevated free fatty acid levels. *Diabetes* 51:2658-2664
45. Horikawa Y, Oda N, Cox NJ et al (2000) Genetic variation in the gene encoding calpain-10 is associated with type 2 diabetes mellitus. *Nat Genet* 26:163-175
46. Marshall C, Hitman GA, Partridge CJ et al (2005) Evidence that an isoform of calpain-10 is a regulator of exocytosis in pancreatic beta-cells. *Mol Endocrinol* 19:213-224
47. Sreenan SK, Zhou YP, Otani K et al (2001) Calpains play a role in insulin secretion and action. *Diabetes* 50:2013-2020
48. Baier LJ, Permana PA, Yang X et al (2000) A calpain-10 gene polymorphism is associated with reduced muscle mRNA levels and insulin resistance. *J Clin Invest* 106:R69-73

49. Carlsson E, Fredriksson J, Groop L, Ridderstrale M (2004) Variation in the calpain-10 gene is associated with elevated triglyceride levels and reduced adipose tissue messenger ribonucleic acid expression in obese Swedish subjects. *J Clin Endocrinol Metab* 89:3601-3605
50. Stumvoll M, Fritsche A, Madaus A et al (2001) Functional significance of the UCSNP-43 polymorphism in the CAPN10 gene for proinsulin processing and insulin secretion in nondiabetic Germans. *Diabetes* 50:2161-2163

TABLES

Table 1. Clinical characteristics of women with and without GDM

Variable	GDM % (n)	Controls % (n)	p value
Age (years)	32.2±0.2 (588)	30.5±0.1 (1189)	<0.0001
Weight gain during pregnancy			
• < 5 kg	11.8 (38/323)	4.0 (32/794)	<0.0001
• 5-10 kg	31 (100/323)	19.1 (152/794)	<0.0001
• 11-15 kg	30.6 (99/323)	39.6 (314/794)	0.005
• > 15 kg	26.6 (86/323)	37.3 (296/794)	0.0007
Smoking	10.7 (35/327)	9.5 (76/802)	0.53
At least one pregnancy before index pregnancy	59.1 (202/342)	52.9 (431/815)	0.053
Twin or triple pregnancies	2.7 (13/474)	1.4 (17/1189)	<0.0001
Insulin treatment during pregnancy	4.9 (13/263)	0.0 (0/429)	<0.0001

Table 2. Genotype and allele distributions and corresponding odds ratios for GDM.

SNP (rs number)	Genotype or allele	GDM n (%)	Controls n (%)	OR (95% CI) for GDM	OR (95% CI) for GDM, recessive model	OR (95% CI) for GDM, dominant model
<i>KCNJ11</i> E23K (rs5219)	EE	185 (31.5)	440 (37.3)			
	EK	310 (52.7)	576 (48.8)	1.28 (1.03–1.60) ^b		
	KK	93 (15.8)	164 (13.9) ^a	1.35 (0.99–1.83) ^c	1.16 (0.88–1.53)	1.3 (1.05–1.60) ^e
	K	496 (42.2)	904 (38.3)	1.17 (1.02–1.35) ^d		
<i>IRS1</i> G972R (rs1801278)	GG	534 (91)	1078 (90.7)			
	GR	49 (8.3)	111 (9.3)	0.89 (0.63–1.27)		
	RR	4 (0.7)	0 (0.0) ^f	Not applicable	Not applicable	0.96 (0.68–1.36)
	R	57 (4.8)	111 (4.7)	1.04 (0.75–1.44)		
<i>UCP2</i> -866G>A (rs659366)	AA	87 (15.0)	164 (13.9)			
	GA	268 (46.4)	607 (51.7)	0.83 (0.62–1.12)		
	GG	223 (38.6)	404 (34.4)	1.04 (0.77–1.41)	1.2 (0.98–1.47)	0.92 (0.69–1.21)
	G	714 (61.8)	1415 (60.2)	1.07 (0.92–1.23)		
<i>CAPN10</i> SNP43 (rs3792267)	AA	52 (9.0)	85 (7.2)			
	GA	220 (38.1)	476 (40.3)	0.76 (0.52–1.11)		
	GG	305 (52.9)	620 (52.5)	0.80 (0.55–1.17)	1.01 (0.83–1.24)	0.78 (0.55–1.12)
	G	830 (71.9)	1716 (72.6)	0.96 (0.82–1.13)		
<i>CAPN10</i> SNP44 (rs2975760)	TT	392 (66.9)	787 (66.7)			
	TC	177 (30.2)	351 (29.7)	1.01 (0.81–1.26)		
	CC	17 (2.9)	43 (3.6)	0.79 (0.45–1.41)	0.79 (0.45–1.40)	0.99 (0.80–1.22)
	C	211 (18.0)	437 (18.5)	0.97 (0.81–1.16)		

^a P-value for difference in genotype frequencies between women with and without GDM, p=0.050

^b P-value for comparison of EK vs. EE between women with and without GDM, p=0.028

^c P-value for comparison of KK vs. EE between women with and without GDM, p=0.056

^d P-value for difference in allele frequencies between women with and without GDM, p=0.027

^e P-value for comparison of KK+EK vs. EE between women with and without GDM, p=0.016

^f P-value for difference in genotype frequencies between women with and without GDM, p=0.014

Table 3. Genotype and allele distributions of the *KCNJ11* E23K polymorphism and corresponding odds ratios for GDM in women without islet autoantibodies or low C-peptide (<0.3 nmol/l).

SNP (rs number)	Genotype or allele	GDM n (%)	Controls n (%)	OR (95% CI) for GDM	OR (95% CI) for GDM, recessive model	OR (95% CI) for GDM, dominant model
<i>KCNJ11</i> E23K (rs5219)	EE	171 (30.8)	439 (37.3)			
	EK	299 (53.9)	574 (48.8)	1.34 (1.07–1.68) ^b		
	KK	85 (15.3)	164 (13.9) ^a	1.33 (0.97–1.83) ^c	1.12 (0.84–1.48)	1.34 (1.08–1.66) ^c
	K	469 (42.2)	902 (38.3)	1.18 (1.02–1.36) ^d		

^a P-value for difference in genotype frequencies between women with and without GDM, p=0.030


^b P-value for comparison of EK vs. EE between women with and without GDM, p=0.011

^c P-value for comparison of KK vs. EE between women with and without GDM, p=0.076

^d P-value for difference in allele frequencies between women with and without GDM, p=0.027

^e P-value for comparison of KK+EK vs. EE between women with and without GDM, p=0.008

Figure 1. Odds ratios and 95% CI for *KCNJ11* E23K polymorphism in women with GDM. The E/E genotype or the E-allele is defined as the reference (i.e., OR=1.0).



ESM Table S1. Primers and probes used for TaqMan allelic discrimination assay

SNP (rs number)	Forward primer (5'→3')	Reverse primer (5'→3')	Probe (5'FAM)	Probe (5'VIC)
<i>KCNJ11</i> E23K (rs5219)	CAGTTGCCTTTCT- TGGACACAAA	CCGAGGAATACG- TGCTGACA	CTGCCGAGCCCAG	CCTGCCAAGCC- CAG
<i>IRS1</i> G972R (rs1801278)	GGGTAGGCCTGC- AAATGCTA	GGTCGAGATGGG- CAGACT	CAGCCCTGGGA -GGT	CAGCCCCGGGA- GGT
<i>UCP2</i> -866G>A (rs659366)	GGGCCTGGTTCG- CCTTTAATT	GCCAGAGGGCC- CAATTGTT	TTCACGCATCAGT- TAC	CACGCGTCAGT- TAC
<i>CAPN10</i> SNP43 (rs3792267)	GCGCTCACGCTT- GCT	CCTCACCAAGTCA- AGGCTTAGC	AAGTAAGGCATTTG- AAG	AAGTAAGGCGT- TTGAAG
<i>CAPN10</i> SNP44 (rs2975760)	GCAGGGCGCTC- ACG	CCTCACCAAGTCA- AGGCTTAGC	CCTTACTTCACAGC- AAG	CCTTACTTCGC- AGCAAG

ESM Table S2. Primers used for template PCR

SNP & (rs number)	Forward primer (5'→3')	Reverse primer (5'→3')	Fragment size (bp)	Annealing temp. (°C)
<i>KCNJ11</i> E23K (rs5219)	GACTCTGCAGTGAG- GCCCTA	ACGTTGCAGTTG- CCTTTCTT	210	58
<i>IRS1</i> G972R (rs1801278)	CTTCTGTCAGGTGT- CCATCC	TGGCGAGGTGT- CCACGTAGC	263	62
<i>UCP2</i> -866G>A (rs659366)	CACGCTGCTTCTG- CCAGGAC	AGGCGTCAGGAG- ATGGACCG	369	63
<i>CAPN10</i> SNP43 (rs3792267) & -SNP44 (rs2975760)	GCTGGCTGGTGA- CATCAGTGC OR AGAGTTTCTGT- GTGTGGGCA	TCAGGTTCCATC- TTTCTGCCAG OR TTCTTCCTCATC- CTCACCAA	475 105	64 55