

Geophysical Research Abstracts
Vol. 18, EGU2016-12324, 2016
EGU General Assembly 2016
© Author(s) 2016. CC Attribution 3.0 License.



Neotectonic fault structures in the Lake Thun area (Switzerland)

Stefano C. Fabbri (1), Marco Herwegh (1), Fritz Schlunegger (1), Christian Hübscher (2), Benedikt J. Weiss (2), Cédric Schmelzbach (3), Heinrich Horstmeyer (3), Kaspar Merz (3), and Flavio S. Anselmetti (4)

(1) Institute of Geological Sciences, University of Bern, 3012 Bern, Switzerland, (2) Institute of Geophysics, University of Hamburg, 20146 Hamburg, Germany, (3) Institute of Geophysics, Dept. of Earth Sciences, ETH Zürich, 8092 Zürich, Switzerland, (4) Institute of Geological Sciences and Oeschger Centre of Climate Change Research, University of Bern, 3012 Bern, Switzerland

Strong historic earthquakes (i.e. intensities $I_0 \geq V$) in Switzerland are well documented by the earthquake catalogue of Switzerland ECOS-09 (e.g. Frutigen, 1729 AD, $M_w=5.2$, $I_0=VI$). Many of these strong events can be recognized paleoseismically by large subaquatic, earthquake-triggered mass movements that occur frequently in Swiss Lakes. Some of these represent the occasional occurrence of even stronger earthquakes (i.e. $M_w \sim 6.5$) in the Alpine region (Strasser et al., 2013), which are expected to produce noticeable surface ruptures. However, convincing evidence for Quaternary displacements with offset surface expressions have scarcely been found (e.g., Wiemer et al., 2009). Applying a multi-disciplinary approach, this study presents potential candidates for such faults in the larger Lake Thun area at the edge of the Alps.

The overdeepened basin of Lake Thun is situated at the northern Alpine front, which extends orthogonally to the general strike direction of the Alpine nappe front. The northern shoreline is predominantly shaped by the front of the Subalpine Molasse, which is in strong contrast to the south western shore built by the structurally higher units of the Middle and Lower Penninic nappes. This pattern with obvious differences of both lake sides suggests a major fault along the lake axis and high tectonic activity during nappe emplacement, i.e. from Eocene times throughout the Late Miocene. The area is dominated today by a strike-slip stress regime with a slight normal faulting component (Kastrup et al., 2004).

As part of a multi-disciplinary study, attempting to find potential neotectonically active fault structures in the Lake Thun area, a 2D ground penetrating radar (GPR) survey was conducted. The aim of the GPR survey was to link observations from a multichannel reflection seismic survey and a multibeam bathymetric survey carried out in Lake Thun with findings in a nearby gravel quarry revealing suspicious deformation features such as rotated gravel clast as well as significantly offset horizons.

The GPR data reveal the occurrence of several morphologic depressions from gypsum cones and clearly dipping reflections. The reflection seismic data set shows prominent reflections, characteristic seismic facies and a few sets of normal and reverse faults in the north western part of the lake basin within the glacio-lacustrine deposits that may point to a transpressional strike-slip regime. A first neotectonic analysis links these prominent lake floor features with geomorphologic patterns from the surrounding landscape, pointing to a potential candidate for a fault that is active in the Quaternary period.