

# 1 **Long-term variations in the net inflow record for Lake** 2 **Malawi**

3

4 *K. Sene<sup>1</sup> & B. Piper<sup>1</sup> & D. Wykeham<sup>1</sup> & R.T. McSweeney<sup>1</sup>*5 *<sup>1</sup>Atkins Limited, Water & Environment, United Kingdom*

6

7 *W. Tych<sup>2</sup> & K. Beven<sup>2</sup>*8 *<sup>2</sup>Lancaster Environment Centre, Lancaster University, United Kingdom*

## 9 **ABSTRACT**

10 Lake Malawi is the third largest lake in Africa and plays an important role in water  
11 supply, hydropower generation, agriculture and fisheries in the region. Lake level  
12 observations started in the 1890s and anecdotal evidence of variations dates back to the  
13 early 1800s. A chronology of lake level and outflow variations is presented together  
14 with updated estimates for the net inflow to the lake. The inflow series and selected  
15 rainfall records were also analysed using an Unobserved Component approach and,  
16 although there was little evidence of long-term trends, there was some indication of  
17 increasing inter-annual variability in recent decades. A weak quasi-periodic behaviour  
18 was also noted with a period of approximately 4-8 years. The results provide useful  
19 insights into the severity of drought and flood events in the region since the 1890s and  
20 the potential for seasonal forecasting of lake levels and outflows.

## 21 **KEYWORDS**

22 Malawi, Lake Malawi, southern Africa, climate, trend, variability, rainfall, lake levels

23

24 **INTRODUCTION**

25 Lake Malawi – with a mean surface area of approximately 28,760 km<sup>2</sup> - is the third  
26 largest lake in Africa and occupies approximately 20% of the land area of Malawi. The  
27 lake is used for water supply, fisheries and navigation and is a major tourist attraction.  
28 The river Shire, which is the sole outflow from the lake, supports extensive areas of  
29 irrigation in the Lower Shire valley together with the water supply to Malawi's second  
30 largest city, Blantyre, as well as being a major tributary of the Zambezi. The  
31 hydropower schemes in the middle reaches of the Shire supply more than 90% of the  
32 national electricity output.

33 There have been several studies of the water balance of the lake and much of the  
34 early work was linked to investigations of the potential for water supply, irrigation and  
35 hydropower. Drayton (1984) provides a useful summary of historical developments up  
36 to the 1980s which included landmark studies by Cochrane (1956), Pike (1964) and  
37 WMO (1976). The latter study was subsequently updated and extended in the early  
38 1980s (WMO 1983) and subsequent research and operational studies include those by  
39 Neuland (1984), Calder *et al.* (1994), Spigel and Coulter (1996), Shela (2000), MoIWD  
40 (2001), Jury and Gwazantini (2002), Kumambala and Ervine (2010) and Lyons *et al.*  
41 (2011).

42 The methods used have varied widely in terms of record lengths, simulation intervals  
43 (daily, monthly, annual), and approaches to infilling and extending rainfall and tributary  
44 flow records and estimating lake evaporation. Typically, the tributary inflow terms are  
45 estimated from records for key flow gauging stations and where necessary scaled up to  
46 the full catchment area using regression relationships. Similarly, lake rainfall estimates  
47 are normally derived from an area average of raingauge values from around the

48 lakeshore and the lake evaporation by averaging Penman estimates of open water  
49 evaporation from lakeshore meteorological stations. Some studies have included  
50 additional terms in the water balance such as a groundwater component (e.g. WMO,  
51 1983, Lyons *et al.*, 2011) or investigated individual components such as the lake rainfall  
52 and evaporation in more detail (e.g. Nicholson and Yin 2002). Several have also used  
53 rainfall-runoff models to explore the sensitivity of lake levels and outflows to factors  
54 such as land-use changes (Calder *et al.*,1994) and climate variations (e.g. WMO,1983,  
55 Kumambala, 2009).

56 An important finding throughout has been the extreme sensitivity of lake levels and  
57 outflows to changes in the net inflow or net basin supply to the lake, which is often  
58 called the ‘free-water’ in studies of Lake Malawi. On account of its length, this record  
59 also provides useful insights into climate variability in the region, particularly in the  
60 early 1900s before raingauge networks were first established (e.g. WMO 1983, MIWD  
61 2001). Here, we use a stochastic signal extraction technique (Young *et al.* 1999) to  
62 explore the trends and interannual variations in this record in more detail. For  
63 comparison, the same technique is applied to indicative updates to previous estimates  
64 for the lake rainfall (WMO 1983). The findings are also compared with the results from  
65 several other studies regarding the variability in lake levels and rainfall in Malawi and  
66 other parts of southern and eastern Africa.

67

68 **THE STUDY AREA**

69 The lake catchment (Fig. 1) has a land-surface area of nearly 100,000 km<sup>2</sup> of which  
70 approximately 67% is in Malawi, 27% in Tanzania and 6% in Mozambique. The main  
71 inflows arise from the rivers Bua, South Rukuru, Dwangwa and Linthipe in Malawi, the  
72 Ruhuhu and Kiwira in Tanzania, and the Songwe, which forms the border between  
73 Malawi and Tanzania. These mainly originate in the highland areas surrounding the lake  
74 which reach elevations of 2500-3000m before dropping down the rift valley escarpment  
75 to the lakeshore, which is typically at an elevation of about 500m. In the Malawi section  
76 of the catchment there are also extensive areas of plateau above the escarpment, which  
77 are typically at an elevation of 1000-1500m.

78 In Malawi the predominant climate-type is temperate (dry winters, hot summers),  
79 with regions of arid savannah and arid steppe in the south (Peel *et al.* 2007). Variations  
80 in both rainfall and river flows are linked to the passage of the Inter-Tropical  
81 Convergence Zone (ITCZ) and intrusions of Atlantic air via the Congo basin.  
82 Additional influences sometimes include the remnants from tropical cyclones in the  
83 Indian Ocean and local convectively driven rainfall associated with the annual arrival of  
84 the ITCZ and the onset of the southeasterly trade winds as it departs towards the north.  
85 These various influences result in a main rainfall season from November to April or  
86 May over much of the lake catchment. Approximately 95% of the annual rainfall  
87 typically falls in that period although there is some evidence of a transition to a mid-  
88 latitude rainfall regime during February at many locations, resulting in a temporary  
89 reduction in rainfall intensity during that month (Nicholson *et al.* 2014). The lake is  
90 large enough to have a local influence on the diurnal atmospheric circulation and  
91 differential heating of the water and land surfaces often results in onshore winds in the

92 afternoon and offshore winds in the morning, with lake rainfall tending to occur in the  
93 late night and/or early morning (e.g. WMO 1983, Nicholson and Yin 2002).

94 Due to topographic influences, the annual rainfall is often greater at the lakeshore  
95 and escarpment (1500-2000mm typically) than over the higher elevation plateau areas  
96 (700-1000mm typically) and can exceed 3000mm near the northwestern part of the lake  
97 due to wind funnelling effects in lakeside valleys (UNDP 1986). Further north and to  
98 the east, in the Tanzanian and Mozambique portions of the catchment, the annual  
99 rainfall is typically in the range 1000-2500mm. The lakeshore and/or escarpment,  
100 plateau areas, highlands and lower Shire rift valley are therefore often considered as  
101 distinct climatic zones in Malawi, although the boundaries and names used differ  
102 between individual studies (e.g. Fry *et al.* 2004, Nicholson *et al.* 2014).

103 The tributary inflows to Lake Malawi follow a similar seasonal pattern to the rainfall,  
104 typically reaching a peak in February or March and then reducing to low or zero values  
105 by the end of the dry season. Several studies have shown that due to these distinct wet  
106 and dry seasons – with little over-year storage - there is a strong correlation between  
107 rainfall and runoff on an annual basis (e.g. WMO, 1983, Drayton, 1984); also that, due  
108 both to the higher rainfall and topographic influences, the contribution to inflows from  
109 the smaller Tanzanian portion of the catchment exceeds that from areas in Malawi and  
110 is typically slightly more than half of the total tributary inflow to the lake (WMO, 1983,  
111 UNDP, 1986).

112 Lake levels have been recorded since the 1890s and some notable events since then  
113 (Table 2) include the near cessation of outflows for more than twenty years up to 1935,  
114 unusually high levels and outflows in the late 1970s and in 1980 which caused flooding

115 of lakeshore communities and areas immediately downstream, and unusually low levels  
116 and outflows associated with a widespread regional drought in the early 1990s. Since  
117 1965, the lake outflows have been controlled at a barrage – the Kamuzu Barrage –  
118 which is situated near Liwonde about 83km downstream from the lake outlet. Some  
119 estimates suggest that by the 1990s the cumulative influence of the temporary bunds  
120 built during construction of the barrage and then during subsequent operations led to  
121 lake levels being up to 0.4-0.8m higher than they would have been otherwise (Drayton,  
122 1984, Shela, 2000).

## 123 **METHODOLOGY**

### 124 Lake water balance

125 Fig. 1 shows the main catchment area for Lake Malawi. For a given time interval the  
126 water balance for the lake can be expressed as:

$$127 \quad \Delta S = P - E + Q_{in} + Q_{GW} - Q_{out} \quad (1)$$

128 where  $\Delta S$  is the change in storage,  $P$  is the lake rainfall,  $E$  the lake evaporation,  $Q_{in}$  and  
129  $Q_{GW}$  are the catchment and groundwater inflows, and  $Q_{out}$  is the lake outflow to the  
130 Shire. Here, all flow terms are expressed in terms of a depth per unit lake area and a  
131 constant area is assumed. This assumption is an approximation but a reasonable one  
132 since based on level-area estimates presented in Lyons *et al.* (2011) at current levels the  
133 area varies by less than 1% per metre rise or fall.

134 Eq. (1) can be rewritten in the form:

$$135 \quad N = \Delta S + Q_{out} = P - E + Q_{in} + Q_{GW} \quad (2)$$

136 where  $N$  is the net inflow, net basin supply or free-water. This expresses the balance  
137 between two terms of which the first is based on levels and outflows whilst the second  
138 is based on quantities which are more difficult to estimate or observe. For example  
139 some observational challenges include the small number of long-term meteorological  
140 stations around the lake, a lack of groundwater observations, the large number of lake  
141 tributaries – some of which are ungauged - and the large spatial variations in rainfall  
142 around the lake catchment.

143 Table 1 illustrates the range of mean values suggested for the terms in the right hand  
144 side of Eq. (1). As might be expected, these types of study usually also show that  
145 estimates for the lake evaporation vary least both seasonally and from year to year. For  
146 example, based on the values presented in WMO (1983), the annual lake evaporation  
147 typically varies over a range within about 4-5% of the mean value, but the  
148 corresponding value for lake rainfall is about 24-25%; likewise the coefficients of  
149 variation for annual values are about 0.02 and 0.14 respectively. However, as can be  
150 seen from the table, the mean values across these studies typically span a wide range,  
151 although this in part reflects the different averaging periods and datasets used.

#### 152 Derivation of the net inflows

153 Given these difficulties, for this study the net inflow was estimated from the lake level  
154 and outflow terms in the water balance. These calculations were performed using  
155 published data up to the 1980s (WMO 1983, UNDP 1986) and more recent records  
156 provided by the Ministry of Irrigation and Water Development (MIWD) in Malawi.  
157 Until 1915, levels were only documented twice per year and for a single gauge, with  
158 monthly values obtained by interpolation; however since then measurements have been

159 made daily at three gauges (Chilumba, Monkey Bay, Nkhata Bay) and an average value  
160 computed, representing the mean lake level (Shela, 2000).

161 Outflows from the lake are usually recorded at the Liwonde gauge which is situated  
162 close to the Kamuzu Barrage. This river gauge – established in 1948 - was the first in  
163 Malawi and the observations are important both for operation of the barrage and for  
164 management of the hydropower and irrigation schemes further downstream. The flow  
165 record is generally considered to be of good quality and the few periods of missing data  
166 were infilled by linear interpolation in the present study. The gauge record is also a  
167 good surrogate for the outflow from the lake since this reach of the Shire is very flat,  
168 only dropping by 1-2m between the lake outlet and the barrage and with only a few  
169 minor tributary inflows, although possibly with some losses due to seepage and  
170 evaporation in Lake Malombe which lies between the lake outlet and Liwonde. An  
171 investigation of these influences (WMO 1983) suggested that on an annual basis they  
172 tend to cancel out and that even the largest seasonal differences have a negligible  
173 influence on flows at Liwonde.

174 For the period before the Liwonde gauge was established, an alternative approach  
175 needs to be used to estimate the lake outflows. Regarding the cessation of flows, some  
176 studies (e.g. WMO 1983) have suggested that outflows first stopped in 1915 but – in  
177 perhaps the most detailed review to date of historical accounts - MIWD (2001) suggest  
178 that this began in 1908. The blockage was possibly caused by sediment washed in  
179 during floods from tributaries downstream from the lake outlet and theories vary  
180 regarding its nature; for example ranging from a distinct sandbar formed at the lake  
181 outlet to more extensive sediment deposition in the river channel further downstream.



182 Lake levels then rose by 3-4m over the following 2-3 decades until outflows resumed in  
183 1935, with the blockage cleared by 1938.

184 This time sequence of events has also been adopted here, re-computing the outflows  
185 in the periods 1899-1908 and 1935-1948 using a weir formula based on the channel bed  
186 (or sill) levels assumed by MIWD (2001). The outflows were assumed to be zero in the  
187 intervening years and the observed values were used from 1948 until 2010, which was  
188 the latest year for which records were available in this study. Comparisons suggested  
189 that, on an annual basis, the results were similar to those reported previously for  
190 1899/00 to 1989/99 in MIWD (2001) and from 1954/55 to 1979/80 in WMO (1983).  
191 Here the notation 1979/80 etc. refers to the Malawi hydrological year which extends  
192 from November to October.

193 To help to assess the sensitivity of the results to these assumptions, for some of the  
194 analyses a second version of the record was used which omitted the period up to 1915 -  
195 when only two lake level readings were made per year - and from 1935 to 1947 when  
196 outflows were estimated from the weir formula. This record is called the partial net  
197 inflow series in the following text. It is also worth noting that, during the time that the  
198 blockage was present, there may have been some flood flows due to overtopping of the  
199 sandbar and/or outflows due to seepage through or beneath it; however these effects  
200 could not be quantified and are therefore an additional source of uncertainty in the  
201 analyses.

## 202 Lake rainfall estimates

203 Whilst the focus in this paper is on the long-term net inflow record, it was also  
204 considered useful to make some comparisons with previous estimates for the lake

205 rainfall. However, as noted earlier there are many challenges in deriving these values; in  
206 particular due to the sparse raingauge coverage in early years and the influence of the  
207 lake on local rainfall.

208 Perhaps some of the most detailed studies to date are those reported by WMO  
209 (1983), which was one of the final outputs from more than a decade of  
210 hydrometeorological studies in Malawi. In that study, the following two long-term  
211 rainfall records were derived:

- 212 • Lake rainfall - monthly values for the period November 1954 to October 1980  
213 derived on the basis of a weighted average of 17 raingauge records from around the  
214 lakeshore, including 4 stations in Tanzania and one on an island in the lake
- 215 • Climate index series - annual values for the years 1920/21 to 1979/90 based on a  
216 weighted average of 10 long-term raingauge records which was derived to provide  
217 an indication of the long-term variability in catchment and lake rainfall

218 Due to limitations on the raingauge data available before the 1950s, the index series was  
219 based only on records from Malawi and, of necessity, made use of records for several  
220 more distant gauges which were not used in the lake rainfall estimation procedure.  
221 Regarding the lake rainfall series, some limitations that were noted included the sparse  
222 nature of the raingauge network in the middle section of the lake due to lakeshore  
223 access difficulties, and the logistical challenges in obtaining rainfall data from islands in  
224 the lake.

225 As part of the present study, the feasibility of extending these records using the same  
226 methodology was investigated based on raingauge records obtained from more recent  
227 studies (e.g. IFAD, 2001) and from the Department of Climate Change and

228 Meteorological Services in Malawi. However, this proved not to be possible; for  
229 example for the climate index series only five of the ten gauges used in the original  
230 study appeared to have more recent data and of those records were only available for  
231 two gauges before the 1950s: Nkhota Kota and Mzimba (Table 3).

232 Instead alternative estimates were derived based on this smaller number of  
233 raingauges and the net inflow record itself. Table 4 summarises the approaches that  
234 were used which were as follows:

- 235 • WMO 1983 climate index (present study) – monthly rainfall values estimated  
236 from the WMO (1983) annual series using a typical seasonal profile
- 237 • Raingauge regression model – a multiple regression relationship between the  
238 WMO (1983) monthly lake rainfall and the records for the Nkhota Kota and  
239 Mzimba gauges
- 240 • Net inflow regression model – a linear regression relationship between the net  
241 inflows and the WMO (1983) monthly lake rainfall record

242 As part of this work double mass and time series comparisons were also made of the  
243 two raingauge records versus that for the only other gauge in the lake catchment with  
244 records dating from the 1920s, at Kasungu, and these checks showed no obvious major  
245 discrepancies.

246 For the regression analyses a Dynamic Linear Regression technique was used (Young et  
247 al. 1999) which is closely linked to the stochastic techniques described in the following  
248 section. For the purpose of estimating annual rainfall values some minor infilling of  
249 monthly values was also performed based on records for nearby gauges, where  
250 available, or long-term mean values.

251 Based on these analyses the mean values for the individual series ranged from about  
252 1414 to 1573mm for the period in common (1954/55 to 1979/80). When compared to  
253 the monthly lake rainfall series, the Nash-Sutcliffe efficiencies were about 0.90 and 0.86  
254 respectively for the raingauge and net inflow regression models and 0.86 for the WMO  
255 1983 climate index series.

#### 256 Investigations of trend and variability

257 There are many approaches to estimating the temporal characteristics of hydrological  
258 records and some commonly used techniques include linear regression (with time), tests  
259 based on sign (e.g. the Mann-Kendall test), subtracting an assumed cyclical component,  
260 and comparisons of mean values for different averaging periods. Some typical  
261 challenges include the limitations of short record lengths, dealing with missing data  
262 values, and the identification of statistically significant behaviour.

263 An approach which avoids many of these problems is to adopt methods based on the  
264 unobserved component signal extraction techniques developed for the analysis of non-  
265 stationary observations. For the analyses of the net inflow and lake rainfall records  
266 derived in this study, the Dynamic Harmonic Regression technique (UC-DHR) of  
267 Young *et al.* (1999) was used and can be considered as an extension of the classical  
268 Fourier series approach which in addition allows for time-varying parameters. This  
269 provides a powerful and computationally efficient technique for data exploration with  
270 few prior assumptions required about the nature or magnitude of any trends or quasi-  
271 periodic behaviour. The method has been used for trend identification, interpolation of  
272 missing data and forecasting for a wide range of environmental and economic

273 applications, including investigations of the impacts of land use change on runoff in the  
274 UK and Malaysia (Chappell *et al.* 2012).

275 The methodology is described in detail in the papers cited so only key details are  
276 provided. In essence though the approach used is to assume a functional form for the  
277 time varying nature of a series involving estimating changing coefficients of a harmonic  
278 regressive model by optimal filtering/smoothing operations using a combination of  
279 Kalman Filter and a Fixed Interval Smoother (KF/FIS). A recursive formulation -  
280 essentially time stepping through the data in both directions – provides both a  
281 mathematically elegant and computationally efficient approach accommodating any  
282 missing data and outliers within the methodological framework. Measures of  
283 uncertainty of the estimation results are an inherent part of the stochastic nature of this  
284 model.

285 In addition to correlation coefficients, additional more complicated performance  
286 measures known as information criteria are used to help identify optimum model  
287 metrics. Other important elements of the method include the assumed variance  
288 parameters of the stochastic model (Noise Variance Ratios in the KF/FIS formulation);  
289 these parameters define the time scale of the parametric variation.

290 Regarding the model formulation, various forms are available and the version used  
291 for this study had the following form:

$$292 \quad y_t = T_t + S_t + e_t \quad (3)$$

293 where  $y_t$  is the observed time series,  $T_t$  is a stochastic trend or low frequency  
294 component,  $S_t$  is a seasonal component, and  $e_t$  is an ‘irregular’ component, arising from

295 factors such as the observation error. This approach is sometimes referred to as spectral  
296 decomposition, as the signal is split into the following three components:

- 297 • A very slow, low frequency trend component  $T_t$
- 298 • The specific periodicity or periodicities (seasonal, diurnal, cyclic – as required in  
299 the model, and their harmonics in  $S_t$ )
- 300 • An unmodelled component  $e_t$  covering the rest of the spectrum, interpreted as  
301 the model residual

302 The seasonal component is represented by a combination of sine and cosine  
303 functions:

$$304 \quad S_t = \sum_{i=1}^N \{a_{i,t} \cos(\omega_i t) + b_{i,t} \sin(\omega_i t)\} \quad (4)$$

305 where  $a_{i,t}$  and  $b_{i,t}$  are stochastic time-varying amplitude parameters and  $\omega_i$ ,  $i=1,2,..N$  are  
306 the fundamental and harmonic frequencies associated with the periodicity in the series,  
307 in this case on an annual or sub-annual basis. Other possibilities – not required here –  
308 include the options to specify a longer-term quasi-cyclical (extra-annual) component  
309 and/or a vector of external input (i.e. exogenous) variables.

310 The extent, if any, to which each term in Eq. (3) is statistically significant is then  
311 assessed using confidence intervals computed as an inherent part of the estimation  
312 procedure, thus providing the vital model/data uncertainty information and allowing for  
313 assessment of the significance of any or all of the components of the model. The input  
314 data can include missing values if required and can be analysed for any desired time  
315 interval, including daily, monthly or annual values.

316

317 **RESULTS**

318 Annual variations in net inflows

319 Fig. 2 shows the estimated values for the annual net inflows using the full record from  
320 1899 to 2010, with the years with the greatest uncertainties in lake levels and outflows  
321 highlighted. Values are expressed as an equivalent depth over the lake surface assuming  
322 a mean surface area of 28760km<sup>2</sup>; as noted previously the change in area per metre rise  
323 or fall is thought to be small (less than 1%). Over this period, the estimated mean annual  
324 inflow value was approximately 0.3m but in some years dropped below zero, most  
325 probably when the losses exceeded the combined rainfall and runoff into the lake. Also,  
326 it can be seen that two of the key events in recent times – the low flows of the early  
327 1990s and the 1979/80 floods – were some of the most extreme in this record, with  
328 comparable dry periods only occurring in 1900/01 and 1948/49, and the high flow  
329 period unmatched.

330 Table 3 summarises these events and a number of others in the history of Lake  
331 Malawi, based on the observational record and earlier traveller's reports of variations in  
332 lake levels during the 19<sup>th</sup> century (UNDP, 1986, Nicholson and Yin, 2001).  
333 Interestingly, there is evidence that lake levels were also exceptionally low in the early  
334 part of the 19<sup>th</sup> century and Nicholson (1998) notes that a drought – defined as  
335 unusually low rainfall – prevailed for most of the period from the start of the century to  
336 the 1860s and was particularly intense in the 1820s and 1830s, affecting major lakes  
337 throughout Africa.

338 More generally, the drought of the early 1990s was widespread in southern Africa  
339 and has been linked to El Niño Southern Oscillation (ENSO) events in the period 1991-

340 1995 (e.g. Jury and Mwafulirwa, 2002). By contrast, the increase in levels in 1997/98 is  
341 thought to have been due to increased rainfall in the eastern catchments of Lake Malawi  
342 and in eastern Africa, which caused increases in lake levels as far north as Ethiopia and  
343 Sudan (Birkett *et al.*, 1999). Again there may have been an El Niño influence since this  
344 tends to cause above normal rainfall in East Africa but droughts in southern Africa  
345 (Nicholson and Selato 2000). Indeed, studies based on reanalyses from atmospheric  
346 models have shown that this event was linked to both ENSO and the Indian Ocean  
347 dipole (e.g. Reason and Jagadheesha 2005).

348 The years 1961 and 1962 also saw exceptional rainfall in East Africa with significant  
349 rises in the levels of lakes such as Lake Victoria; however, although there was also an  
350 increase in the net inflow series for Lake Malawi, this was significantly less than for the  
351 1979/80 event. During the years 2002 and 2005 there were also major droughts in  
352 Malawi (World Bank, 2009) but in terms of the net inflow do not appear particularly  
353 abnormal on an annual basis, although this may mask seasonal variations. It is also  
354 worth noting that, during the 2001/02 growing season, crop damage from short-lived  
355 heavy rainfall events may also have been a factor in the food shortages which occurred.

#### 356 Trends and variability in net inflows

357 The long-term variations in flows are also of interest and a first step in applying Eq. (4)  
358 was to select the fundamental frequency and harmonic periods to use. Following  
359 inspection of the autoregressive spectrum, intervals of 1 year and 6, 4, 3 and 2.4 months  
360 were identified. The Nash-Sutcliffe efficiency of the resulting model was about 0.87 for  
361 the full series and 0.89 for the partial series and the corresponding values for the  
362 coefficient of determination were about 0.89 and 0.90.



363 From the annual time series of net inflows (Fig. 2) there is the visual impression of  
364 an increasing trend, although perhaps with a return towards average values since the  
365 unusually dry period in the 1990s. However, when using monthly values, for the full  
366 series the model (Eq. 3) suggested a sustained positive trend until the 1930s and then  
367 another increase in the period leading up to the unusually wet years of the late-1970s.  
368 This was then followed by a precipitous fall to the 1990s and then a subsequent increase  
369 in the following years. The partial record showed similar variations. However in neither  
370 case were the changes significant when compared with 95% confidence intervals. The  
371 estimates for the trend slope, shown for the full series in Fig. 3, illustrate an additional  
372 point, which is that the rate-of-change in the trend is rarely stable and sustained changes  
373 can occur over periods of years or even decades, reflecting the long periods of drought  
374 and above average rainfall which occur in this region. Again the partial series had a  
375 similar response.

376 The model also provides estimates for the seasonal components in net inflows and  
377 Fig. 4 shows the estimated amplitudes for the three largest terms (Annual and 6 and 4  
378 months) based on the full net inflow series. As might be expected, given the distinct wet  
379 and dry seasons around the lake, the response is dominated by the annual component.  
380 For the parts of the record in which there is most confidence (i.e. based on the partial  
381 record) the model suggests that the calendar years with the largest annual amplitudes  
382 were 1950, 1963, 1979, 1989 and 2001 whilst the lowest values were in 1953, 1966,  
383 1967 and 1991.

384 Although there is always a danger of seeing periodic behaviour when there is none,  
385 the annual amplitudes do sometimes seem to alternate between high and low periods,  
386 with increasing variability since the 1940s. For example, considering the main turning

387 points in the record, the highest ‘peaks’ and ‘dips’ seem to be clustered around intervals  
388 of around 4-8 years, as shown later. In comparison, in a study of storage variations  
389 alone for Lake Malawi, Jury and Gwazantini (2002) found a biennial oscillation of 2-2.6  
390 years and a weaker oscillation of around 5.6 years. These periods are typical of those  
391 often reported for the El Niño Southern Oscillation although northern Malawi is thought  
392 to lie near a transition zone between the separate regions of influence in southern and  
393 eastern Africa mentioned earlier (e.g. Jury and Mwafulirwa 2002). There are also  
394 indications that cold (La Niña) events affect rainfall in southern Africa (e.g. Nicholson  
395 and Selato 2000) together with influences from the Indian Ocean (e.g. Saji *et al.* 1999,  
396 Nicholson 2007, Manatsa *et al.* 2011, Jury 2013) although the interactions between  
397 these various mechanisms remain an active area for research.

#### 398 Trends and variability in lake rainfall

399 Similar techniques were used to analyse the long-term lake rainfall records. Again  
400 monthly values were considered and for convenience a logarithmic transformation was  
401 used in the analyses.

402 Since combining the series might mask underlying signals, the records derived in the  
403 present study were initially analysed separately, with similar results for all three series.  
404 For the amplitudes, the annual component was again by far the largest and again there  
405 seemed to be little evidence of an increasing or decreasing trend in the periods of record  
406 either from the trend slope results or the trend values. As for the net inflows, the late  
407 1970s again appear as a high rainfall period and the early 1990s as a low rainfall period.

408 The lack of any definite trend has also been found in other studies of rainfall in  
409 Malawi and surrounding regions using different datasets and techniques. For example,

410 for the period 1960-2001, Ngongondo *et al.* 2011 found a roughly equal split between  
411 an increasing or decreasing trend in annual rainfall for the 42 raingauge records  
412 considered in Malawi, although this was only statistically significant for three of those  
413 stations. Similarly, based on an analysis of records for 71 raingauges in Malawi,  
414 including locations outside the lake catchment, Nicholson *et al.* (2014) found no long-  
415 term trends in the period 1900-2010, although noted that rainfall in the northern  
416 lakeshore and plateau areas was generally below normal in the 1990s and 2000s. Some  
417 differences were also noted in both the interannual variability and spatial coherence in  
418 records between the early and later parts of the rainfall season, which were attributed to  
419 long-term changes in atmospheric circulation.

420 In contrast, for the southern highlands of Tanzania, including parts of the Lake  
421 Malawi catchment, in an analysis for 16 raingauge records from 1970-2010 Mbululo  
422 and Nyihirani (2012) found that the wettest years were 1977/78, 1978/79, 1984/85,  
423 1988/89 and 1997/98 whilst the driest years were 1976/77, 1987/88, 1990/00, 2002/03  
424 and 2005/06. It therefore appears that there are some differences in high and low rainfall  
425 years when compared to those for Malawi, perhaps indicating a different rainfall  
426 response in this part of the lake catchment; however there were insufficient long-term  
427 records to investigate this aspect further.

428 As for the net inflow analyses, the annual amplitude values also provided some  
429 useful insights into quasi-cyclical behaviour, and a similar pattern was exhibited in all  
430 three series; in particular there appeared to be unusually low amplitudes ('dips') in  
431 hydrological years 1968, 1983, 1991 and 1990 in all three series and high values  
432 ('peaks') in 1956 and 1978.

433 This effect was less apparent in the individual rainfall records, although there were  
434 some periods with high or low values at two or more raingauges; for example lows were  
435 experienced in 1967 and 1968 and highs in 1979 and lows in 1999 for two of the three  
436 gauges. The irregular components of the rainfall series – as defined by Eq. (3) – also  
437 suggested a change in pattern towards more extreme values in more recent years for the  
438 Nkhota Khota and Kasungu gauges but the results were more mixed for the Mzimba  
439 gauge. So, although there might be some signs of increasing variability in recent  
440 decades, this did not appear to be a general result, based on this small sample of gauge  
441 records.

442 To provide a more quantitative estimate for this cyclical behaviour, typical turning  
443 points were identified manually and the time intervals between them estimated. A  
444 similar exercise was also performed for the net inflow amplitude series (in Fig. 4) and  
445 Fig. 5 shows the results of these analyses, which cover about 100 turning points in total.  
446 The distributions for the net inflows and lake rainfall were generally similar and the  
447 ranges spanned were 2-6 and 2-7 years for the lake rainfall ‘peaks’ and ‘dips’  
448 respectively and 2-10 and 3-8 years for the corresponding values for the net inflows.

449 Although subjective, this again illustrates a possible linkage to phenomena occurring  
450 on timescales of a few years, such as the El Niño Southern Oscillation or Indian Ocean  
451 Dipole. Here, before performing this analysis, the individual lake rainfall series were  
452 combined into a single annual record which, although not a statistically homogenous  
453 series, still provides some information on the relative magnitudes of rainfall in different  
454 periods, and whether dry or wet years tend to occur in succession. This series was  
455 constructed as follows, again using the terminology defined in Table 4 (and shown here  
456 as *period – series used*):

- 457 • 1899/00-1919/20 – Net inflow regression model (present study)
- 458 • 1920/21-1953/54 - WMO 1983 annual index (present study)
- 459 • 1954/55-1979/80 - WMO 1983 lake rainfall
- 460 • 1980/81-2008/09 – Rainfall regression model (present study)

461 For exploring long-term variations it is also convenient to plot the annual values for this  
462 series (Fig 6). Here Figure 6(a) shows a comparison of this combined record with the  
463 net inflows, standardised in terms of the mean values and standard deviations, and  
464 Figure 6(b) shows the rainfall series itself, in terms of the percentage departures from  
465 the mean.

466 In general terms Figure 6(a) shows a close correspondence between the standardised  
467 rainfall and net inflow series, although with some notable exceptions, such as in the late  
468 1920s and in 1983/84 and 1992/93. This helps to confirm the value of the net inflow as  
469 an indicator of regional rainfall and at a more basic level, adds confidence in the  
470 underlying records used to calculate these values. The differences that are observed  
471 could be a real-effect and/or related to errors in lake levels, outflows and/or individual  
472 raingauge records; for example, the net inflow also responds to variations in evaporation  
473 and catchment runoff which may vary in different ways to the lake rainfall in some  
474 years. In these comparisons values for the period 1899/00-1919/20 should of course be  
475 ignored since the rainfall estimates are based on the net inflows in those years (and were  
476 also ignored when considering the turning points summarised in Figs. 5(a) and 5(b)).

477 From Figure 6(b), it is also interesting that some of the most notable events in the  
478 observational records for levels and outflows appear to have been caused by rainfall  
479 shortfalls or excesses that were not extreme in terms of magnitude, but did occur over a  
480 period of years. From the records available it therefore appears that major changes in

481 levels and outflows tend to occur from prolonged periods of above or below average  
482 rainfall, rather than single unusually dry or wet years. However there is always the  
483 potential for an extreme rainfall event in an individual year to lead to a rapid rise or fall  
484 in levels.

## 485 **DISCUSSION AND CONCLUSIONS**

486 These results illustrate a number of interesting features regarding the long-term  
487 variations in the net inflows to Lake Malawi and the rainfall in its catchment area. In  
488 particular, in the 20<sup>th</sup> century, the most extreme periods in the observational record to  
489 2010 appear to have been the dry years of the early 1990s and the high inflows during  
490 the 1979/80 floods. Some other notably low inflow years were 1900/01 and 1948/49  
491 although it is of interest that the blockage at the lake outlet in the early 1900s seems to  
492 have resulted from a sustained period of low rainfall and inflows rather than from any  
493 one particular event.

494 Based on the model outputs, overall there seems to have been a slight but statistically  
495 insignificant increasing trend in the net inflows since the start of observations.  
496 However, this has been swamped by periods of low and high inflows, which can last for  
497 a decade or more in some cases. Other complicating factors may also have played a role  
498 such as changes in land use and water abstractions on the tributaries flowing into the  
499 lake. These are difficult to quantify although it is worth noting the lake catchment area  
500 remains largely rural with few major irrigation or dam schemes to date, although with  
501 widespread clearance of natural vegetation for agricultural and other purposes (e.g.  
502 Chavula *et al.* 2011). There was also little discernible trend in the rainfall records  
503 although with some evidence of increasing variability in recent decades.

504        Regarding the seasonal component of net inflows, as expected the model suggested  
505        that this was dominated by the annual contribution. There was also some evidence that  
506        the highest amplitudes seem to recur at intervals of about 4-8 years. As noted earlier  
507        these are typical of the timescales which are often cited for the El Niño Southern  
508        Oscillation and other quasi-periodic variations in the Pacific and Indian Oceans. This  
509        raises the interesting possibility of improving seasonal forecasts for the net inflows and  
510        hence lake levels and outflows based on ocean and atmospheric conditions or indices  
511        linked to these phenomena, such as the Southern Oscillation Index (e.g. Ropelewski and  
512        Jones, 1987) and the Dipole Mode Index (e.g. Saji *et al.* 1999). For example, for the  
513        lake storage alone, Jury and Gwazantini (2002) found that a regression approach based  
514        on sea surface temperatures and pressures and upper zonal winds could provide  
515        potentially useful results, and Jury (2014) – in investigations of a naturalized outflow  
516        record - found evidence that it should be possible to anticipate lake level changes by  
517        about two months for some choices of global climate variables.

518        Although this would be the most direct approach, another possibility would be to  
519        forecast net inflows from estimates for the individual terms in the water balance. This  
520        would entail using downscaled medium- to long-range meteorological forecasts for the  
521        region to estimate the lake rainfall combined with rainfall-runoff models for the  
522        tributary inflows and possibly an energy budget model for the lake evaporation.  
523        However some potential challenges in model calibration include major gaps in the flow  
524        observations for some sub-catchments and the large spatial variations in rainfall and  
525        runoff around the catchment. Previous studies have also suggested some enhancement  
526        of lake rainfall due to local variations in atmospheric circulation resulting from the

527 temperature differences between the lake surface and the surrounding land, as has been  
528 observed on some other large lakes, such as Lake Victoria in East Africa.

529 In contrast, due to the large storage capacity of the lake, the net inflow represents an  
530 accumulation of these factors, helping to integrate or smooth out these effects. The  
531 results presented here also suggest that it varies in a similar way to the lake rainfall,  
532 providing another option for estimating that parameter in the first half of the 20<sup>th</sup>  
533 century, when few raingauge records were available. This then allows insights into the  
534 nature of variations in regional rainfall during the period in which lake outflows ceased,  
535 and for the previous decade.

536 Regarding forecasting techniques, both statistical and dynamical seasonal forecasting  
537 approaches have been used operationally in southern Africa since the 1990s, particularly  
538 for commercial agriculture operations (e.g. Jury 2013). For Lake Malawi, given the  
539 many uncertainties in observations and models, a probabilistic approach would be  
540 desirable and it could also be useful to update the net inflow estimates using data  
541 assimilation techniques based on near real-time observations of lake levels and  
542 outflows. For shorter-range forecasts, there might also be advantages in using daily or  
543 10-day (decadal) values rather than monthly values, although the flow routing effects of  
544 the lake storage would become more apparent at these timescales. The application of  
545 this approach could then provide a more risk-based basis to decision-making for a  
546 number of applications, including water supply, hydropower, and irrigation operations.

#### 547 **ACKNOWLEDGEMENTS**

548 Several of the hydrological datasets used in this paper were reviewed as part of the  
549 World Bank funded Malawi Water Resources Investment Strategy for the Ministry of  
550 Irrigation and Water Development in Malawi (2010-2011). We are grateful to the



551 Ministry both for providing data and for expert advice on the quality and history of  
552 gauging station records. The Department of Climate Change and Meteorological  
553 Services in Malawi also provided updates to some of the raingauge records used in the  
554 analyses. The WMO (1983) study was performed by Dr Chris Kidd and provides a  
555 valued contribution to understanding of the hydrology of Lake Malawi.

556

## 557 REFERENCES

558 Birkett, C., Murtugudde, R. & Allan, T. 1999 Indian Ocean climate event brings floods  
559 to East Africa's lakes and the Sudd Marsh. *Geophys. Res. Letters* **26(8)**, 1031-1034.

560 Calder, I.R., Hall, R.L., Bastable, H.G., Gunston, H.M., Shela, O., Chirwa, A., Kafundu,  
561 R. 1994 The impacts of land use change on water resources in sub-Saharan Africa: a  
562 modelling study of Lake Malawi. *Journal of Hydrology* **170**, 123-135.

563 Chappell, N.A., Tych, W. 2012 Identifying step changes in single streamflow and  
564 evaporation records due to forest cover change. *Hydrol. Process* **26**, 100–116.

565 Chavula, G., Brezonik, P., Bauer, M. 2011 Land Use and Land Cover Change (LULC)  
566 in the Lake Malawi Drainage Basin, 1982-2005. *International Journal of Geosciences*  
567 **2**, 172-178.

568 Cochrane, N.J. 1957 Lake Nyasa and the Shire River. *Proc. Instn Civ. Engrs* **8**, 363-  
569 382.

570 Drayton, R.S. 1984 Variations in the level of Lake Malawi. *Hydrological Sciences*  
571 *Journal* **29**, 1-3.

572 Fry, M.J., Houghton-Carr, H.A., Folwell S.S., Butao Uka, Z. 2004 Development and  
573 implementation of GIS water resources software In Southern Africa FRIEND Phase II  
574 2000-2003 report (Eds: Meigh J, Fry M), UNESCO, Paris.

575 IFAD 2001 Smallholder Flood Plains Development Programme (SFPDP). Water  
576 Resources Assessment Study (WRAS). International Fund for Agricultural  
577 Development / Ministry of Agriculture and Irrigation.

578 Jury, M.R., Gwazantini, M.E. 2002 Climate variability in Malawi, part 2: sensitivity and  
579 prediction of lake levels. *International Journal of Climatology*, **22(11)**, 1303-1312.

580 Jury, M.R., Mwfulirwa, N.D. 2002 Climate variability in Malawi, part 1: dry summers,  
581 statistical associations and predictability. *International Journal of Climatology*, **22(11)**,  
582 1289–1302.

583 Jury, M.R. 2013 Climate prediction experiences in southern Africa 1990–2005 and key  
584 outcomes. *Natural Hazards*, **65**, 1883-1894.

585 Jury, M.R. 2014 Malawi's Shire River Fluctuations and Climate. *Journal of*  
586 *Hydrometeorology*, *October 2014*, 2039-2049.

- 587 Kumambala, P. 2009 Sustainability of Water Resources Development for Malawi with  
588 Particular Emphasis on North and Central Malawi. PhD thesis, University of Glasgow,  
589 Glasgow.
- 590 Kumambala, P.N., Ervine, A. 2010 Water Balance Model of Lake Malawi and its  
591 Sensitivity to Climate Change. *The Open Hydrology Journal* **4**, 152-162.
- 592 Lyons, R.P., Kroll, N., Scholz, C.A. 2011 An energy balance hydrologic model for the  
593 Lake Malawi Rift Basin, East Africa. *Global and Planetary Change* **75**, 83-97.
- 594 Manatsa, D., Matarira, C.H., Mukwada, G. 2011 Relative impacts of ENSO and Indian  
595 Ocean dipole/zonal mode on east SADC rainfall. *International Journal of Climatology*  
596 **31(4)**, 558–577.
- 597 Mbululo, Y., Nyihirani, F. 2012 Climate Characteristics over Southern Highlands  
598 Tanzania. *Atmospheric and Climate Sciences*, **2012(2)**, 454-463.
- 599 MIWD 2001 Integrated Water Resources Management System for the Lake Malawi and  
600 Shire River: Lake Malawi Level Control. Pre-feasibility study, Norconsult.
- 601 Neuland, H. 1984 Abnormal High Water Levels of Lake Malawi? – An attempt to  
602 assess the future behaviour of the lake water levels. *Geojournal* **9.4**, 323-334.
- 603 Ngongondo, C., Xu, C-Y., Gottschalk, L. 2011 Evaluation of spatial and temporal  
604 characteristics of rainfall in Malawi: a case of data scarce region. *Theor Appl Climatol*.  
605 **106**, 79–93.
- 606 Nicholson, S.E. 1998 Fluctuations of Rift Valley Lakes Malawi and Chilwa during  
607 historical times: a synthesis of geological, archaeological and historical information. In:  
608 Environmental Change and Response in East African Lakes (J.T. Lehman, ed.), Kluwer  
609 Academic Publishers, Dordrecht, pp. 207–231.
- 610 Nicholson, S.E. 2007. The nature of rainfall variability in Africa South of the Equator.  
611 *International Journal of Climatology*, **6(5)**, 515–530.
- 612 Nicholson, S.E., Selato, J.C. 2000 The influence of La Niña on African rainfall. *Int. J.*  
613 *Climatol.* **20**, 1761-1776.
- 614 Nicholson, S.E., Yin, X. 2001 Rainfall Conditions In Equatorial East Africa During The  
615 Nineteenth Century As Inferred From The Record Of Lake Victoria. *Climatic Change*  
616 **48**, 387–398.
- 617 Nicholson, S.E., Yin, X. 2002 Mesoscale patterns of rainfall, cloudiness and  
618 evaporation over the Great Lake of East Africa. In ‘The East African Great Lakes:  
619 Limnology, Paleoclimatology and Biodiversity’ (Eds E.O.Odada and D.O.Olago),  
620 Kluwer Academic Publishers, The Netherlands, pp. 93-119
- 621 Nicholson, S.E., Klotter, D., Chavula, G. 2014 A detailed rainfall climatology for  
622 Malawi, Southern Africa. *International Journal of Climatology* **34(2)**, 315-325.
- 623 Peel, M.C., Finlayson, B.L., McMahon, T.A. 2007 Updated world map of the Köppen-  
624 Geiger climate classification. *Hydrol. Earth Syst. Sci.* **11**, 1633–1644.
- 625 Pike, J.G. 1964 The Hydrology of Lake Nyasa. *Journal of the Institution of Water*  
626 *Engineers and Scientists* **18**, 542-564.

- 627 Reason, C.J.C., Jagadheesha, D. 2005 A model investigation of recent ENSO impacts  
628 over southern Africa. *Meteorol Atmos Phys* **89**, 181-205
- 629 Ropelewski, C.F. Jones, P.D. 1987 An extension of the Tahiti-Darwin Southern  
630 Oscillation Index. *Monthly Weather Review* **115**, 2161-2165.
- 631 Saji, N., Goswami, B., Vinayachandran, P., Yamagata, T. 1999 A dipole mode in the  
632 tropical Indian Ocean. *Nature* **401**, 360-363.
- 633 Shela, O. 2000 Naturalisation of Lake Malawi levels and Shire river flows: Challenges  
634 of Water Resources Research and Sustainable Utilisation of the Lake Malawi-Shire  
635 River System. Proc. 1st WARFSA/WaterNet Symposium: Sustainable Use of Water  
636 Resources, Maputo, 1-2 November 2000.
- 637 Spigel RH, Coulter GW (1996) Comparison of Hydrology and Physical Limnology of  
638 the East African Great Lakes: Tanganyika, Malawi, Victoria, Kivu and Turkana (with  
639 reference to some North American Great Lakes). In 'The Limnology, Climatology and  
640 Paleoclimatology of the East African Lakes' (Eds.TC Johnson. EO Odada), Gordon and  
641 Breach Publishers, pp. 103-140
- 642 UNDP 1986 Annex 2B, National Water Resources Master Plan. United Nations  
643 Development Programme Projects MLW-79-015/MLW-84-003.
- 644 WMO 1976 A Water Resources Assessment of Lake Malawi. World Meteorological  
645 Organisation, Geneva.
- 646 WMO 1983. A Water Resources Evaluation of Lake Malawi and the Shire River. World  
647 Meteorological Organization, Report No. MLW/77/012, Geneva.
- 648 World Bank 2009 Malawi: Economic Vulnerability and Disaster Risk Assessment.  
649 Drought and Flood Risk Atlas.
- 650 Young, P.C., Pedregal, D.J., Tych, W. (1999) Dynamic harmonic regression. *Journal of*  
651 *Forecasting* **18**: 369–394.

Table 1 – Examples of estimates for the annual water balance of Lake Malawi

Reference	Period	P (mm)	E (mm)	Q <sub>in</sub> (mm)	Q <sub>GW</sub> (mm)
WMO (1976) in Drayton (1984)	1953-74	1350	1610	653	-
WMO (1983)	1954-79	1414	2264	1000	380
Neuland (1984)	1954-79	1374	1605	693	-
Spigel and Coulter (1996)	Not stated	1350	1610	650	0
Nicholson and Yin (2002)	1956-80	1350	~1700-1900	-	-
Kumambala (2009)	1975-90	1272	1695	400	-
Lyons <i>et al.</i> (2011)	1992-07	955	1665	-	Negligible

Table 2 – Some key events which have influenced the levels and outflows for Lake Malawi (WMO, 1983, Drayton, 1984, UNDP, 1986, Shela, 2000, MoIWD, 2001 and other sources)

Period	Description	Period	Description
1800-1809	Levels were "...so low that local inhabitants	1900-1909	Lake levels dropped with the outflow stopped by a sandbar in 1908 (MoIWD 2001)
1810-1819	traversed dry land where a deep lake now resides" and the Ruhuhu tributary	1910-1919	No outflow. Minimum level reached in 1915 after which values rose by nearly 1m in the remainder of the decade
1820-1829	"...was completely desiccated at some time early in the century".	1920-1929	No outflow. Levels rose by nearly 2m over the decade
1830-1839	Levels may have been about 465m at the start of the century (Nicholson and Yin 2001)	1930-1939	Levels rose by about 2.5m from 1930 to a peak in 1937. Outflows resumed from 1935
1840-1849	By mid-century "Lake Malawi had risen about	1940-1949	Country-wide drought in 1948/49. The lake level was about 1.5m below the 1937 peak
1850-1859	6m and maintained this level throughout the next few decades" (Nicholson and Yin 2001)	1950-1959	Temporary bund in place at the outlet from the lake from October 1956 to July 1957
1860-1869		1960-1969	Temporary bund placed across the Shire at Liwonde in 1965 during construction of the Kamuzu Barrage, which was also commissioned in 1965. Outflows regulated from that time
1870-1879	Lake level high in 1873 (~475m; Pike, in WMO 1983), but falling in the remainder of the decade	1970-1979	Peak annual levels of about 477m reached in the years 1978, 1979 and 1980 with inundation of lakeshore areas and high flows in the Shire
1880-1889	Lake level high in 1882 (~474m; Pike, in WMO 1983) but falling in the remainder of decade	1980-1989	
1890-1899	Lake level about 470m in 1890 but rising to the mid-1890s then falling again (Pike, in WMO 1983)	1990-1999	Levels declined by about 2m from 1989 to 1997 affecting flows in the Shire and hydropower generation, in part through temporary changes to the barrage operating rules
		2000-09	Unusual rainfall patterns in the 2001/02 crop season caused both drought and flooding. There was also a country-wide drought following rainfall deficits in the 2004/05 wet season. However, lake levels varied within a range of about 1m in this decade

Table 3 – Summary of raingauge records used in the analyses

Name	Climate zone	Approximate elevation (m)	Period selected	Approximate mean annual rainfall (mm)	Description
Kasungu Boma	Plateau	1036	1925-2009	800	Moved to Kasungu airport in 1983
Mzimba	Plateau	1350	1933-2009	870	Long established gauge in a plateau region to the west of Lake Malawi
Nkhota Kota	Lakeshore	500	1922-2009	1500	Long established gauge near the lakeshore in the northwest part of the lake

Table 4 – Summary of lake rainfall and rainfall index series discussed in the text

Series	Hydrological year (Nov-Oct)	Type	Basis of approach
WMO 1983 lake rainfall	1954/55-1979/80	Monthly lake rainfall estimates	Weighted average of 17 raingauge records of which 12 were around the lakeshore in Malawi and 4 along the Tanzanian lakeshore, with the remaining gauge on Likoma Island in the Malawi part of the lake. In the weighting scheme used, the gauge records from Malawi accounted for about 80% of the total
WMO 1983 climate index	1920/21-1979/80	Annual index series	Weighted average of 10 raingauge records, all from Malawi, of which 4 were used in the above estimation procedure and the remainder were of necessity from locations more distant from the lake, but within or near the lake catchment. Approximately two-thirds of the contribution to total values was from the following 4 gauges: Nkhota Kota, Livingstonia, Karonga and Chintech
WMO 1983 climate index (present study)	1920/21-1979/80	Monthly index series	The annual WMO (1983) values disaggregated to monthly values using a seasonal profile. The profile for the Nkhota Kota gauge was used since a comparison with the WMO 1983 lake rainfall series showed this to be the most representative record, when compared with those for the Mzimba and Kasungu gauges. To help with infilling missing periods in the lake rainfall, the profile for the period to 1953/54 was used
Raingauge regression model (present study)	1933/34-2008/09	Monthly index series	A fixed parameter multiple regression relationship developed between the scaled logarithms of the Nkhota Kota and Mzimba records and the WMO 1983 lake rainfall record
Net inflow regression model (present study)	1899/00-2008/09	Monthly index series	A fixed parameter linear regression relationship between the net inflow record and the logarithm of the WMO 1983 lake rainfall record, with any negative estimated rainfall values set to zero for the purpose of this approximate analysis; the net effect of this assumption was to change the mean lake rainfall estimate by about 2-3-%




Figure 1 – Location map




Figure 2 – Estimated annual net inflows for 1900-2008; the periods in which the lake outflows were estimated and/or levels only recorded twice per year are shown as a dotted line






Figure 3 – Estimated trend slopes and confidence intervals for the full monthly net inflow estimates from 1899-2009







Figure 4 – Estimated amplitudes of the annual, 6-monthly and 4-monthly components for the full monthly net inflows from 1899-2009




(a)



(b)




(c)




(d)

Figure 5 – Comparison of the frequencies of occurrence of peaks and dips in the annual amplitude series for the lake rainfall and net inflow series (a) lake rainfall (peaks) (b) lake rainfall (dips) (c) net inflow (peaks) (d) net inflow (dips)



(a)



(b)

Figure 6 (a) comparison of the standardised net inflow and combined lake rainfall series  
 (b) annual percentage departures from the mean for the combined lake rainfall series