



RESEARCH ARTICLE

Isolation of ‘*Candidatus Nitrosocosmicus franklandus*’, a novel ureolytic soil archaeal ammonia oxidiser with tolerance to high ammonia concentration

Laura E. Lehtovirta-Morley[†], Jenna Ross[†], Linda Hink, Eva B. Weber, Cécile Gubry-Rangin, Cécile Thion[‡], James I. Prosser* and Graeme W. Nicol[‡]

Institute of Biological and Environmental Sciences, Cruickshank Building, St Machar Drive, University of Aberdeen, Aberdeen AB24 3UU, UK

*Corresponding author: Institute of Biological and Environmental Sciences, University of Aberdeen, Cruickshank Building, St Machar Drive, Aberdeen AB24 3UU, UK. Tel: +0044-1224-273254; E-mail: j.prosser@abdn.ac.uk

[†]Joint first authors.

[‡]Present address: Environmental Microbial Genomics Group, Laboratoire Ampere, École Centrale de Lyon, Université de Lyon, 69134 Ecully cedex, France.

One sentence summary: This study describes ecophysiological characteristics of the first cultivated representative of a soil archaeal group performing an important biogeochemical process, ammonia oxidation, with significance for niche specialisation and archaeal phylogeny.

Editor: Riks Laanbroek

ABSTRACT

Studies of the distribution of ammonia oxidising archaea (AOA) and bacteria (AOB) suggest distinct ecological niches characterised by ammonia concentration and pH, arising through differences in substrate affinity and ammonia tolerance. AOA form five distinct phylogenetic clades, one of which, the ‘*Nitrososphaera* sister cluster’, has no cultivated isolate. A representative of this cluster, named ‘*Candidatus Nitrosocosmicus franklandus*’, was isolated from a pH 7.5 arable soil and we propose a new cluster name: ‘*Nitrosocosmicus*’. While phylogenetic analysis of *amoA* genes indicates its association with the *Nitrososphaera* sister cluster, analysis of 16S rRNA genes provided no support for a relative branching that is consistent with a ‘sister cluster’, indicating placement within a lineage of the order *Nitrososphaerales*. ‘*Ca. N. franklandus*’ is capable of ureolytic growth and its tolerances to nitrite and ammonia are higher than in other AOA and similar to those of typical soil AOB. Similarity of other growth characteristics of ‘*Ca. N. franklandus*’ with those of typical soil AOB isolates reduces support for niche differentiation between soil AOA and AOB and suggests that AOA have a wider physiological diversity than previously suspected. In particular, the high ammonia tolerance of ‘*Ca. N. franklandus*’ suggests potential contributions to nitrification in fertilised soils.

Keywords: Thaumarchaeota; *Nitrososphaera* sister cluster; soil; ammonia inhibition; *Nitrosocosmicus*; nitrification

INTRODUCTION

Soil nitrification, the sequential oxidation of ammonia to nitrite and nitrate, is generally limited by the activity of ammonia oxidisers (AO), which perform the first step in this process. Traditionally, ammonia oxidation was thought to be

dominated by ammonia oxidising bacteria (AOB), but the cultivation of autotrophic ammonia oxidising archaea (AOA) from marine (Könneke et al. 2005) and subsequently terrestrial (Lehtovirta-Morley et al. 2011; Tourna et al. 2011) environments led to a reassessment of the microbial ecology of soil nitrification. Potential differences in archaeal and bacterial physiology

Received: 23 December 2015; Accepted: 9 March 2016

© FEMS 2016. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1. Characteristics of the ‘model’ AOA (*N. maritimus*) and AOB (*N. europaea*), six AOA soil isolates or enrichments and a range of soil AOB. Some data are not available for certain strains and cell yield and activity data for *Nitrosotalea* strains are not directly comparable when presented in terms of NH_3 . Cell yield and specific cell activity of C13 calculated using cell and *amoA* gene concentrations are labelled ^a and ^b, respectively.

Organism	Size (μm)	μ_{max} (h^{-1})	Cell yield (cells μM^{-1} NH_3)	Specific cell activity (fmol NO_2^- cell $^{-1}$ h^{-1})	Inhibitory NH_4^+ conc. (mM)	Inhibitory HNO_2 conc. (μM)	References
<i>Nitrosopumilus maritimus</i> SCM1	0.17–0.22 \times 0.5–0.9	0.033	5×10^4	0.53	2	0.028	Könneke et al. (2005)
<i>Nitrososphaera viennensis</i> EN76	0.6–0.8 diam	0.024			20	0.79	Stieglmeier et al. (2014a)
<i>Nitrosoarchaeum koreensis</i> MY1	0.3–0.5 \times 0.6–1.0		1.1×10^5	2.5	20	0.50	Jung et al. (2011)
<i>Nitrosotenuis chungbukensis</i> MY2	0.2 \times 0.7		3.4×10^5		20	0.50	Jung et al. (2014)
<i>Nitrosotalea devanaterrea</i> Nd1	0.33 \times 0.89	0.011	4.5×10^5	0.072	50	0.91–3.5	Lehtovirta-Morley et al. (2011)
<i>Nitrosotalea</i> sp. Nd2	0.2 \times 0.7	0.025	4×10^5	0.065		1.61–5.7	Lehtovirta-Morley et al. (2014)
<i>Nitrosocosmicus franklandus</i> (C13)	1.1 diam	0.024	^a 7.60×10^3 / ^b 1.29×10^5	^a 2.02 ^b 0.58	>100 mM	1.58	This study
<i>Nitrosomonas europaea</i> ATCC 19718	0.8–1.1 \times 1.0–1.7	0.052–0.066	$4.61\text{--}6.44 \times 10^3$	11	>400	31.6	Prosser (1989); Hunik, Meijer and Trammer (1992); Koops et al. (2006)
Soil AOB (data collated from approximately 25 strains)	0.3–0.8 \times 1.0–8.0	0.005–0.044	$1.38\text{--}10.6 \times 10^3$	4–23	7–50		Prosser (1989); Koops and Pommerening-Röser (2001); Koper et al. (2010); Prosser and Nicol (2012)

presented the possibility of niche differentiation between AOB and AOA. This is exemplified by the high affinity for ammonia of the marine isolate *Nitrosopumilus maritimus* (Martens-Habben et al. 2009), which explains the dominance of AOA in oligotrophic sea water, and by acidophilic growth of the soil isolates, *Nitrosotalea devanaterrea* and *Nitrosotalea* sp. Nd2, which provides an explanation for nitrification in acid soils (Lehtovirta-Morley et al. 2011, 2014). There is limited evidence for greater sensitivity of AOA to ammonia, with inhibition of growth at micromolar concentrations of ammonia, while AOB typically grow at millimolar concentrations (Koper et al. 2010; Prosser and Nicol 2012). In addition, differences in effects of organic compounds have been reported within and between AOA and AOB (Lehtovirta-Morley et al. 2011, 2014; Sayavedra-Soto and Arp 2011; Tourna et al. 2011) and there is evidence for greater nitrite sensitivity in *N. maritimus* (presented in terms of nitrous acid in Table 1). Cell mass and specific cell activity are approximately one order of magnitude lower in cultivated soil AOA than soil AOB but maximum specific growth rates are similar (Prosser and Nicol 2012; Table 1).

Characterisation of soil AOA and comparison with AOB are limited by difficulties in isolation of pure cultures, due to low specific growth rates of AO and their susceptibility to contamination by faster growing heterotrophs. Pure cultures of AOB have been available since their first isolation from soil by Frankland and Frankland (1890), although few strains have been subjected to detailed physiological analysis. In contrast, only nine AOA have been isolated to date: *N. maritimus* strains SCM1 (from a salt water aquarium) (Könneke et al. 2005), NAO2 and NAO6 (from marine surface water) (Elling et al. 2015), *N. devanaterrea* Nd1 and *Nitrosotalea* sp. Nd2 (from acidic soils) (Lehtovirta-Morley et al. 2011, 2014), *Nitrososphaera viennensis* EN76 (from garden soil) (Tourna et al. 2011), *N. gargensis* Ga9.2 (from a hot spring) (Palatinzky et al. 2015) and *Nitrosopumilus* strains PS0 and HCA1 (closely related to *N. maritimus* SCM1, isolated from coastal waters) (Qin et al. 2014). Only three of these isolates are from soil, but two soil

enrichments have also been characterised, *Nitrosoarchaeum koreensis* MY1 (Jung et al. 2011) and *Nitrosotenuis chungbukensis* MY2 (Jung et al. 2014).

AOA and AOB use the enzyme ammonia monooxygenase (AMO) in the first step of ammonia oxidation, whose three subunits, A, B, C, are encoded by *amoA*, *amoB* and *amoC* genes, respectively. The *amoA* gene has been used extensively as a marker in diversity analyses, and initial studies showed that AOA are widely distributed in most environments (soil, marine waters and sediments, freshwater, geothermal springs) (e.g. Francis et al. 2005; Reigstad et al. 2008). In the soil, AOA typically outnumber AOB (e.g. Leininger et al. 2006) and the relative abundance of phylogenetic groups within the AOA varies in different soils (Gubry-Rangin et al. 2011).

A phylogenetic study (Pester et al. 2011) of publically available archaeal *amoA* gene sequences placed all within one of five major clusters. Four of these clusters have characterised, cultured representatives (the *Nitrosopumilus*, *Nitrosotalea*, *Nitrosocaldus* and *Nitrososphaera* clusters). The fifth has a distinct, but specific association with the *Nitrososphaera* cluster, it was termed the ‘*Nitrososphaera* sister’ cluster (Pester et al. 2011) and its phylogenetic divergence occurred early in thaumarchaeotal diversification (Gubry-Rangin et al. 2015). Sequences within this group are widely distributed in soil and other environments, but typically do not comprise the major proportion of *amoA* gene sequences in high throughput sequencing studies of soil (e.g. Gubry-Rangin et al. 2011). Pester et al. (2011) found *amoA* gene sequences representative of this group in seven of nine soils sampled from four different continents, comprising up to 13% of all archaeal *amoA* gene sequences. In addition, a genome sequence of an unpublished soil enrichment culture has been deposited in GenBank (accession number CP012850.1) and has provisionally been named ‘*Candidatus Nitrosocosmicus oleophilus*’, with the proposed genus name describing ‘a widely distributed AO’ (Rhee, pers. comm.).

Table 2. Details of primers and PCR cycling conditions used in this study.

Target gene	Primer	Sequence (5'-3')	Reference	Cycling conditions
Bacterial 16S rRNA	27f 1492r	AGAGTTTGGATCMTGGCTCAG GYYACCTTGTTACGACTT	Lane (1991) Nicol et al. (2008)	95°C 5 min; 10 cycles of 94°C 30 s, 55°C 30 s, 72°C 2 min; 25 cycles of 92°C 30 s, 55°C 30 s, 72°C 2 min; 72°C 5 min
Fungal 18S rRNA	Fung5f FF390r	GGGAACCAGGACTTTTAC GAGGTCTCGTTCGTTATCG	Smit et al. (1999) Vainio and Hantula (2000)	95°C 5 min; 10 cycles of 94°C 30 s, 55°C 30 s, 72°C 50 s; 25 cycles of 92°C 30 s, 55°C 30 s, 72°C 50 s; 72°C 5 min
Archaeal 16S rRNA	A109F 1492r	ACKGCTCAGTAACACGT GYYACCTTGTTACGACTT	Grosskopf, Janssen and Liesack (1998) Nicol et al. (2008)	95°C 5 min; 10 cycles of 94°C 30 s, 55°C 30 s, 72°C 2 min; 25 cycles of 92°C 30 s, 55°C 30 s, 72°C 2 min; 72°C 5 min
Thaumarchaeal <i>amoA</i>	CrenamoA23f CrenamoA616r	ATGGTCTGGCTWAGACG GCCATCCATCTGTATGTCCA	Tourna et al. (2008) Tourna et al. (2008)	95°C 5 min; 10 cycles of 94°C 30 s, 55°C 30 s, 72°C 1 min; 25 cycles of 92°C 30 s, 55°C 30 s, 72°C 1 min; 72°C 5 min
Nitrosotalea sp. Nd2 <i>amoA</i> gene cluster	Nd2F1 Nd2R2	GATACTTGCAGTGATACCTACC CCAGATATTCTTGTTCAACAGAGG	This study This study	95°C 5 min; 35 cycles of 94°C 30 s, 52°C 30 s, 72°C 4 min; 72°C 10 min

The aim of this study was to characterise a novel AOA belonging to the *Nitrososphaera* sister cluster (Pester et al. 2011), isolated from a near-neutral pH agricultural soil, and to consider the implications for niche differentiation of AOA and AOB in soil.

MATERIALS AND METHODS

Culture enrichment and isolation

Sandy loam soil of pH 7.5 was sampled from an agricultural plot on the Scottish Rural College's Craibstone estate, Aberdeen (grid reference NJ872104). AO enrichment cultures were obtained by inoculation of 0.5 g soil in 50 ml 'fresh water medium' (FWM), containing NaCl (1 g l⁻¹), MgCl₂·6H₂O (0.4 g l⁻¹), CaCl₂·2H₂O (0.1 g l⁻¹), KH₂PO₄ (0.2 g l⁻¹), KCl (0.5 g l⁻¹), NaHCO₃ (0.168 g l⁻¹), NH₄Cl (0.027 g l⁻¹) and 1 ml l⁻¹ additions of the following solutions: modified non-chelated trace elements (Könneke et al. 2005), Fe-NaEDTA (7.5 mM) (Tourna et al. 2011) and 50 mg l⁻¹ streptomycin sulphate. Medium was adjusted to pH 7.5 at room temperature and filter-sterilised through a bottle-top filter (0.22 µm, Merck Millipore, Watford) into 100-ml Duran bottles (Sigma-Aldrich, Gillingham, UK). Soil suspensions were initially incubated statically at 37°C in the dark, monitored for nitrite accumulation and subcultured (2% inoculum) when nitrite concentration increased exponentially. Nitrite and ammonium concentrations were determined colorimetrically using Griess reagent and indophenol method, respectively, as previously described (Lehtovirta-Morley et al. 2014). Reactions and absorbance measurements were performed in clear polystyrene 96-well microplates (Greiner Bio-One, Stonehouse, UK) and measured using an Infinite F50 Microplate Reader and Magellan reader control and data analysis software (Tecan, Männedorf, Switzerland).

The presence of bacteria in enrichment cultures was determined by spread-inoculating 1 ml of culture onto Tryptone Soy, Nutrient and LB Agar plates (Sigma-Aldrich, Gillingham, UK) at

5%, 10% and 25% of standard concentrations, respectively, and 1.5% w/v agar. Agar plates were incubated in the dark at 30°C or 40°C. In addition, culture purity was assessed by PCR amplification of bacterial and fungal SSU rRNA genes. Cells were pelleted from liquid culture before extracting DNA using a standard bead-beating protocol with SDS extraction buffer and phenol:chloroform:isoamyl alcohol (Lehtovirta-Morley et al. 2011). PCR was performed using BIOTAQ polymerase and dNTP mix (Bioline, London, UK) according to manufacturer's instruction (see Table 2 for details of primers and PCR cycling conditions).

To eliminate bacteria from enrichment cultures, their antibiotic sensitivity was determined by spread-inoculating 0.5 ml of an exponentially growing enrichment on FWM agar plates containing antibiotic discs and incubating at 40°C in the dark. Antibiotic discs were prepared by permeating sterile paper with 50 mg l⁻¹ stock solutions (diluted in H₂O) of ampicillin, carbenicillin, gentamycin, kanamycin, clindamycin, penicillin or rifampicin and dried in the dark at ambient temperature (Lehtovirta-Morley et al. 2014). Antibiotics inhibiting bacterial contaminants were then added to FWM at a final concentration of 50 mg l⁻¹, in an attempt to purify the enrichment culture.

Culture maintenance and preservation

The purified archaeal strain, initially termed C13, was routinely cultured in FWM as described above but with the addition of 1 ml l⁻¹ of the following solutions: vitamin solution (excluding C₅H₁₄NO⁺) (Lehtovirta-Morley et al. 2011), Se-We solution (Widdel and Bak 1992) and 10 ml l⁻¹ of HEPES buffer (1 M HEPES, 0.6 M NaOH), with or without clindamycin (50 mg l⁻¹) and ampicillin (100 mg l⁻¹). All solutions were added prior to filter sterilisation.

C13 was cryopreserved and stored at -80°C using dimethylsulfoxide (DMSO) (VWR, Lutterworth, UK) as a cryoprotectant. DMSO was diluted to a concentration of 14% (v/v) using FWM

and filter-sterilised (0.22- μm filter) before the addition of 0.5 ml DMSO solution to 0.5 ml of an exponentially growing culture in a sterile 1.5 ml microcentrifuge tube. Tubes were agitated for 5 min at 25°C in the dark to allow the DMSO to penetrate the cells before storage at -80°C. To resuscitate cells, samples were thawed at 25°C and immediately centrifuged at 4°C at 1200 $\times g$ for 10 min to minimise the cytotoxic effects of DMSO. Supernatant was then removed, pelleted cells were resuspended in 1 ml sterile FWM and centrifuged and washed three times, to remove residual DMSO, before final resuspension in 10 ml FWM. Growth of cells preserved for 3 weeks was detectable after resuscitation for 4 days.

Physiological characterisation

Unless otherwise stated, all physiological characteristics were determined during batch cultivation, with static incubation of triplicate cultures in FWM in the dark at 40°C as described above for enrichment cultures. The effects of ammonium concentration were determined in FWM supplemented with 0, 1, 2, 5, 10, 20, 50 or 100 mM NH_4Cl and inhibition by nitrite concentration by supplementation with 0, 1, 2, 5, 10, 20 or 50 mM NaNO_2 . Ureolytic activity was investigated in ammonia-free FWM supplemented with 1 mM urea. Growth was also determined during incubation at temperatures ranging from 15°C to 60°C (5°C intervals). The effect of initial pH value in the range of 5–9 (0.5 pH intervals) was investigated in unbuffered medium containing a reduced concentration of 100 μM NH_4Cl , to minimise the decrease in pH decline due to ammonia oxidation, and in medium buffered with 5 mM MES (2-(*N*-morpholino)ethanesulfonic acid) or 10 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid). As MES and HEPES have pK_a values of 5.91 and 7.31, respectively, at 40°C, neither possessed the buffering capacity to fulfil the full range of pH tested in this study, and MES buffer was used at pH 5.5–7 and HEPES buffer at pH 7–8.5.

Cell counts and microscopy

Cells were enumerated microscopically in 1 ml samples of liquid culture, fixed with 5% formaldehyde (final concentration; w/v) and stored at 4°C. Each sample was repeatedly passed through a syringe and needle (0.4 mm diameter) to disperse aggregated cells. Cells were then diluted and stained with 2 μg DAPI (4, 6-diamidino-2-phenylindole), incubated for 10–15 min at 4°C in the dark, filtered onto a black 0.2- μm polycarbonate membrane and washed with PBS. Dried filters were placed on a glass slide mounted with an antifadent (Citifluor AF2, Citifluor Ltd, Leicester, UK) and counted under oil immersion using an Olympus BH-2 microscope with a U-MWU2 fluorescence light source (Olympus, Southend-on-Sea, UK) at $\times 1000$ magnification. For scanning electron microscopy, cells were fixed and gold-sputtered as previously described (Lehtovirta-Morley *et al.* 2014) and images taken using an Evo MA 10 scanning electron microscope (Carl Zeiss, London, UK).

Abundance of *amoA* genes

Growth of strain C13 was also determined by quantification of *amoA* genes. DNA was extracted as described above, except that 20 μg glycogen was added as co-precipitant during DNA precipitation. qPCR amplification was performed in a BioRad MyIQ Single-Color Real-Time PCR Detection System (Hertfordshire, UK) using amplification-conditions and cycling-parameters described in Thion and Prosser (2014), except that 0.6 μM of each

primer was used. A PCR product containing the *amo* gene cluster of *Nitrosotalea* sp. Nd2 (see Table 2 for details of primers and PCR cycling conditions) was used as a standard at dilutions giving 10^2 – 10^8 *amoA* genes. The amplification efficiency was 97% and the r^2 value 0.99. Positive amplification was confirmed by melting curve analysis and agarose gel electrophoresis.

DOPE-FISH analysis

Purity of strain C13 was also determined by DOPE-FISH (double labelling of oligonucleotide probes for fluorescence in situ hybridisation) using archaeal ARCH915 probe (Stahl and Amann 1991) labelled with Cy3 and bacterial EUB338 probe (Daims *et al.* 1999) labelled with fluorescein. One ml of stationary phase culture was washed twice with PBS, concentrated in 100 μl of PBS and fixed with 96% ethanol (1:1 v/v). Before applying the cells to a teflon-covered microscope slide, cells in the fixed culture were dispersed by passing through a syringe and needle as described above. After dehydrating the cells on a microscope slide, 1 μl of each doubly labelled FISH probe (50 ng μl^{-1}) was added to 10 μl of hybridisation buffer using 30% stringency (0.9 M NaCl, 20 mM Tris-HCl, 30% (v/v) formamide and 0.01% SDS) (Stoecker *et al.* 2010). Cells were incubated at 46°C for 4 h in the dark, after which the microscope slide was transferred to the pre-warmed washing buffer (112 mM NaCl, 20 mM Tris-HCl and 5 mM EDTA) and incubated for 15 min at 48°C. Cells were then resuspended in 100 μl of pre-warmed washing buffer and incubated at 48°C for 20 min. Finally, the microscope slide was dried and counter-stained with DAPI in an antifadent oil (Citifluor AF1, Citifluor Ltd, Leicester, UK) and samples were visualised using a Zeiss Imager M2 fluorescent microscope (Carl Zeiss, London, UK).

Phylogenetic analysis

PCR was performed with primer sets targeting the archaeal *amoA* gene (CrenamoA23f/CrenamoA616r) and 16S rRNA gene (A109f/1492r) (Table 2). PCR products were purified using a Macherey–Nagel NucleoSpin PCR Clean-up Kit (Fisher Scientific, Loughborough, UK) and sequenced, with assembled 16S rRNA and *amoA* gene sequences deposited in GenBank with accession numbers KU290365 and KU290366, respectively. Sequences were aligned with those from all previously cultivated AOA and a selection of environmental clones using the ClustalW function within the BioEdit software (Hall 1999) before removing regions of ambiguous alignment. Phylogenetic analyses were performed using maximum-likelihood with General Time Reversible-correction (PhyML; Guindon and Gascuel 2003), Bayesian (MrBayes (Ronquist *et al.* 2012), parsimony (MEGA5 (Tamura *et al.* 2011)) and Tamura's 3-parameter pairwise distance analysis (MEGA5) for 16S rRNA gene analysis, and Jones-Taylor-Thornton (JTT)-corrected maximum-likelihood (PhyML), Bayesian (MrBayes), parsimony (MEGA5) and JTT-corrected pairwise distance (MEGA5) analyses of inferred translated amino acid sequences for *amoA* genes. Where appropriate, analyses used estimated variable sites only with gamma-distributed site variation and bootstrap support was calculated 1000 times for ML, parsimony, distance analyses, with Bayesian analysis performed with 100 000 iterations, sampling frequencies of 10, with an average standard deviation <0.01.

Statistical analysis

Maximum specific growth rate, μ_{max} , was calculated as the gradient of semi-logarithmic plots of nitrite, cell or *amoA* gene




Figure 1. Typical DOPE-FISH images of stationary-phase C13 culture (a, b) using archaeal ARCH915 probe (red), bacterial EUB338 probe (green) and counterstained with DNA DAPI stain (blue). Scanning electron micrographs of individual cells (c) and aggregates (d) of archaeal ammonia oxidiser strain C13.

concentration vs time during exponential growth. The effect of treatments on μ_{max} was analysed by one-way ANOVA and linear regression analysis was performed on μ_{max} using Excel. Cell activity was determined for each replicate culture by minimising the sum of squares of differences between nitrite concentrations determined experimentally and those predicted by assumption of constant specific cell activity during exponential growth using the Solver routine in Excel, with μ_{max} and initial cell concentration or *amoA* gene concentration determined experimentally for each replicate culture.

RESULTS

Strain isolation

Enrichment cultures containing Thaumarchaeota, identified by amplification and sequencing of *amoA* genes, were obtained by inoculation of inorganic (FWM) medium containing 0.5 mM NH_4^+ with a sandy loam soil, pH 7.5. One culture containing a single *amoA* gene clonal population (designated C13) was selected for purification. Growth of heterotrophic bacteria was observed on Nutrient, Tryptone Soy and LB agar, but was eliminated after seven successive subcultures in FWM containing 500 mg l⁻¹ kanamycin. Bacterial contamination was, however, detected by PCR amplification of 16S rRNA genes and by growth on solid FWM.

Heterotrophic contamination was reduced further by supplementation of medium with one of five antibiotics (ampicillin, carbenicillin, gentamycin, clindamycin and rifampicin) to which bacterial contaminants were sensitive. Nitrite was produced in medium containing each antibiotic except gen-

tamycin, but was not detectable in subsequent subcultures, presumably because of a requirement for compounds produced by co-cultured bacteria. Growth was restored by the addition of spent (filter-sterilised) growth medium (1:10 ratio) and the addition of clindamycin and ampicillin. After a further eight subcultures with addition of spent medium and antibiotics, bacterial growth was not detected on FWM agar plates incubated for two weeks. Subsequent growth occurred in the absence of spent medium with no detectable amplification of bacterial 16S rRNA genes. The purity of strain C13 was also confirmed by DOPE-FISH which revealed the presence of archaeal cells, but never bacteria (Fig. 1a, b).

Physiological characteristics

The influence of ammonia concentration on growth was determined by cultivation of strain C13 in medium containing 0–100 mM NH_4^+ (Fig. 2a). μ_{max} was greatest (0.025 [s.e. 0.0002] h⁻¹) at 5 mM, and lower at 1 mM. μ_{max} decreased with increasing ammonium concentration, but growth was possible at the highest concentration tested, 100 mM, at 0.0035 [s.e. 0.0002] h⁻¹. Nitrite was tolerated at initial concentrations of 0–10 mM NO_2^- ($p = 0.13$), but growth was completely inhibited by 20 mM and 50 mM NO_2^- , with no detectable increase in nitrite concentration after incubation for 21 days (Fig. 2b).

Strain C13 is mesophilic and grew at 30°C–45°C (Fig. 2c), with no detectable increase in cell abundance or nitrite concentration outside this temperature range after incubation for 21 days. Growth was optimal at 40°C, with a μ_{max} of 0.027 [s.e. 0.00006] h⁻¹. The isolate is neutrophilic, growing in the pH range 6–8.5 (Fig. 2d), and no detectable increase in cell abundance or nitrite




Figure 2. The influence of (a) initial ammonium concentration, (b) initial nitrite concentration, (c) temperature and (d) pH on maximum specific growth rate of archaeal ammonia oxidiser isolate in batch culture. Growth in (d) was in unbuffered FWM or FWM buffered with MES or HEPES. Error bars are standard errors of means from triplicate cultures but are often smaller than the symbol size.

concentration occurred outside this pH range during incubation for 21 days. Growth in unbuffered FWM was optimal at pH 7, with a maximum specific growth rate of 0.025 [s.e. 0.0002] h⁻¹ (Fig. 2d), but pH in this medium varied during growth, increasing to 7.5 after incubation for 4 days, and subsequently decreasing to 7.3 when ammonium was completely utilised. Growth in FWM buffered by addition of MES or HEPES was optimal at pH 7.5 (Fig. 2d), but the μ_{max} was lower than in unbuffered medium, at 0.016 [s.e. 0.00006] h⁻¹, presumably due to inhibition by buffer.

C13 is ureolytic and grew on medium containing 1 mM urea as the sole source of ammonia (Fig. 3a) with a μ_{max} of 0.025 s.e. 0.0005 h⁻¹. Ammonia concentration increased from 65 μ M at day 0 (carried over in inoculum) to >2 mM during incubation for 6 days and was stoichiometrically converted to nitrite (Fig. 3b). μ_{max} estimated as specific rates of increase in nitrite, cell and *amoA* concentration differed slightly (Fig. 3c, Table 3), although differences between μ_{max} values are statistically significant (t-test, $p > 0.05$). *amoA* gene concentrations were approximately one order of magnitude greater than cell concentrations, and cell activities and cell yields calculated using both sets of data are presented in Table 3.

Morphology

Cells are irregularly shaped cocci with a diameter of 0.96 μ m (standard error = 0.021 μ m, range = 0.43–1.59 μ m, $n = 66$) (Fig. 1c). Cells frequently occurred as chains of four cells or in irregular clusters, often with substantial extracellular material that was also observed detached from cells. These irregularly shaped aggregates varied in diameter from 7 to 15 μ m (Fig. 1d).

Phylogenetic analyses

Phylogenetic analysis of 16S rRNA and *amoA* genes placed C13 within a well-supported, monophyletic lineage associated with organisms found in enrichment cultures and cloned sequences from various soil-based studies, with 16S rRNA and *amoA* genes sharing 99% and 92% identity, respectively, with those from the unpublished genome of ‘*Ca. Nitrosocosmicus oleophilus*’ and an enrichment culture obtained from arctic soil (Alves et al. 2013). It was however, distinct from all cultured *Nitrososphaera* strains, with closest 16S rRNA similarity to *N. viennensis* at 95% identity (Fig. 4). While analysis of both genes shows a clear phylogenetic association with *Nitrososphaera*-like organisms, differences were observed when associating strain C13 (and closely related organisms) within a separate, sister cluster to the *Nitrososphaera*. Analysis of derived *AmoA* protein sequences demonstrated that C13 was placed in the clade designated by Pester et al. (2011) as the *Nitrososphaera* sister cluster (sharing 91%–99% identity with the 14 sequences analysed in that study), and distinct from those designated as belonging to the *Nitrososphaera* cluster. We thus propose the new clade delineation ‘*Nitrosocosmicus*’ cluster for these sequences. However, analysis of 16S rRNA gene sequences did not provide support for a separate lineage affiliation, with C13 being placed within a lineage that is associated with other members of the *Nitrososphaerales* (Stieglmeier et al. 2014a).

DISCUSSION

The thaumarchaeotal strain C13 is a mesophilic, neutrophilic autotrophic archaeal AO that grows within the temperature




Figure 3. (a) Changes in ammonium and nitrite concentration during growth of strain C13 during batch growth on FWM with urea as nitrogen source; (b) changes in ammonium and nitrite concentrations and (c) semilogarithmic plots of nitrite (\square), cell (Δ) and *amoA* (\diamond) concentrations during growth on FWM with 2 mM ammonium chloride. Error bars are standard errors of means from triplicate cultures but are often smaller than the symbol size.

Table 3. Growth characteristics of strain C13 during batch growth at 40°C in HEPES-buffered FWM at pH 7.5 with initial ammonia concentration of 2 mM and determined on the basis of changes in concentrations of nitrite, cells and *amoA* genes. Means and standard errors are calculated from triplicate cultures.

Basis for calculations	μ_{max} (h^{-1})	Specific cell activity ($\text{fmol NO}_2^- \text{ cell}^{-1} \text{ h}^{-1}$)	Cell yield ($\text{cells } \mu\text{M}^{-1} \text{ NH}_3$)
Nitrite concentration	0.0237 (s.e. 0.0001)		
Cell concentration	0.0248 (s.e. 0.0011)	2.02 (s.e. 0.19)	7.60×10^3 (s.e. 0.19×10^3)
<i>amoA</i> concentration	0.0239 (s.e. 0.0004)	0.58 (s.e. 0.17)	1.29×10^5 (s.e. 0.097×10^5)


range 30°C–45°C and pH range 6–8.5. In these respects, it is similar to the soil AOA *N. viennensis* (28°C–45°C, pH 6–8.5) (Stieglmeier et al. 2014b) and *N. chungbukensis* MY2 (25°C–40°C, pH 6.5–8) (Jung et al. 2014). It also utilises urea as a source of ammonia, in common with several other members of the ‘Nitrososphaera’ lineage that possess genes encoding urease and urea transporters and *N. viennensis* and *N. gargensis* that can grow on urea (Tourna et al. 2011; Spang et al. 2012; Zhalina et al. 2014).

Cells of strain C13 are larger than those of *N. viennensis* (0.6–0.8 μm diam) (Stieglmeier et al. 2014a) and other soil AOA isolates, with an estimated cell volume of 0.7 μm^3 . Cells were associated with extracellular polymeric material that led to aggregate formation and may facilitate biofilm formation, which has been shown to protect AOB from deleterious effects of low pH and nitrification inhibitors (de Boer et al. 1991; Powell and Prosser 1991; Allison and Prosser 1993). μ_{max} values were within the range found for other neutrophilic soil AOA (Table 1). Cell

concentration was approximately one order of magnitude lower than *amoA* gene concentration, and similar differences have been reported in *N. maritimus* (Nakagawa and Stahl 2013). Cell concentration may be underestimated by incomplete dispersion of cells in aggregates, while *amoA* gene concentration may be overestimated by extracellular DNA following cell lysis. Cell yield and activity estimated using *amoA* gene concentrations are more consistent with those for other soil AOA and with *N. koreensis* MY1, respectively.

Strain C13 was more tolerant to ammonia than other soil AOA (Table 1), although the mechanisms for ammonia tolerance are unknown, and growth was possible at 100 mM NH_4^+ , at which μ_{max} was reduced to approximately 10% of that at optimal ammonium concentration. Strain C13 was also more tolerant to inhibition by nitrite than other neutrophilic soil AOA and grew at initial concentrations up to 20 mM NO_2^- . At pH 7.5, this is equivalent to 1.58 μM HNO_2 , which is believed to cause

(a) 16S rRNA



(b) AmoA




Figure 4. Maximum-likelihood phylogenetic analysis of (a) 16S rRNA genes and (b) derived AmoA protein sequences of isolated archaeon strain C13 with sequences from other cultivated organisms, metagenomes or cloned environmental sequences. Clade names are as described by Stieglmeier et al. (2014a) (16S rRNA) or Pester et al. (2011) (AmoA). Circles describing support represent the most conservative value from four methods used (bootstrap (ML, distance, parsimony) or posterior probabilities (Bayesian)) and the scale bar represents 0.05 changes per nucleotide or amino acid position. Accession number and environmental source are given in parentheses. If an accession number describes a genome or genomic fragment containing both a 16S rRNA and *amoA* gene, details are provided in the 16S rRNA tree only.

inhibition, and nitrous acid tolerance of strain C13 is therefore similar to that of *N. devanaterrea* and *Nitrosotalea* sp. Nd2, greater than that of other cultivated soil AOA and one order of magnitude greater and lower than *N. maritimus* and *Nitrosomonas europaea*, respectively (Table 1).

Comparisons of the 'model' AOA and AOB, *N. maritimus* and *N. europaea* ATCC 19718, have led to the view that AOA are smaller, have lower growth rates and specific cell activity and lower tolerance to ammonia and nitrite than AOB; these differences have further been suggested to drive niche differentiation between AOA and AOB. While these distinctions are evident for the model AOA and AOB (Table 1), and exist across the breadth of AOA and AOB, they are less marked when considering only soil isolates (Table 1). Soil AOA and AOB have a similar range for μ_{max} and are therefore likely to compete equally for ammonia when it is not growth-limiting. AOA are generally smaller and consequently have lower specific cell activity and greater cell yield than AOB, but this, in itself, provides no obvious competitive advantage. However, strain C13 extends the range of ammonia tolerance beyond that of characterised cultivated soil AOB (Prosser 1989; Koper et al. 2010), reducing support for selective inhibition of AOA at high ammonia concentration, and suggesting a potential role for AOA in nitrification in fertilised soils. Care is therefore required when using information from a limited number of isolates to explain ecological phenomena and the differences between physiological characteristics of *Nitrosotalea* isolates (Lehtovirta-Morley et al. 2014) highlight physiological diversity between strains that are very closely related in terms of 16S rRNA gene sequences. In addition, culti-

vation conditions will select strains that may not be representative of dominant and active members of natural communities and adaptation and selection will increase with continued cultivation. In this respect, C13 did not grow at 25°C and growth was optimal at 40°C, which is not typical of its environmental origin.

Phylogenetic analysis of predicted AmoA amino acid sequences places strain C13 in the *Nitrososphaera* sister cluster (Pester et al. 2011), a lineage that is closely related to, but distinct from, the *Nitrososphaera* cluster. However, the same relative branching order was not resolved in 16S rRNA gene phylogenies, with strain C13 belonging to clade placed within, and not basal to, the order *Nitrososphaerales*, a lineage that is generally referred to as 'Group 1.1b' in environmental surveys (e.g. Nicol et al. 2008). These contrasting topologies are consistent with a recent detailed comparison by Vico Oton et al. (2015) of thaumarchaeal 16S rRNA (clusters O and M) and *amoA* (C13.1 and C13.2) gene phylogenies. However, as strain C13 shares only 94%–95% 16S rRNA gene identity with other cultivated *Nitrososphaera* representatives, C13 probably represents a novel genus within the *Nitrososphaerales*.

This paper therefore reports the discovery of an AO that grows at neutral pH, but whose growth and activity are similar to those of soil AOB enrichments and isolates at neutral pH. Phylogenetic analysis of 16S rRNA and *amoA* genes places the organism within the *Nitrososphaerales* but distinct from previously isolated representatives of this order. We propose the following candidate status:

'*Nitrosocosmicus franklandus*' gen. et sp. nov.

Etymology. nitrosus (Latin masculine adjective): nitrous; cosmicus (Latin masculine adjective): cosmopolitan; Frankland (English noun): after Percy and Grace Faraday Frankland. The name reflects its ability to oxidise ammonia to nitrite, its widespread distribution in the environment, and provides a dedication to the authors of the first publication to describe isolation of organisms associated with ammonia oxidising activity (Frankland and Frankland 1890).

Source. Agricultural soil of pH 7.5, Craibstone, Aberdeen, Scotland, UK.

Description. A chemolithoautotrophic AO of the kingdom Thaumarchaeota, appearing as irregular cocci with a diameter 1.1 µm.

In accordance with other members of Nitrososphaerales, *N. franklandus* is an ammonia oxidising archaeon, isolated from a terrestrial environment, demonstrating the characteristic irregular coccoid cell morphology associated with this order (Stieglmeier et al. 2014a). As with *N. viennensis*, it is able to grow on urea but it can tolerate higher ammonium concentrations, equivalent to those tolerated by oligotrophic bacterial AOs (Tourna et al. 2011). *Nitrosocosmicus franklandus* represents an ecologically relevant strain of Thaumarchaeota with the potential to compete successfully with ammonium oxidising bacteria in fertilised soils where ammonium concentrations are high. The discovery of *N. franklandus* reduces support for niche differentiation between AOA and AOB in soil, with the possible exception in acid soils, and suggests that physiological diversity within each group may be more important in determining the composition of natural communities.

ACKNOWLEDGEMENTS

The authors would like to thank Mr Kevin Mackenzie and Mrs Gillian Milne (University of Aberdeen) for technical support with scanning electron microscopy, and Dr Robin Walker for access to the Woodlands Field experimental plots at the SRUC, Craibstone Estate, Aberdeen.

FUNDING

This work was financially supported by Natural Environmental Research Council (standard grants NE/I027835/1 and NE/L006286/1 and fellowship NE/J019151/1), EC Marie Curie ITN NORA, Grant Agreement No. 316472, the AXA Research Fund and the Centre for Genome Enabled Biology and Medicine, University of Aberdeen.

Conflict of interest. None declared.

REFERENCES

Allison SM, Prosser JI. Ammonia oxidation at low pH by attached populations of nitrifying bacteria. *Soil Biol Biochem* 1993;25:935–41.

Alves RJ, Wanek W, Zappe A et al. Nitrification rates in Arctic soils are associated with functionally distinct populations of ammonia-oxidising archaea. *ISME J* 2013;7:1620–31.

de Boer W, Klein Gunnewiek PJA, Veenhuis M et al. Nitrification at low pH by aggregated autotrophic bacteria. *Appl Environ Microb* 1991;57:3600–4.

Daims H, Bruhl A, Amann R et al. The domain-specific probe EUB338 is insufficient for the detection of all bacteria: devel-

opment and evaluation of a more comprehensive probe set. *Syst Appl Microbiol* 1999;22:434–4.

Elling FJ, Könneke M, Mussmann M et al. Influence of temperature, pH, and salinity on membrane lipid composition and TEX86 of marine planktonic thaumarchaeal isolates. *Geochim Cosmochim Acta* 2015;151:238–55.

Francis CA, Roberts KJ, Beman JM et al. Ubiquity and diversity of ammonia-oxidising archaea in water columns and sediments of the ocean. *P Natl Acad Sci USA* 2005;102:14683–8.

Frankland PF, Frankland GC. The nitrifying process and its specific ferment. Part I. *Philos T Roy Soc B* 1890;181:107–28.

Grosskopf R, Janssen PH, Liesack W. Diversity and structure of the methanogenic community in anoxic rice paddy soil microcosms as examined by cultivation and direct 16S rRNA gene sequence retrieval. *Appl Environ Microb* 1998;64:960–9.

Gubry-Rangin C, Hai B, Quince C et al. Niche specialisation of terrestrial archaeal ammonia oxidisers. *P Natl Acad Sci USA* 2011;108:21206–11.

Gubry-Rangin C, Kratsch C, Williams TA et al. Coupling of diversification and pH adaptation during the evolution of terrestrial Thaumarchaeota. *P Natl Acad Sci USA* 2015;112:9370–75.

Guindon S, Gascuel O. PHYML: a simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* 2003;52:696–704.

Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl Acid Symp Ser* 1999;41:95–8.

Hunik JH, Meijer HJG, Tramper J. Kinetics of *Nitrosomonas europaea* at extreme substrate, product and salt concentrations. *Appl Microbiol Biot* 1992;37:802–7.

Jung M-Y, Park S-J, Kim S-J et al. A mesophilic, autotrophic, ammonia-oxidising archaeon of thaumarchaeal group i.1a cultivated from a deep oligotrophic soil horizon. *Appl Environ Microb* 2014;80:3645–55.

Jung M-Y, Park S-J, Min D et al. Enrichment and characterisation of an autotrophic ammonia-oxidising archaeon of mesophilic crenarchaeal group I.1a from an agricultural soil. *Appl Environ Microb* 2011;77:8635–47.

Könneke M, Bernhard AE, de la Torre JR et al. Isolation of an autotrophic ammonia-oxidising marine archaeon. *Nature* 2005;437:543–46.

Koops H-P, Pommerening-Röser A. Distribution and ecophysiology of the nitrifying bacteria emphasizing cultured species. *FEMS Microbiol Ecol* 2001;72:1–9.

Koops H-P, Purkhold U, Pommerening-Röser A et al. The lithoautotrophic ammonia-oxidising bacteria. In: Dworkin M, Falkow S, Rosenberg E et al. (eds). *The Prokaryotes*, vol 6, 3rd ed. New York: Springer, 2006, 778–811.

Koper TE, Stark JM, Habteselassie MY et al. Nitrification exhibits Haldane kinetics in an agricultural soil treated with ammonium sulfate or dairy-waste compost. *FEMS Microbiol Ecol* 2010;74:316–22.

Lane DJ. 16/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M (eds). *Nucleic Acid Techniques in Bacterial Systematics*. New York: John Wiley and Sons, 1991, 115–75.

Lehtovirta-Morley LE, Ge C, Ross J et al. Characterisation of terrestrial acidophilic archaeal ammonia oxidisers and their inhibition and stimulation by organic compounds. *FEMS Microbiol Ecol* 2014;89:542–52.

Lehtovirta-Morley LE, Stoecker K, Vilcinskas A et al. Cultivation of an obligate acidophilic ammonia oxidiser from a nitrifying acid soil. *P Natl Acad Sci USA* 2011;108:15892–7.

- Leininger S, Urich T, Schloter M et al. Archaea predominate among ammonia-oxidising prokaryotes in soils. *Nature* 2006;**442**:806–9.
- Martens-Habbena W, Berube PM, Urakawa H et al. Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature* 2009;**461**:976–9.
- Nakagawa T, Stahl DA. Transcriptional response of the archaeal ammonia oxidiser *Nitrosopumilus maritimus* to low and environmentally relevant ammonia concentrations. *Appl Environ Microb* 2013;**79**:6911–6.
- Nicol GW, Leininger S, Schleper C et al. The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia-oxidising archaea and bacteria. *Environ Microbiol* 2008;**10**:2966–78.
- Palatinszky M, Herbold, Jehmlich N et al. Cyanate as an energy source for nitrifiers. *Nature* 2015;**524**:105–8.
- Pester M, Rattei T, Flechl S et al. *amoA*-based consensus phylogeny of ammonia-oxidising archaea and deep sequencing of *amoA* genes from soils of four different geographic regions. *Environ Microbiol* 2011;**14**:525–39.
- Powell SJ, Prosser JI. Protection of *Nitrosomonas europaea* colonising clay minerals from inhibition by nitrapyrin. *J Gen Microbiol* 1991;**137**:1923–29.
- Prosser JI. Autotrophic nitrification in bacteria. *Adv Microb Physiol* 1989;**30**:125–81.
- Prosser JI, Nicol GW. Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. *Trends Microbiol* 2012;**20**:523–31.
- Qin W, Amin SA, Martens-Habbena W et al. Marine ammonia-oxidising archaeal isolates display obligate mixotrophy and wide ecotypic variation. *P Natl Acad Sci USA* 2014;**111**:12504–9.
- Reigstad LJ, Richter A, Daims H et al. Nitrification in terrestrial hot springs of Iceland and Kamchatka. *FEMS Microbiol Ecol* 2008;**64**:167–74.
- Ronquist F, Teslenko M, van der Mark P et al. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* 2012;**61**:539–42.
- Sayavedra-Soto LA, Arp DJ. Ammonia-oxidising bacteria: their biochemistry and molecular biology. In: Ward BB, Arp DJ, Klotz MG (eds). *Nitrification*. Washington DC: ASM Press, 2011, 11–37.
- Smit E, Leeflang P, Glandorf B et al. Analysis of fungal diversity in the wheat rhizosphere by sequencing of cloned PCR-amplified genes encoding 18S rRNA and temperature gradient gel electrophoresis. *Appl Environ Microb* 1999;**65**:2614–21.
- Spang A, Poehlein A, Offre P et al. The genome of the ammonia oxidising candidate *Nitrososphaera gargensis*: insights into metabolic versatility and environmental adaptations. *Environ Microbiol* 2012;**14**:3122–45.
- Stahl DA, Amann R. Development and application of nucleic acid probes. In: Stackebrandt E, Goodfellow M (eds). *Nucleic Acid Techniques in Bacterial Systematics*. Chichester, UK: John Wiley & Sons Ltd, 1991, 205–48.
- Stieglmeier M, Klingl A, Alves RJE et al. *Nitrososphaera viennensis* sp. nov., an aerobic and mesophilic ammonia-oxidising archaeon from soil and member of the archaeal phylum Thaumarchaeota. *Int J Syst Evol Microbiol* 2014a;**64**:2738–52.
- Stieglmeier M, Mooshammer M, Kitzler B et al. Aerobic nitrous oxide production through N-nitrosating hybrid formation in ammonia-oxidising archaea. *ISME J* 2014b;**8**:1135–46.
- Stoecker K, Doringner C, Daims H et al. Double labeling of oligonucleotide probes for fluorescence in situ hybridisation (DOPE-FISH) improves signal intensity and increases rRNA accessibility. *Appl Environ Microb* 2010;**76**:922–6.
- Tamura K, Peterson D, Peterson N et al. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011;**28**:2731–9.
- Thion C, Prosser JI. Differential response of non-adapted ammonia-oxidising archaea and bacteria to drying–rewetting stress. *FEMS Microbiol Ecol* 2014;**90**:380–9.
- Tourna M, Freitag TE, Nicol GW et al. Growth, activity and temperature responses of ammonia oxidising archaea and bacteria in soil microcosms. *Environ Microbiol* 2008;**10**:1357–64.
- Tourna M, Stieglmeier M, Spang A et al. *Nitrososphaera viennensis*, an ammonia oxidising archaeon from soil. *P Natl Acad Sci USA* 2011;**108**:8420–5.
- Vainio EJ, Hantula J. Direct analysis of wood-inhabiting fungi using denaturing gradient gel electrophoresis of amplified ribosomal DNA. *Mycol Res* 2000;**104**:927–36.
- Vico Oton E, Quince C, Nicol GW et al. Phylogenetic congruence and ecological coherence in terrestrial Thaumarchaeota. *ISME J* 2015;**10**:85–96.
- Widdel F, Bak F. Gram-negative mesophilic sulfate-reducing bacteria. In: Ballows A, Trüper HG, Dworkin M, Harder W, Schleifer K-H (eds). *The Prokaryotes. A Handbook on the Biology of Bacteria: Ecophysiology, Isolation, Identification, Application* 2nd edn. New York: Springer, 1992, 3352–78.
- Zhalnina K, de Quadros PD, FAO Camargo et al. Drivers of archaeal ammonia-oxidising communities in soil. *Front Microbiol* 2014;**3**:210.