

# Opposing gamete plastic responses to thermal environments in *Tribolium castaneum*

**NERC** SCIENCE OF THE ENVIRONMENT



Ram Vasudeva (r.vasudeva@uea.ac.uk)  
School of Biological Sciences, University of East Anglia, Norwich, NR4 7TJ







## INTRODUCTION

- Female and Male gametes represent good examples of common sexual traits under divergent selection and optima<sup>1,2,3,4</sup>.
- Using a poikilothermic insect model, I examine how gamete sizes develop following controlled variation in the thermal environment.
- Resulting plasticity could inform on 1) adaptive significance of gamete size, and 2) degree of intralocus conflict within gamete optima.
- Using 30°C and 38°C lab selection lines following >45 generations of selection, I measure genetic and phenotypic plasticity in sperm and egg sizes when exposed to 30°C and 38°C thermal environments.

## METHODS



## RESULTS



## CONCLUSIONS

- Clear differences in egg and sperm sizes between 30°C and 38°C thermal regimes, however the experimental switches show that this is driven by short-term developmental plasticity<sup>5</sup>.
- Gamete responses go in opposite directions: warmer conditions = bigger eggs but smaller sperm.
- The opposing responses suggest opposing optima<sup>6,7</sup>.
- Ongoing work is testing the adaptive significance of variation in both traits for their thermal regime.

## REFERENCES

- 1) Gavrillets (2000) *Nature*.| 2) Morrow & Gage (2001) *J. Zool.*| 3) Chapman (2006) *Curr. Biol.*| 4) Bonduriansky & Chenoweth (2008) *Trend.Eco.Evol.*| 5) Moczek (2015) *Heredity.*| 6) Lupold *et al.* (2016) *Nature.*| 7) Fox *et al.* (1997) *Am. Nat.*

## ACKNOWLEDGEMENTS

Members of Team Tribolium UEA : Matt Gage, Kris Sales, Jo Godwin, Alyson Lumley, Conrad Gillett and Laura Hebberecht Lopez.