


SCIENTIFIC REPORTS

OPEN

Long-term increases in soil carbon due to ecosystem fertilization by atmospheric nitrogen deposition demonstrated by regional-scale modelling and observations

E. Tipping ¹, J. A. C. Davies², P. A. Henrys¹, G. J. D. Kirk ³, A. Lilly ⁴, U. Dragosits⁵, E. J. Carnell ⁵, A. J. Dore⁵, M. A. Sutton⁵ & S. J. Tomlinson⁵

Fertilization of nitrogen (N)-limited ecosystems by anthropogenic atmospheric nitrogen deposition (N_{dep}) may promote CO_2 removal from the atmosphere, thereby buffering human effects on global radiative forcing. We used the biogeochemical ecosystem model N14CP, which considers interactions among C (carbon), N and P (phosphorus), driven by a new reconstruction of historical N_{dep} to assess the responses of soil organic carbon (SOC) stocks in British semi-natural landscapes to anthropogenic change. We calculate that increased net primary production due to N_{dep} has enhanced detrital inputs of C to soils, causing an average increase of 1.2 kgCm^{-2} (c. 10%) in soil SOC over the period 1750–2010. The simulation results are consistent with observed changes in topsoil SOC concentration in the late 20th Century, derived from sample-resample measurements at nearly 2000 field sites. More than half (57%) of the additional topsoil SOC is predicted to have a short turnover time (c. 20 years), and will therefore be sensitive to future changes in N_{dep} . The results are the first to validate model predictions of N_{dep} effects against observations of SOC at a regional field scale. They demonstrate the importance of long-term macronutrient interactions and the transitory nature of soil responses in the terrestrial C cycle.

Soil organic matter (SOM) is a key ecosystem component, both as a store of carbon that can exchange with the atmosphere thereby affecting climate¹, and because of its key soil functional roles in water and heat retention, nutrient cycling, and sorption of contaminants². Its source is predominantly plant material (litterfall, decaying roots, root exudates), the supply of which might increase where net primary production (NPP) is enhanced by the fertilizing effect of atmospheric N deposition^{3–5}. The latter originates from a wide range of emission sources, and comprises oxidised N (NO_x) and reduced N (NH_x). Whereas NO_x is mainly due to combustion processes such as those occurring in power generation, industry and motorised transport (including shipping), most ammonia (NH_3) emissions originate from agricultural practices including the management of livestock manures and slurries, and mineral fertiliser application⁶.

Whereas there is convincing evidence from both experimental manipulations and field observations that forest growth responds positively to N_{dep} ^{7–9}, evidence for a consequent SOC response comes mainly from short-term experimental studies, almost all at high N inputs compared with ambient values⁹, and from modelled estimates^{10,11}. One field study is that of Kirby *et al.*¹² who reported that for British woodlands, increases in the SOM of mineral and organo-mineral soils between 1971 and 2001 were positively correlated with changes in modelled N_{dep} . Data for other ecosystem types are sparse¹⁰. To explore possible long-term and widespread SOC response to N_{dep} in both forest and non-forest ecosystems, we conducted a combined modelling and data analysis of the soils of British semi-natural ecosystems comprising broadleaf woodland, unimproved grasslands, and shrublands, which account for a total area of $6.8 \times 10^4 \text{ km}^2$, i.e. 32.5% of Great Britain (Fig. 1). These ecosystems are well-suited to

¹Centre for Ecology and Hydrology, Lancaster, LA1 4AP, UK. ²Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK. ³Cranfield University, Bedford, MK43 0AL, UK. ⁴James Hutton Institute, Aberdeen, AB15 8QH, UK. ⁵Centre for Ecology and Hydrology, Edinburgh, EH26 0QB, UK. Correspondence and requests for materials should be addressed to E.T. (email: et@ceh.ac.uk)




Figure 1. Maps showing the occurrence of the three semi-natural habitat types (grey scale) and sampling locations (circles). The fractional land cover refers to individual $5\text{ km} \times 5\text{ km}$ grid cells used in simulation modelling. The maps were derived and plotted by aggregating land cover mapped at $25\text{ m} \times 25\text{ m}$ resolution in the UK Land Cover Map 2007³³, using ArcGIS 10.4 (<http://desktop.arcgis.com/en/arcmap/>) (accessed 09/11/2016).

a large-scale evaluation in view of the wide range of N_{dep} (1st to 99th percentile $0.2\text{--}3.6\text{ gNm}^{-2}\text{ a}^{-1}$ in 2010) and the wealth of data available from repeated soil sampling. The modelling was done by combining the ecosystem model N14CP¹³ with a new high-resolution spatio-temporal N_{dep} dataset covering the period since 1750, produced through atmospheric emission, transport and deposition modelling.

Results

We calculated how SOC pools evolved in Great Britain over the Holocene using the ecosystem model N14CP which links the plant-soil C, N and phosphorus (P) cycles, and has been parameterised and tested with plot-scale data for sites across N Europe¹³ (see Methods). The model was driven by national climate, vegetation and soil datasets, modelled atmospheric deposition (see Methods) and estimated vegetation history. It was implemented for $5\text{ km} \times 5\text{ km}$ grid cells. A key modelling assumption is that NPP depends on a single limiting factor, temperature, rainfall, N supply or P supply; for these British sites, all locations were found to have been continuously N limited. In blind comparisons with data, the simulations produced reasonable estimates of SOC pools (see Methods).

The changes in anthropogenic N_{dep} (Fig. 2, top row) caused increases in NPP (Fig. 2, middle row), by an average of $110\text{ gCm}^{-2}\text{ a}^{-1}$ (50%), and therefore a greater supply of organic C to the soil and enhanced SOC pools (Fig. 2, bottom row). The additional amounts of SOC from 1750 to 2010 were 21, 38 and 20 Mt for broadleaf woodland, unimproved grassland and shrubland respectively, equivalent to an average increment of 1.2 kgCm^{-2} or an average annual accumulation of $4.6\text{ gCm}^{-2}\text{ a}^{-1}$. From an analysis of model performance (see Methods), it can be shown that uncertainty in these SOC increases arises primarily from uncertainty in N_{dep} , and it is reasonable to assume an error of c. 30% in the estimated values. The extra SOC was spread through four model pools, fast (mean residence time 1 yr), slow (c. 20 yr), passive (c. 1000 yr) and subsoil (c. 2000 yr), with more than half (57%) accumulating in the slow pool (Fig. 3). The average calculated ratio of SOC gained to N_{dep} for all the soils considered over the entire period was 8.2 gCgN^{-1} , at the lower end of the range suggested by de Vries *et al.*¹⁰. Enrichment of the ecosystems by N_{dep} was calculated to affect total denitrification (i.e. gaseous loss of both nitrous oxide, N_2O , and dinitrogen, N_2), such that over the period 1750–2010, instead of 32 gNm^{-2} being lost to the atmosphere, the total was 40 gNm^{-2} , a net increase of 8 gNm^{-2} .

The model predictions of increased SOC were tested using data for 1911 field sites (Fig. 1) where topsoils had been sampled and then resampled, on average 27 years later, during the period 1959 to 2010 (Tables 1 and 2; see Methods). The measured SOC variable was concentration, expressed as % by weight. At each site, the model-simulated change in SOC pool (gCm^{-2}) was converted to a simulated change in SOC concentration (%C) by assuming that the average simulated SOC concentration was the same as the average of the measured values. This took into account both concentration change and soil thickening that could arise from increased organic matter input (see Methods). To evaluate the model's prediction of SOC change, we compared values of R_{SF} , the ratio of topsoil SOC concentration at the second sampling to that at the first (see Methods), transforming the data logarithmically to reduce the skew in the data and hence the influence of extreme values. We combined data for soils under grass and shrubs into a single data set, and analysed them separately from those for the woodland soils.




Figure 2. Simulated changes in atmospheric nitrogen deposition (N_{dep}), net primary productivity (NPP) and soil organic carbon (SOC) over the period 1700–2010 for semi-natural land areas of Great Britain. The central line is the mean, ranges are 5 and 95%-iles. Dashed lines show mean results with N_{dep} set to zero.




Figure 3. Simulated soil organic carbon (SOC) pools in 1800 and 2000; average values weighted by area of broadleaf woodland, grassland, and shrubland in Great Britain. Key: fast (F, mean residence time 1 yr), slow (S, c. 20 yr), passive (P, c. 1000 yr) and subsoil (Sub, c. 2000 yr). The values above the pairs of bars show the fractional contributions to the total increase in SOC.

Over the full range of SOC concentration (0.6–53%), simulated and observed average R_{SF} values for woodland soils are similar, and the observed value is significantly ($P < 0.01$) greater than 1.0 (Table 3a), indicating significant increases in SOC concentrations over the observation period. The confidence intervals for the observations are inevitably wider than those for the modelled values, owing to the large scatter in the observed data, most of which results from local spatial soil heterogeneity, which is not captured by the model. For all soils under unimproved grassland and shrubland, an average value of R_{SF} greater than 1.0 was predicted by the N14CP model, whereas the observations give a value slightly, although not significantly, less than 1.0. A clearer result is obtained by restricting the analysis to soils with relatively low SOC concentrations, since in such soils changes in the C pool are mainly seen as changes in SOC concentration, rather than as soil thickening. When this is done (Table 3b), simulated and observed average R_{SF} values remain similar for woodland soils, but now there is better agreement for soils under unimproved grassland and shrubland, and the observed means for both vegetation groups are significantly greater than 1.0.

Data source	Locations ^a	Year of 1 st sample	Year of 2 nd sample	Mean time period, yr	Number of sites			
					Broadleaf woodland	Unimproved grassland	Shrubland	Total
National Soil Inventory	E, W	1978–1983	2003	22	95	199	54	348
Countryside Survey 1	E, S, W	1978	1998	20	11	137	56	204
Countryside Survey 2	E, S, W	1978	2007	29	13	146	63	222
British Woodland Survey	E, S, W	1971	2001	30	1106	—	—	1106
National Soil Inventory of Scotland & Rare Soils	S	1959–1988	2007–2010	27	—	13	18	31
Total					1225	495	191	1911

Table 1. Summary of topsoil resampling sites. ^aE- England, S- Scotland, W- Wales.

	Sampling	Broadleaf woodland	Unimproved grassland	Shrubland
National Soil Inventory	1	2.2–17.4	2.9–47.2	4.5–54.1
	2	2.1–10.5	2.9–42.2	4.4–47.0
Countryside Survey 1	1	2.2–11.8	2.8–43.5	9.5–51.6
	2	3.5–19.7	3.6–48.9	15.1–53.0
Countryside Survey 2	1	2.2–10.1	2.8–43.5	5.2–51.7
	2	3.1–20.2	2.9–38.3	10.5–52.3
British Woodland Survey	1	3.4–24.4	—	—
	2	3.5–26.5	—	—
National Soil Inventory of Scotland & Rare Soils	1	—	4.5–48.9	12.6–49.9
	2	—	6.2–47.5	13.7–48.1

Table 2. Topsoil SOC data, 5–95 percentiles of %C.

	Calculated	Observed
<i>(a) all soils</i>		
broadleaf	1.052 ± 0.003	1.053 ± 0.035**
grass & shrub	1.022 ± 0.002	0.989 ± 0.044 ^{NS}
<i>(b) soils with [SOC] < 10%</i>		
broadleaf	1.054 ± 0.004	1.052 ± 0.037***
grass & shrub	1.032 ± 0.004	1.052 ± 0.046*

Table 3. Calculated and observed values of R_{SP} , the ratio of the soil organic carbon (SOC) concentration at the second sampling to that at the first, with 95% confidence intervals intervals, for topsoils of Great Britain, based on sampling between 1959 and 2010 (Table 1). For the observed ratios, significant difference from 1.0 is indicated: NS not significant, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

A second test of the model is whether the data show significant spatial differences in relation to N_{dep} . Again, this is best considered only for the soils with SOC < 10%, to focus on conditions where variations in SOC pool are reflected by variations in SOC concentration. Regression slopes of simulated and observed $\log_{10}R_{SF}$ against N_{dep} (Table 4) are in agreement, although the observed slope for soils under improved grassland and shrubland is not significant.

Discussion

The present study used a large dataset for British semi-natural soils, which we generated by combining several smaller datasets (Table 1). Previous evaluations of SOC change using the individual datasets gave less clear-cut results, including decrease or increase in SOC depending upon the range of SOC concentration considered^{14, 15} and little or no change^{12, 16, 17}. Our results are more definite because of (a) the larger sample size, (b) the omission of coniferous woodland which is mostly newly-planted and managed, (c) the use of R_{SF} as the diagnostic variable, and (d) our focus on soils with relatively low SOC concentrations.

Overall, the observations provide strong quantitative support for the model prediction of a general increase in the SOC contents of these soils, and the role of N_{dep} . The signal is stronger for woodland soils than those under

	<i>n</i>	Calculated	Observed
broadleaf	864	+0.0091 ± 0.0032	+0.0112 ± 0.0060***
grass & shrub	338	+0.0055 ± 0.0023	+0.0056 ± 0.0104 ^{NS}

Table 4. Calculated and observed slopes of spatial regressions of $\log_{10} R_{SF}$ (ratio of SOC concentration in second to first soil sampling) against N_{dep} ($gNm^{-2} a^{-1}$) in 2000 at 5 km grid resolution across Great Britain. The number of sites is indicated by *n*, error values are 95% confidence intervals. Whether the observed slope is significantly different from zero is shown by NS (not significant) and *** ($P \leq 0.001$).

grass and shrubs, because the scavenging effects of trees enhance N_{dep} inputs¹⁸ so that trees receive proportionally greater extra N. Moreover, the fewer data for soils under unimproved grassland and shrubland make it harder to achieve statistical significance.

The underlying cause of the net accumulation of SOC simulated by the N14CP model is the enrichment of the ecosystem by atmospherically-deposited N, which enhances the flux of plant available N, so that, because N is the limiting factor for plant growth, there is an increase in NPP. In turn, litter production and the supply of carbon to the soil rise, enlarging all three topsoil SOC pools and also subsoil SOC. In the ecosystem transition brought about by N deposition, the SOC pools grow in inverse proportion to their turnover rates, but because the slow pool is relatively large (Fig. 3) it shows the largest absolute SOC increase (Fig. 3). The soil organic N pools grow, more-or-less in parallel with the SOC, so that the simulated soil C:N ratio remains about the same throughout the enrichment¹³. The model omits other possible large-scale determinants of SOC change such as the effects of increased N on litter decomposition rates^{9, 19, 20}, fertilization by increasing CO_2 ²¹, changes in forest management^{12, 22}, and reductions in NPP due to ozone toxicity²³, which may mean that at least in the long term these effects are quantitatively less important than fertilization by N_{dep} , and/or that their effects largely cancel each other. Although the parameterisation includes the effects of pH on SOM turnover, the parameter values are such that there is little pH dependence, and consequently little effect on SOC of recent widespread increasing pH in British soils^{12, 16, 24} is simulated.

Possible climatic effects are included in N14CP via the effect of temperature on maximum NPP, but more importantly by the assumed positive temperature dependences of metabolic reactions, notably soil organic matter turnover rates. Temperature effects alone can be gauged by the simulations for zero N_{dep} in Fig. 2 (dashed lines in the middle and bottom panels). Minor increases in NPP are predicted owing to greater N availability resulting from faster organic matter turnover, but the resulting increases in detrital additions to soil organic matter are more than offset by increased turnover rates, leading to slight declines in SOC. However these effects are small in comparison with the large decreases in SOC (mean $\sim 70 gCm^{-2} a^{-1}$) reported over the period 1987/88 to 2011 for 24 forest sites in the German Alps²⁵, and attributed to faster SOC turnover due to increased summer temperature (mean $0.49 K decade^{-1}$) and changes in snow cover and freeze-thaw cycles associated with climate change. A loss rate of such magnitude due to temperature change cannot be simulated with N14CP; our calculated rate of SOC accumulation (mainly due to N_{dep}) of $13 g m^{-2} a^{-1}$ for broadleaf woodland over the period 1971–2007 is more compatible with the results of a study²⁶ covering the whole of Germany (1800 resampled sites) for the period 1987/1992 to 2006/2008, in which an increased forest SOC of $41 g m^{-2} a^{-1}$ was found.

As well as its effects on SOC, fertilization by N_{dep} can also promote the storage of organic carbon in woody biomass^{9–11}. However for Great Britain such storage is limited firstly by the relatively small area of broadleaf woodland, c. 20% of the total area of the semi-natural ecosystems considered here, and secondly because of the removal and burning of wood. Carbon storage in grass and shrub biomass is inconsequential. Therefore the effects of N_{dep} on carbon storage in Great Britain are mainly through SOC. The estimated increased SOC storage calculated here totals 78 Mt, which is equivalent to about 0.5% of the total anthropogenic emissions of CO_2-C from Great Britain over the same period (1750–2010). The high population density of Great Britain means that its anthropogenic CO_2-C emissions per unit area are relatively large, and its small area of woodland limits biomass storage, and therefore at bigger scales, European or global, the proportional effect of N_{dep} on C storage is likely to have been appreciably greater.

The accumulated SOC cannot be regarded as permanently sequestered. To maintain present levels, or to increase them, continual inputs of litter at the same rate are needed, which means that the elevated NPP must be maintained. Future declines in N_{dep} would eventually lead to decreased NPP, the litter input would be reduced, and CO_2-C losses by decomposition would become greater than litter C inputs, with a net return of CO_2-C to the atmosphere. As noted above, much (57%) of the C calculated to have been gained by the soils studied here over the period 1750 to 2010 is mainly in the 20-year pool (Fig. 3), which means that the soils respond fairly quickly to changes in N_{dep} . This is shown by the fairly rapid accumulation of SOC since 1750, and by the recent lower rate of increase in modelled SOC, reflecting reductions in N_{dep} since 1990 (Fig. 2). In terms of greenhouse gas balance, the release of N_2O from the soils as part of the increased denitrification that accompanies N enrichment due to N_{dep} will have partially offset the CO_2-C sequestration, but this too depends upon changes in N_{dep} . These dynamic relationships, and the transitory nature of soil responses, need to be factored into predictions of long term changes in the regional and global cycles of C and N.

Methods

Modelling long-term spatially-distributed N_{dep} . Atmospheric nitrogen deposition (N_{dep}) was modelled for a series of time steps between 1800 and the present (2010), at a $5 km \times 5 km$ grid resolution. The N deposition maps were derived by modelling the magnitude and spatial distribution of emissions of oxides of nitrogen (NO_x)

and ammonia (NH_3) for the historical time period, and running an atmospheric chemistry and transport model (FRAME; Fine Resolution Atmospheric Multi-pollutant Exchange) to estimate total N deposition for the years 1800, 1900, 1950, 1970, 1990 and 2010. Sulphur dioxide (SO_2) emissions, chemistry and deposition were also modelled for the same years, as they impact on the chemical transformations in the atmosphere and therefore the lifetime and transport distance of nitrogen-containing gases and particles before deposition.

Emissions of NO_x , NH_3 and SO_2 originate from a wide range of sources. Oxidised N and SO_2 emissions are mainly due to combustion processes such as those occurring in the power generation and industry sectors, as well as from domestic combustion and motorised transport (including shipping etc.). Ammonia, on the other hand, mainly originates from agricultural sources such as livestock manures and slurries as well as mineral fertiliser application (83% of total 2014 UK NH_3 emissions⁶), with waste processing, domestic pets and horses and natural sources etc. also contributing. Combustion sources are also estimated to contribute to NH_3 emissions significantly on a global scale^{27,28}, though for historical emissions for the UK the main uncertainty concerns fuel source emissions from domestic coal burning^{29,30}. Spatial datasets are normally produced by combining activity data (i.e. emission source populations or process locations and activities) with emission factors (i.e. average source strength of emissions per unit of activity data).

The emission maps were derived through spatial modelling, using a combination of spatial and non-spatial data, and a number of different approaches for different emission source sectors. For agricultural NH_3 emissions, the AENEID model^{31,32} was used, which combines agricultural livestock and crop statistics with land cover and agricultural practice data to derive high spatial resolution emission maps^{33–38}. The model was adapted for different historic periods, to account for changes over time in the zonal aggregation and livestock and crop categories recorded in the available agricultural statistics parish, county, rural district level datasets, which also differed between the four countries of the UK, i.e. England, Wales, Scotland, Northern Ireland^{39,40}. In terms of emission factors, the latest UK estimates for 2010 and back-cast to 1990 from the national inventory (NAEI) were used, with data for earlier years adapted to historic conditions, taking into account fertiliser N input to crops and grassland^{41,42}, N content in livestock excreta^{43–47}, milk yield of dairy cows (highly correlated with N excretion rates⁴⁷), changing livestock weights (e.g. increases in size due to animal breeding and feeding advances etc. ref. 48). For activity statistics and emission factors for combustion sources, available modern national inventory estimates and back-cast timelines from the NAEI were used for the more recent period (2010, 1990, 1970), with historic time series of power station fuel use, road transport statistics, human population census data, etc. used to extend the time series back to 1800^{49–58}.

Substantial NH_3 emissions from domestic coal combustion had previously been estimated for the UK²⁹ at 40, 110, 80 and 5 kt of $\text{NH}_3\text{-N}$ for 1850, 1900, 1950 and 1990, respectively, while recognizing a shortage of NH_3 emission measurements from this source. For 1900, this source amounted to around 70% of total estimated UK NH_3 emissions. To address this uncertainty, a specific campaign of measurements of N emissions from domestic coal burning was conducted (to be reported elsewhere), which was not able to support these large estimates. Measured emission rates of NO_x in this campaign were consistent with the inventory estimates, and the latter were used, because they are based on a larger dataset. For NH_3 , a much lower emission rate of 4.5 g $\text{NH}_3\text{-N}$ per tonne of coal burnt was used, based on the new measurements, which is consistent with a total UK $\text{NH}_3\text{-N}$ emission from coal of 0.15 kt $\text{NH}_3\text{-N}$ for 1900. While much smaller estimates of NH_3 emission from this source were supported by these measurements as compared with ref. 29, it is recognized that burning sources of NH_3 emission remain a major source of uncertainty^{30,59}.

Long-term land-cover changes were incorporated based on the current UK Land Cover Map 2007³³ and the mapping by Dudley Stamp and colleagues in the 1930s⁶⁰, with changes such as large areas of afforestation (e.g. Kielder Forest, northern England) influencing emission distributions and N deposition regionally over the historic period. For all datasets, careful checking and gap filling was necessary, taking account of historic changes in boundary maps (e.g. county boundaries), and a key consideration was the development of a consistent multi-year gridded dataset that was suitable for analysis, interpretation and comparisons across the time period.

The emission maps, together with land cover and meteorological data, were used as input to the UK FRAME atmospheric chemistry transport model to generate estimates of nitrogen and sulphur deposition. FRAME is a Lagrangian model run at a 1 km⁶¹ or 5 km⁶² grid resolution over the British Isles which simulates the emissions, transport, chemical transformation of gases and particles as well as dry and wet deposition of nitrogen and sulphur compounds. The boundary conditions for air concentrations used in the regional simulation were generated from a European scale simulation run with a 50 km grid resolution using a time series of European emission maps for 1900–2010 (ref. 63 and Simpson D., Chalmers University of Technology, Gothenburg, Sweden, pers. comm.). The performance of the FRAME model has been evaluated by comparison with measurements of gas and aerosol concentrations as well as concentrations in precipitation from national monitoring networks^{61,64}. The model was able to satisfy the criteria of fitness for purposes required for use in policy applications and has been used for future emissions scenario modelling and the assessment of impacts on natural ecosystems^{65,66}. For this study, long-term average meteorological conditions were used rather than specific annual data, to keep the long-term trends comparable, rather than having to take specific annual conditions into account when interpreting a 200+ year time series, with the model run at a 5 km grid resolution. The deposition datasets were compared against measurement based data⁶⁷ (wet deposition, gaseous and aerosol concentrations) for the recent period where national monitoring networks exist (1990 and 2010), with a good fit. However, for the historic period (1800 to 1950), suitable measurements are very scarce²⁹.

The gridded population estimates (1801–1951) created for use here are based on data provided through www.VisionofBritain.org.uk, using historical material which has been re-districted by the Linking Censuses through Time system, created as part of ESRC AwardH507255151, by Danny Dorling, David Martin and Richard Mitchell. The gridded population estimates (1971–2011) are based on data provided with the support of the ESRC and JISC, and use boundary material which is copyright of the Crown and the EDLINE Consortium data provided

with the support of the ESRC and JISC. The agricultural livestock and crop maps for 1970–2010 were created with data from the EDINA AgCensus database <http://edina.ac.uk/agcensus/> at Edinburgh University Data Library and DEFRA for England, The Welsh Assembly Government for Wales, and The Scottish Government (formerly SEERAD) for Scotland.

Ecosystem modelling with N14CP. The N14CP ecosystem model¹³ simulates the inter-linked cycles of C, N and P in plant–soil systems, starting at the onset of the Holocene, on a quarter-year time step. Four types of vegetation are considered, namely broadleaved woodland, coniferous woodland, herbs (principally rough, neutral, calcareous and acid grasslands) and shrubs (heather, heather grassland and montane habitats), each of which has two end-member stoichiometric compositions.

NPP is calculated on the basis of a limiting factor, potentially temperature, rainfall, available N or available P. For each location and year, total rainfall and mean temperature values are first used to estimate NPP without nutrient limitation. Plant N and P demands to achieve this NPP are then compared with calculated available N and P from SOM turnover and stored plant pools. If the available nutrients cannot meet the NPP demand, then whichever has the lower relative availability is considered to be the limiting nutrient.

SOC is generated by the entry into the soil of plant detritus comprising above-ground leaves and twigs, dead roots and root exudates. Larger woody components (boles and branches) are assumed to decompose on the soil surface, releasing CO₂ to the atmosphere but N and P to the soil. The SOM is partitioned among pools with turnover rates of c. 1 year (fast), c. 20 years (slow) and c. 1000 years (passive); the exact turnover rates depend upon temperature. The model is driven by climate, atmospheric deposition and soil type, and simple estimates are made about vegetation history. Key assumptions about the behaviour of N are (a) that N fixation depends upon the availability of P, and (b) that there is down-regulation of N fixation by anthropogenic N_{dep}, which means that during the last 200–300 years N_{dep} has become the dominant source of nutrient N to natural and semi-natural ecosystems in many parts of northern Europe, including Great Britain. The model was calibrated¹³ with data for semi-natural ecosystem sites of northern Europe, and with generic ¹⁴C data to constrain the contributions of the fast, slow and passive pools.

N14CP was applied to British semi-natural sites on a 5 km × 5 km grid. For the period 1910 to 2011, the model was driven by seasonal temperatures and precipitation⁶⁸. Prior to this, long-term gridded climate averages (1910–1940) were modified by historical anomalies⁶⁹. For the period from 1750, spatially-resolved N_{dep} and S_{dep} were obtained from the deposition model described above with values interpolated linearly between 1750 and the deposition model years of 1800, 1900, 1950, 1970 and 2010. Base cation deposition over time was estimated from values for 2012 from the CBED model⁶⁷, combined with an historical anomaly¹³ to provide time-series inputs. The UK Land Cover Map 2007 (LCM)³³ was processed to provide fractional covers on this grid scale for the three broad habitat types examined here, i.e. broadleaf woodland, grasslands and shrubs. It was assumed that areas classified today as broadleaf woodland have been so since forests succeeded post-glaciation grasslands, broadly estimated to have been at 6000 BC. Contemporary grasslands were assumed to have been grasslands since forest clearance, dates for which are estimated from data across GB⁷⁰. Shrublands were assumed always to have been shrublands. An initial weatherable P pool (gPm⁻²) was set for each grid square on the basis of two soil types ((i) podzols and rankers, (ii) other soils)¹³, the fractional coverage of each being calculated from gridded soil survey data from the National Soil Resources Institute⁷¹ and the James Hutton Institute (JHI)⁷². Model outputs, on an annual basis, were the pools of SOC and nutrients (biomass C, N and P, topsoil and subsoil SOC, SON and SOP, P sorbed to soil surfaces), and their fluxes (NPP, N fixation, CO₂, N lost via denitrification, DOC, DON, DOP, and dissolved inorganic N and P).

Model performance. To constrain the model parameters and test the model's performance, we previously¹³ conducted a detailed calibration using plot-scale data for 44 semi-natural ecosystem sites (woodland, grassland, shrubland) of northern Europe, and blind-tested this non-site specific parameter set against a further 44 sites. A comprehensive sensitivity analysis considering parameter interactions identified the parameters which had most influence on the multiple outputs of interest, including NPP, soil C, N and P, and dissolved nutrients. These parameters were calibrated in a combination of multiple local and global searches using NPP, soil C, N and P pools and fluxes, and generic radiocarbon data, and then the model was tested using the same output variables. The results showed that the model successfully simulates and predicts average ecosystem behaviour, but in the absence of site-specific information, e.g. P weathering rate, it cannot distinguish the behaviours of individual sites.

In addition to the site-scale testing¹³, evidence that N14CP can provide reliable predictions is as follows. (1) The model found that N limitation is widespread which agrees with literature data⁷³. (2) The model predicts an increase of c. 30 gC in above-ground unmanaged woodland biomass per g N_{dep}⁻¹, in agreement with the range of 15–40 gC g N⁻¹ estimated for European forests¹⁰. (3) A general assessment of model performance for Great Britain was performed by comparing modelled topsoil C pools with values estimated for a depth of 15 cm from measured SOC concentrations reported by the Countryside Survey for Great Britain (CS)^{16,74} (see below), by estimating soil bulk density (BD) from %C using an equation derived from CS data ($BD = 1.29e^{-0.206\%C} + 2.51e^{-0.003\%C} - 2.057$)¹⁶. The average simulated topsoil C pool over all the sites on both sampling dates was 6430 gCm⁻² (standard deviation, sd, 1500 gCm⁻²), in reasonable agreement with the average pool to a depth of 15 cm estimated from the measurements, 7580 gCm⁻² (sd 1320 gCm⁻²). For soils under different vegetation types, the simulated average soil pools (gCm⁻²) were 5701 (broadleaf woodland), 6842 (grassland) and 9897 (shrubland), in the same sequence as the observed values of 7440, 7505 and 8620 respectively. The average modelled soil C:N ratio for the 398 CS sites with measurements of both C and N was 15.6 (sd 1.2) sampled in 1998 and 2007, in agreement with the measured ratio of 16.2 (sd 7.9). (4) The model is tested further in the present work, by comparison of simulated and observed changes in SOC concentration in the late 20th Century (see Results, Tables 3 and 4).

We explored how uncertainty in key input drivers affects model output by conducting a simple sensitivity analysis. In this analysis, we ran N14CP for 9 randomly chosen example locations covering the 3 vegetation types at the 20th, 50th and 80th percentile of cumulative N_{dep} loading between 1750 and 2010. We considered the sensitivity of the soil carbon outputs at these locations to variation in vegetation history, soil type and N_{dep} inputs. The contemporary vegetation at a given location is relatively reliable, but vegetation history is less certain. Therefore we ran the model for different histories, including changes to forest clearance dates and succession (see SI for further details). As described in the preceding section, the fraction coverage of soil types in the grid cells determines the initial P weathering condition. To determine sensitivity to this condition, we varied the soil type between the minimum and maximum P_{weath0} values. Finally, to investigate sensitivity to N_{dep} , we varied the input magnitude for each example by $\pm 30\%$, which encompasses the national scale variation suggested by an inter-comparison of different atmospheric chemistry transport models⁶⁴. In the sensitivity simulations, we considered the one-at-a-time effects of changes to these inputs and the combined effects, requiring 243 model runs.

The results of the sensitivity analysis are shown in Figures S1 and S2. We find that the three input drivers examined have an influence on the magnitude of contemporary carbon stores to varying extents, with shrub locations broadly displaying the greatest sensitivity, which is clearly seen in S1. However, vegetation history and soil type have less influence on the dynamic trends i.e. they have little effect on the change in soil organic carbon over the last ~200 years, which is the focus here. We can conclude that uncertainties in these input drivers do not alter the conclusion that N_{dep} has been a major driver of soil carbon increases for these ecosystem types. The N_{dep} sensitivity results show that the model's estimates for change in SOC are most sensitive to this input.

Soils data. Data were compiled for soils under broadleaf woodland, unimproved grasslands and shrubs, since these ecosystems are generally long-established in the UK. None of them is entirely natural, because of relatively light management, but none is significantly fertilized, and they are commonly referred to as “semi-natural”. Conifer sites were excluded because they are predominantly managed woodland undergoing recent (20th Century) land use change. Ombrotrophic peats were excluded because they are nearly exclusively composed of organic matter, and therefore changes in soil C pools cannot be revealed by changes in soil C concentration. Only results referring to the same vegetation on both sampling occasions were used, so as to minimise errors due to site mis-identification. In all cases, soils were sampled from the surface to a measured depth (15 cm in 98% of cases), and any coarse litter on the surface was removed before sampling. The average time difference between samplings was 27 years. Site locations, sampling dates, and numbers of samples are summarized in Table 1, and SOC concentrations in Table 2.

Data from the National Soil Inventory (NSI) came from surveys carried out in England and Wales¹⁴. The samples were taken to 15 cm depth, bulked from 25 separate soil cores at each site. For both first and second samplings, organic carbon was analysed by the Walkley-Black method at low concentrations (<15% C), and by loss-on-ignition (LOI) for soils with greater %C. Countryside Survey data were obtained from the Centre for Ecology & Hydrology^{16,74,75}. Single 15 cm deep core samples were taken where possible, shallower samples in cases where a full core could not be taken. Samples from each campaign were analysed for LOI, which was converted to %C using a factor of 0.55⁷⁵. The British Woodland Survey (BWS) soil sampling campaigns¹² were conducted with the same protocols as CS. Data from the National Soil Inventory of Scotland (NSIS) were taken from sites first sampled in 1978–88 and resampled in 2007–9 by JHI staff. Additionally, we included data from samples taken from soils rare in a Scottish context, initially sampled between 1959 and 1981 and resampled in 2010. The Scottish data referred to excavated pits with sampling by soil horizon, and therefore not to a specified depth. However, sampling depths for the first and second samplings were very similar; on both sampling occasions the range of topsoil depths was from 10 to 24 cm, with an average of 17 cm.

Estimation of SOC concentrations from simulated SOC pools. We compared simulated and measured changes in SOC in terms of SOC concentration, [SOC] (%C), which was the measured and reported variable in the soil surveys (see above). To relate [SOC] to the simulated SOC pool (Q_{SOC} , gCm^{-2}), we assumed that the topsoil mineral matter pool, Q_{MM} (gm^{-2}), which is not modelled with N14CP, was constant, and made the approximation that the topsoil was homogeneous. The mean Q_{SOC} during the period between two samplings is given by

$$Q_{\text{SOC,mean}} = (Q_{\text{SOC,1}} + Q_{\text{SOC,2}})/2 \quad (1)$$

and the mean simulated SOC concentration (%) by

$$[\text{SOC}]_{\text{sim,mean}} = 100 \times Q_{\text{SOC,mean}} / \{(Q_{\text{SOC,mean}}/f_C) + Q_{\text{MM}}\} \quad (2)$$

where f_C is the fraction of SOM that is SOC (taken to be 0.55 in this work). Rearrangement of equation (2) gives

$$Q_{\text{MM}} = \{100 \times Q_{\text{SOC,mean}} / [\text{SOC}]_{\text{sim,mean}}\} - \{Q_{\text{SOC,mean}}/f_C\} \quad (3)$$

which allows the value of Q_{MM} to be calculated by assuming that $[\text{SOC}]_{\text{sim,mean}}$ is equal to the mean of the measured values of [SOC], i.e. $[\text{SOC}]_{\text{obs,1}}$ and $[\text{SOC}]_{\text{obs,2}}$, at the two samplings. The simulated values of [SOC] at the first and second samplings are

$$[\text{SOC}]_{\text{sim,1}} = 100 \times Q_{\text{SOC,1}} / \{(Q_{\text{SOC,1}}/f_C) + Q_{\text{MM}}\} \quad (4)$$

$$[\text{SOC}]_{\text{sim,2}} = 100 \times Q_{\text{SOC,2}} / \{(Q_{\text{SOC,2}}/f_C) + Q_{\text{MM}}\} \quad (5)$$

We used the variable R_{SF} (i.e. $[SOC]_2/[SOC]_1$) to assess change in SOC, to take into account the two ways that a soil can respond to extra C, which are by increasing in C concentration and by thickening. Thus if a soil has low %C, i.e. is high in mineral matter, any additional C mostly goes to increasing the C concentration, whereas a soil with high %C and little mineral matter will respond by thickening, with little or no change in C concentration¹⁷. If SOC (%) is used as the diagnostic variable, results for the latter type of soil exert a disproportionate influence in the statistical analysis, but the ratio approach avoids this.

Statistical Methods. Change in SOC was evaluated by modelling the ratio of values at two time points (R_{SF}) on the logarithmic scale, using a linear mixed effects model⁷⁶. A mixed effects model estimates global fixed effects across the whole system whilst accounting for differences within it by allowing additional sources of error to be included in the model specification. These so called random effects were used here to account for potential differences arising from pooling data across different sites. Similarly data from different habitats could result in random deviances from the global average that are not consistent, i.e. they have the same variance, across habitats. $\log_{10}R_{SF}$ was therefore modelled simply using an intercept term to estimate the mean overall change, but incorporating random effects representing different sampling units from which the data are obtained. Analyses were carried out in the R statistical computing environment⁷⁷.

To test whether the mean $\log_{10}R_{SF}$ differs significantly from zero (i.e. whether mean R_{SF} differs from 1.0) one would naturally want to test whether the fitted intercept in the model deviates from zero. However, because the $\log_{10}R_{SF}$ data cannot be expected to conform to a normal distribution, or any other exponential family distribution, the estimated variance parameters output directly from the model would not be robust enough for hypothesis testing. We therefore adopted a bootstrapping⁷⁸ approach to test the null hypothesis that $\log_{10}R_{SF} = 0$. We produced random multiple pseudo-sample sets by re-ordering pairs of SOC values repeatedly (1000 times) and re-estimating the mean $\log_{10}R_{SF}$. The mixed model described above was then fitted in exactly the same way to each of these sets to produce 1000 estimates of mean $\log_{10}R_{SF}$ against which the observed change from the original data could be compared. The probability (p) value was then obtained as the percentile amongst the pseudo $\log_{10}R_{SF}$ values corresponding to the observed $\log_{10}R_{SF}$.

Testing whether there was a significant relationship between $\log_{10}R_{SF}$ and N_{dep} was carried out in a similar manner. Linear mixed effects models were used as described above with the addition of N_{dep} as a fixed effect term in the model as well as the intercept term. The random effects structure remained the same and a similar bootstrap procedure was adopted based on the estimated coefficient between $\log_{10}R_{SF}$ and N_{dep} .

References

- Schlesinger, W. H. & Bernhardt, E. S. *Biogeochemistry. An Analysis of Global Change* (Elsevier, 2013).
- Jandl, R. *et al.* Current status, uncertainty and future needs in soil organic carbon monitoring. *Sci. Tot. Environ* **468–469**, 376–383, doi:10.1016/j.scitotenv.2013.08.026 (2014).
- Townsend, A. R., Braswell, B. H., Holland, E. A. & Penner, J. E. Spatial and temporal patterns in terrestrial carbon storage due to deposition of fossil fuel nitrogen. *Ecol. Appl* **6**, 806–814, doi:10.2307/2269486 (1996).
- Butterbach-Bahl, K. *et al.* *The European Nitrogen Assessment* (eds Sutton M.A. *et al.*) Ch. 6, 99–125 (Cambridge University Press, 2014).
- Reay, D. S., Dentener, F., Smith, P., Grace, J. & Feely, R. A. Global nitrogen deposition and carbon sinks. *Nature Geosci* **1**, 430–437, doi:10.1038/ngeo230 (2008).
- UK National Atmospheric Emission Inventory (NAEI) <http://www.naei.org.uk> (Date of access 09/11/2016) (2014).
- De Vries, W., Reinds, G. J., Gundersen, P. & Sterba, H. The impact of nitrogen deposition on carbon sequestration in European forests and forest soils. *Global Change Biol* **12**, 1151–1173, doi:10.1111/gcb.2006.12.issue-7 (2006).
- Kahle, H.-P. (ed.) *Causes and Consequences of Forest Growth Trends in Europe* (Brill, 2008).
- Frey, S. D. *et al.* Chronic nitrogen additions suppress decomposition and sequester soil carbon in temperate forests. *Biogeochem* **121**, 305–316, doi:10.1007/s10533-014-0004-0 (2014).
- De Vries, W. *et al.* The impact of nitrogen deposition on carbon sequestration by European forests and heathlands. *For. Ecol. Manag* **258**, 1814–1823, doi:10.1016/j.foreco.2009.02.034 (2009).
- De Vries, W., Du, E. & Butterbach-Bahl, K. Short and long-term impacts of nitrogen deposition on carbon sequestration by forest ecosystems. *Current Opinion Environ. Sustain* **9–10**, 90–104, doi:10.1038/srep12625 (2015).
- Kirby, K. J. *et al.* Long term ecological change in British woodland (1971–2001). A re-survey and analysis of change based on the 103 sites in the Nature Conservancy ‘Bunce 1971’ woodland survey. *English Nature Research Report Number 653* (English Nature, 2005).
- Davies, J. A. C. *et al.* Long-term P weathering and recent N deposition control contemporary plant-soil C, N and P. *Glob. Biogeochem. Cycles* **30**, 231–249, doi:10.1002/gbc.v30.2 (2016).
- Bellamy, P. H., Loveland, P. J., Bradley, R. I., Lark, R. M. & Kirk, G. J. D. Carbon losses from all soils across England and Wales 1978–2003. *Nature* **437**, 245–248, doi:10.1038/nature04038 (2005).
- Kirk, G. J. D. & Bellamy, P. H. Analysis of changes in organic carbon in mineral soils across England and Wales using a simple single-pool model. *Eur. J. Soil Sci* **61**, 406–411, doi:10.1111/ejs.2010.61.issue-3 (2010).
- Reynolds, B. *et al.* Countryside Survey: National ‘Soil Change’ 1978–2007 for Topsoils in Great Britain—Acidity, Carbon, and Total Nitrogen Status. *Vadose Zone J* **12**, 0, doi:10.2136/vzj2012.0114 (2013).
- Chapman, S. J. *et al.* Comparison of soil carbon stocks in Scottish soils between 1978 and 2009. *Eur. J. Soil. Sci.* **64**, 455–465, doi:10.1111/ejss.12041 (2013).
- Sutton, M. A., Fowler, D. & Moncrieff, J. B. The exchange of atmospheric ammonia with vegetated surfaces. 1. Unfertilised vegetation. *Quart. J. Roy. Met. Soc.* **119**, 1023–1045, doi:10.1002/(ISSN)1477-870X (1993).
- Whittinghill, K. A., Currie, W. S., Zak, D. R., Burton, A. J. & Pregitzer, K. S. Anthropogenic N Deposition increases soil C storage by decreasing the extent of litter decay: Analysis of field observations with an ecosystem model. *Ecosystems* **15**, 450–461, doi:10.1007/s10021-012-9521-7 (2012).
- Janssens, I. A. *et al.* Reduction of forest soil respiration in response to nitrogen deposition. *Nature Geoscience* **3**, 315–322, doi:10.1038/ngeo844 (2010).
- Walker, A. P. *et al.* Predicting long-term carbon sequestration in response to CO₂ enrichment: How and why do current ecosystem models differ? *Global Biogeochem. Cycles* **29**, 476–495, doi:10.1002/2014GB004995 (2015).
- Ciais, P. *et al.* Carbon accumulation in European forests. *Nature Geosci* **1**, 425–429, doi:10.1038/ngeo233 (2008).
- Ashmore, M. R. Assessing the future global impacts of ozone on vegetation. *Plant, Cell and Environment* **28**, 949–964, doi:10.1111/pce.2005.28.issue-8 (2005).

24. Kirk, G. J. D., Bellamy, P. B. & Lark, R. M. Changes in soil pH across England and Wales in response to decreased acid deposition. *Glob. Change Biol* **16**, 3111–3119 (2010).
25. Prieztzel, J., Zimmermann, L., Schubert, A. & Christophel, D. Organic matter losses in German Alps forest soils since the 1970s most likely caused by warming. *Nature Geosci.* Published online (2016).
26. Grüneberg, E., Ziche, D. E. & Wellbrock, N. Organic carbon stocks and sequestration rates of forest soils in Germany. *Glob. Change Biol.* **20**, 2644–2662, doi:10.1111/gcb.2014.20.issue-8 (2014).
27. Bouwman, A. F. *et al.* A global high resolution emission inventory for ammonia. *Glob. Biogeochem. Cycles* **11**, 561–587, doi:10.1029/97GB02266 (1997).
28. Sutton, M. A. *et al.* Toward a climate-dependent paradigm of ammonia emission & deposition. *Phil. Trans. Roy. Soc. (Ser. B)* **368**, 20130166 (2013).
29. Fowler, D. *et al.* A Chronology of Nitrogen Deposition in the UK between 1900 and 2000. *Water, Air and Soil Pollution: Focus* **4**, 9–23, doi:10.1007/s11267-004-3009-1 (2004).
30. Sutton, M. A., Erismann, J. W., Dentener, F. & Moeller, D. Ammonia in the environment: from ancient times to the present. *Environ. Pollut.* **156**, 583–604, doi:10.1016/j.envpol.2008.03.013 (2008).
31. Dragosits, U., Sutton, M. A., Place, C. J. & Bayley, A. A. Modelling the spatial distribution of agricultural ammonia emissions in the UK. *Environmental Pollution* **102**(S1), 195–203, doi:10.1016/S0269-7491(98)80033-X (1998).
32. Hellsten, S. *et al.* Modelling seasonal dynamics in the UK ammonia emission inventory. *Water Air and Soil Pollution: Focus* **7**, 3–11, doi:10.1007/s11267-006-9087-5 (2007).
33. Morton, D. *et al.* *Final Report for LCM2007 – the New UK Land Cover Map.* (CS Technical Report No. 11/07 NERC/Centre for Ecology & Hydrology, 2011).
34. Dowling, M. W., Clarkson, L. A., Kennedy, L. & Crawford, E. M. *Database of Irish Historical Statistics: Agricultural Statistics 1911–1973* (1998).
35. Turner, M. E. Counting sheep: waking up to new estimates of livestock numbers in England c 1800. *Agric. History Rev* **46**(2), 142–161 (1998).
36. Clarkson, L. A., Kennedy, L., Crawford, E. M. & Dowling, M. W. *Database of Irish Historical Statistics: Population, 1821–1911* (UK Data Service. SN: 3578, 1997) <http://dx.doi.org/10.5255/UKDA-SN-3578-1> (2014).
37. Thompson, F. M. L. Nineteenth-century horse sense. *Economic Hist. Rev* **29**, 60–81, doi:10.2307/2594507 (1976).
38. FAO. *Report on the 1950 World Census of Agriculture* (FAO, 1955).
39. EDINA. *Agricultural Census database (AgCensus)* <http://edina.ac.uk/agcensus/> (Date of access 09/11/2016) (2014).
40. Vision of Britain Historic agricultural data. <http://www.visionofbritain.org.uk> (Date of access 09/11/2016) (2014).
41. Cooke, G. W. Changes in fertiliser use in the UK from 1950 to 1980. *Proc. Fertiliser Society* **190** (1980).
42. Szpak, P., Millaire, J. E., White, C. D. & Longstaffe, F. J. Influence of seabird guano and camelid dung fertilization on the nitrogen isotopic composition of field-grown maize (*Zea mays*). *J. Archaeolog. Sci* **39**, 3721–3740, doi:10.1016/j.jas.2012.06.035 (2012).
43. Wood, T. B. Losses in making and storing farmyard manure. *J. Agric. Sci* **2**, 207–215, doi:10.1017/S002185960001283 (1907).
44. Crowther, B. Y. C. & Woodman, H. E. A study of nitrogen metabolism in a dairy cow. *J. Agric. Sci* **2**, 207–215 (1917).
45. Broster, W. H., Tuck, V. J., Smith, T. & Johnson, V. W. Experiments on the nutrition of the dairy heifer: VII. Observations on the effect of the energy intake on the utilization of protein in growth and in lactation. *J. Agric. Sci* **72**, 13–30, doi:10.1017/S0021859600020372 (1969).
46. Pike, I. H. The effect on nitrogen utilization in the sow of variation in dietary protein concentration and pattern of feeding in pregnancy. *J. Agric. Sci* **74**, 209–215, doi:10.1017/S0021859600022826 (1970).
47. Mitchell, B. R. *British Historical Statistics* (Cambridge University Press, 1998).
48. Turner, M. E., Beckett, J. V. & Afton, B. *Agricultural Production, Output, and Productivity, 1700–1914* In *Farm Production in England 1700–1914* (Oxford University Press, 2001).
49. Brenkert, A. L., Smith, A. J., Kim, S. H. & Pitcher, H. M. *Molde documentation for the MiniCAM* (Pacific Northwest National Laboratory, 2003).
50. Department of Energy and Climate Change. *1948 Power Stations in Great Britain (DECC, 2012)* <https://www.decc.gov.uk/en/content/cms/statistics/publications/dukes/dukes.aspx> (Date of access 09/11/2016) (2014).
51. Department of Energy and Climate Change. *Historical electricity data: 1920 to 2014 (DECC, 2013)* <https://www.gov.uk/government/statistical-data-sets/historical-electricity-data-1920-to-2011> (Date of access 09/11/2016) (2014).
52. Department of Energy and Climate Change. *Historical coke and breeze data: coal carbonized and coke and breeze produced at coke ovens 1920 to 2013 (DECC, 2014)* <https://www.gov.uk/government/statistical-data-sets/historical-coke-and-breeze-data-coal-carbonized-and-coke-and-breeze-produced-at-coke-ovens-1920-to-2011> (Date of access 09/11/2016) (2014).
53. Department of Energy and Climate Change. *Historical gas data: gas production and consumption and fuel input 1920 to 2013. (DECC 2014)* <https://www.gov.uk/government/statistical-data-sets/historical-gas-data-gas-production-and-consumption-and-fuel-input-1882-to-2011> (Date of access 09/11/2016) (2014).
54. Department of Energy and Climate Change. *Historical coal data: coal production, availability and consumption 1853 to 2014. (DECC, 2015)* <https://www.gov.uk/government/statistical-data-sets/historical-coal-data-coal-production-availability-and-consumption-1853-to-2011> (Date of access 09/11/2016) (2015).
55. Flinn, M. W. *The History of The British Coal Industry* Volume 2, 1700–1830 *The Industrial Revolution* (Clarendon Press, 1984).
56. Lee, D. S. & Longhurst, J. W. S. Estimates of emissions of SO₂, NO_x, HCl and NH₃ from a densely populated region of the UK. *Environ. Pollut.* **79**, 37–44, doi:10.1016/0269-7491(93)90175-N (1993).
57. Olivier, J. G. J. *et al.* Sectoral emission inventories of greenhouse gases for 1990 on a per country basis as well as on 1° × 1°. *Environ. Sci. Pol* **2**, 241–263 (1999).
58. Tsagatakis, I., Jephcote, C., Brace, S. & Passant, N. *UK Emission Mapping Methodology: A Report of the National Atmospheric Emission Inventory 2013* (Ricardo Energy & Environment, 2015).
59. R'Honi, Y. *et al.* Exceptional emissions of NH₃ and HCOOH in the 2010 Russian wildfires. *Atmos. Chem. Phys.* **13**, 4171–4181, doi:10.5194/acp-13-4171-2013 (2013).
60. Stamp, L. D. (ed.). *The Land of Britain. The Report of the Land Utilisation Survey of Britain* (Geographical Publications, 1937).
61. Hallsworth, S. *et al.* The role of indicator choice in quantifying the threat of atmospheric ammonia to the 'Natura 2000' network. *Environ Sci Policy* **13**, 671–687, doi:10.1016/j.envsci.2010.09.010 (2010).
62. Dore, A. J. *et al.* The influence of model grid resolution on estimation of national scale nitrogen deposition and exceedance of critical loads. *Biogeosc* **9**, 1597–1609, doi:10.5194/bg-9-1597-2012 (2012).
63. Simpson, D. *et al.* Impacts of climate and emission changes on nitrogen deposition in Europe: A multi-model study. *Atmos. Chem. Physics* **14**, 6995–7017, doi:10.5194/acp-14-6995-2014 (2014).
64. Dore, A. J. *et al.* Evaluation of the performance of different atmospheric chemical transport models and inter-comparison of nitrogen and sulphur deposition estimates for the UK. *Atmos. Environ.* **119**, 131–143, doi:10.1016/j.atmosenv.2015.08.008 (2015).
65. Dore, A. J. *et al.* Modelling the atmospheric transport and deposition of sulphur and nitrogen over the United Kingdom and assessment of the influence of SO₂ emissions from international shipping. *Atmos. Environ.* **41**, 2355–2367, doi:10.1016/j.atmosenv.2006.11.013 (2007).
66. Matejko, M. *et al.* The influence of long term trends in pollutant emissions on deposition of sulphur and nitrogen and exceedance of critical loads in the United Kingdom. *Environ. Sci. Pol* **12**, 882–896, doi:10.1016/j.envsci.2009.08.005 (2009).

67. Werner, M. *et al.* Modelling of base cation emissions, concentrations and deposition in the UK. *Atmos. Chem. Phys. Discuss.* **10.9**, 21989–22018, doi:[10.5194/acpd-10-21989-2010](https://doi.org/10.5194/acpd-10-21989-2010) (2010).
68. Perry, M. & Hollis, D. The generation of monthly gridded datasets for a range of climatic variables over the UK. *Internat. J. Climatol.* **25.8**, 1041–1054, doi:[10.1002/\(ISSN\)1097-0088](https://doi.org/10.1002/(ISSN)1097-0088) (2005).
69. Mauri, A., Davis, B. A. S., Collins, P. M. & Kaplan, J. O. The climate of Europe during the Holocene: a gridded pollen-based reconstruction and its multi-proxy evaluation. *Quatern. Sci. Rev.* **112**, 109–127, doi:[10.1016/j.quascirev.2015.01.013](https://doi.org/10.1016/j.quascirev.2015.01.013) (2015).
70. Roberts, N. *The Holocene: An Environmental History* (Wiley, 2013).
71. National Soil Resources Institute. Soilscales. <http://www.landis.org.uk/soilscales> (Date of access 09/11/2016) (2013).
72. Soil Survey of Scotland Staff. *Soil Maps of Scotland at a Scale of 1:250 000* (Macaulay Institute for Soil Research, 1981).
73. LeBauer, D. S. & Treseder, K. K. Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology* **89**, 371–379, doi:[10.1890/06-2057.1](https://doi.org/10.1890/06-2057.1) (2008).
74. Emmett, B. A. *et al.* Countryside Survey Soils Report from 2007. Countryside Survey Technical Report No. 9/07 (Centre for Ecology & Hydrology, 2010).
75. Emmett, B. A. *et al.* Countryside Survey Soils Manual. Countryside Survey Technical Report No. 04/07 (Centre for Ecology & Hydrology, 2008).
76. Pinheiro, J. C. & Bates, D. M. *Mixed-Effects Models in S and S-PLUS* (Springer, 2000).
77. R Development Core Team. *R: A Language and Environment for Statistical Computing* (2011).
78. Efron, B. & Tibshirani, R. J. *An Introduction to the Bootstrap* (CRC Press, 1994).

Acknowledgements

This research was funded by the UK Natural Environment Research Council Macronutrient Cycles Programme (LTLS project, grants NE/J011533/1 and NE/J011703/1) and by the Rural & Environment Science & Analytical Services Division of the Scottish Government. We thank E.C. Rowe (CEH Bangor) for helpful comments.

Author Contributions

E.T. coordinated the study, wrote the initial draft, contributed to the ecosystem modelling and statistical analysis. J.A.C.D. led the ecosystem modelling. P.A.H. led the statistical analysis. G.J.D.K. and A.L. collaborated on soil data interpretation. U.D., E.J.C., S.J.T. and A.J.D. performed the emissions and deposition modelling; M.A.S. coordinated input from atmospheric measurements and contributed to the modelling strategy. All authors contributed to the final version.

Additional Information

Supplementary information accompanies this paper at doi:[10.1038/s41598-017-02002-w](https://doi.org/10.1038/s41598-017-02002-w)

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017