



Barber, J., & Itasaki, N. (2016). Expression of Prolyl hydroxylases 2 and 3 in chick embryos. *Gene Expression Patterns*, 21(2), 97-102. DOI: 10.1016/j.gep.2016.04.001

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.gep.2016.04.001](https://doi.org/10.1016/j.gep.2016.04.001)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <http://www.sciencedirect.com/science/article/pii/S1567133X16300163>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms.html>

Expression of Prolyl hydroxylases 2 and 3 in chick embryos

Jake Barber and Nobue Itasaki
Faculty of Health Sciences, University of Bristol
Southwell Street, Bristol BS2 8EJ, UK

Correspondence:

Nobue Itasaki
nobue.itasaki@bristol.ac.uk

Key words:

Prolyl Hydroxylase
EGLN
Hypoxia
HIF
chick embryo

Abstract

Hypoxic cellular response is crucial for normal development as well as in pathological conditions in order to tolerate low oxygen. The response is mediated by Hypoxia Inducible Factors (HIFs), where the α -subunit of HIF is stabilised and able to function only in low oxygen. Prolyl hydroxylases (PHDs) are oxygen dependent dioxygenase enzymes that hydroxylate HIF- α leading to HIF degradation. Thus PHDs function as an oxygen sensor for the function of HIFs. Here we describe the mRNA expression pattern of PHDs in chick embryos. Up to embryonic day 2, PHDs are weak without specific localisation, whereas from day 3 localised expression was observed in the eye, branchial arches and dermomyotome. Later in the limb development PHDs were expressed in the perichondral mesenchyme, excluded from the developing limb cartilages.

Introduction

Tolerance to hypoxia is essential in embryonic development (Dunwoodie, 2009) as well as in ischemic conditions in adults (Semenza, 2014). Hypoxic cellular response is mostly exerted via the activity of Hypoxia Inducible Factor (HIF), a heterodimeric complex composed of an oxygen-sensitive HIF- α and a constitutively expressed HIF- β subunit (Wang et al., 1995). As a transcription factor, HIF targets over 100 genes to counteract the reduction in oxygen levels at the cellular, local tissue and systemic level (Liu et al., 2012). The main role of HIFs includes stimulating angiogenesis with increased VEGF transcription, promoting erythropoiesis, enhancing glucose uptake and glycolysis and engaging programmed cell death for adaptation (Biju et al., 2004; Haase, 2013; Lu et al., 2002; Shweiki et al., 1992). Hypoxia has also been linked to disease processes such as cardiac abnormalities (Bishop and Ratcliffe, 2015) and tumour metastasis (Semenza, 2013).

The primary means of controlling HIF activity in response to oxygen concentration is oxygen-dependent hydroxylation of specific proline residues of the α -subunit, which promotes an association with the von Hippel-Lindau ubiquitin E3 ligase and subsequent destruction of HIF- α (Bruick and McKnight, 2001). HIF prolyl hydroxylation is catalysed by prolyl hydroxylase domain (PHD) enzymes 1-3, also known as Egl-2, -1 and -3 respectively, that are members of the iron and oxygen dependent, 2-oxoglutarate-dependent dioxygenase family (Kaelin and Ratcliffe, 2008). All three PHDs are similar at the C-terminus containing prolyl-4-hydroxylase-sequence, but PHD2 has a zinc finger at its N-terminus absent from the other PHDs (Place and Domann, 2013). PHDs act as oxygen sensors that keep HIF at low levels during normoxia. On the contrary, when oxygen is not available, PHDs do not hydroxylate HIF- α , thus allowing HIF- α proteins to be stabilised and function

(Semenza, 2014). PHDs are expressed in different cell types and tissues, and contribute to the physiological regulation of HIF- α (Appelhoff et al., 2004; Berra et al., 2003; Lieb et al., 2002). PHDs are transcriptionally up-regulated by HIF- α and therefore involved in a feedback loop with HIF during prolonged hypoxia or rapid degradation upon re-oxygenation (Loor and Schumacker, 2008; Speer et al., 2013).

The importance of HIF-1 α in embryogenesis was demonstrated by targeted deletion that caused a significant decrease in vascularisation and abnormal head fold formation manifest at 8.5-9.5 dpc (Iyer et al., 1998; Ryan et al., 1998). With regard to PHDs, PHD2 knock-outs in mouse leads to embryonic lethality at 12.5-14.5 dpc, whereas deletion of PHD1 or PHD3 does not affect viability (Takeda et al., 2006). The major cell differentiation processes during embryogenesis where the HIF pathway is likely involved are vasculogenesis and chondrogenesis. In addition, epithelial to mesenchymal transition (EMT) may also be initiated by HIF, as HIF directly regulates *Snail1* expression (Imai et al., 2003; Luo et al., 2011).

The expression of HIF-1 α and HIF-2 α in embryos has been described in detail in various species (de Beaucourt and Coumailleau, 2007; Etchevers, 2003; Köblitz et al., 2015; Ota et al., 2007). However, description on PHD expression is largely limited to adult tissues and cell lines (Appelhoff et al., 2004; Lieb et al., 2002); the only description in embryos is in *Xenopus*, where PHDs are expressed in the brain, branchial arches, otic vesicles and pronephros (Han et al., 2012). PHD3 is also expressed in the heart and somites (Han et al., 2012). However, as aqueous animals *Xenopus* may have roles and regulatory mechanisms for the HIF pathway different from amniotes during embryogenesis. Here we describe expression pattern of PHDs in chick embryos focusing on early developmental stages. It should be noted that, in

contrast to other vertebrates that have three PHD genes, chick embryos appear to have only two; PHD2 and PHD3.

Results and Discussion

Gene homology search on Ensembl (<http://www.ensembl.org>) showed that chicks have only two PHDs; PHD2 (EGLN1) and PHD3 (EGLN3) (Fig. 1). Chick PHD3 showed a similarity in both size and amino acid sequences (%) to those of other species, whereas chick PHD2 is smaller (316 amino acids) compared to PHD1 or PHD2 of other species (≥ 400 amino acids). Thus chicks appear to have unique sets of PHDs compared to other vertebrates. The avian species that are sequenced on Ensembl are Chicken, Turkey, Duck, Flycatcher and Zebra Finch. Among those, Chicken and Turkey have exclusively EGLN1 and 3, whereas Duck, Flycatcher and Zebra Finch have EGLN1, EGLN3 and another gene called P4H-TM (prolyl 4-hydroxylase, transmembrane, which may also be called PHD4/EGLN4 (Koivunen et al., 2007). PHD4/EGLN4 is found in most of vertebrate species such as Humans, Mouse, Xenopus and Zebrafish. None of the avian species mentioned above have PHD1 (EGLN2) in the Ensembl search. Therefore, the lack of PHD1 appears to be common in avian species.

Chick embryos at Hamburger and Hamilton (Hamburger and Hamilton, 1951) stages 8-13 showed faint and broad staining of both PHD2 and PHD3 (Fig. 2A-F). To note, at this stage *HIF1- α* is expressed in embryo proper, in a broad manner similarly to PHDs, whereas *HIF2- α* is in extraembryonic tissues (Ota et al., 2007).

To examine whether the expression of PHD2/3 changes in response to availability of oxygen or by stabilisation of HIF, the expression was compared between *in ovo* and *ex ovo* cultured embryos, along with the ones cultured *ex ovo* with dimethyloxalyl glycine (DMOG) (Fig. 1G-L). The difference of staining between three conditions was marginal in both PHD2 and PHD3, although, there was a trend that *ex ovo* cultured embryos showed slightly weaker staining of PHD3 compared to

that of *in ovo* (n=4/6; no apparent difference for PHD2, n=0/6) whereas DMOG treatment caused stronger staining compared to control *ex ovo* cultures (n=4/6 for PHD2, n=5/6 for PHD3).

At stage 19, localised staining of PHD2 was seen in the telencephalon, dorsal mesencephalon, the eye and optic stalk, the maxillary process and 1st and 2nd pharyngeal arches (Fig. 3A,B). A periodic expression in the somite was also observed (Fig. 3A). Transverse sections of the trunk showed PHD2 expression in the apical side of epithelialized somites (Fig. 3G-I) and in the dermomyotome in more differentiated somites (Fig. 3F). The expression of PHD3 was similar to PHD2 in the cranial region at this stage, however, the expression in the somites was not prominent (Fig. 3C-E).

By stage 21, localised expression of PHD2 in the eye, the maxillary process and 1st and 2nd pharyngeal arches had become prominent (Fig. 4A). It has also become noticeable that the heart is relatively weak in staining. This is in agreement with the localised expression of HIF-1 α in the developing heart (Nanka et al., 2006).

The expression pattern of PHD2 in somites had also become distinct by stage 21 (Fig 4B,E,F). Histological sections showed that the expression is localised to the ventral side of the dermomyotome (Fig. 4E) and further localised to the medial part in more differentiated (anterior) levels (Fig. 4F). In contrast, PHD3 did not show such specific expression in somites, while the expression in the eyes and branchial arches are maintained (Fig. 3C,D,G).

The developing limb buds showed a broad expression of both PHD2 and PHD3 (Fig. 4H-K). At stage 27, when the digital cartilages have become prominent, both PHD2 and PHD3 were expressed in the inter-digital mesenchyme, particularly at the edges of the cartilage, with clear exclusion from the cartilage (Fig. 5A-G). This expression was also observed at stage 32 when the digits are elongated (Fig. 5H,I).

There was no obvious difference between the expression of PHD2 and PHD3 in the developing limb. Histological section showed the PHD-expression is in the limb mesenchyme underneath the surface ectoderm (Fig. 5J).

Methods and methods

Chick embryos were purchased from local farm and incubated at 38°C in humidified incubator. Embryos were staged according to Hamburger and Hamilton (Hamburger and Hamilton, 1951). In situ hybridisation was performed as described previously (Acloque et al., 2008). Cartilage staining was performed as described (Bronner-Fraser, 2008). In aligning the protein structure of PHDs, mouse PHD structures were based on; PHD1, AF453879; PHD2, NM_053207; PHD3, NM_028133. Chick PHD2 and PHD3 were based on XM_001231253 and XM_421233, respectively. Xenopus PHD structures were based on; PHD1 NM_001167742; PHD2, NM_001093091; PHD3, NM_001127012. Chick PHD2 and PHD3 probes, antisense and sense, were synthesized using Chick EST clone 400i24 (880 bp) and 680g23 (490 bp), respectively. The PHD2 clone was chosen based on the matching with XM_001231253, including a part of the coding region and 285 base pair of 3'UTR. The PHD3 clone was chosen based on the matching to both XM_421233 and XM_004941757, that are identical at 1-229 amino acids whereas the C-terminal 12 and 20 amino acids, respectively, are different.

Ex ovo culture was performed on agar plate as described (Chapman et al., 2001) with either 0.1% DMSO or 2mM DMOG (Cayman Chemical) dissolved in the agar. The amount of DMSO was the same for both *ex ovo* control and DMOG cultures. Embryos were taken out of the shell at HH stages 5-7, cultured in ambient air on the agar plate and further incubated for 24 hours before fixing.

REFERENCES

- Acloque, H., Wilkinson, D.G. and Nieto, M.A., 2008. In situ hybridization analysis of chick embryos in whole-mount and tissue sections. *Methods Cell Biol.* 87, 169-85.
- Appelhoff, R.J., Tian, Y.-M., Raval, R.R., Turley, H., Harris, A.L., Pugh, C.W., Ratcliffe, P.J. and Gleadle, J.M., 2004. Differential function of the prolyl hydroxylases PHD1, PHD2, and PHD3 in the regulation of hypoxia-inducible factor. *Journal of Biological Chemistry.* 279, 38458-38465.
- Berra, E., Benizri, E., Ginouves, A., Volmat, V., Roux, D. and Pouyssegur, J., 2003. HIF prolyl-hydroxylase 2 is the key oxygen sensor setting low steady-state levels of HIF-1alpha in normoxia. *EMBO J.* 22, 4082-90.
- Biju, M.P., Neumann, A.K., Bensinger, S.J., Johnson, R.S., Turka, L.A. and Haase, V.H., 2004. Vhlh gene deletion induces Hif-1-mediated cell death in thymocytes. *Mol Cell Biol.* 24, 9038-47.
- Bishop, T. and Ratcliffe, P.J., 2015. HIF hydroxylase pathways in cardiovascular physiology and medicine. *Circ Res.* 117, 65-79.
- Bronner-Fraser, M., 2008. *Avian embryology*, 2nd ed. ed. Elsevier/Academic Press, Amsterdam ; London.
- Bruick, R.K. and McKnight, S.L., 2001. A conserved family of prolyl-4-hydroxylases that modify HIF. *Science.* 294, 1337-40.
- Chapman, S.C., Collignon, J., Schoenwolf, G.C. and Lumsden, A., 2001. Improved method for chick whole-embryo culture using a filter paper carrier. *Dev Dyn.* 220, 284-9.
- de Beaucourt, A. and Coumailleau, P., 2007. Molecular cloning and characterization of the *Xenopus* hypoxia-inducible factor 1alpha (xHIF1alpha). *J Cell Biochem.* 102, 1542-52.
- Dunwoodie, S.L., 2009. The role of hypoxia in development of the Mammalian embryo. *Developmental cell.* 17, 755-773.
- Etchevers, H.C., 2003. Early expression of hypoxia-inducible factor 1alpha in the chicken embryo. *Gene Expr Patterns.* 3, 49-52.
- Haase, V.H., 2013. Regulation of erythropoiesis by hypoxia-inducible factors. *Blood Rev.* 27, 41-53.
- Hamburger, V. and Hamilton, H.L., 1951. A series of normal stages in the development of the chick embryo. *Journal of Morphology.* 88, 49-92.
- Han, D., Wen, L. and Chen, Y., 2012. Molecular Cloning of *phd1* and Comparative Analysis of *phd1*, 2, and 3 Expression in *Xenopus laevis*. *The Scientific World Journal.* 2012, 689287.
- Imai, T., Horiuchi, A., Wang, C., Oka, K., Ohira, S., Nikaido, T. and Konishi, I., 2003. Hypoxia attenuates the expression of E-cadherin via up-regulation of SNAIL in ovarian carcinoma cells. *Am J Pathol.* 163, 1437-47.
- Iyer, N.V., Kotch, L.E., Agani, F., Leung, S.W., Laughner, E., Wenger, R.H., Gassmann, M., Gearhart, J.D., Lawler, A.M., Yu, A.Y. and Semenza, G.L., 1998. Cellular and developmental control of O₂ homeostasis by hypoxia-inducible factor 1 alpha. *Genes Dev.* 12, 149-62.
- Kaelin, W.G. and Ratcliffe, P.J., 2008. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Molecular cell.* 30, 393-402.
- Köblitz, L., Fiechtner, B., Baus, K., Lussnig, R. and Pelster, B., 2015. Developmental Expression and Hypoxic Induction of Hypoxia Inducible Transcription Factors in the Zebrafish. *PloS one.* 10, e0128938.

- Koivunen, P., Tiainen, P., Hyvarinen, J., Williams, K.E., Sormunen, R., Klaus, S.J., Kivirikko, K.I. and Myllyharju, J., 2007. An endoplasmic reticulum transmembrane prolyl 4-hydroxylase is induced by hypoxia and acts on hypoxia-inducible factor alpha. *J Biol Chem.* 282, 30544-52.
- Lieb, M.E., Menzies, K., Moschella, M.C., Ni, R. and Taubman, M.B., 2002. Mammalian EGLN genes have distinct patterns of mRNA expression and regulation. *Biochem Cell Biol.* 80, 421-6.
- Liu, W., Shen, S.M., Zhao, X.Y. and Chen, G.Q., 2012. Targeted genes and interacting proteins of hypoxia inducible factor-1. *Int J Biochem Mol Biol.* 3, 165-78.
- Loor, G. and Schumacker, P.T., 2008. Role of hypoxia-inducible factor in cell survival during myocardial ischemia-reperfusion. *Cell Death Differ.* 15, 686-90.
- Lu, H., Forbes, R.A. and Verma, A., 2002. Hypoxia-inducible factor 1 activation by aerobic glycolysis implicates the Warburg effect in carcinogenesis. *J Biol Chem.* 277, 23111-5.
- Luo, D., Wang, J., Li, J. and Post, M., 2011. Mouse snail is a target gene for HIF. *Mol Cancer Res.* 9, 234-45.
- Nanka, O., Valasek, P., Dvorakova, M. and Grim, M., 2006. Experimental hypoxia and embryonic angiogenesis. *Dev Dyn.* 235, 723-33.
- Ota, K., Nagai, H. and Sheng, G., 2007. Expression and hypoxic regulation of hif1 α and hif2 α during early blood and endothelial cell differentiation in chick. *Gene Expression Patterns.* 7, 761-766.
- Place, T.L. and Domann, F.E., 2013. Prolyl-hydroxylase 3: Evolving Roles for an Ancient Signaling Protein. *Hypoxia.* 2013, 13-17.
- Ryan, H.E., Lo, J. and Johnson, R.S., 1998. HIF-1 alpha is required for solid tumor formation and embryonic vascularization. *EMBO J.* 17, 3005-15.
- Semenza, G.L., 2013. HIF-1 mediates metabolic responses to intratumoral hypoxia and oncogenic mutations. *J Clin Invest.* 123, 3664-71.
- Semenza, G.L., 2014. Oxygen sensing, hypoxia-inducible factors, and disease pathophysiology. *Annu Rev Pathol.* 9, 47-71.
- Shweiki, D., Itin, A., Soffer, D. and Keshet, E., 1992. Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature.* 359, 843-5.
- Speer, R.E., Karuppagounder, S.S., Basso, M., Sleiman, S.F., Kumar, A., Brand, D., Smirnova, N., Gazaryan, I., Khim, S.J. and Ratan, R.R., 2013. Hypoxia-inducible factor prolyl hydroxylases as targets for neuroprotection by "antioxidant" metal chelators: From ferroptosis to stroke. *Free Radic Biol Med.* 62, 26-36.
- Takeda, K., Ho, V.C., Takeda, H., Duan, L.J., Nagy, A. and Fong, G.H., 2006. Placental but not heart defects are associated with elevated hypoxia-inducible factor alpha levels in mice lacking prolyl hydroxylase domain protein 2. *Mol Cell Biol.* 26, 8336-46.
- Wang, G.L., Jiang, B.H., Rue, E.A. and Semenza, G.L., 1995. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc Natl Acad Sci U S A.* 92, 5510-4.

Figure legends

Fig. 1. Schematic drawing of mouse chick and *Xenopus* PHD proteins. The shaded region shows prolyl-4-hydroxylase alpha subunit containing (P4Hc) sequence. Numbers indicate amino acid positions.

Fig. 2. *PHD2* and *PHD3* expression in chick embryos at stages 8-13.

(A-F) HH stages 8 (A, D) and 13 (B,C,E,F) showing *PHD2* (A,B) and *PHD3* (D,E) expression with antisense probe, and negative controls with sense probes (C,F). There is no distinct localisation of expression at these stages. (B) and (C) are stained in parallel hence comparable, so are (E) and (F). Scale bar, 500 μm .

(G-L) HH stage 11-12 embryos incubated *in ovo* (G,J), *ex ovo* with control DMSO (H,K) or *ex ovo* with DMOG (I,L), stained with *PHD2* (G-I) or *PHD3* (J-L). DMOG treated embryos show slightly stronger staining compared to control embryos cultured *ex ovo* with DMSO (H,I; K,L). Embryo cultured *ex ovo* with DMSO shows a slightly weaker staining of *PHD3* compared to *in ovo* cultured embryo (J,K).

Fig. 3. *PHD2* and *PHD3* expression in chick embryos at stage 19.

(A-E) HH stage 19 embryos stained with *PHD2* (A,B) and *PHD3* (C-E). Localised staining of *PHD2* is seen in the telencephalon, dorsal mesencephalon, the eye and optic stalk (arrowhead), the maxillary process and 1st and 2nd pharyngeal arches (arrows), as well as in somites (small arrowheads). (B) is a coronal section crossing the midbrain and pharyngeal arches. ph1; pharyngeal arch 1. The expression pattern of *PHD3* (C) is similar to *PHD2* except the absence in somites. (D) and (E) were stained in parallel with either antisense (D) or sense (E) probes of *PHD3*. Scale bars in (B), 200 μm ; (C), 100 μm .

(F-I) Transverse sections of the trunk showing PHD2 expression in the dermomyotome (F) and in the apical side of epithelialized somites (G-J). Scale bar, 100 μm .

Fig. 4. *PHD2* and *PHD3* expression in chick embryos at stages 19-21.

(A-D) Whole mount embryos stained with PHD2 (A,B) and PHD3 (C,D). Both PHD2 and PHD3 show localised expression in the eye, the maxillary process and 1st and 2nd pharyngeal arches (A,C), however, expressions in somites are seen only for PHD2 (B) and not PHD3 (D). Scale bar, 500 μm .

(E-G) Histological sections at trunk levels showing PHD2 (E,F) and PHD3 (G) expression. (F,G) are more anterior than (E). PHD2 expression is localised to the ventral side of the dermomyotome (E) or the medial part of it in more anterior levels (F). PHD3 does not show prominent expression in somites (G). Scale bar, 200 μm .

(H-K) Whole mount trunk region of stage 19 embryos at the forelimb level stained with PHD2 (H) and PHD3 (J). (I,K) are with sense probes of each. Scale bar, 200 μm .

Fig. 5. *PHD2* and *PHD3* expression in chick limb buds.

(A-G) Whole mount (A,C) and section (B,D) of PHD2 (A,B) and PHD3 (C,D) expression at stage 27, showing expression in the inter-digital mesenchyme excluding from the cartilage. Alcian blue staining is shown in (E) for comparison. (F,G) are staining with antisense and sense probes of PHD2, respectively. Scale bars, 500 μm .

(H-J) Whole mount (H,I) and section (J) of PHD2 (H) and PHD3 (I,J) expression at stage 32. Digital cartilages are negative for both PHD2 and PHD3. (J) shows that PHD3 expression is excluded from the surface ectoderm (bracket). There was no

obvious difference between the expression of PHD2 and PHD3 in the developing limb. Scale bars in (H), 500 μm ; (J), 25 μm .




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5