



Kennedy, E., Heron, J., & Munafò, M. (2017). Substance use, criminal behaviour and psychiatric symptoms following childhood traumatic brain injury: findings from the ALSPAC cohort. *European Child and Adolescent Psychiatry*. DOI: 10.1007/s00787-017-0975-1

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1007/s00787-017-0975-1](https://doi.org/10.1007/s00787-017-0975-1)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Springer at <http://link.springer.com/article/10.1007%2Fs00787-017-0975-1>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms.html>

Supplementary Table 1. Descriptive statistics for participants with injury information included in analyses and participants excluded from analyses due to missing injury information.

	Included (<i>n</i> =11412)	Excluded (<i>n</i> =4033)	p value*
	N (%)	N (%)	
Male	5849 (51.3)	1786 (51.8)	0.553
Social class IV – V ^a	4111 (41.9)	926 (53.6)	<0.001
Rented subsidised housing	1208 (11.4)	731 (25.0)	<0.001
Mother completed secondary school	6496 (62.1)	1587 (78.1)	<0.001
Maternal daily smoking	3034 (28.2)	763 (41.2)	<0.001
Maternal daily alcohol use	1408 (13.1)	103 (5.6)	<0.001
Three or more early life events ^b	5797 (54.4)	330 (27.1)	<0.001
	M (SD)	M (SD)	
Maternal age at birth (years)	28.51 (4.76)	26.16 (5.23)	<0.001
Bonding at 8 months ^c	28.23 (3.65)	28.46 (3.87)	0.040
Positive parenting experience at 21 months ^d	20.78 (2.74)	20.94 (2.90)	0.145
Negative parenting experience at 21 months ^d	5.99 (1.52)	6.00 (1.58)	0.850

Injury from birth to age 16 years (data present *n* = 11412; data missing *n* = 4033); * *p* values calculated using chi square or analysis of variance; ^a highest social class of either parent is skilled non-manual or lower occupation based on the Registrar General's classification of occupations; ^b parent-reported questionnaire relating to upsetting events in the child's life completed when offspring was 6, 30, 42 and 81 months old; ^c parent-report questionnaire completed when offspring was 8 months old; ^d positive and negative parenting experiences based on parent-completed questionnaire when offspring was 21 months old.

Supplementary Table 2 Descriptive statistics for covariates on complete case sample for all covariates and all substance use (alcohol, tobacco, cannabis) measures.

	No Injury (n=1,363)	TBI (n=207)	OI (n=504)	p value*
	N (%)	N (%)	N (%)	
Male	553 (40.6)	112 (54.1)	258 (51.2)	<0.001
Social Class IV – V ^a	450 (33.0)	65 (31.4)	168 (33.3)	0.878
Rented subsidised housing	69 (5.1)	10 (4.8)	18 (3.6)	0.575
Mother completed secondary school	657 (48.2)	93 (44.9)	256 (50.8)	0.338
Maternal daily smoking	241 (17.7)	45 (21.7)	92 (18.3)	0.371
Maternal daily alcohol use	208 (15.3)	36 (17.4)	73 (14.5)	0.619
Three or more early life events ^b	764 (56.0)	132 (63.8)	284 (56.3)	0.120
	M (SD)	M (SD)	M (SD)	
Maternal age at birth (years)	29.84 (4.29)	29.42 (4.26)	29.61 (4.49)	0.600
Bonding at 8 months ^c	27.93 (3.57)	28.06 (3.20)	27.99 (3.49)	0.844
Positive parenting experience at 21 months ^d	5.95 (1.39)	5.94 (1.34)	5.87 (1.34)	0.518
Negative parenting experience at 21 months ^d	20.84 (2.67)	20.55 (2.80)	20.90 (2.58)	0.272

*TBI: traumatic brain injury; OI: orthopaedic injury; * p values calculated using chi square or analysis of variance; ^a highest social class of either parent is skilled non-manual or lower occupation based on the Registrar General's classification of occupations; ^b parent-reported questionnaire relating to upsetting events in the child's life completed when offspring was 6, 30, 42 and 81 months old; ^c parent-report questionnaire completed when offspring was 8 months old; ^d positive and negative parenting experiences based on parent-completed questionnaire when offspring was 21 months old.*

Supplementary Table 3 Associations between traumatic brain injury and orthopaedic injuries from birth to age 16 years and substance use at age 17 years on complete case sample

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol^{a*}				
<i>n</i>	<i>n</i> = 2074	<i>n</i> = 2074	<i>n</i> = 2074	<i>n</i> = 2074
TBI vs no Injury	1.54 (1.15 - 2.06)	1.51 (1.13 - 2.03)	1.48 (1.10 - 2.00)	1.31 (0.94 - 1.82)
OI vs no Injury	0.88 (0.72 - 1.09)	0.87 (0.70 - 1.07)	0.87 (0.70 - 1.08)	0.77 (0.61 - 0.98)
TBI vs OI	1.74 (1.26 - 2.41)	1.75 (1.26 - 2.42)	1.71 (1.23 - 2.38)	1.69 (1.17 - 2.45)
<i>Omnibus p</i>	0.541	0.408	0.412	0.080
Tobacco^{b**}				
<i>n</i>	<i>n</i> = 2074	<i>n</i> = 2074	<i>n</i> = 2074	<i>n</i> = 2074
TBI vs no Injury	1.36 (0.98 - 1.89)	1.43 (1.02 - 1.99)	1.37 (0.98 - 1.93)	1.09 (0.74 - 1.62)
OI vs no Injury	1.09 (0.85 - 1.39)	1.12 (0.87 - 1.43)	1.13 (0.88 - 1.46)	1.15 (0.86 - 1.55)
TBI vs OI	1.25 (0.86 - 1.81)	1.27 (0.88 - 1.85)	1.21 (0.83 - 1.77)	0.95 (0.61 - 1.47)
<i>Omnibus p</i>	0.341	0.238	0.227	0.331
Cannabis^{c**}				
<i>n</i>	<i>n</i> = 2074	<i>n</i> = 2074	<i>n</i> = 2074	<i>n</i> = 2074
TBI vs no Injury	1.60 (1.18 - 2.16)	1.56 (1.15 - 2.11)	1.51 (1.11 - 2.05)	1.23 (0.87 - 1.74)
OI vs no Injury	1.10 (0.88 - 1.37)	1.09 (0.87 - 1.37)	1.09 (0.87 - 1.37)	1.02 (0.79 - 1.33)
TBI vs OI	1.46 (1.04 - 2.03)	1.43 (1.02 - 2.00)	1.39 (0.99 - 1.95)	1.20 (0.82 - 1.77)
<i>Omnibus p</i>	0.200	0.236	0.254	0.718

complete cases had no missing data for the exposure, outcome or covariates. *TBI*: traumatic brain injury; *OI*: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 4 Associations between traumatic brain injury from birth to age 16 years, with no additional orthopaedic injury, and substance use at age 17 years compared to orthopaedic injury

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol^{a*}				
<i>n</i>	<i>n</i> = 3564	<i>n</i> = 3148	<i>n</i> = 2778	<i>n</i> = 1992
TBI only vs OI	1.48 (1.10 - 1.99)	1.44 (1.05 - 1.98)	1.57 (1.13 - 2.18)	1.98 (1.28 - 3.09)
<i>Omnibus p</i>	0.056	0.313	0.306	0.081
Tobacco^{b**}				
<i>n</i>	<i>n</i> = 2991	<i>n</i> = 2642	<i>n</i> = 2326	<i>n</i> = 1992
TBI only vs OI	1.14 (0.78 - 1.66)	1.12 (0.75 - 1.68)	1.04 (0.67 - 1.62)	0.96 (0.56 - 1.63)
<i>Omnibus p</i>	0.094	0.072	0.078	0.374
Cannabis^{c**}				
<i>n</i>	<i>n</i> = 3843	<i>n</i> = 3384	<i>n</i> = 2978	<i>n</i> = 1992
TBI only vs OI	1.20 (0.88 - 1.62)	1.09 (0.79 - 1.52)	1.08 (0.76 - 1.53)	1.14 (0.71 - 1.82)
<i>Omnibus p</i>	0.007	0.078	0.109	0.588

Sample size reduces per adjustment as the participants who are missing covariate data get excluded *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 5 Associations between traumatic brain injury from birth to age 16 years, with no additional orthopaedic injury, and substance use at age 17 years compared to orthopaedic injury on complete case sample

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol ^{a*}				
<i>n</i>	<i>n</i> = 1992	<i>n</i> = 1992	<i>n</i> = 1992	<i>n</i> = 1992
TBI only vs OI	1.81 (1.22 - 2.68)	1.82 (1.22 - 2.70)	1.81 (1.22 - 2.70)	1.98 (1.28 - 3.09)
Omnibus <i>p</i>	0.435	0.328	0.344	0.081
Tobacco ^{b**}				
<i>n</i>	<i>n</i> = 1992	<i>n</i> = 1992	<i>n</i> = 1992	<i>n</i> = 1992
TBI only vs OI	1.08 (0.68 - 1.71)	1.12 (0.71 - 1.78)	1.10 (0.69 - 1.76)	0.96 (0.56 - 1.63)
Omnibus <i>p</i>	0.446	0.324	0.302	0.374
Cannabis ^{c**}				
<i>n</i>	<i>n</i> = 1992	<i>n</i> = 1992	<i>n</i> = 1992	<i>n</i> = 1992
TBI only vs OI	1.24 (0.82 - 1.87)	1.23 (0.81 - 1.86)	1.22 (0.80 - 1.85)	1.14 (0.71 - 1.82)
Omnibus <i>p</i>	0.319	0.355	0.365	0.588

complete cases had no missing data for the exposure, outcome or covariates. *TBI*: traumatic brain injury; *OI*: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 6 Associations between traumatic brain injury and orthopaedic injuries from birth to age 16 years and criminal behaviours at age 17 years on complete case sample

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 2115	<i>n</i> = 2115	<i>n</i> = 2115	<i>n</i> = 2115
TBI vs no Injury	1.82 (1.30 - 2.54)	1.62 (1.15 - 2.28)	1.58 (1.13 - 2.23)	1.29 (0.09 - 1.88)
OI vs no Injury	1.71 (1.34 - 2.18)	1.57 (1.22 - 2.01)	1.56 (1.22 - 2.00)	1.67 (1.27 - 2.19)
TBI vs OI	1.06 (0.74 - 1.53)	1.03 (0.72 - 1.49)	1.01 (0.70 - 1.47)	0.77 (0.52 - 1.16)
Omnibus <i>p</i>	<0.001	<0.001	<0.001	<0.001
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 2077	<i>n</i> = 2077	<i>n</i> = 2077	<i>n</i> = 2077
TBI vs no Injury	1.73 (1.20 - 2.48)	1.52 (1.05 - 2.21)	1.49 (1.02 - 2.17)	1.17 (0.77 - 1.77)
OI vs no Injury	1.15 (0.87 - 1.53)	1.01 (0.75 - 1.35)	1.02 (0.76 - 1.36)	1.03 (0.75 - 1.42)
TBI vs OI	1.50 (1.00 - 2.25)	1.51 (1.00 - 2.30)	1.46 (0.96 - 2.23)	1.14 (0.71 - 1.81)
Omnibus <i>p</i>	0.158	0.707	0.678	0.765

Complete cases had no missing data for the exposure, outcome or covariates *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 7 Associations between traumatic brain injury from birth to age 16 years, with no additional orthopaedic injury, and criminal behaviour at age 17 years compared to orthopaedic injury

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 3719	<i>n</i> = 3283	<i>n</i> = 2886	<i>n</i> = 2031
TBI only vs OI	1.14 (0.81 - 1.60)	1.15 (0.80 - 1.66)	1.18 (0.81 - 1.73)	0.76 (0.46 - 1.26)
<i>Omnibus p</i>	<0.001	0.001	0.001	<0.001
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 3657	<i>n</i> = 3228	<i>n</i> = 2844	<i>n</i> = 1995
TBI only vs OI	0.94 (0.63 - 1.40)	0.86 (0.55 - 1.34)	0.90 (0.57 - 1.44)	0.80 (0.44 - 1.47)
<i>Omnibus p</i>	0.001	0.096	0.102	0.930

Sample size reduces per adjustment as the participants who are missing covariate data get excluded *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 8 Associations between traumatic brain injury from birth to age 16 years, with no additional orthopaedic injury, and criminal behaviour at age 17 years compared to orthopaedic injury on complete case sample

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 2031	<i>n</i> = 2031	<i>n</i> = 2031	<i>n</i> = 2031
TBI only vs OI	0.94 (0.60 - 1.46)	0.95 (0.60 - 1.49)	0.95 (0.60 - 1.50)	0.76 (0.46 - 1.26)
<i>Omnibus p</i>	<0.001	<0.001	<0.001	<0.001
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 1995	<i>n</i> = 1995	<i>n</i> = 1995	<i>n</i> = 1995
TBI only vs OI	1.02 (0.60 - 1.74)	1.03 (0.60 - 1.79)	1.02 (0.59 - 1.76)	0.80 (0.44 - 1.47)
<i>Omnibus p</i>	0.299	0.956	0.921	0.930

Complete cases had no missing data for the exposure, outcome or covariates *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 9 Associations between traumatic brain injury and orthopaedic injuries from birth to age 16 years and psychiatric symptoms based on the Strengths and Difficulties Questionnaire at age 17 years on complete case sample

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 4493	<i>n</i> = 4493	<i>n</i> = 4493
TBI vs no Injury	1.64 (1.11 – 2.43)	1.72 (1.16 – 2.55)	1.62 (1.08 – 2.41)
OI vs no Injury	1.08 (0.79 – 1.48)	1.10 (0.80 – 1.50)	1.07 (0.78 – 1.47)
TBI vs OI	1.52 (0.97 - 2.36)	1.57 (1.01 - 2.45)	1.51 (0.96 - 2.37)
<i>Omnibus p</i>	0.391	0.340	0.445
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 4483	<i>n</i> = 4483	<i>n</i> = 4483
TBI vs no Injury	0.92 (0.62 – 1.37)	0.88 (0.59 – 1.31)	0.85 (0.57 – 1.26)
OI vs no Injury	0.84 (0.64 – 1.11)	0.81 (0.61 – 1.07)	0.79 (0.60 – 1.05)
TBI vs OI	1.10 (0.71 - 1.71)	1.09 (0.70 - 1.69)	1.07 (0.68 - 1.67)
<i>Omnibus p</i>	0.206	0.127	0.090

Complete cases had no missing data for the exposure, outcome or covariates *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 10 Associations between traumatic brain injury from birth to age 16 years, with no additional orthopaedic injury, and psychiatric symptoms at age 17 years compared to orthopaedic injury

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 2437	<i>n</i> = 4818	<i>n</i> = 4328
TBI only vs OI	1.51 (0.96 - 2.38)	1.74 (1.07 - 2.84)	1.80 (1.08 - 3.00)
Omnibus <i>p</i>	0.201	0.285	0.458
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 5427	<i>n</i> = 4806	<i>n</i> = 4316
TBI only vs OI	1.17 (0.75 - 1.83)	1.17 (0.71 - 1.93)	1.15 (0.68 - 1.96)
Omnibus <i>p</i>	0.829	0.121	0.100

Sample size reduces per adjustment as the participants who are missing covariate data get excluded *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 11 Associations between traumatic brain injury from birth to age 16 years, with no additional orthopaedic injury, and psychiatric symptoms at age 17 years compared to orthopaedic injury on complete case sample

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 4328	<i>n</i> = 4328	<i>n</i> = 4328
TBI only vs OI	1.76 (1.06 - 2.91)	1.85 (1.12 - 3.07)	1.80 (1.08 - 3.00)
Omnibus <i>p</i>	0.412	0.366	0.458
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 4316	<i>n</i> = 4316	<i>n</i> = 4316
TBI only vs OI	1.15 (0.68 - 1.95)	1.16 (0.68 - 1.96)	1.15 (0.68 - 1.96)
Omnibus <i>p</i>	0.217	0.133	0.100

Complete cases had no missing data for the exposure, outcome or covariates *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 12 Association between traumatic brain injury and orthopaedic injuries from birth to age 11 years and substance use at age 17 years

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol^{a*}				
<i>n</i>	<i>n</i> = 3188	<i>n</i> = 2812	<i>n</i> = 2381	<i>n</i> = 1788
TBI vs no Injury	1.26 (0.89 - 1.78)	1.28 (0.89 - 1.85)	1.37 (0.93 - 2.02)	1.13 (0.68 - 1.88)
OI vs no Injury	1.16 (0.97 - 1.38)	1.10 (0.91 - 1.33)	1.13 (0.92 - 1.38)	0.81 (0.61 - 1.07)
TBI vs OI	1.09 (0.75 - 1.57)	1.16 (0.78 - 1.72)	1.22 (0.81 - 1.85)	1.40 (0.80 - 2.44)
Omnibus <i>p</i>	0.072	0.231	0.266	0.167
Tobacco^{b**}				
<i>n</i>	<i>n</i> = 2675	<i>n</i> = 2364	<i>n</i> = 2084	<i>n</i> = 1788
TBI vs no Injury	1.35 (0.88 - 2.08)	1.31 (0.83 - 2.07)	1.15 (0.69 - 1.92)	1.00 (0.53 - 1.87)
OI vs no Injury	1.01 (0.80 - 1.28)	1.02 (0.79 - 1.32)	1.00 (0.76 - 1.32)	0.83 (0.58 - 1.19)
TBI vs OI	1.34 (0.84 - 2.15)	1.28 (0.78 - 2.12)	1.16 (0.66 - 2.01)	1.20 (0.60 - 2.39)
Omnibus <i>p</i>	0.786	0.739	0.940	0.328
Cannabis^{c**}				
<i>n</i>	<i>n</i> = 3436	<i>n</i> = 3023	<i>n</i> = 2668	<i>n</i> = 1788
TBI vs no Injury	1.61 (1.14 - 2.28)	1.44 (0.99 - 2.08)	1.45 (0.98 - 2.15)	1.47 (0.88 - 2.47)
OI vs no Injury	1.17 (0.98 - 1.41)	1.12 (0.92 - 1.37)	1.10 (0.89 - 1.36)	1.03 (0.76 - 1.40)
TBI vs OI	1.38 (0.95 - 2.00)	1.28 (0.86 - 1.91)	1.32 (0.86 - 2.02)	1.43 (0.81 - 2.52)
Omnibus <i>p</i>	0.041	0.162	0.254	0.671

Sample size reduces per adjustment as the participants who are missing covariate data get excluded *TBI*: traumatic brain injury; *OI*: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 13 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and substance use at age 17 years compared to orthopaedic injury

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol ^{a*}				
<i>n</i>	<i>n</i> = 3144	<i>n</i> = 2776	<i>n</i> = 2454	<i>n</i> = 1762
TBI only vs OI	1.37 (0.89 - 2.11)	1.40 (0.89 - 2.20)	1.47 (0.91 - 2.37)	1.64 (0.87 - 3.13)
Omnibus <i>p</i>	0.059	0.207	0.243	0.181
Tobacco ^{b**}				
<i>n</i>	<i>n</i> = 2640	<i>n</i> = 2333	<i>n</i> = 2055	<i>n</i> = 1762
TBI only vs OI	1.30 (0.74 - 2.27)	1.31 (0.73 - 2.35)	1.16 (0.61 - 2.21)	0.98 (0.43 - 2.20)
Omnibus <i>p</i>	0.851	0.774	0.968	0.275
Cannabis ^{c**}				
<i>n</i>	<i>n</i> = 3391	<i>n</i> = 2986	<i>n</i> = 2635	<i>n</i> = 1762
TBI only vs OI	1.53 (1.00 - 2.36)	1.43 (0.91 - 2.25)	1.47 (0.91 - 2.37)	1.51 (0.79 - 2.87)
Omnibus <i>p</i>	0.044	0.164	0.259	0.214

Sample size reduces per adjustment as the participants who are missing covariate data get excluded *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 14 Associations between traumatic brain injury and orthopaedic injuries from birth to age 11 years and criminal behaviour at age 17 years

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 3325	<i>n</i> = 2931	<i>n</i> = 2584	<i>n</i> = 1818
TBI vs no Injury	1.26 (0.82 - 1.94)	1.15 (0.72 - 1.83)	1.25 (0.77 - 2.02)	0.97 (0.53 - 1.79)
OI vs no Injury	1.30 (1.05 - 1.62)	1.31 (1.03 - 1.66)	1.37 (1.06 - 1.75)	1.78 (1.30 - 2.45)
TBI vs OI	0.97 (0.61 - 1.53)	0.88 (0.53 - 1.44)	0.91 (0.54 - 1.53)	0.54 (0.28 - 1.04)
<i>Omnibus p</i>	0.013	0.025	0.012	0.001
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 3275	<i>n</i> = 2886	<i>n</i> = 2549	<i>n</i> = 1790
TBI vs no Injury	1.44 (0.90 - 2.28)	1.20 (0.73 - 1.98)	1.31 (0.78 - 2.21)	1.17 (0.62 - 2.22)
OI vs no Injury	1.34 (1.05 - 1.70)	1.28 (0.99 - 1.67)	1.27 (0.96 - 1.69)	1.15 (0.79 - 1.69)
TBI vs OI	1.06 (0.66 - 1.76)	0.94 (0.55 - 1.60)	1.03 (0.59 - 1.81)	1.02 (0.50 - 2.06)
<i>Omnibus p</i>	0.012	0.056	0.078	0.432

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 15 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and criminal behaviour at age 17 years compared to orthopaedic injury

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 3285	<i>n</i> = 2898	<i>n</i> = 2555	<i>n</i> = 1792
TBI only vs OI	1.08 (0.64 - 1.83)	1.00 (0.58 - 1.75)	1.07 (0.60 - 1.91)	0.65 (0.31 - 1.34)
<i>Omnibus p</i>	0.012	0.021	0.010	<0.001
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 3235	<i>n</i> = 2852	<i>n</i> = 2519	<i>n</i> = 1764
TBI only vs OI	1.14 (0.65 - 2.00)	0.98 (0.53 - 1.79)	1.08 (0.57 - 2.05)	1.04 (0.47 - 2.29)
<i>Omnibus p</i>	0.012	0.055	0.080	0.434

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 16 Associations between traumatic brain injury and orthopaedic injuries from birth to age 11 years and psychiatric symptoms based on the Strengths and Difficulties Questionnaire at age 17 years

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 4923	<i>n</i> = 4372	<i>n</i> = 3937
TBI vs no Injury	2.20 (1.37 – 3.53)	2.33 (1.41 – 3.85)	1.90 (1.11 - 3.26)
OI vs no Injury	1.05 (0.76 – 1.44)	0.99 (0.69 – 1.41)	0.96 (0.66 - 1.39)
TBI vs OI	2.10 (1.23 - 3.57)	2.35 (1.33 - 4.17)	1.98 (1.08 - 3.65)
<i>Omnibus p</i>	0.442	0.656	0.884
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 4912	<i>n</i> = 4359	<i>n</i> = 3924
TBI vs no Injury	1.51 (0.95 – 2.39)	1.30 (0.79 – 2.13)	0.99 (0.57 - 1.21)
OI vs no Injury	1.01 (0.77 – 1.31)	0.91 (0.68 – 1.22)	0.89 (0.65 - 1.21)
TBI vs OI	1.51 (0.91 - 2.49)	1.42 (0.83 - 2.45)	1.12 (0.61 - 2.05)
<i>Omnibus p</i>	0.780	0.666	0.456

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 17 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and psychiatric symptoms at age 17 years compared to orthopaedic injury

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 4854	<i>n</i> = 4310	<i>n</i> = 3880
TBI only vs OI	2.65 (1.48 - 4.74)	3.04 (1.64 - 5.64)	2.77 (1.45 - 5.31)
Omnibus <i>p</i>	0.450	0.651	0.826
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 4842	<i>n</i> = 4296	<i>n</i> = 3866
TBI only vs OI	1.58 (0.88 - 2.84)	1.47 (0.78 - 2.77)	1.28 (0.64 - 2.54)
Omnibus <i>p</i>	0.828	0.639	0.481

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 18 Associations between traumatic brain injury and orthopaedic injuries from birth to age 11 years and psychiatric symptoms based on the Development and Well-Being Assessment (DAWBA) at age 15 years

DAWBA	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Externalising Behaviour ^{a*}			
<i>n</i>	<i>n</i> = 3994	<i>n</i> = 3515	<i>n</i> = 3112
TBI vs no Injury	2.25 (1.32 – 3.81)	2.35 (1.35 – 4.11)	1.83 (0.98 - 3.41)
OI vs no Injury	0.85 (0.57 – 1.27)	0.90 (0.59 – 1.39)	0.87 (0.55 - 1.38)
TBI vs OI	2.65 (1.43 - 4.91)	2.61 (1.36 - 4.98)	2.11 (1.03 - 4.32)
<i>Omnibus p</i>	0.779	0.970	0.796
ODD ^{b**}			
<i>n</i>	<i>n</i> = 3983	<i>n</i> = 3506	<i>n</i> = 3105
TBI vs no Injury	2.28 (1.28 – 4.07)	2.42 (1.31 – 4.55)	1.78 (0.89 - 3.58)
OI vs no Injury	0.81 (0.51 – 1.28)	0.98 (0.61 – 1.56)	0.98 (0.60 - 1.61)
TBI vs OI	2.82 (1.42 - 5.59)	2.48 (1.22 - 5.02)	1.82 (0.82 - 4.01)
<i>Omnibus p</i>	0.673	0.730	0.851
CD ^{c**}			
<i>n</i>	<i>n</i> = 3982	<i>n</i> = 3505	<i>n</i> = 3104
TBI vs no Injury	1.09 (0.34 – 3.55)	0.83 (0.20 – 3.53)	0.72 (0.17 - 3.10)
OI vs no Injury	0.66 (0.31 - 1.40)	0.78 (0.36 – 1.70)	0.85 (0.39 - 1.86)
TBI vs OI	1.66 (0.44 - 6.33)	1.06 (0.22 - 5.10)	0.94 (0.17 - 4.13)
<i>Omnibus p</i>	0.301	0.523	0.636
ADHD ^{d**}			
<i>n</i>	<i>n</i> = 3994	<i>n</i> = 3515	<i>n</i> = 3112
TBI vs no Injury	3.15 (1.07 – 9.28)	2.86 (0.94 – 8.70)	3.02 (0.97 - 9.36)
OI vs no Injury	1.42 (0.62 - 3.21)	0.91 (0.34 – 2.47)	0.57 (0.16 - 1.98)
TBI vs OI	2.23 (0.66 - 7.48)	3.14 (0.82 - 12.02)	5.28 (1.14 - 24.51)
<i>Omnibus p</i>	0.274	0.891	0.623

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI*: traumatic brain injury; *OI*: orthopaedic injury; * logistic regression; ^a externalising disorder symptoms based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years;

^b ODD: oppositional defiant disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^c CD: conduct disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^d ADHD: attentional defiant hyperactivity disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Table 19 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and psychiatric symptoms based on the Development and Well-Being Assessment (DAWBA) at age 15 years compared to orthopaedic injury

DAWBA	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Externalising Behaviour ^{a*}			
<i>n</i>	<i>n</i> = 3931	<i>n</i> = 3460	<i>n</i> = 3063
TBI only vs OI	2.34 (1.11 - 4.92)	2.20 (1.00 - 4.83)	1.99 (0.85 - 4.64)
Omnibus <i>p</i>	0.609	0.833	0.698
ODD ^{b**}			
<i>n</i>	<i>n</i> = 3920	<i>n</i> = 3451	<i>n</i> = 3056
TBI only vs OI	2.76 (1.25 - 6.12)	2.29 (0.99 - 5.28)	1.94 (0.79 - 4.79)
Omnibus <i>p</i>	0.553	0.876	0.901
CD ^{c**}			
<i>n</i>	<i>n</i> = 3919	<i>n</i> = 3450	<i>n</i> = 3055
TBI only vs OI	1.69 (0.35 - 8.06)	0.83 (0.10 - 6.77)	0.67 (0.08 - 5.60)
Omnibus <i>p</i>	0.294	0.512	0.652
ADHD			
<i>n</i>	<i>n</i> = 3931	<i>n</i> = 3460	<i>n</i> = 3063
TBI only vs OI	1.68 (0.35 - 8.03)	2.53 (0.48 - 13.39)	4.03 (0.65 - 25.14)
Omnibus <i>p</i>	0.345	0.990	0.514

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI*: traumatic brain injury; *OI*: orthopaedic injury; ^a logistic regression; ^a externalising disorder symptoms based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^b ODD: oppositional defiant disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^c CD: conduct disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^d ADHD: attentional defiant hyperactivity disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 20 Association between traumatic brain injury and orthopaedic injuries from birth to age 11 years and substance use at age 17 years on complete case sample

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol ^{a*}				
<i>n</i>	<i>n</i> = 1788	<i>n</i> = 1788	<i>n</i> = 1788	<i>n</i> = 1788
TBI vs no Injury	1.25 (0.80 - 1.96)	1.24 (0.79 - 1.94)	1.18 (0.75 - 1.86)	1.13 (0.68 - 1.88)
OI vs no Injury	0.88 (0.69 - 1.13)	0.88 (0.69 - 1.12)	0.87 (0.68 - 1.12)	0.81 (0.61 - 1.07)
TBI vs OI	1.42 (0.87 - 2.31)	1.41 (0.86 - 2.30)	1.35 (0.82 - 2.21)	1.40 (0.80 - 2.44)
<i>Omnibus p</i>	0.429	0.388	0.360	0.167
Tobacco ^{b**}				
<i>n</i>	<i>n</i> = 1788	<i>n</i> = 1788	<i>n</i> = 1788	<i>n</i> = 1788
TBI vs no Injury	1.09 (0.64 - 1.87)	1.08 (0.63 - 1.85)	0.99 (0.57 - 1.72)	1.00 (0.53 - 1.87)
OI vs no Injury	0.90 (0.66 - 1.21)	0.91 (0.68 - 1.24)	0.91 (0.67 - 1.23)	0.83 (0.58 - 1.19)
TBI vs OI	1.22 (0.68 - 2.19)	1.18 (0.65 - 2.13)	1.09 (0.60 - 2.00)	1.20 (0.60 - 2.39)
<i>Omnibus p</i>	0.518	0.596	0.532	0.328
Cannabis ^{c**}				
<i>n</i>	<i>n</i> = 1788	<i>n</i> = 1788	<i>n</i> = 1788	<i>n</i> = 1788
TBI vs no Injury	1.45 (0.92 - 2.29)	1.47 (0.92 - 2.32)	1.37 (0.86 - 2.18)	1.47 (0.88 - 2.47)
OI vs no Injury	1.03 (0.79 - 1.33)	1.05 (0.80 - 1.36)	1.04 (0.80 - 1.36)	1.03 (0.76 - 1.40)
TBI vs OI	1.41 (0.86 - 2.33)	1.40 (0.85 - 2.32)	1.32 (0.79 - 2.19)	1.43 (0.81 - 2.52)
<i>Omnibus p</i>	0.659	0.557	0.615	0.671

Complete cases had no missing data for the exposure, outcome or covariates. TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 21 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and substance use at age 17 years compared to orthopaedic injury on complete case sample

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol ^{a*}				
<i>n</i>	<i>n</i> = 1762	<i>n</i> = 1762	<i>n</i> = 1762	<i>n</i> = 1762
TBI only vs OI	1.72 (0.98 - 3.04)	1.70 (0.96 - 3.02)	1.61 (0.91 - 2.87)	1.64 (0.87 - 3.13)
Omnibus <i>p</i>	0.453	0.412	0.383	0.181
Tobacco ^{b**}				
<i>n</i>	<i>n</i> = 1762	<i>n</i> = 1762	<i>n</i> = 1762	<i>n</i> = 1762
TBI only vs OI	1.18 (0.59 - 2.35)	1.17 (0.58 - 2.34)	1.07 (0.53 - 2.17)	0.98 (0.43 - 2.20)
Omnibus <i>p</i>	0.495	0.577	0.511	0.275
Cannabis ^{c**}				
<i>n</i>	<i>n</i> = 1762	<i>n</i> = 1762	<i>n</i> = 1762	<i>n</i> = 1762
TBI only vs OI	1.58 (0.89 - 2.81)	1.58 (0.88 - 2.81)	1.47 (0.82 - 2.63)	1.51 (0.79 - 2.87)
Omnibus <i>p</i>	0.672	0.568	0.622	0.214

Complete cases had no missing data for the exposure, outcome or covariates. TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 22 Associations between traumatic brain injury and orthopaedic injuries from birth to age 11 years and criminal behaviour at age 17 years on complete case sample

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 1818	<i>n</i> = 1818	<i>n</i> = 1818	<i>n</i> = 1818
TBI vs no Injury	1.28 (0.74 - 2.21)	1.12 (0.64 - 1.96)	1.06 (0.60 - 1.86)	0.97 (0.53 - 1.79)
OI vs no Injury	1.57 (1.18 - 2.08)	1.57 (1.18 - 2.10)	1.57 (1.17 - 2.09)	1.78 (1.30 - 2.45)
TBI vs OI	0.82 (0.46 - 1.46)	0.71 (0.39 - 1.29)	0.68 (0.37 - 1.23)	0.54 (0.28 - 1.04)
<i>Omnibus p</i>	0.002	0.002	0.003	0.001
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 1790	<i>n</i> = 1790	<i>n</i> = 1790	<i>n</i> = 1790
TBI vs no Injury	1.50 (0.85 - 2.63)	1.24 (0.69 - 2.22)	1.17 (0.65 - 2.11)	1.17 (0.62 - 2.22)
OI vs no Injury	1.11 (0.79 - 1.54)	1.06 (0.75 - 1.50)	1.07 (0.76 - 1.51)	1.15 (0.79 - 1.69)
TBI vs OI	1.35 (0.73 - 2.51)	1.16 (0.61 - 2.21)	1.09 (0.57 - 2.08)	1.02 (0.50 - 2.06)
<i>Omnibus p</i>	0.422	0.654	0.635	0.432

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 23 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and criminal behaviour at age 17 years compared to orthopaedic injury on complete case sample

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 1792	<i>n</i> = 1792	<i>n</i> = 1792	<i>n</i> = 1792
TBI only vs OI	0.95 (0.49 - 1.84)	0.83 (0.42 - 1.62)	0.80 (0.40 - 1.57)	0.65 (0.31 - 1.34)
<i>Omnibus p</i>	0.001	0.002	0.002	<0.001
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 1764	<i>n</i> = 1764	<i>n</i> = 1764	<i>n</i> = 1764
TBI only vs OI	1.48 (0.74 - 2.99)	1.25 (0.61 - 2.58)	1.16 (0.56 - 2.42)	1.04 (0.47 - 2.29)
<i>Omnibus p</i>	0.431	0.650	0.634	0.434

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 24 Associations between traumatic brain injury and orthopaedic injuries from birth to age 11 years and psychiatric symptoms based on the Strengths and Difficulties Questionnaire at age 17 years on complete case sample

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 3937	<i>n</i> = 3937	<i>n</i> = 3937
TBI vs no Injury	2.14 (1.26 – 3.63)	2.17 (1.28 - 3.70)	1.90 (1.11 - 3.26)
OI vs no Injury	0.98 (0.68 – 1.41)	0.99 (0.68 – 1.43)	0.96 (0.66 - 1.39)
TBI vs OI	2.19 (1.20 - 3.98)	2.20 (1.21 - 4.01)	1.98 (1.08 - 3.65)
<i>Omnibus p</i>	0.752	0.708	0.884
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 3924	<i>n</i> = 3924	<i>n</i> = 3924
TBI vs no Injury	1.17 (0.68 - 2.02)	1.10 (0.63 - 1.90)	0.99 (0.57 - 1.21)
OI vs no Injury	0.92 (0.68 - 1.25)	0.91 (0.67 - 1.24)	0.89 (0.65 - 1.21)
TBI vs OI	1.26 (0.70 - 2.29)	1.20 (0.66 - 2.19)	1.12 (0.61 - 2.05)
<i>Omnibus p</i>	0.684	0.605	0.456

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 25 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and psychiatric symptoms at age 17 years compared to orthopaedic injury on complete case sample

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 3880	<i>n</i> = 3880	<i>n</i> = 3880
TBI only vs OI	3.02 (1.60 - 5.69)	3.05 (1.61 - 5.76)	2.77 (1.45 - 5.31)
Omnibus <i>p</i>	0.716	0.667	0.826
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 3866	<i>n</i> = 3866	<i>n</i> = 3866
TBI only vs OI	1.42 (0.72 - 2.80)	1.36 (0.69 - 2.69)	1.28 (0.64 - 2.54)
Omnibus <i>p</i>	0.695	0.623	0.481

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 26 Associations between traumatic brain injury and orthopaedic injuries from birth to age 11 years and psychiatric symptoms based on the Development and Well-Being Assessment (DAWBA) at age 15 years on complete case sample

DAWBA	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Externalising Behaviour ^{a*}			
<i>n</i>	<i>n</i> = 3112	<i>n</i> = 3112	<i>n</i> = 3112
TBI vs no Injury	2.19 (1.20 - 4.00)	2.11 (1.15 - 3.88)	1.83 (0.98 - 3.41)
OI vs no Injury	0.88 (0.56 - 1.39)	0.88 (0.55 - 1.38)	0.87 (0.55 - 1.38)
TBI vs OI	2.49 (1.24 - 5.01)	2.41 (1.19 - 4.87)	2.11 (1.03 - 4.32)
<i>Omnibus p</i>	0.894	0.866	0.796
ODD ^{b**}			
<i>n</i>	<i>n</i> = 3105	<i>n</i> = 3105	<i>n</i> = 3105
TBI vs no Injury	2.11 (1.07 - 4.17)	2.04 (1.03 - 4.04)	1.78 (0.89 - 3.58)
OI vs no Injury	0.98 (0.60 - 1.60)	0.98 (0.60 - 1.60)	0.98 (0.60 - 1.61)
TBI vs OI	2.15 (0.99 - 4.67)	2.08 (0.96 - 4.54)	1.82 (0.82 - 4.01)
<i>Omnibus p</i>	0.797	0.820	0.851
CD ^{c**}			
<i>n</i>	<i>n</i> = 3104	<i>n</i> = 3104	<i>n</i> = 3104
TBI vs no Injury	0.91 (0.22 - 3.82)	0.90 (0.21 - 3.80)	0.72 (0.17 - 3.10)
OI vs no Injury	0.83 (0.38 - 1.81)	0.83 (0.38 - 1.81)	0.85 (0.39 - 1.86)
TBI vs OI	1.09 (0.23 - 5.19)	1.08 (0.23 - 5.17)	0.94 (0.17 - 4.13)
<i>Omnibus p</i>	0.641	0.636	0.636
ADHD ^{d**}			
<i>n</i>	<i>n</i> = 3112	<i>n</i> = 3112	<i>n</i> = 3112
TBI vs no Injury	3.69 (1.23 - 11.12)	3.39 (1.11 - 10.37)	3.02 (0.97 - 9.36)
OI vs no Injury	0.62 (0.18 - 2.13)	0.60 (0.18 - 2.07)	0.57 (0.16 - 1.98)
TBI vs OI	5.94 (1.31 - 26.81)	5.62 (1.23 - 25.65)	5.28 (1.14 - 24.51)
<i>Omnibus p</i>	0.758	0.702	0.623

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; * logistic regression; ^a externalising disorder symptoms based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^b ODD: oppositional defiant disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^c CD: conduct disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^d ADHD: attentional defiant hyperactivity disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 27 Associations between traumatic brain injury from birth to age 11 years, with no additional orthopaedic injury, and psychiatric symptoms based on the Development and Well-Being Assessment (DAWBA) at age 15 years compared to orthopaedic injury on complete case sample

DAWBA	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Externalising Behaviour ^{a*}			
<i>n</i>	<i>n</i> = 3063	<i>n</i> = 3063	<i>n</i> = 3063
TBI only vs OI	2.25 (0.98 - 5.17)	2.31 (1.01 - 5.32)	1.99 (0.85 - 4.64)
Omnibus <i>p</i>	0.759	0.747	0.698
ODD ^{b**}			
<i>n</i>	<i>n</i> = 3056	<i>n</i> = 3056	<i>n</i> = 3056
TBI only vs OI	2.25 (0.93 - 5.44)	2.28 (0.94 - 5.52)	1.94 (0.79 - 4.79)
Omnibus <i>p</i>	0.863	0.880	0.901
CD ^{c**}			
<i>n</i>	<i>n</i> = 3055	<i>n</i> = 3055	<i>n</i> = 3055
TBI only vs OI	0.81 (0.10 - 6.54)	0.88 (0.11 - 7.12)	0.67 (0.08 - 5.60)
Omnibus <i>p</i>	0.617	0.624	0.652
ADHD			
<i>n</i>	<i>n</i> = 3063	<i>n</i> = 3063	<i>n</i> = 3063
TBI only vs OI	4.38 (0.72 - 26.52)	4.77 (0.78 - 29.23)	4.03 (0.65 - 25.14)
Omnibus <i>p</i>	0.585	0.565	0.514

Complete cases had no missing data for the exposure, outcome or covariates. *TBI*: traumatic brain injury; *OI*: orthopaedic injury; ^a logistic regression; ^a externalising disorder symptoms based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^b ODD: oppositional defiant disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^c CD: conduct disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years; ^d ADHD: attentional defiant hyperactivity disorder based on the Development and Well-being Assessment (DAWBA) self-reported at age 15 years

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 28 Associations between traumatic brain injury and orthopaedic injuries from age 12 to age 16 years and substance use at 17 years

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol ^{a*}				
<i>n</i>	<i>n</i> = 2926	<i>n</i> = 2580	<i>n</i> = 2263	<i>n</i> = 1649
TBI vs no Injury	1.71 (1.28 - 2.27)	1.59 (1.17 - 2.15)	1.72 (1.25 - 2.37)	1.41 (0.93 - 2.15)
OI vs no Injury	1.06 (0.83 - 1.35)	0.98 (0.75 - 1.27)	0.98 (0.74 - 1.31)	0.70 (0.47 - 1.04)
TBI vs OI	1.61 (1.13 - 2.31)	1.62 (1.11 - 2.38)	1.76 (1.17 - 2.63)	2.03 (1.17 - 3.53)
Omnibus <i>p</i>	0.116	0.432	0.319	0.282
Tobacco ^{b**}				
<i>n</i>	<i>n</i> = 2488	<i>n</i> = 2193	<i>n</i> = 1923	<i>n</i> = 1649
TBI vs no Injury	1.56 (1.11 - 2.19)	1.67 (1.16 - 2.41)	1.71 (1.15 - 2.52)	1.15 (0.72 - 1.86)
OI vs no Injury	1.50 (1.13 - 2.00)	1.61 (1.17 - 2.21)	1.76 (1.25 - 2.48)	2.00 (1.29 - 3.09)
TBI vs OI	1.04 (0.68 - 1.58)	1.04 (0.66 - 1.63)	0.97 (0.60 - 1.57)	0.58 (0.32 - 1.06)
Omnibus <i>p</i>	0.001	0.000	0.000	0.003
Cannabis ^{c**}				
<i>n</i>	<i>n</i> = 3172	<i>n</i> = 2788	<i>n</i> = 2439	<i>n</i> = 1649
TBI vs no Injury	1.49 (1.11 - 1.99)	1.32 (0.97 - 1.81)	1.36 (0.98 - 1.88)	1.14 (0.74 - 1.76)
OI vs no Injury	1.32 (1.03 - 1.68)	1.23 (0.95 - 1.60)	1.28 (0.96 - 1.72)	1.04 (0.69 - 1.58)
TBI vs OI	1.13 (0.79 - 1.62)	1.08 (0.73 - 1.58)	1.06 (0.70 - 1.60)	1.09 (0.62 - 1.92)
Omnibus <i>p</i>	0.004	0.051	0.034	0.702

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 29 Associations between traumatic brain injury from age 12 to age 16 years, with no additional orthopaedic injury, and substance use at age 17 years compared to orthopaedic injury

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol ^{a*}				
<i>n</i>	<i>n</i> = 2890	<i>n</i> = 2548	<i>n</i> = 2232	<i>n</i> = 1625
TBI only vs OI	1.59 (1.09 - 2.32)	1.58 (1.06 - 2.37)	1.76 (1.14 - 2.71)	2.41 (1.35 - 4.29)
Omnibus <i>p</i>	0.172	0.540	0.403	0.364
Tobacco ^{b**}				
<i>n</i>	<i>n</i> = 2455	<i>n</i> = 2164	<i>n</i> = 1895	<i>n</i> = 1625
TBI only vs OI	0.95 (0.60 - 1.49)	0.89 (0.54 - 1.47)	0.82 (0.48 - 1.40)	0.59 (0.31 - 1.13)
Omnibus <i>p</i>	0.002	0.001	<0.001	0.002
Cannabis ^{c**}				
<i>N</i>	<i>n</i> = 3135	<i>n</i> = 2755	<i>n</i> = 2407	<i>n</i> = 1625
TBI only vs OI	1.06 (0.72 - 1.56)	0.997 (0.66 - 1.51)	0.97 (0.62 - 1.51)	1.12 (0.61 - 2.03)
Omnibus <i>p</i>	0.008	0.070	0.051	0.714

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 30 Associations between traumatic brain injury and orthopaedic injuries from age 12 to 16 years and criminal behaviour at age 17 years

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 3079	<i>n</i> = 2718	<i>n</i> = 2372	<i>n</i> = 1681
TBI vs no Injury	2.05 (1.50 - 2.80)	1.88 (1.34 - 2.63)	1.99 (1.40 - 2.82)	1.52 (0.97 - 2.39)
OI vs no Injury	1.89 (1.44 - 2.45)	1.47 (1.09 - 1.97)	1.53 (1.11 - 2.11)	1.49 (0.99 - 2.24)
TBI vs OI	1.09 (0.74 - 1.60)	1.28 (0.85 - 1.94)	1.30 (0.83 - 2.01)	1.02 (0.58 - 1.79)
<i>Omnibus p</i>	<0.001	0.001	<0.001	0.022
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 3024	<i>n</i> = 2668	<i>n</i> = 2339	<i>n</i> = 1651
TBI vs no Injury	1.74 (1.22 - 2.48)	1.43 (0.96 - 2.13)	1.51 (1.00 - 2.29)	1.21 (0.72 - 2.03)
OI vs no Injury	1.59 (1.17 - 2.17)	1.09 (0.77 - 1.54)	1.12 (0.77 - 1.64)	0.86 (0.52 - 1.42)
TBI vs OI	1.09 (0.70 - 1.70)	1.31 (0.80 - 2.15)	1.35 (0.80 - 2.27)	1.41 (0.72 - 2.77)
<i>Omnibus p</i>	<0.001	0.360	0.252	0.756

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 31 Associations between traumatic brain injury from age 12 to age 16 years, with no additional orthopaedic injury, and criminal behaviour at age 17 years compared to orthopaedic injury

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 3043	<i>n</i> = 2682	<i>n</i> = 2341	<i>n</i> = 1657
TBI only vs OI	0.94 (0.62 - 1.42)	1.09 (0.70 - 1.71)	1.09 (0.67 - 1.75)	0.85 (0.46 - 1.58)
<i>Omnibus p</i>	<0.001	0.002	0.002	0.041
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 2988	<i>n</i> = 2636	<i>n</i> = 2308	<i>n</i> = 1627
TBI only vs OI	0.80 (0.48 - 1.31)	0.90 (0.51 - 1.58)	0.90 (0.50 - 1.64)	0.92 (0.43 - 1.99)
<i>Omnibus p</i>	0.002	0.662	0.538	0.435

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 32 Associations between traumatic brain injury and orthopaedic injuries from age 12 to 16 years and psychiatric symptoms based on the Strengths and Difficulties Questionnaire at age 17 years

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 4460	<i>n</i> = 3939	<i>n</i> = 3536
TBI vs no Injury	1.21 (0.74 – 1.96)	1.41 (0.84 – 2.38)	1.39 (0.81 - 2.39)
OI vs no Injury	1.39 (0.92 - 2.12)	1.46 (0.92 - 2.31)	1.37 (0.84 - 2.24)
TBI vs OI	0.87 (0.47 - 1.59)	0.97 (0.50 - 1.85)	1.02 (0.51 - 2.02)
<i>Omnibus p</i>	0.095	0.058	0.126
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 4452	<i>n</i> = 3929	<i>n</i> = 3526
TBI vs no Injury	0.87 (0.55 – 1.38)	0.75 (0.45 – 1.25)	0.72 (0.42 - 1.24)
OI vs no Injury	0.87 (0.57 - 1.32)	0.56 (0.34 - 0.93)	0.55 (0.32 - 0.95)
TBI vs OI	1.00 (0.55 - 1.81)	1.35 (0.67 - 2.69)	1.30 (0.62 - 2.74)
<i>Omnibus p</i>	0.421	0.014	0.018

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 33 Associations between traumatic brain injury from age 12 to age 16 years, with no additional orthopaedic injury, and psychiatric symptoms at age 17 years compared to orthopaedic injury

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 4404	<i>n</i> = 3890	<i>n</i> = 3490
TBI only vs OI	1.01 (0.54 - 1.87)	1.13 (0.58 - 2.19)	1.18 (0.58 - 2.38)
Omnibus <i>p</i>	0.065	0.042	0.096
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 4396	<i>n</i> = 3880	<i>n</i> = 3480
TBI only vs OI	1.17 (0.64 - 2.14)	1.58 (0.78 - 3.21)	1.51 (0.71 - 3.22)
Omnibus <i>p</i>	0.560	0.024	0.029

Sample size reduces per adjustment as the participants who are missing covariate data get excluded. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 34 Associations between traumatic brain injury and orthopaedic injuries from age 12 to age 16 years and substance use at 17 years on complete case sample

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol^{a*}				
<i>n</i>	<i>n</i> = 1649	<i>n</i> = 1649	<i>n</i> = 1649	<i>n</i> = 1649
TBI vs no Injury	1.76 (1.22 - 2.54)	1.72 (1.19 - 2.49)	1.71 (0.60 - 1.20)	1.41 (0.93 - 2.15)
OI vs no Injury	0.88 (0.63 - 1.24)	0.84 (0.59 - 1.18)	0.85 (0.60 - 1.20)	0.70 (0.47 - 1.04)
TBI vs OI	2.00 (1.24 - 3.21)	2.05 (1.27 - 3.31)	2.02 (1.24 - 3.27)	2.03 (1.17 - 3.53)
<i>Omnibus p</i>	0.721	0.964	0.915	0.282
Tobacco^{b**}				
<i>n</i>	<i>n</i> = 1649	<i>n</i> = 1649	<i>n</i> = 1649	<i>n</i> = 1649
TBI vs no Injury	1.55 (1.04 - 2.31)	1.70 (1.13 - 2.55)	1.69 (1.12 - 2.55)	1.15 (0.72 - 1.86)
OI vs no Injury	1.55 (1.08 - 2.22)	1.60 (1.10 - 2.31)	1.68 (1.15 - 2.44)	2.00 (1.29 - 3.09)
TBI vs OI	1.00 (0.60 - 1.65)	1.06 (0.64 - 1.77)	1.01 (0.60 - 1.69)	0.58 (0.32 - 1.06)
<i>Omnibus p</i>	0.004	0.002	0.001	0.003
Cannabis^{c**}				
<i>n</i>	<i>n</i> = 1649	<i>n</i> = 1649	<i>n</i> = 1649	<i>n</i> = 1649
TBI vs no Injury	1.71 (1.17 - 2.48)	1.64 (1.12 - 2.39)	1.64 (1.12 - 2.41)	1.14 (0.74 - 1.76)
OI vs no Injury	1.26 (0.89 - 1.78)	1.18 (0.83 - 1.68)	1.18 (0.83 - 1.69)	1.04 (0.69 - 1.58)
TBI vs OI	1.35 (0.84 - 2.19)	1.39 (0.85 - 2.26)	1.39 (0.85 - 2.27)	1.09 (0.62 - 1.92)
<i>Omnibus p</i>	0.035	0.094	0.092	0.702

Complete cases had no missing data for the exposure, outcome or covariates. TBI: traumatic brain injury; OI: orthopaedic injury; * logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 35 Associations between traumatic brain injury from age 12 to age 16 years, with no additional orthopaedic injury, and substance use at age 17 years compared to orthopaedic injury on complete case sample

Substance Use	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Alcohol ^{a*}				
<i>n</i>	<i>n</i> = 1625	<i>n</i> = 1625	<i>n</i> = 1625	<i>n</i> = 1625
TBI only vs OI	2.07 (1.25 - 3.43)	2.13 (1.28 - 3.55)	2.09 (1.25 - 3.49)	2.41 (1.35 - 4.29)
Omnibus <i>p</i>	0.823	0.952	0.997	0.364
Tobacco ^{b**}				
<i>n</i>	<i>n</i> = 1625	<i>n</i> = 1625	<i>n</i> = 1625	<i>n</i> = 1625
TBI only vs OI	0.83 (0.47 - 1.44)	0.89 (0.51 - 1.56)	0.84 (0.48 - 1.49)	0.59 (0.31 - 1.13)
Omnibus <i>p</i>	0.011	0.006	0.003	0.002
Cannabis ^{c**}				
<i>n</i>	<i>n</i> = 1625	<i>n</i> = 1625	<i>n</i> = 1625	<i>n</i> = 1625
TBI only vs OI	1.16 (0.69 - 1.95)	1.21 (0.72 - 2.05)	1.21 (0.71 - 2.06)	1.12 (0.61 - 2.03)
Omnibus <i>p</i>	0.081	0.164	0.163	0.714

Complete cases had no missing data for the exposure, outcome or covariates. TBI: traumatic brain injury; OI: orthopaedic injury; * logistic regression; **generalised ordinal regression; ^a alcohol measured using the Alcohol Use Disorder Identification Test (AUDIT); ^b tobacco measured using the Fagerström Test for Nicotine Dependence; ^c cannabis measured using the Cannabis Abuse Screening Test.

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use and crime variables

Supplementary Table 36 Associations between traumatic brain injury and orthopaedic injuries from age 12 to 16 years and criminal behaviour at age 17 years on complete case sample

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 1681	<i>n</i> = 1681	<i>n</i> = 1681	<i>n</i> = 1681
TBI vs no Injury	2.22 (1.49 - 3.30)	2.02 (1.35 - 3.02)	2.01 (1.34 - 3.01)	1.52 (0.97 - 2.39)
OI vs no Injury	2.01 (1.40 - 2.88)	1.59 (1.10 - 2.30)	1.59 (1.10 - 2.30)	1.49 (0.99 - 2.24)
TBI vs OI	1.10 (0.67 - 1.81)	1.27 (0.76 - 2.10)	1.27 (0.76 - 2.11)	1.02 (0.58 - 1.79)
<i>Omnibus p</i>	<0.001	0.001	0.001	0.022
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 1651	<i>n</i> = 1651	<i>n</i> = 1651	<i>n</i> = 1651
TBI vs no Injury	1.89 (1.22 - 2.94)	1.75 (1.11 - 2.76)	1.74 (1.10 - 2.76)	1.21 (0.72 - 2.03)
OI vs no Injury	1.26 (0.81 - 1.95)	0.90 (0.57 - 1.42)	0.91 (0.57 - 1.43)	0.86 (0.52 - 1.42)
TBI vs OI	1.51 (0.84 - 2.70)	1.94 (1.06 - 3.55)	1.92 (1.05 - 3.53)	1.41 (0.72 - 2.77)
<i>Omnibus p</i>	0.063	0.751	0.728	0.756

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 37 Associations between traumatic brain injury from age 12 to age 16 years, with no additional orthopaedic injury, and criminal behaviour at age 17 years compared to orthopaedic injury on complete case sample

Criminal Behaviour	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)
Offences ^{a**}				
<i>n</i>	<i>n</i> = 1657	<i>n</i> = 1657	<i>n</i> = 1657	<i>n</i> = 1657
TBI only vs OI	0.87 (0.50 - 1.50)	1.03 (0.59 - 1.79)	1.03 (0.59 - 1.80)	0.85 (0.46 - 1.58)
<i>Omnibus p</i>	<0.001	0.004	0.004	0.041
Trouble with the Police ^{b**}				
<i>n</i>	<i>n</i> = 1627	<i>n</i> = 1627	<i>n</i> = 1627	<i>n</i> = 1627
TBI only vs OI	0.91 (0.47 - 1.79)	1.20 (0.60 - 2.41)	1.19 (0.59 - 2.38)	0.92 (0.43 - 1.99)
<i>Omnibus p</i>	0.277	0.744	0.767	0.435

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; **generalised ordinal regression; ^a offences measured by self-report questionnaire at age 17 years; ^b trouble with the police measured by self-report questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Model 3: As Model 2 with additional adjustment for substance use variables

Supplementary Table 38 Associations between traumatic brain injury and orthopaedic injuries from age 12 to 16 years and psychiatric symptoms based on the Strengths and Difficulties Questionnaire at age 17 years on complete case sample

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 3536	<i>n</i> = 3536	<i>n</i> = 3536
TBI vs no Injury	1.31 (0.77 – 2.24)	1.41 (0.83 – 2.41)	1.39 (0.81 - 2.39)
OI vs no Injury	1.35 (0.83 - 2.19)	1.37 (0.84 – 2.24)	1.37 (0.84 - 2.24)
TBI vs OI	0.97 (0.49 - 1.91)	1.03 (0.52 - 2.03)	1.02 (0.51 - 2.02)
<i>Omnibus p</i>	0.147	0.116	0.126
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 3526	<i>n</i> = 3526	<i>n</i> = 3526
TBI vs no Injury	0.76 (0.44 – 1.30)	0.71 (0.41 – 1.22)	0.72 (0.42 - 1.24)
OI vs no Injury	0.62 (0.36 - 1.06)	0.55 (0.32 - 0.94)	0.55 (0.32 - 0.95)
TBI vs OI	1.22 (0.59 - 2.55)	1.29 (0.62 - 2.70)	1.30 (0.62 - 2.74)
<i>Omnibus p</i>	0.053	0.015	0.018

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)

Supplementary Table 39 Associations between traumatic brain injury from age 12 to age 16 years, with no additional orthopaedic injury, and psychiatric symptoms at age 17 years compared to orthopaedic injury on complete case sample

SDQ	Unadjusted OR (95% CI)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
Conduct problems ^{a*}			
<i>n</i>	<i>n</i> = 3490	<i>n</i> = 3490	<i>n</i> = 3490
TBI only vs OI	1.13 (0.56 - 2.25)	1.2 (0.60 - 2.42)	1.18 (0.58 - 2.38)
Omnibus <i>p</i>	0.111	0.088	0.096
Peer Problems ^{b*}			
<i>n</i>	<i>n</i> = 3480	<i>n</i> = 3480	<i>n</i> = 3480
TBI only vs OI	1.41 (0.66 - 2.98)	1.51 (0.71 - 3.21)	1.51 (0.71 - 3.22)
Omnibus <i>p</i>	0.079	0.025	0.029

Complete cases had no missing data for the exposure, outcome or covariates. *TBI: traumatic brain injury; OI: orthopaedic injury; *logistic regression; ^a conduct problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years; ^b peer problems based on parent-completed Strengths and Difficulties Questionnaire at age 17 years.*

Unadjusted: Injuries from birth to age 16 years with main substance use variable in each analysis

Model 1: As unadjusted with additional adjustment for pre-birth confounders (mother's age at birth, mother's education at birth, social class and gender)

Model 2: As Model 1 with additional adjustment for childhood confounders (early life events, parental bonding, positive and negative parenting experiences, maternal alcohol use and maternal tobacco smoking)