

Kent Academic Repository

Full text document (pdf)

Citation for published version

Alharbi, Naif Khalaf and Padron-Regalado, Eriko and Thompson, Craig P and Kupke, Alexandra and Wells, Daniel and Sloan, Megan A and Grehan, Keith and Temperton, Nigel J. and Lambe, Teresa and Warimwe, George and Becker, Stephan and Hill, Adrian V S and Gilbert, Sarah C (2017) ChAdOx1 and MVA based vaccine candidates against MERS-CoV elicit neutralising

DOI

<https://doi.org/10.1016/j.vaccine.2017.05.032>

Link to record in KAR

<http://kar.kent.ac.uk/61976/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

1 **ChAdOx1 and MVA based Vaccine Candidates against MERS-CoV Elicit Neutralising Antibodies and**
2 **Cellular Immune Responses in Mice**

3 Naif Khalaf Alharbi^{1,2}, Eriko Padron Regalado¹, Craig P. Thompson^{1,3}, Alexandra Kupke^{4,5}, Daniel Wells¹,
4 Megan A. Sloan¹, Keith Grehan⁶, Nigel Temperton⁶, Teresa Lambe¹, George Warimwe¹, Stephan
5 Becker^{4,5}, Adrian V. S. Hill¹, and Sarah C. Gilbert¹

6

7 ¹The Jenner Institute, University of Oxford, Oxford, OX3 7DQ, UK

8 ²King Abdullah International Medical Research Center, Riyadh, Saudi Arabia

9 ³Department of Zoology, University of Oxford, Oxford, UK

10 ⁴Institute of Virology, Philipps University of Marburg, Marburg, Germany

11 ⁵German Center for Infection Research, TTU Emerging Infections, Germany

12 ⁶Viral Pseudotype Unit, School of Pharmacy, University of Kent, Chatham Maritime, Kent, ME4 4TB, UK

13 **Abstract**

14 The Middle East respiratory syndrome coronavirus (MERS-CoV) has infected more than 1900 humans,
15 since 2012. The syndrome ranges from asymptomatic and mild cases to severe pneumonia and death.
16 The virus is believed to be circulating in dromedary camels without notable symptoms since the 1980s.
17 Therefore, dromedary camels are considered the only animal source of infection. Neither antiviral drugs
18 nor vaccines are approved for veterinary or medical use despite active research on this area. Here, we
19 developed four vaccine candidates against MERS-CoV based on ChAdOx1 and MVA viral vectors, two
20 candidates per vector. All vaccines contained the full-length spike gene of MERS-CoV; ChAdOx1 MERS
21 vaccines were produced with or without the leader sequence of the human tissue plasminogen activator
22 gene (tPA) where MVA MERS vaccines were produced with tPA, but either the mH5 or F11 promoter
23 driving expression of the spike gene. All vaccine candidates were evaluated in a mouse model in prime
24 only or prime-boost regimens. ChAdOx1 MERS with tPA induced higher neutralising antibodies than
25 ChAdOx1 MERS without tPA. A single dose of ChAdOx1 MERS with tPA elicited cellular immune
26 responses as well as neutralising antibodies that were boosted to a significantly higher level by MVA
27 MERS. The humoral immunogenicity of a single dose of ChAdOx1 MERS with tPA was equivalent to two
28 doses of MVA MERS (also with tPA). MVA MERS with mH5 or F11 promoter induced similar antibody
29 levels; however, F11 promoter enhanced the cellular immunogenicity of MVA MERS to significantly
30 higher magnitudes. In conclusion, our study showed that MERS-CoV vaccine candidates could be
31 optimised by utilising different viral vectors, various genetic designs of the vectors, or different regimens
32 to increase immunogenicity. ChAdOx1 and MVA vectored vaccines have been safely evaluated in camels
33 and humans and these MERS vaccine candidates should now be tested in camels and in clinical trials.

34 **Introduction**

35 Middle East respiratory syndrome (MERS) is caused by a novel betacoronavirus (MERS-CoV) that was
36 isolated in late 2012 in Saudi Arabia (1). The syndrome (MERS) is described as a viral infection that
37 causes fever, cough, and/or shortness of breath and to a lesser extent gastrointestinal symptoms such
38 as diarrhea (2). Severe disease from MERS-CoV infection can cause respiratory failure and organ failure,
39 and cases can be fatal, especially in patients with co-morbidities such as diabetes and cardiac
40 complications. However, the infection can be asymptomatic or mild in many cases (3-7). MERS-CoV has
41 spread to 27 countries and infected more than 1900 humans with a mortality rate of 40% (2).
42 Dromedary camels, especially juveniles, contract the infection and shed the virus, without notable
43 symptoms of disease; this is now known to have been occurring since the early 1980s (8-13). The
44 mechanism of camel to human transmission is still not clear, but several primary cases have been
45 associated with camel contact, which is considered an important risk factor (14-16). Therefore, camels
46 are being considered an intermediate host and one of the sources of MERS-CoV infection (8-13). Other
47 livestock animals such as sheep, goats, cows, chicken, and horses have proved seronegative in many
48 studies (17-20). Further, these animals did not productively contract MERS-CoV when they were
49 inoculated experimentally (21, 22). Therefore, to date, dromedary camels are the only confirmed animal
50 reservoir. There is currently no approved vaccine against MERS-CoV for camels or humans despite active
51 vaccine research and development. A number of vaccine candidates have been developed using various
52 platforms and regimens and have been tested in several animal models (23). Viral vectors are potent
53 platform technologies that have been utilised to develop vaccines against malaria, tuberculosis,
54 influenza, HIV, HCV, Ebola, and many viral pathogens. These vectors include adenoviruses, poxviruses,
55 yellow fever viruses, and alphaviruses (24, 25), and they are preferred for their ability to induce cellular
56 immune responses in addition to humoral immunity. Here, we report development of MERS-CoV
57 vaccine candidates that are based on two different viral vectors: Chimpanzee Adenovirus, Oxford

58 University #1 (ChAdOx1) (26) and Modified Vaccinia virus Ankara (MVA) (27, 28). Each viral vector was
59 developed by generating two alternative versions, resulting in four vaccine candidates that all encode
60 the same complete MERS-CoV spike gene (S). The two ChAdOx1 based vaccines were produced with or
61 without the signal peptide of the human tissue plasminogen activator gene (tPA) at the N terminus.
62 Previous studies have shown that encoding tPA upstream of recombinant antigens enhanced
63 immunogenicity, although results differed depending on the antigens employed. The tPA encoded
64 upstream of influenza A virus nucleoprotein, in a DNA vector, enhanced both cellular and humoral
65 immune responses in mice (29, 30), whereas the same leader sequence resulted in increased humoral
66 sequences but decreased cellular responses to HIV Gag (30). The two MVA based vaccines were
67 produced with either the mH5 or F11 poxviral promoter driving antigen expression, both including the
68 tPA sequence at the N terminus of MERS-CoV Spike protein. Previously, we reported the ability of the
69 strong early F11 promoter to enhance cellular immunogenicity of vaccine antigen candidates for malaria
70 and influenza, as compared to utilising p7.5 or mH5 early/late promoters which resulted in a lower level
71 of gene expression immediately after virus infection of target cells, but higher levels at a later stage (31).
72 Here, we continue to assess the F11 promoter in enhancing cellular immunogenicity, and to investigate
73 its ability to impact on humoral immune responses. The four vaccine candidates were evaluated in a
74 number of different regimens in mouse models that showed a single dose of ChAdOx1 MERS inducing
75 higher cellular and humoral immunogenicity than a single dose of MVA MERS, or equivalent to two
76 doses of MVA MERS. ChAdOx1 based vaccines have been tested in different animal models, including
77 camels (32), and in human clinical trials and proved safe and immunogenic (33). Therefore, based on our
78 data, ChAdOx1 MERS can be readily developed for use as a MERS vaccine in humans. Furthermore,
79 utilising ChAdOx1 MERS for camel vaccination can serve the one-health approach whereby blocking
80 MERS-CoV transmission in camels is expected to prevent human infections.

81 **Materials and methods**

82 **Transgene and shuttle vector cloning**

83 The spike (S) gene of MERS-CoV camel isolate (Genbank accession number: KJ650098.1) was synthesised
84 by GeneArt Gene Synthesis (Thermo Fisher Scientific). The S transgene was then cloned into four shuttle
85 plasmid vectors following In-Fusion cloning (Clontech). Two plasmids contained the S transgene within
86 the E1 homologous region of ChAdOx1, driven by the human cytomegalovirus major immediate early
87 promoter (IE CMV) that includes intron A. One of the ChAdOx1 shuttle plasmids was designed to include
88 the tPA signal sequence upstream of the transgene sequence while the second plasmid did not contain
89 the tPA. The ChAdOx1 shuttle plasmids contained the S transgene within Gateway® recombination
90 cassettes. To construct MVA MERS, one of the shuttle plasmids for MVA was designed to have the
91 upstream and downstream (flanks) of the *F11L* ORF as homologous sequence arms. Inserting the S
92 transgene within these arms enabled the utilisation of the endogenous F11 promoter, which is part of
93 the right homologous arm, while deleting the native F11L ORF. This resulted in the shuttle vector for
94 generation of F11-MVA MERS (F11 shuttle vector). The mH5 promoter sequence was subcloned
95 upstream of the S transgene; and this mH5-S transgene was then subcloned into the F11 shuttle vector.
96 This resulted in the shuttle vector for generation of mH5-MVA MERS (F11/mH5 shuttle vector). mH5-
97 MVA MERS contained the mH5 promoter at the *F11L* locus, however, the endogenous F11 promoter is
98 intact and located upstream of the mH5 promoter. The endogenous F11 promoter could not be replaced
99 with the mH5 since it is part of the essential upstream ORF.

100 **Immunostaining for Transgene Expression**

101 The ChAdOx1 shuttle plasmid, described above, was used to validate the expression of MERS-CoV spike
102 protein *in vitro*. An African green monkey kidney cell line (Vero cells) was seeded into 6-well plate to
103 80% confluence. Then the plasmid DNA was transfected into Vero cells using Lipofectamine® 2000

104 (Thermo Fisher Scientific) following manufacturer's instruction. Twenty four hours after transfection,
105 cells were fixed, permeabilised, and immunostained using a rabbit polyclonal anti-MERS-CoV spike
106 antibody, following standard protocols. DAPI stain was used to label nuclei.

107 **Construction of recombinant ChAdOx1 and MVA encoding MERS-CoV S antigens**

108 The ChAdOx1 MERS vaccines were prepared by Gateway® recombination between the ChAdOx1
109 destination DNA BAC vector (described in (26)) and entry plasmids containing the coding sequence for
110 MERS-CoV spike gene (ChAdOx1 shuttle vectors explained above), according to standard protocols.
111 ChAdOx1 MERS genomes were then derived in HEK293A cell lines (Invitrogen, Cat. R705-07), the
112 resultant viruses were purified by CsCl gradient ultracentrifugation as previously described (34). The
113 titres were determined on HEK293A cells using anti-hexon immunostaining assay based on the
114 QuickTiter™ Adenovirus Titer Immunoassay kit (Cell Biolabs Inc). For MVA MERS vaccines chicken
115 embryo fibroblast cells (CEFs) were infected with MVA parental virus that encodes dsRed marker instead
116 of the native F11L ORF and transfected with MVA shuttle plasmids containing MERS-CoV spike gene
117 (explained above) to allow recombination with the MVA genome and deletion of dsRed marker whilst
118 keeping the F11 promoter sequence. Recombinant MVA expressing MERS-CoV S protein was purified by
119 plaque-picking and fluorescent selection using the sorting function of CyCLONE robotic module of a
120 MoFlo Flow cytometer (Dako Cytomation, Denmark) as previously described (31). F11-MVA MERS and
121 mH5-MVA MERS were confirmed to lack the native *F11L* ORF (and the dsRed marker), and contain
122 MERS-CoV S by PCR (identity and purity PCR screening). The sequence of the S transgene amplified from
123 these vaccines was confirmed. The recombinant viruses (vaccines) were amplified in 1500 cm²
124 monolayers of CEFs cells, partially purified over sucrose cushions and titrated in CEFs cells according to
125 standard practice, and purity and identity were again verified by PCR.

126 **Mouse immunogenicity**

127 Female BALB/c mice (Harlan, UK) aged 6 to 8 weeks were immunised intramuscularly (i.m.) in the upper
128 leg (total volume 50 μ L) with a total of 10^8 IU of ChAdOx1 MERS with or without tPA or with a total of
129 10^6 pfu of either F11-MVA MERS or mH5-MVA MERS. For induction of short-term anaesthesia, animals
130 were anaesthetised using vaporised IsoFloH. In prime only regimens, mice were vaccinated with
131 ChAdOx1 with blood samples taken at 14 days post immunisation (d.p.i) or 28 d.p.i. for serum isolation;
132 and spleens were collected at 28 d.p.i. In heterologous prime-boost regimens, mice were vaccinated
133 with ChAdOx1 MERS and boosted with MVA MERS at 28 d.p.i; mice were bled at 28 d.p.i. (post-prime) or
134 42 d.p.i (14 days post-boost) for serum isolation, and spleens were collected at 42 d.p.i. In homologous
135 regimens, mice were vaccinated with MVA MERS and boosted with MVA MERS at 21 d.p.i; mice were
136 bled on 21 d.p.i. (post-prime) or 42 d.p.i (post-boost) for serum isolation and spleens were collected at
137 42 d.p.i.

138 **ELISpot, ICS, and flow cytometry**

139 Splenocytes were harvested for analysis by IFN- γ ELISpot or intracellular cytokine staining (ICS) and flow
140 cytometry as previously described (35, 36), using re-stimulation with 2 μ g/mL S291 MERS-CoV S-specific
141 peptide (VYDTIKYYSIIPHSI); for vaccine cellular immunogenicity (37)); or 1 μ g/mL E3 and F2(G) MVA
142 vector-specific peptides (38) (for anti-MVA immune responses). In the absence of peptide re-
143 stimulation, the frequency of IFN- γ^+ cells, which was typically 0.1% by flow cytometry or less than 50 SFC
144 by ELISpot, was subtracted from tested re-stimulated samples.

145 **ELISA**

146 2 μ g/ml with capturing antigen (S1 recombinant protein from MyBioSource, CA, USA) were used to coat
147 ELISA plates, and standard endpoint ELISA protocol was followed, as previously described (39). Sera
148 were prepared in a 10-fold serial dilution in PBS/T and then 50 μ l were plated in duplicate wells. Serum
149 from a naïve BALB/c mouse was included as a negative control. Goat anti-mouse total IgG conjugated to

150 alkaline phosphatase (Sigma) and PNPP tablet (20 mg p-nitrophenylphosphate, SIGMA) substrate were
151 used in the assay.

152 **MERSpp Neutralisation assay**

153 MERS pseudotyped viral particles (MERSpp) were produced and titrated using Huh7.5 cell line as
154 described previously (40). For the MERSpp neutralization assay, serum samples were serially diluted in
155 96-well white plates (Nunc). A standard concentration of the MERSpp were added to the wells and
156 plates were incubated for 1 h at 37 °C. After incubation, Huh7.5 cells (10,000 cells per well) were added
157 to the plate in duplicates. Following 48 h incubation, cells were lysed and luciferase activity was
158 measured. IC90 neutralisation titres were calculated for each mouse serum sample using GraphPad
159 Prism.

160 **Virus neutralisation assay**

161 Induction of virus-neutralising antibodies was confirmed according to previously published protocols
162 (37, 41). Briefly, mouse serum samples were tested for their capacity to neutralise MERS-CoV (EMC
163 isolate) infections *in vitro* with 100 50% tissue culture infective doses (TCID₅₀) in Huh-7 cells. Sera of non-
164 immunised mice served as negative control.

165 **Statistical analysis**

166 GraphPad Prism (GraphPad software) was used for statistical analysis and to plot data.

167 **Ethics statement**

168 All animal procedures were performed in accordance with the terms of the UK Animals (Scientific
169 Procedures) Act (ASPA) for the project licenses 30/2414 or 30/2889 and were approved by the
170 University of Oxford Animal Care and Ethical Review Committee. All mice were housed for at least 7

171 days for settlement prior to any procedure in the University animal facility, Oxford, UK under Specific
172 Pathogen Free (SPF) conditions.

173 **Results**

174 **Construction and antigen expression of MERS-CoV vaccine candidates**

175 The spike gene from a camel isolate (Camel/Qatar_2_2014 MERS-CoV isolate, GenBank accession
176 number KJ650098.1) was cloned into four shuttle vectors that facilitate homologous recombination with
177 the genome of ChAdOx1 or MVA. Four recombinant viral vectors, two ChAdOx1 and two MVA, were
178 derived as described in the materials and methods. ChAdOx1 based vaccine candidates were generated
179 with or without the signal peptide of the human tissue plasminogen activator gene (tPA). The spike
180 transgene expression in ChAdOx1 MERS vaccine candidates is under the control of the human
181 cytomegalovirus major immediate early promoter (CMV IE) that includes intron A. In MVA MERS vaccine
182 candidates, the tPA was also inserted upstream of the spike transgene, which was under the control of
183 either the ectopic mH5 promoter or the endogenous F11 promoter (Figure 1A). All of our MERS-CoV
184 vaccine candidates contain the same codon-optimized spike transgene. The expression of the newly
185 synthesized transgene was first tested by transfection of an African green monkey kidney cell line (Vero
186 cells) with the adenovirus shuttle vector, and immunofluorescence staining of the transfected cells
187 (Figure 1B and 1C). This was performed to confirm the expression of the codon optimized spike
188 transgene in mammalian cells. The level of transgene expression from the four vaccine candidates was
189 not evaluated *in vitro*. We have previously reported that differences in MVA promoter activity
190 detectable *in vitro* does not correlate with *in vivo* immunogenicity (31), and that only *in vivo* expression
191 correlates with the *in vivo* immunogenicity.

192 **Humoral Immunogenicity of ChAdOx1 based MERS-CoV vaccine candidates**

193 To evaluate humoral immune responses to ChAdOx1 MERS with or without tPA, BALB/c mice were
194 vaccinated with 1×10^8 IU of ChAdOx1 intramuscularly. Serum samples from 14 and 28 d.p.i. were
195 collected and evaluated by ELISA. Both vaccine candidates induced a high level of S1-specific antibodies
196 (mean endpoint titre (Log_{10}) = 4.8 with tPA, 4.7 without tPA), unlike the control vaccine, ChAdOx1
197 encoding enhanced green fluorescent protein (ChAdOx1-eGFP, mean endpoint titre (Log_{10}) = 1). These
198 antibody levels were similar between the two candidates (with or without tPA) at day 14. However, at
199 28 d.p.i. ChAdOx1 MERS with tPA induced significantly higher S1-specific antibodies than ChAdOx1
200 MERS without tPA (mean endpoint titre (Log_{10}) = 5.13 with tPA, 4.6 without tPA, Figure 2A). Serum
201 samples from day 28 were selected for MERSpp neutralisation assay. Serum antibodies induced by
202 ChAdOx1 MERS with tPA showed significantly higher neutralisation activity than without tPA (mean titre
203 IC_{90} (Log_{10}) = 2.8 with tPA, 2.2 without tPA; Figure 2B). In order to confirm that the pseudotyped virus
204 neutralisation assay was producing biologically relevant results, serum samples from mice immunised
205 with ChAdOx1 MERS with tPA were also tested in a neutralisation assay utilising wildtype MERS virus.
206 This assay confirmed the neutralisation activity of mouse antibodies (nAb) with a median of 360 VNT
207 (Virus Neutralization Test antibody titre; Figure 2C). We therefore continued to evaluate ChAdOx1 MERS
208 with tPA in addition to generating MVA MERS vaccine candidates with tPA.

209 **Cellular Immunogenicity of ChAdOx1 based MERS-CoV vaccine candidates**

210 Having established the utility of tPA in ChAdOx1 MERS vaccines (referred to as ChAdOx1 MERS in the
211 rest of this report) at increasing humoral responses, spleens were collected at 28 d.p.i. from immunised
212 BALB/c mice. Splenocytes were processed to evaluate cellular immune responses to ChAdOx1 MERS in
213 ELISpot and Intracellular cytokine staining (ICS). Peptide S291, described by others (37), was used to re-
214 stimulate the cells in both assays and ELISpot data showed a high level of IFN- γ secreting splenocytes

215 (Median = 1300 SFU/10⁶ splenocytes; Figure 3A). ICS data confirmed the IFN- γ secreting CD8⁺
216 splenocytes also secreted TNF- α and IL-17 (Figure 3B).

217 **Immunogenicity of Heterologous ChAdOx1 and MVA vaccination against MERS-CoV**

218 To evaluate humoral immune responses to heterologous prime-boost vaccination, BALB/c mice were
219 immunised with ChAdOx1 MERS vaccine and boosted with one of two different MVA MERS vaccine
220 candidates four weeks later. The MVA based candidates differ in the promoters that controls the
221 transgene expression: F11-MVA MERS utilises the endogenous strong early F11 promoter and mH5-
222 MVA MERS utilises the ectopic early/late mH5 promoter. Serum samples from 28 d.p.i. (post-prime) or
223 42 d.p.i. (post-boost) were collected and evaluated by ELISA and MERSpp neutralisation assay. At 28
224 d.p.i. ChAdOx1 MERS induced similar levels of S1-specific antibodies and nAb as observed previously
225 (Figure 4A and B). At 42 d.p.i. S1-specific antibodies were boosted to a higher level (mean endpoint titre
226 (Log_{10}) = 5 by ChAdOx1 MERS boosted to 5.8 by mH5-MVA MERS or 5.9 by F11-MVA MERS); Figure 4A)
227 with nAb also enhanced to a statistically significant level (mean titre IC_{90} (Log_{10}) = 2.87 by ChAdOx1
228 MERS boosted to 3.3 by mH5-MVA MERS or 3.5 by F11-MVA MERS; Figure 4B). There was no difference
229 in antibody levels induced using either the F11 or mH5 promoter in the MVA.

230 At 42 d.p.i. splenocytes were also processed to evaluate cellular immune responses to ChAdOx1 MERS
231 MVA MERS prime-boost vaccination in ELISpot and ICS as shown in Figure 3. The T cell responses to
232 MERS S were boosted by the MVA vaccinations; in the ICS experiments, F11-MVA and mH5-MVA
233 boosted the percentage of IFN- γ ⁺ splenic CD8⁺ T cells to 7.3 and 5.2% respectively (Figure 4D) whereas
234 the percentage was 2.5% after ChAdOx1 MERS prime in Figure 3B. The percentage of TNF- α ⁺ splenic
235 CD8⁺ T cells were also increased by MVA boost (comparing Figure 3B and 4D). Utilising the F11 promoter
236 resulted in a trend towards greater cell-mediated immunogenicity (Figure 4C and D). Splenocytes were
237 also re-stimulated with MVA backbone-specific E3 and F(G)2 peptides and evaluated in ICS. Both MVA

238 based vaccines induced similar responses to E3 or to F(G)2 peptides, 2 weeks after MVA vaccination
239 (Figure 4E and F). This similarity confirmed the efficiency of vaccine titration, vaccination, and sample
240 processing because responses to each of those peptides are not expected to be different unless there is
241 variation in the doses administered or sample preparation. Overall, MVA MERS vaccines were able to
242 boost the humoral and cellular immune responses to ChAdOx1 MERS prime vaccination. There was no
243 difference between the F11 and mH5 promoter in the resulting antibody titres after ChAdOx1
244 prime/MVA boost, but there was a trend towards increased cellular immunogenicity when the F11
245 promoter was used.

246 **Immunogenicity of Homologous MVA vaccination against MERS-CoV**

247 To evaluate humoral immune responses to a homologous MVA MERS prime-boost vaccination, two
248 groups of BALB/c mice were immunised with F11-MVA MERS or mH5-MVA MERS and boosted with the
249 same vaccine after three weeks. Serum samples from 21 d.p.i. (post-prime) or 42 d.p.i. (post-boost)
250 were collected and evaluated in ELISA and MERSpp neutralisation assays. At 21 d.p.i. F11-MVA MERS
251 and mH5-MVA induced similar levels of S1-specific antibodies (mean endpoint titre (Log_{10}) = 3.2 and 2.8
252 respectively; Figure 5A). At 42 d.p.i. S1-specific antibody levels had increased to 4.7 and 4.8 respectively
253 (Figure 5A). The titres of nAb (MERS pp assay) were also similar for both vaccines (mean titre IC_{90} (Log_{10})
254 = 2.71 (F11-MVA MERS) and 2.76 respectively; Figure 5B). Utilising different promoters in MVA vectors
255 did not result in differences in the induced antibody levels. However, at 42 d.p.i. IFN- γ secreting
256 splenocytes induced by F11-MVA MERS were statistically significantly higher than those of mH5-MVA
257 MERS ((Median = 525 and 249 SFU/ 10^6 splenocytes, respectively, Figure 5C). Both MVA vaccines induced
258 similar vector-specific immune responses as expected (Figure 5D and E).

259 Discussion

260 Vaccines against MERS-CoV have been developed and tested in a number of animal models (including
261 non-human primates (42-44) and camels (45)) as well as in human clinical trials (46). All vaccine
262 candidates focused on the spike antigen because it contains the receptor-binding domain used for cell
263 entry by the virus, against which neutralising antibodies may be induced, and it is conserved. Therefore,
264 the improvement of MERS-CoV vaccines focuses on platform and vaccination regimens rather than
265 antigen selection and optimisation. Here, we focused on using the same antigen (transgene) to develop
266 a vaccine against MERS-CoV, and to assess different vectors, different versions of each vector, and
267 different vaccination regimens. We generated a number of MERS-CoV vaccine candidates based on the
268 same codon optimized spike transgene and ensured its expression *in vitro* before we evaluated the
269 humoral and cellular immunogenicity in a pre-clinical BALB/c mouse model. ChAdOx1 based vaccine
270 candidates were produced with or without tPA. The tPA signal peptide was predicted to enhance the
271 humoral immunogenicity of encoded vaccine antigens, based on previous reports (29). Our data
272 supported this hypothesis and showed a significant increase in the S1-specific antibody levels at 28 d.p.i.
273 The level of neutralising antibodies was also increased when tPA was utilised. However, ChAdOx1 MERS
274 without tPA was still a potent vaccine candidate, inducing a high level of both S1-specific binding
275 antibodies and MERS-CoV neutralising antibodies. Neutralisation activity of mouse serum antibodies
276 was assayed by using MERS-CoV pseudotyped viral particles (MERSpp), an approach used by a number
277 of researchers for other human pathogens such as HIV, Influenza, and HCV to overcome the necessity of
278 handling BSL-3 viruses (40). Additionally, we confirmed the ability of serum samples from vaccinated
279 mice to neutralise live MERS virus. We therefore selected ChAdOx1 MERS with tPA (simply referred to
280 ChAdOx1 MERS) for further evaluation.

281 ChAdOx1 MERS also induced cellular responses for MERS S, with polyfunctional CD8⁺ T cells detected in
282 the spleen of immunized mice. This supports the potency of the ChAdOx1 viral vector in inducing T
283 cellular immunity, observed previously in animal models (26, 32, 47) as well as in humans (33). Following
284 ChAdOx1 prime/MVA boost, MVA significantly boosted the neutralizing antibody titres to higher levels.
285 No difference in humoral immunity was found when either the F11 or mH5 promoter was used.
286 Regarding the promoter effect on MVA cellular immunogenicity, we have previously reported that
287 utilising the F11 promoter enhanced malaria and influenza antigens in MVA (31). Here, we again report
288 that F11-MVA MERS induced higher T cell responses than mH5-MVA MERS in a homologous prime-boost
289 MVA MERS vaccination.

290 All of our vaccine candidates induced humoral (with nAb) and cellular immune (with polyfunctional CD8⁺
291 T cell) responses against MERS-CoV spike antigen. Modest effects on immunogenicity of different
292 versions of the vaccines were noted, with the use of the tPA leader sequence in ChAdOx1, and the use
293 of the F11 promoter in MVA producing small increases in immunogenicity compared to no leader
294 sequence, or the mH5 promoter. The protective level of either antibodies or cellular immunity required
295 to counter MERS-CoV infection in humans or in animal models is not yet defined, despite some efforts
296 (48-51). The ideal vaccine would provide rapid onset of immunity and complete protective efficacy after
297 a single dose, with a long duration of immunity. Complete protective efficacy of one dose of ChAdOx1
298 expressing the external glycoprotein of Rift Valley Fever Virus has been demonstrated in multiple
299 species and it is already known that ChAdOx1 RVF is highly immunogenic in camels (32). To date, the
300 only vaccine against MERS to be tested in camels is an MVA vectored vaccine (41) which was protective
301 in hDPP4 transgenic mice immunized with a homologous prime/boost regimen (37) but in camels
302 required two doses given both intranasally and intramuscularly to provide partial protection and
303 reduction of virus shedding (45). Here we find that a single dose of ChAdOx1 MERS is as immunogenic as
304 two doses of MVA MERS, suggesting that this regimen should be tested for protective efficacy in camels.

305 However if this is not completely protective, administration of MVA MERS as a heterologous boost
306 should be considered next. In our hands one dose of MVA resulted in an endpoint titre of 3 logs, two
307 doses of MVA produced 4.7 logs, one dose of ChAdOx1 produced 5 logs, and ChAdOx1/MVA prime
308 boost produced 5.9 logs. If a single dose of ChAdOx1 MERS is not protective and a two dose regimen is
309 required, ChAdOx1/MVA would be more likely to provide complete protection than MVA/MVA.

310 ChAdOx1 MERS should now be evaluated for immunogenicity and efficacy in larger animal species,
311 including both camels and humans.

312 **Figure legends**

313 **Figure 1: Construction of MERS-CoV vaccine candidates**

314 **A:** schematic representation of ChAdOx1 and MVA based vaccines, each encodes the same MERS-CoV spike gene (MERS-CoV S).
315 The S gene was inserted into the E1 region of ChAdOx1 genome or into the *F11L* locus of MVA genome. tPA: Human tissue
316 plasminogen activator (tPA) signal peptide sequence. IE CMV: The human cytomegalovirus major immediate early promoter.
317 mH5 and F11: Poxviral promoters. LHA: left homology arm sequence. RHA: right homology arm sequence. **B:** The expression of
318 spike transgene, cloned into a plasmid vector, was validated by transfection into an African green monkey kidney cell line (Vero
319 cells) and confirmed by immunostaining. C: Untransfected cells control. Green colour represents detection of the spike protein.
320 Blue colour represents nuclei by staining nucleic acid with DAPI.

321 **Figure 2: Antibody responses to ChAdOx1 MERS vaccine candidates.**

322 BALB/c mice (n = 6) were immunised with a single injection of ChAdOx1 MERS that either encodes or lacks tPA signal peptide,
323 intramuscularly at 1×10^8 IU. A control group of mice were immunised with ChAdOx1 expressing eGFP instead of MERS-CoV S
324 gene. Serum samples were collected at 14 and 28 days post immunisation (d.p.i.). S1-binding antibodies were detected at both
325 time points by ELISA (A) and neutralisation activity of the antibodies were confirmed by MERS-CoV pseudotyped viral particles
326 (MERSpp) neutralisation assay (B) or neutralisation assay (C). Individual data points are shown with line as the median. Data are
327 representative of two independent experiments. Statistical significance by Kruskal–Wallis test is shown.

328 **Figure 3: Cellular immune responses to ChAdOx1 MERS vaccine candidate.**

329 BALB/c mice (n = 6) were immunised with a single injection of ChAdOx1 MERS that encodes tPA signal peptide intramuscularly
330 at 1×10^8 IU. Twenty eight days post-immunisation, IFN- γ *ex vivo* ELISpot (A) or Intracellular Cytokine Staining (ICS (B)), were
331 performed to determine the percentage of splenic IFN- γ secreting CD4⁺ and CD8⁺ after *in vitro* re-stimulation with a MERS-CoV
332 S-specific peptide. Individual data points are shown with line as the median (A) or error bars as the SD (B). Data are
333 representative of two independent experiments.

334 **Figure 4: Humoral and cellular immunogenicity of heterologous ChAdOx1 MERS and MVA MERS** 335 **vaccination.**

336 BALB/c mice (n = 6) were immunised with ChAdOx1 MERS that encodes tPA signal peptide, intramuscularly at 1×10^8 IU. At 28
337 d.p.i. mice were boosted with MVA MERS at 1×10^6 pfu. MVA MERS candidates either contain mH5 or F11 promoter for
338 transgene expression. Serum samples were collected at 28 (post-prime) and 42 (post-boost) d.p.i. S1-binding antibodies were
339 detected at both time points by ELISA (A) and neutralisation activity of serum antibodies at 42 d.p.i. were confirmed by MERSpp
340 neutralisation assay (B). At 42 d.p.i, IFN- γ *ex vivo* ELISpot (C) or Intracellular Cytokine Staining (ICS (D)) were performed to
341 determine the percentage of CD8⁺ IFN- γ ⁺ splenocytes after *in vitro* re-stimulation with a MERS-CoV S-specific peptide. ICS of
342 splenocytes re-stimulated with MVA-specific peptides (F(G)2 and E3) was also performed (E and F). Individual data points are
343 shown with line as the median. Data are representative of two independent experiments. Statistical significance by Kruskal–
344 Wallis test is shown. Symbols are closed squares (■) for ChAdOx1 prime responses, open circles (○) for mH5-MVA boost
345 responses, and closed circles (●) for F11-MVA boost responses.

346 **Figure 5: Humoral and cellular immunogenicity of homologous MVA MERS vaccination.**

347 BALB/c mice (n = 6) were immunised with MVA MERS at 1×10^6 pfu, intramuscularly, in a homologous prime-boost vaccination
348 with three-weeks interval. MVA MERS candidates either contain mH5 or F11 promoter for transgene expression. Serum
349 samples were collected at 21 (post-prime) and 42 (post-boost) d.p.i. S1-binding antibodies were detected at both time points by
350 ELISA (A) and neutralisation activity of serum antibodies at 42 d.p.i. were confirmed by MERSpp neutralisation assay (B). At 42
351 d.p.i splenocytes were processed and re-stimulated with a MERS-CoV S-specific peptide (CD8⁺ T cell specific) for IFN- γ *ex vivo*
352 ELISpot (C). ICS of splenocytes re-stimulated with MVA-specific peptides (F(G)2 and E3) was also performed (D and E) as was
353 performed in figure 4. Individual data points are shown with line as the median. Data are representative of two independent
354 experiments. Statistical significance by Kruskal–Wallis test is shown. Symbols are open circles (○) for mH5-MVA and closed
355 circles (●) for F11-MVA.

356 **Conflict of interest**

357 SCG is a co-founder of, consultant to and shareholder in Vaccitech plc which is developing vectored influenza and MERS
358 vaccines.

359 **References**

- 360 1. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel
361 coronavirus from a man with pneumonia in Saudi Arabia. *The New England journal of medicine*.
362 2012;367(19):1814-20.
- 363 2. World Health Organisation. Middle East respiratory syndrome coronavirus (MERS-CoV) 2016.
364 Available from: <http://www.who.int/emergencies/mers-cov/en/>. [accessed 20.02.2017].
- 365 3. Lessler J, Salje H, Van Kerkhove MD, Ferguson NM, Cauchemez S, Rodriquez-Barraquer I, et al.
366 Estimating the Severity and Subclinical Burden of Middle East Respiratory Syndrome Coronavirus
367 Infection in the Kingdom of Saudi Arabia. *American journal of epidemiology*. 2016;183(7):657-63.
- 368 4. Al Hammadi ZM, Chu DK, Eltahir YM, Al Hosani F, Al Mulla M, Tarnini W, et al. Asymptomatic
369 MERS-CoV Infection in Humans Possibly Linked to Infected Dromedaries Imported from Oman to United
370 Arab Emirates, May 2015. *Emerging infectious diseases*. 2015;21(12):2197-200.
- 371 5. Oboho IK, Tomczyk SM, Al-Asmari AM, Banjar AA, Al-Mugti H, Aloraini MS, et al. 2014 MERS-CoV
372 outbreak in Jeddah--a link to health care facilities. *N Engl J Med*. 2015;372(9):846-54.
- 373 6. Who Mers-Cov Research G. State of Knowledge and Data Gaps of Middle East Respiratory
374 Syndrome Coronavirus (MERS-CoV) in Humans. *PLoS currents*. 2013;5.
- 375 7. Memish ZA, Zumla AI, Assiri A. Middle East respiratory syndrome coronavirus infections in
376 health care workers. *The New England journal of medicine*. 2013;369(9):884-6.
- 377 8. Muller MA, Corman VM, Jores J, Meyer B, Younan M, Liljander A, et al. MERS coronavirus
378 neutralizing antibodies in camels, Eastern Africa, 1983-1997. *Emerging infectious diseases*.
379 2014;20(12):2093-5.
- 380 9. Perera RA, Wang P, Gomaa MR, El-Shesheny R, Kandeil A, Bagato O, et al. Seroepidemiology for
381 MERS coronavirus using microneutralisation and pseudoparticle virus neutralisation assays reveal a high
382 prevalence of antibody in dromedary camels in Egypt, June 2013. *Euro surveillance : bulletin Europeen
383 sur les maladies transmissibles = European communicable disease bulletin*. 2013;18(36):pii=20574.
- 384 10. Hemida MG, Perera RA, Al Jassim RA, Kayali G, Siu LY, Wang P, et al. Seroepidemiology of Middle
385 East respiratory syndrome (MERS) coronavirus in Saudi Arabia (1993) and Australia (2014) and
386 characterisation of assay specificity. *Euro surveillance : bulletin Europeen sur les maladies transmissibles
387 = European communicable disease bulletin*. 2014;19(23).
- 388 11. Gutierrez C, Tejedor-Junco MT, Gonzalez M, Lattwein E, Renneker S. Presence of antibodies but
389 no evidence for circulation of MERS-CoV in dromedaries on the Canary Islands, 2015. *Euro surveillance :
390 bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin*.
391 2015;20(37).
- 392 12. Ali MA, Shehata MM, Gomaa MR, Kandeil A, El-Shesheny R, Kayed AS, et al. Systematic, active
393 surveillance for Middle East respiratory syndrome coronavirus in camels in Egypt. *Emerging microbes &
394 infections*. 2017;6(1):e1.
- 395 13. van Doremalen N, Hijazeen ZS, Holloway P, Al Omari B, McDowell C, Adney D, et al. High
396 Prevalence of Middle East Respiratory Coronavirus in Young Dromedary Camels in Jordan. *Vector borne
397 and zoonotic diseases*. 2017;17(2):155-9.

- 398 14. Alraddadi BM, Watson JT, Almarashi A, Abedi GR, Turkistani A, Sadran M, et al. Risk Factors for
399 Primary Middle East Respiratory Syndrome Coronavirus Illness in Humans, Saudi Arabia, 2014. *Emerging*
400 *infectious diseases*. 2016;22(1):49-55.
- 401 15. Gossner C, Danielson N, Gervelmeyer A, Berthe F, Faye B, Kaasik Aaslav K, et al. Human-
402 Dromedary Camel Interactions and the Risk of Acquiring Zoonotic Middle East Respiratory Syndrome
403 Coronavirus Infection. *Zoonoses and public health*. 2016;63(1):1-9.
- 404 16. Funk AL, Goutard FL, Miguel E, Bourgarel M, Chevalier V, Faye B, et al. MERS-CoV at the Animal-
405 Human Interface: Inputs on Exposure Pathways from an Expert-Opinion Elicitation. *Frontiers in*
406 *veterinary science*. 2016;3:88.
- 407 17. Reusken CB, Ababneh M, Raj VS, Meyer B, Eljarah A, Abutarbush S, et al. Middle East Respiratory
408 Syndrome coronavirus (MERS-CoV) serology in major livestock species in an affected region in Jordan,
409 June to September 2013. *Euro surveillance : bulletin Europeen sur les maladies transmissibles =*
410 *European communicable disease bulletin*. 2013;18(50):20662.
- 411 18. Hemida MG, Perera RA, Wang P, Alhammadi MA, Siu LY, Li M, et al. Middle East Respiratory
412 Syndrome (MERS) coronavirus seroprevalence in domestic livestock in Saudi Arabia, 2010 to 2013. *Euro*
413 *surveillance : bulletin Europeen sur les maladies transmissibles = European communicable disease*
414 *bulletin*. 2013;18(50):20659.
- 415 19. Alagaili AN, Briese T, Mishra N, Kapoor V, Sameroff SC, Burbelo PD, et al. Middle East respiratory
416 syndrome coronavirus infection in dromedary camels in Saudi Arabia. *mBio*. 2014;5(2):e00884-14.
- 417 20. Reusken CB, Haagmans BL, Muller MA, Gutierrez C, Godeke GJ, Meyer B, et al. Middle East
418 respiratory syndrome coronavirus neutralising serum antibodies in dromedary camels: a comparative
419 serological study. *The Lancet Infectious diseases*. 2013;13(10):859-66.
- 420 21. Vergara-Alert J, van den Brand JM, Widagdo W, Munoz Mt, Raj S, Schipper D, et al. Livestock
421 Susceptibility to Infection with Middle East Respiratory Syndrome Coronavirus. *Emerging infectious*
422 *diseases*. 2017;23(2):232-40.
- 423 22. Adney DR, Brown VR, Porter SM, Bielefeldt-Ohmann H, Hartwig AE, Bowen RA. Inoculation of
424 Goats, Sheep, and Horses with MERS-CoV Does Not Result in Productive Viral Shedding. *Viruses*.
425 2016;8(8).
- 426 23. Alharbi NK. Vaccines against Middle East respiratory syndrome coronavirus for humans and
427 camels. *Reviews in medical virology*. 2016.
- 428 24. Draper SJ, Heeney JL. Viruses as vaccine vectors for infectious diseases and cancer. *Nat Rev*
429 *Microbiol*. 2010;8(1):62-73.
- 430 25. Tao D, Barba-Spaeth G, Rai U, Nussenzweig V, Rice CM, Nussenzweig RS. Yellow fever 17D as a
431 vaccine vector for microbial CTL epitopes: protection in a rodent malaria model. *The Journal of*
432 *experimental medicine*. 2005;201(2):201-9.
- 433 26. Dicks MD, Spencer AJ, Edwards NJ, Wadell G, Bojang K, Gilbert SC, et al. A novel chimpanzee
434 adenovirus vector with low human seroprevalence: improved systems for vector derivation and
435 comparative immunogenicity. *PloS one*. 2012;7(7):e40385.
- 436 27. Gomez CE, Perdiguero B, Garcia-Arriaza J, Esteban M. Clinical applications of attenuated MVA
437 poxvirus strain. *Expert review of vaccines*. 2013;12(12):1395-416.
- 438 28. Gilbert SC. Clinical development of Modified Vaccinia virus Ankara vaccines. *Vaccine*.
439 2013;31(39):4241-6.
- 440 29. Luo M, Tao P, Li J, Zhou S, Guo D, Pan Z. Immunization with plasmid DNA encoding influenza A
441 virus nucleoprotein fused to a tissue plasminogen activator signal sequence elicits strong immune
442 responses and protection against H5N1 challenge in mice. *Journal of virological methods*. 2008;154(1-
443 2):121-7.

444 30. Wallace A, West K, Rothman AL, Ennis FA, Lu S, Wang S. Post-translational intracellular
445 trafficking determines the type of immune response elicited by DNA vaccines expressing Gag antigen of
446 Human Immunodeficiency Virus Type 1 (HIV-1). *Hum Vaccin Immunother.* 2013;9(10):2095-102.

447 31. Alharbi NK, Spencer AJ, Salman AM, Tully CM, Chinnakannan SK, Lambe T, et al. Enhancing
448 cellular immunogenicity of MVA-vectored vaccines by utilizing the F11L endogenous promoter. *Vaccine.*
449 2016;34(1):49-55.

450 32. Warimwe GM, Gesharisha J, Carr BV, Otieno S, Otingah K, Wright D, et al. Chimpanzee
451 Adenovirus Vaccine Provides Multispecies Protection against Rift Valley Fever. *Sci Rep.* 2016;6:20617.

452 33. Antrobus RD, Coughlan L, Berthoud TK, Dicks MD, Hill AV, Lambe T, et al. Clinical assessment of a
453 novel recombinant simian adenovirus ChAdOx1 as a vectored vaccine expressing conserved Influenza A
454 antigens. *Molecular therapy : the journal of the American Society of Gene Therapy.* 2014;22(3):668-74.

455 34. Cottingham MG, Carroll F, Morris SJ, Turner AV, Vaughan AM, Kapulu MC, et al. Preventing
456 spontaneous genetic rearrangements in the transgene cassettes of adenovirus vectors. *Biotechnol*
457 *Bioeng.* 2012;109(3):719-28.

458 35. Sridhar S, Reyes-Sandoval A, Draper SJ, Moore AC, Gilbert SC, Gao GP, et al. Single-dose
459 protection against *Plasmodium berghei* by a simian adenovirus vector using a human cytomegalovirus
460 promoter containing intron A. *J Virol.* 2008;82(8):3822-33.

461 36. Cottingham MG, Andersen RF, Spencer AJ, Saurya S, Furze J, Hill AV, et al. Recombination-
462 mediated genetic engineering of a bacterial artificial chromosome clone of modified vaccinia virus
463 Ankara (MVA). *PLoS one.* 2008;3(2):e1638.

464 37. Volz A, Kupke A, Song F, Jany S, Fux R, Shams-Eldin H, et al. Protective efficacy of recombinant
465 Modified Vaccinia virus Ankara (MVA) delivering Middle East Respiratory Syndrome coronavirus spike
466 glycoprotein. *Journal of virology.* 2015.

467 38. Tschärke DC, Woo WP, Sakala IG, Sidney J, Sette A, Moss DJ, et al. Poxvirus CD8+ T-cell
468 determinants and cross-reactivity in BALB/c mice. *Journal of virology.* 2006;80(13):6318-23.

469 39. Draper SJ, Moore AC, Goodman AL, Long CA, Holder AA, Gilbert SC, et al. Effective induction of
470 high-titer antibodies by viral vector vaccines. *Nature medicine.* 2008;14(8):819-21.

471 40. Grehan K, Ferrara F, Temperton N. An optimised method for the production of MERS-CoV spike
472 expressing viral pseudotypes. *MethodsX.* 2015;2:379-84.

473 41. Song F, Fux R, Provacía LB, Volz A, Eickmann M, Becker S, et al. Middle East respiratory
474 syndrome coronavirus spike protein delivered by modified vaccinia virus Ankara efficiently induces virus-
475 neutralizing antibodies. *Journal of virology.* 2013;87(21):11950-4.

476 42. Muthumani K, Falzarano D, Reuschel EL, Tingey C, Flingai S, Villarreal DO, et al. A synthetic
477 consensus anti-spike protein DNA vaccine induces protective immunity against Middle East respiratory
478 syndrome coronavirus in nonhuman primates. *Science translational medicine.* 2015;7(301):301ra132.

479 43. Lan J, Yao Y, Deng Y, Chen H, Lu G, Wang W, et al. Recombinant Receptor Binding Domain
480 Protein Induces Partial Protective Immunity in Rhesus Macaques Against Middle East Respiratory
481 Syndrome Coronavirus Challenge. *EBioMedicine.* 2015;2(10):1438-46.

482 44. Wang L, Shi W, Joyce MG, Modjarrad K, Zhang Y, Leung K, et al. Evaluation of candidate vaccine
483 approaches for MERS-CoV. *Nature communications.* 2015;6:7712.

484 45. Haagmans BL, van den Brand JM, Raj VS, Volz A, Wohlsein P, Smits SL, et al. An orthopoxvirus-
485 based vaccine reduces virus excretion after MERS-CoV infection in dromedary camels. *Science.*
486 2016;351(6268):77-81.

487 46. Pellerin C. Army Scientists Begin First MERS Vaccine Clinical Trial. DoD News, Defense Media
488 Activity. 2016.


489 47. Warimwe GM, Lorenzo G, Lopez-Gil E, Reyes-Sandoval A, Cottingham MG, Spencer AJ, et al.
490 Immunogenicity and efficacy of a chimpanzee adenovirus-vectored Rift Valley fever vaccine in mice.
491 *Virology journal.* 2013;10:349.

- 492 48. Wang W, Wang H, Deng Y, Song T, Lan J, Wu G, et al. Characterization of anti-MERS-CoV
493 antibodies against various recombinant structural antigens of MERS-CoV in an imported case in China.
494 *Emerging microbes & infections*. 2016;5(11):e113.
- 495 49. Min CK, Cheon S, Ha NY, Sohn KM, Kim Y, Aigerim A, et al. Comparative and kinetic analysis of
496 viral shedding and immunological responses in MERS patients representing a broad spectrum of disease
497 severity. *Scientific reports*. 2016;6:25359.
- 498 50. Park WB, Perera RA, Choe PG, Lau EH, Choi SJ, Chun JY, et al. Kinetics of Serologic Responses to
499 MERS Coronavirus Infection in Humans, South Korea. *Emerging infectious diseases*. 2015;21(12):2186-9.
- 500 51. Corman VM, Albarrak AM, Omrani AS, Albarrak MM, Farah ME, Almasri M, et al. Viral Shedding
501 and Antibody Response in 37 Patients With Middle East Respiratory Syndrome Coronavirus Infection.
502 *Clin Infect Dis*. 2016;62(4):477-83.


503

Figure 1
[Click here to download high resolution image](#)

A



B



C




Figure 2
[Click here to download high resolution image](#)






Figure 3
[Click here to download high resolution image](#)





Figure 4
[Click here to download high resolution image](#)




Figure 5
[Click here to download high resolution image](#)

