

Competition between movement plans increases motor variability: evidence of a shared resource for movement planning

Running head: A shared resource for movement planning

Leonie Oostwoud Wijdenes^{1,2}, Richard B. Ivry³ & Paul M. Bays^{1,4}

¹Institute of Neurology, University College London, London, UK; ²Radboud University Nijmegen, Donders Institute for Brain, Cognition and Behaviour, Nijmegen, The Netherlands; ³Institute of Cognitive and Brain Sciences, University of California Berkeley, USA; ⁴University of Cambridge, Department of Psychology, Cambridge, UK

Corresponding author: Leonie Oostwoud Wijdenes, Radboud University Nijmegen, Donders Institute for Brain, Cognition and Behaviour, Box 9104, 6500 HE Nijmegen, The Netherlands, l.oostwoudwijdenes@donders.ru.nl

Author Contributions

P.M.B, L.O.W. and R.B.I. conception and design of research; L.O.W. performed experiments; L.O.W. analyzed data; L.O.W., P.M.B. and R.B.I interpreted results of experiments; L.O.W. prepared figures; L.O.W. and P.M.B. drafted manuscript; L.O.W., P.M.B. and R.B.I edited and revised manuscript; L.O.W., P.M.B. and R.B.I approved final version of manuscript.

Abstract

Do movement plans, like representations in working memory, share a limited pool of resources? If so, the precision with which each individual movement plan is specified should decrease as the total number of movement plans increases. To explore this, human participants made speeded reaching movements toward visual targets. We examined if preparing one movement resulted in less variability than preparing two movements. The number of planned movements was manipulated in a delayed response cueing procedure that limited planning to a single target (Exp. 1) or hand (Exp. 2), or required planning of movements toward two targets (or with two hands). For both experiments, initial movement direction variability was higher in the two-plan than in the one-plan condition, demonstrating a cost associated with planning multiple movements, consistent with the limited resource hypothesis. In a third experiment we show that the advantage in initial variability of preparing a single movement was present only when the trajectory could be fully specified. This indicates that the difference in variability between one and two plans reflects the specification of full motor plans, not a general preparedness to move. The precision cost related to concurrent plans represents a novel constraint on motor preparation, indicating that multiple movements cannot be planned independently, even if they involve different limbs.

New & Noteworthy

Various lines of evidence indicate that multiple movements can be prepared in parallel. We show that preparing more than one movement comes with a cost: a movement plan is more variable if it is prepared simultaneously with another plan. This suggests that the

representations of movement plans share a common neural resource, and implies that the number of alternative plans is constrained by noise.

Keywords

Movement planning, motor control, parallel encoding, action, reaching

Introduction

A simple decision, such as selecting which of two identical cartons of milk to grab from the supermarket shelf, involves both perceptual processes, to determine the locations of the cartons, and motor processes, to prepare the appropriate movement (Song and Nakayama 2009). Neurophysiology studies with non-human primates indicate that choosing between two possible actions is not a serial process, whereby one of the two actions is selected and then the selected action is planned and executed, but rather a parallel process in which both actions are prepared simultaneously (Cisek and Kalaska 2005; Cisek 2006). These studies suggest that the final movement is the outcome of a competition between the motor plans. This idea has been motivated by evidence suggesting parallel processing during action preparation (Kornblum et al. 1990; Prescott et al. 1999; Cisek and Kalaska 2005; Cisek 2006) and formalized in the affordance competition hypothesis (Cisek 2007; Cisek and Kalaska 2010).

Studies with humans also indicate that we are capable of simultaneously planning multiple movements. Behavioral evidence comes from studies examining the effects of non-cued targets on movement selection (Gallivan et al. 2015), external perturbations that require rapid switches to an alternative plan (Nashed et al. 2014), and tasks in which participants are forced to move before the target is fully specified (Ghez et al. 1997; Chapman et al.

2010; Stewart et al. 2014; Haith et al. 2015; Gallivan et al. 2016). Neurophysiologically, signatures of parallel planning are evident in studies showing the modulatory effects of target separation on neural activity (Grent-'t-Jong et al. 2014) as well as cortical inhibition of non-selected movements (Duque et al. 2010; Klein-Flugge and Bestmann 2012). The level at which movement plans compete has been the subject of considerable discussion. Hypotheses range from competition at the level of motor goals (Wong et al. 2014; Haith et al. 2015) to competition at the level of biomechanically specified movements (Cos et al. 2014; Gallivan et al. 2015) and movement control policies (Gallivan et al. 2016).

Simultaneous preparation of multiple movements has been suggested to enable the motor system to respond fast and flexibly to environmental changes (Nashed et al. 2014; Gallivan et al. 2016; Klapetek et al. 2016). However, response selection studies show that if the number of response options increases, the reaction time also increases (Hick 1952; Rosenbaum 1980). This suggests that initiating one prepared movement takes less time than initiating one out of two prepared movements. When two movements need to be performed simultaneously, response time increases can mainly be ascribed to indirect cueing of movements and to a lesser extent to spatial incompatibility of the responses (Diedrichsen et al. 2001; Heuer and Klein 2006; Oliveira and Ivry 2008). Thus, the preparation and execution of multiple movements simultaneously is associated with response time-related costs.

Here we ask if there are also precision costs related to the simultaneous preparation of multiple movements. A central tenet of cognitive psychology is that the brain is limited in its capacity for performing multiple tasks simultaneously (Moray 1967), with an architecture that allows the system to prioritize among these tasks. These constraints are encapsulated in the concept of a shared central resource. Early theoretical and quantitative models formalized this concept into a relation between the amount of resource dedicated to a goal and the chance of a correct response (Kahneman 1973; Shaw 1978;

Norman and Shallice 1980). More recently, work on visual working memory has highlighted the idea that resource sharing has consequences for the precision with which information is internally represented (Palmer 1990; Bays and Husain 2008; Zhang and Luck 2008; van den Berg et al. 2012; Ma et al. 2014). Formally, these models describe the variability with which visual items are represented in memory by a power law: As the number of items increases, the variability associated with the representation of each item increases.

By analogy, if the internal representations of planned movements share a common resource, planning variability should increase as the number of simultaneously-planned movements increases. In speeded movements, this increase in planning noise may be observable as an increase in movement variability early in the trajectory (recognizing that feedforward or feedback control processes may influence latter aspects of the trajectory). We tested this prediction by comparing movement variability in conditions requiring one or two movement plans, in two different contexts: 1) Within a hand (like moving with one hand to pick up the milk carton on the left or the one on the right) and 2) Between different hands (like using the left or right hand to pick up a single carton of milk). We found that movement plans at both levels of competition were more variable if two movements were prepared at the same time than if only one movement was prepared, consistent with the concept of a limited resource for movement planning.

Method

Participants and apparatus

In total, 64 volunteers (21 males, 43 females) aged 19-36 years ($M = 23.9$ years) participated in one of three experiments (16 in Exp. 1; 16 in Exp. 2; 32 in Exp. 3). Sample

size for all experiments was fixed before testing. All participants were naïve to the research question, had normal hearing ability, and normal or corrected-to-normal visual acuity. Five participants were left-handed, the other 59 were right-handed. The study was approved by the UCL Research Ethics Committee.

Participants were seated with their chin on a rest and their index finger(s) touched a surface, slanted 30° from the horizontal plane. Via a mirror, stimuli displayed on a 21" CRT monitor (screen refresh rate 130 Hz) were reflected into the same plane as the movement surface, although participants could not see their hands. Index finger movement was tracked at 133.3 Hz with an electromagnetic tracking device (3D Guidance trakSTAR, Ascension Technology Corporation, Burlington VT, USA).

Experiment 1

Participants triggered the start of each trial by placing the index finger of their preferred hand on a starting point (crosshair) with the aid of visual feedback (a black circle presented at the fingertip location, 0.4 cm diameter, Fig. 1a). After stable finger placement, visual feedback was removed and four tones were played at 600 ms intervals over speakers (Creative Inspire T10), placed to the left and right of the set-up. Together with the first tone, two potential movement targets were presented. The left target location was randomly drawn from a 30° arc with a 13.33 cm radius from the starting point. The center of the arc was 45° counterclockwise with respect to the mid-sagittal plane. The right target was always located 90° in the clockwise direction around the starting point with respect to the left target. Targets were red circles, 1 cm in diameter, both of which turned green at the same time as the third tone. To verify if movements were prepared in advance of the third tone, in 20% of the trials the targets changed position ("jumped") at the same time as the third tone. The new target locations were randomly drawn from the 30° target arc and

were at least 10° away from the original locations. Both targets changed by the same distance in the same direction (clockwise or counterclockwise) so that the targets were always 90° apart.

In this first experiment, the laterality of the tones (left or right) specified which of the two targets was the movement goal. In the *early cue* block (50% of trials) all tones were played from a single speaker, thus specifying the goal early in the trial. In the *late cue* block (50% of trials) the first two tones were played from both speakers and only the last two tones were played from either the left or right speaker (Fig. 1c). In the *late cue* condition, left and right targets were randomly interleaved. In the *early cue* condition, trials to the left and right target were blocked and the blocks were counterbalanced across participants. *Early cued* trials were blocked to maximize the chance that participants would plan a single movement. Participants were instructed not to move before the third tone (the *imperative*) and to hit the specified target by the time of the fourth tone.

The first 10 trials for each cue condition were practice trials during which visual feedback of the index finger position was provided throughout the movement. For the test trials, participants received feedback about the end position of their finger at the end of each response, as well as information concerning whether they were successful on that trial. Success required an endpoint landing inside the target area within 200 ms of the fourth tone. If the movement was initiated before the third tone the trial was aborted and repeated with new parameters (1.7% of trials repeated). Each participant completed 450 trials in total.

Experiment 2

In the second experiment, the tones cued which hand participants had to move toward a single target. There were two starting positions (one for each hand), located 8 cm apart, and a single target, with the location randomly drawn from a 30° arc with a 12.75 cm radius centered halfway between the two starting points (Fig. 1b). The laterality of the tones (left or right) specified which of the two hands to move to the target, with *early* and *late* cued trials tested in separate blocks. Trials were aborted and repeated with new parameters if the movement was initiated before the third tone or if the wrong hand moved (7.2% of trials repeated). All other features of the design were the same as in Exp. 1.

Experiment 3

To disentangle the influences of hand selection and movement planning on movement variability, we incorporated uncertainty about the target location in Experiment 3. We varied the timing of target presentation relative to cue information. The experimental configuration was the same as Exp. 2, but there were no target jumps. In half the trials, the target was displayed at the first tone (*target early*), and in the other half it was displayed at the third tone (*target late*). *Target early* and *late* trials were blocked and counterbalanced between participants. For 16 participants the hand was always cued early (*hand early*; Exp. 3A), and for 16 other participants the hand was always cued late (*hand late*; Exp. 3B). Again, trials were aborted and repeated with new parameters if the movement was initiated before the third tone or if the wrong hand moved (3.3% of trials repeated). Each participant completed 300 trials in total.

Analysis

The fingertip position time series were filtered with a low-pass second-order bidirectional Butterworth filter at 10 Hz. Movement speed was obtained by differentiating these series in the two-dimensional movement plane. Movement initiation was defined as the last sample before peak speed in which the speed was < 2 cm/sec, movement end was defined as the first sample after peak speed in which the speed was < 2 cm/sec, and movement time was defined as the time between movement initiation and movement end. Response time was defined as the time between the third tone and movement initiation. Movement direction was measured as the angle of the fingertip position with respect to the starting position at the time of peak speed (Messier and Kalaska 1999; Churchland et al. 2006a). On average, peak speed occurred at 44% (SEM = 0.4%) of the total movement duration (175 ms, SEM = 4 ms after movement initiation). Direction error was defined as the deviation of movement direction from a straight line toward the target. Movement bias was operationalized as the mean direction error and movement variability as the standard deviation of the direction error.

Practice trials, trials for which there was an error with movement registration, and trials for which offline analysis showed that the online registration had failed to detect movement of the wrong hand or toward the wrong target, or failed to detect that the initial fingertip position was outside the start position, were rejected from analysis ($< 7\%$ of all trials). The data for one participant in Exp. 1 was discarded as he initially moved in a direction in between the two targets on a large number of trials (behavior not seen in other participants).

Movement variability, movement bias, response time and movement time, across items and conditions, were examined with ANOVA and post-hoc t-tests. We applied a Welch degrees of freedom modification for samples of unequal variance. To assess if response time and peak velocity influenced movement variability, we calculated the standard deviation of the initial movement directions for each quartile of the data, based on slowest

to fastest response time or peak velocity for each condition and each participant. Linear regression slopes were fit to the distributions of early- and late-cued no-jump trials for each participant, and we determined if the regression coefficients were different from zero with a one-sample t-test. The same procedure was used with response time and absolute movement bias to analyze if they were correlated.

Results

To examine the influence of the number of movement plans on movement variability, we asked participants to generate rapid reaching movements to one of two visual targets (Fig. 1a). In the *early cue* block, target information was presented well in advance of movement initiation (Fig. 1c), allowing participants to prepare a single movement. In the *late cue* block, this information was only presented at the time movement initiation was signaled; as such, participants should be motivated to prepare movements to both targets given the temporal requirements of the task.

To confirm that movements were prepared in advance of the imperative, the target unpredictably changed location (jumped) at the time of the imperative on 20% of the trials. If participants were waiting until the imperative to plan their movement, this manipulation should not lead to an increase in movement variability. A target (*left or right*) x cue (*early or late*) x jump (*no-jump or jump*) repeated measures ANOVA showed a main effect for jump, indicating that the initial direction of movement was more variable if the target changed position than if it did not ($F(1, 14) = 20.08, p < 0.001$). There was also a significant interaction between cue and jump ($F(1, 14) = 21.56, p < 0.001$), revealing that the increase in movement variability due to the target jump was actually greater for the *early cue* than for the *late cue* (Fig. 2a, empty bars). Pairwise comparisons showed that target jumps increased movement variability in both the *early* ($t(29) = 6.02, p < 0.001$) and *late cue*

conditions ($t(29) = 2.65, p = 0.013$). The increase in movement variability for the target jump condition confirms that participants prepared movements in advance of the imperative.

The key result of this experiment is shown in Fig. 2a (filled bars): initial movement direction in the *early cue* condition was significantly less variable than in the *late cue* condition ($t(14) = 4.31, p < 0.001$; *no-jump* trials only). A target (*left or right*) x target in previous trial (*same side or other side*) repeated measures ANOVA showed that within the *late cue* block for the no jump trials movement variability after the target switched sides ($M = 4.0$ deg) was not different from the variability when the target was on the same side ($M = 4.4$ deg) ($F(1,14) = 2.10, p = 0.169$), indicating that it is unlikely that the difference between the *early* and *late cue* condition is due to the blocked design. Thus, we can conclude that preparing two movements at the same time results in more variable movements than preparing one movement, suggesting a precision cost related to the concurrent planning of multiple movements that could be explained by movement plans sharing a common resource.

The concurrent preparation of multiple movements did not affect movement bias. Fig. 3 shows the probability density distributions of direction errors at the time of peak speed. A target (*left or right*) x cue (*early or late*) x jump (*no-jump or jump*) repeated measures ANOVA showed a main effect for target, indicating that movements towards the left target ($M = 2.7$ deg) were 2.5 degrees more to the right than movements to the right target ($M = 0.2$ deg) ($F(1,14) = 8.74, p = 0.010$). Thus participants were aiming slightly less eccentric than straight to the target. Note that if participants were aiming in between the two targets, their initial direction would be 45 and -45 degrees for the left right target respectively, which was never observed (Fig. 3). There were no other main effects and no interaction effects (all $p > 0.166$), indicating that the movements were similar in the *early* and *late cue* conditions. Movement bias was not correlated with response time. The regression

coefficients for the relation between response time and movement bias were not significantly different from zero for either *early* or *late* cued *no-jump* trials (early: $M = 6.0 \times 10^{-7}$, $t(14) = 0.26$, $p = 0.797$; late: $M = -5.1 \times 10^{-6}$, $t(14) = -1.86$, $p = 0.085$).

Differences between the cue conditions in movement variability were accompanied by differences in response time. A target (*left* or *right*) x cue (*early* or *late*) x jump (*no-jump* or *jump*) repeated measures ANOVA showed main effects for cue and jump (Fig. 2b). Movements were initiated longer after the imperative if the target was cued *late* than if it was cued *early* ($F(1,14) = 22.86$, $p < 0.001$). As illustrated in Fig. 2c, response time was uncorrelated with movement variability: Regression coefficients for the relation between reaction time and movement variability within the *early* and *late* cued *no-jump* trials were not significantly different from zero (early: $M = -2.7 \times 10^{-5}$, $t(14) = 2.07$, $p = 0.057$; late: $M = -4.4 \times 10^{-5}$, $t(14) = 1.60$, $p = 0.131$). Thus, response variability was larger in the *late* cue condition than in the *early* cue condition even when response times are equated. The same was true for peak speed: the regression coefficients for the relation between peak speed and movement variability were not significantly different from zero for either *early* or *late* cued *no-jump* trials (early: $M = -8.5 \times 10^{-5}$, $t(14) = 0.58$, $p = 0.573$; late: $M = -1.0 \times 10^{-4}$, $t(14) = 0.75$, $p = 0.466$).

The main effect of jump reflects the fact that response times were longer if the target changed position than if it did not ($F(1,14) = 14.39$, $p = 0.002$). Together with the quick response times in the *no-jump* trials (205–270 ms), this provides further evidence that participants planned their movements in advance of the imperative. Moreover, in the *early-cue jump* trials reaction time was negatively correlated with movement variability (regression coefficient $M = -3.9 \times 10^{-4}$, $t(14) = 5.65$, $p < 0.001$). A target jump in the *early* cue condition resulted in almost twice as much initial movement variability as *no-jump*, most likely the result of movements with very short response times being initiated in the direction of the target before it jumped. As response times in the *late* cue condition were

longer, the increase in variability is also smaller. This suggests that although movement initiation was slightly delayed in response to a target jump, it was not delayed until the movement plan was completely updated toward the new target position (Haith et al. 2015). This interpretation is in line with the idea that these two processes operate, at least to some extent, independently (Oostwoud Wijdenes et al. 2011).

Competing plans between hands

The results of Exp. 1 suggest that movement plans share a common resource. However, two visual targets presented at the same time will also compete for representational resources in the brain (Kastner and Ungerleider 2000). Prior to a reach, visuospatial attention is directed to the reach target(s) (e.g. Baldauf and Deubel 2010). It is possible that the increase in variability that we observed in Exp. 1 when the target was uncertain is due to noisier representations of the target locations (i.e., divided visuospatial attention). This would not be observed if there was no target uncertainty. In contrast, if the observed variability is related to competing movement plans, variability should also increase when a single target representation is associated with two movement plans, i.e., when the movement may be performed with either the left or right hand (Oliveira et al. 2010). In Exp. 2, participants were presented with a single target and, in separate blocks, were either provided with an *early* cue or a *late* cue specifying the response hand (Fig. 1b).

The results show that planning movements concurrently with two different hands (Exp. 2) has very similar consequences to planning two movements with one hand (Exp. 1). The initial direction of responses was more variable in the *late* cue condition than in the *early* cue condition for the *no-jump* trials ($t(15) = 3.86, p = 0.002$; Fig. 4a, filled bars). Thus we find the same precision-related cost of planning two movements at the same time.

As in Exp. 1, the jump trials confirm that the participants planned their responses in advance of the imperative. The hand (*left* or *right*) x cue x jump repeated measures ANOVA showed that the initial movement direction was more variable for *jump* than for *no-jump* trials ($F(1,15) = 43.45, p < 0.001$). The interaction between cue and jump was also significant, indicating that the effect of the target jump was again larger for *early* than for *late* cue trials ($F(1,15) = 19.91, p < 0.001$). Post-hoc comparisons showed that *jump* trials were more variable than *no-jump* trials for *early*-cued ($t(31) = 7.34, p < 0.001$) and *late*-cued trials ($t(31) = 4.21, p < 0.001$). Thus movements were planned before the imperative, and movement execution was more precise for one planned movement than for two movements planned at the same time.

The movement bias results were also replicated in Exp. 2 (Fig. 5). A hand (*left* or *right*) x cue (*early* or *late*) x jump (*no-jump* or *jump*) repeated measures ANOVA showed a main effect for hand. Movements with the right hand ($M = 1.93$ deg) were more biased towards the right and movements with the left hand were biased to the left ($M = -1.14$ deg) ($F(1,15) = 40.08, p < 0.001$). The interaction effect between hand and jump was close to significant ($F(1,14) = 3.79, p = 0.071$), indicating that the bias for both hands was closer to zero for jump trials than for no-jump trials. The other effects were not significant (all $p > 0.189$). The regression coefficients for the relation between response time and movement bias were also not significantly different from zero for either *early* or *late* cued *no-jump* trials (early: $M = 1.1 \times 10^{-6}, t(15) = 0.69, p = 0.501$; late: $M = 3.1 \times 10^{-7}, t(15) = 0.11, p = 0.915$).

Responses were initiated faster in the *early* than in the *late* cue condition and also faster if the target did not jump than if it jumped ($F(1,15) = 102.9, p < 0.001$ and $F(1,15) = 13.47, p = 0.002$, respectively; see also Fig. 4b). Response times were shorter and peak speeds were lower in the *early* cue condition than in the *late* cue condition. There was no correlation between timing parameters and movement variability in the *no-jump* conditions (regression coefficients for response time [early: $M = -6.0 \times 10^{-6}, t(15) = 0.36, p = 0.726$;

late $M = 3.7 \times 10^{-5}$; $t(15) = 1.32$, $p = 0.206$] and peak speed [early: $M = 6.4 \times 10^{-5}$; $t(15) = 0.36$, $p = 0.722$; late $M = 2.3 \times 10^{-4}$; $t(15) = 1.20$, $p = 0.249$] were not significantly different from zero). Thus, as in Exp. 1, the target jump resulted in delayed movement initiation and higher movement variability.

General preparedness

Our central hypothesis is that the increased variability in two-plan conditions is due to competition between motor plans. We assume that these plans entail some specification of desired trajectories, either in egocentric or joint space (Shadmehr and Mussa-Ivaldi 1994; Gallivan et al. 2015). An alternative possibility is that the differences in variability reflect a more general preparedness of the hand to move that doesn't require trajectory specification, such as hand selection. To examine if hand preparedness influences movement variability, we conducted a third experiment in which we manipulated the timing of the *target* presentation while the hand was cued early (Exp. 3A) or late (Exp. 3B). Given that a trajectory cannot be specified until the target is known, we predicted that the increase in variability associated with multiple plans would only be observed when the target was presented in advance of the imperative signal (*early*), and would not be found when the target was presented at the time of the imperative signal (*late*).

When the target was presented *early*, movements were more variable if the hand was cued *late* than if the hand was cued *early*, replicating the findings of Exp. 2 (between-participants, Welch two sample t-test: $t = 2.21$, $p = 0.031$; Fig. 4a, filled bars). The critical test in this experiment involves the conditions in which the target was presented at the time of the imperative (*late*), given our assumption that this would preclude movement planning. Here there was no effect of hand-cue timing: variability was similar for the hand *early* and hand *late* conditions when the target was presented *late* (Welch two sample t-test: $t = 0.06$, $p = 0.950$; Fig. 4a, hatched bars). Moreover, variability in these conditions

was greater than in the *target early/hand early* condition (*target late/hand early* - paired-samples t-test: $t(31) = 2.36, p = 0.025$; *target late/hand late* - Welch two sample t-test: $t = 2.15, p = 0.036$) suggesting that movement variability is reduced when the forthcoming movement can be completely specified.

Although this was not a straightforward reaction time task, the response time data were congruent with what one would expect based on results of speeded response paradigms (Rosenbaum 1980). Responses were initiated earlier if the target was presented *early* than if it was presented *late*, an effect observed for both hand cued *early* ($t(31) = 3.44, p = 0.002$) and hand cued *late* ($t(31) = 2.21, p = 0.034$), see Fig. 4b. Response times were also faster if the hand was cued *early* than if it was cued *late* (target *early*: $t(36.40) = 5.91, p < 0.001$; target *late*: $t(46.33) = 6.97, p < 0.001$). Thus, in line with previous findings, responses were faster when more information was specified before the imperative, whereas late specification of the hand delayed responses more than late specification of target position.

Discussion

We examined if movement plans share a common resource, by testing if planning variability was affected by the number of movements prepared concurrently. We compared initial movement variability in an *early cue* condition, where preparation should be limited to a single movement, to a *late cue* condition, in which we assumed participants would have to prepare two different movements, given the constraints on reaction time. As predicted, the initial movement direction in the *early cue* condition was less variable than in the *late cue* condition, regardless of whether the two plans corresponded to two different movements of the same hand (Exp. 1; Fig. 2), or movements with the two different hands (Exp. 2; Fig. 4).

To confirm that participants' engaged in advance planning, we included control conditions in which the target changed position at the time of the imperative. If the process of planning a movement in all conditions began only at the imperative, the previous location of the target would be irrelevant and there would be no cost in the target jump conditions. In contrast to this prediction, the initial movement variability and response time increased in the target jump conditions (Figs. 2a and 3a empty bars). Thus, it appears that, consistent with our assumption, movement preparation began before the imperative. The increase in variability and response time observed when the target jumped is likely due to the need to update plans. For very short response times, movements may be initially directed toward the old target location, or to locations between the old and the new target (Van Sonderen and Denier van der Gon 1991; Haith et al. 2015). The increase in variability was larger in the *early cue* condition because here movements were initiated earlier on average, and therefore when target location was more uncertain.

Our results demonstrate a precision cost related to simultaneously preparing multiple movements. This precision cost resembles the cost observed in studies of visual working memory (Ma et al. 2014; Bays 2015), although the increase in variability observed in the present study is somewhat smaller than typically observed in studies of visual working memory. In the working memory studies, variability in recall is found to increase steadily and in a continuous manner as the number of items in memory increases (Palmer 1990; Wilken and Ma 2004; Bays and Husain 2008; Bays et al. 2009; van den Berg et al. 2012). Moreover, salient or goal-relevant items are stored with increased precision, but at a cost to the memorability of other items in memory (Bays and Husain 2008; Bays et al. 2011; Gorgoraptis et al. 2011). These results are consistent with the hypothesis that working memory is constrained by a limited resource that is shared between the representations.

Our results point to a similar constraint for multiple competing movement plans when there are multiple task-relevant targets (Cisek 2006; Chapman et al. 2010; Duque et al. 2010;

Gallivan et al. 2015), or when movements to a single target can be accomplished with either hand (Oliveira et al. 2010). We suggest that the neural mechanisms that cause the representational noise to increase for multiple items in working memory also impacts the fidelity of the representation of multiple movement plans. That is, we propose that neurons face similar energetic constraints when representing multiple visual features or movements, and that the way in which this problem is resolved is similar for the two systems. The variability increase for two competing movement plans may be less than for two competing items in working memory because for reaching, resources can be dynamically reallocated after the late cue. It is also possible that there is a different power law constant for movement planning resources than for working memory resources. Additionally, other factors than the amount of resources dedicated to the movement plan, for which the power law relationship does not hold, might affect movement variability, for example online movement corrections.

A neural model in which feature representations are based on population coding, with the limited shared resource construct implemented by normalization of population activity, can reproduce the specific distributions of errors observed in visual working memory (Bays 2014). As the number of items increases, the firing rates associated with each item decrease, and it becomes harder to distinguish spikes representing individual items from noise; thus working memory variability increases. This model is supported by neurophysiological findings showing that the firing rates correlated with features of remembered items decreases as the number of items increases (Buschman et al. 2011). In addition, in human fMRI, the amplitude of the BOLD response associated with target representations in working memory decreases as the number of represented items increases from one to two (Sprague et al. 2014). Mirroring the findings on firing rates for items in working memory, non-human primate electrophysiology studies show that average firing rates decrease and firing rate variability increases when the number of

response alternatives increases, (Churchland et al. 2011; Churchland and Ditterich 2012). This is consistent with our hypothesis that the same neural constraints are responsible for the increase in planning noise that we observed when multiple plans are prepared simultaneously.

We hypothesize that competition for resources among movement plans takes place in areas associated with movement preparation, such as the premotor cortex and posterior parietal cortex (PPC) (Roland et al. 1980; Snyder et al. 1997; Cui and Andersen 2007). Areas in the intraparietal sulcus and in the premotor cortex have been shown to represent actions independent of the effector that will execute the movement (Medendorp 2004; Gallivan et al. 2013), making them possible loci for competition between motor plans. Top-down allocation of resources could reflect processing in a domain-general area such as prefrontal cortex.

The results of Exp. 3 were especially interesting in showing that there was no difference in variability between *early cue* and *late cue* conditions when the target position was unknown before the imperative (Fig. 4). This result suggests that competition does not arise at the level of a general readiness to move. Instead, the precision benefit for planning a single movement appears to be present only when a specific movement can be prepared. However, the exact level of specification remains to be established: Resources might be distributed across high level representations of motor goals (Wong et al. 2014) or low level fully-specified movement trajectories (Cos et al. 2014; Gallivan et al. 2015).

One might wonder why movements were not more variable in the conditions in which the target was presented late compared to when the target was specified early but the hand was specified late. In the former, it was not possible to plan anything before the imperative, whereas in the latter, there were two competing plans. We speculate that this is related to

the differences in response times, with movement initiation delayed until variance drops below a criterion value (Churchland et al. 2006b).

Besides uncovering a precision related cost associated with concurrently planned movements, we also replicated the well-known response time related cost. Response time was aided by early specification of parameters of the movement. Responses were initiated sooner after the imperative in the *early cue* than in the *late cue* condition, and also sooner in the *target presented early* than *late* condition (Figs. 2b, 4b and 6b). This is in accordance with seminal studies showing that reaction times are faster if there are fewer possible targets, and if more information is specified before the imperative (Hick 1952; Goodman and Kelso 1980; Rosenbaum 1980). Identification of the side that the tone was coming from might have additionally affected the response times in the *late cue* condition. The relationship between the number of cues and reaction time has been formalized in a capacity-sharing model with similarities to a shared resource account (Pellizzer and Hedges 2003).

In previous studies in which the target was specified only *after* the imperative, it has been reported that the movement initially follows a path that is the average of movement paths toward all of the possible targets (Chapman et al. 2010; Stewart et al. 2014). In our study, movement averaging could in theory have contributed to the observed variability in Exp. 1. However, at odds with this hypothesis, movement biases were not affected by cue condition. It is also not consistent with the results of Exp. 2 where the difference between movement plans was based on the number of candidate hands (1 or 2) for the forthcoming response.

In summary, we propose that the level of planning noise, manifest in the form of movement variability (Afshar and Shenoy 2006; Churchland et al. 2006a; van Beers 2009; Chaisanguanthum et al. 2014), is a function of the amount of resources dedicated to a

motor plan. Similar to working memory, resources may be allocated flexibly, depending on the number of alternative movement plans. Importantly, our results indicate a precision constraint on the ability to prepare multiple movements simultaneously. If too many movements are prepared at the same time, each representation will suffer from high levels of noise. Therefore, preparing a small number of plans at one time may be optimal for generating fast, accurate movements.

Grants

This research was supported by the Wellcome Trust and the Van Coeverden Adriani Stichting.

Disclosures

No conflicts of interest, financial or otherwise, are declared by the authors.

References

- Afshar A, Shenoy KV.** A central source of movement variability. *Neuron* 52: 1085–1096, 2006.
- Baldauf D, Deubel H.** Attentional landscapes in reaching and grasping. *Vision Research* 50: 999–1013, 2010.
- Bays PM.** Noise in neural populations accounts for errors in working memory. *J Neurosci* 34: 3632–3645, 2014.
- Bays PM.** Spikes not slots: noise in neural populations limits working memory. *Trends in Cognitive Sciences* (July 2015). doi: 10.1016/j.tics.2015.06.004.
- Bays PM, Catalao RFG, Husain M.** The precision of visual working memory is set by allocation of a shared resource. *Journal of Vision* 9: 7–7, 2009.
- Bays PM, Gorgoraptis N, Wee N, Marshall L, Husain M.** Temporal dynamics of encoding, storage, and reallocation of visual working memory. *Journal of Vision* 11: 6–6, 2011.
- Bays PM, Husain M.** Dynamic Shifts of Limited Working Memory Resources in Human Vision. *Science* 321: 851–854, 2008.
- van Beers RJ.** Motor learning is optimally tuned to the properties of motor noise. *Neuron* 63: 406–417, 2009.
- van den Berg R, Shin H, Chou W-C, George R, Ma WJ.** Variability in encoding precision accounts for visual short-term memory limitations. *Proc Natl Acad Sci USA* 109: 8780–8785, 2012.
- Buschman TJ, Siegel M, Roy JE, Miller EK.** Neural substrates of cognitive capacity limitations. *Proceedings of the National Academy of Sciences* 108: 11252–11255, 2011.
- Chaisanguanthum KS, Shen HH, Sabes PN.** Motor variability arises from a slow random walk in neural state. *J Neurosci* 34: 12071–12080, 2014.
- Chapman CS, Gallivan JP, Wood DK, Milne JL, Culham JC, Goodale MA.** Reaching for the unknown: Multiple target encoding and real-time decision-making in a rapid reach task. *Cognition* 116: 168–176, 2010.
- Churchland AK, Ditterich J.** New advances in understanding decisions among multiple alternatives. *Current Opinion in Neurobiology* 22: 920–926, 2012.
- Churchland AK, Kiani R, Chaudhuri R, Wang X-J, Pouget A, Shadlen MN.** Variance as a Signature of Neural Computations during Decision Making. *Neuron* 69: 818–831, 2011.
- Churchland MM, Afshar A, Shenoy KV.** A central source of movement variability. *Neuron* 52: 1085–1096, 2006a.

- Churchland MM, Yu BM, Ryu SI, Santhanam G, Shenoy KV.** Neural variability in premotor cortex provides a signature of motor preparation. *J Neurosci* 26: 3697–3712, 2006b.
- Cisek P.** Integrated neural processes for defining potential actions and deciding between them: a computational model. *J Neurosci* 26: 9761–9770, 2006.
- Cisek P.** Cortical mechanisms of action selection: the affordance competition hypothesis. *Philosophical Transactions of the Royal Society B: Biological Sciences* 362: 1585–1599, 2007.
- Cisek P, Kalaska JF.** Neural correlates of reaching decisions in dorsal premotor cortex: specification of multiple direction choices and final selection of action. *Neuron* 45: 801–814, 2005.
- Cisek P, Kalaska JF.** Neural mechanisms for interacting with a world full of action choices. *Annu Rev Neurosci* 33: 269–298, 2010.
- Cos I, Duque J, Cisek P.** Rapid prediction of biomechanical costs during action decisions. *Journal of Neurophysiology* 112: 1256–1266, 2014.
- Cui H, Andersen RA.** Posterior Parietal Cortex Encodes Autonomously Selected Motor Plans. *Neuron* 56: 552–559, 2007.
- Diedrichsen J, Hazeltine E, Kennerley S, Ivry RB.** Moving to directly cued locations abolishes spatial interference during bimanual actions. *Psychol Sci* 12: 493–498, 2001.
- Duque J, Lew D, Mazzocchio R, Olivier E, Ivry RB.** Evidence for Two Concurrent Inhibitory Mechanisms during Response Preparation. *Journal of Neuroscience* 30: 3793–3802, 2010.
- Gallivan JP, Barton KS, Chapman CS, Wolpert DM, Flanagan JR.** Action plan co-optimization reveals the parallel encoding of competing reach movements. *Nature Communications* 6: 7428, 2015.
- Gallivan JP, Logan L, Wolpert DM, Flanagan JR.** Parallel specification of competing sensorimotor control policies for alternative action options. *Nature Neuroscience* (January 11, 2016). doi: 10.1038/nn.4214.
- Gallivan JP, McLean DA, Flanagan JR, Culham JC.** Where One Hand Meets the Other: Limb-Specific and Action-Dependent Movement Plans Decoded from Preparatory Signals in Single Human Frontoparietal Brain Areas. *Journal of Neuroscience* 33: 1991–2008, 2013.
- Ghez C, Favilla M, Ghilardi MF, Gordon J, Bermejo R, Pullman S.** Discrete and continuous planning of hand movements and isometric force trajectories. *Exp Brain Res* 115: 217–233, 1997.
- Goodman D, Kelso JA.** Are movements prepared in parts? Not under compatible (naturalized) conditions. *J Exp Psychol Gen* 109: 475–495, 1980.
- Gorgoraptis N, Catalao RFG, Bays PM, Husain M.** Dynamic Updating of Working Memory Resources for Visual Objects. *Journal of Neuroscience* 31: 8502–8511, 2011.

- Greent't-Jong T, Oostenveld R, Jensen O, Medendorp WP, Praamstra P.** Competitive interactions in sensorimotor cortex: oscillations express separation between alternative movement targets. *Journal of Neurophysiology* 112: 224–232, 2014.
- Haith AM, Huberdeau DM, Krakauer JW.** Hedging your bets: intermediate movements as optimal behavior in the context of an incomplete decision. *PLoS Comput Biol* 11: e1004171, 2015.
- Heuer H, Klein W.** The influence of movement cues on intermanual interactions. *Psychological Research Psychologische Forschung* 70: 229–244, 2006.
- Hick WE.** On the rate of gain of information. *Quarterly Journal of Experimental Psychology* 4: 11–26, 1952.
- Kahneman D.** *Attention and effort*. Englewood Cliffs, NJ: Prentice-Hall, 1973.
- Kastner S, Ungerleider LG.** Mechanisms of visual attention in the human cortex. *Annu Rev Neurosci* 23: 315–341, 2000.
- Klapetek A, Jonikaitis D, Deubel H.** Attention allocation before antisaccades. *Journal of Vision* 16: 11, 2016.
- Klein-Flugge MC, Bestmann S.** Time-Dependent Changes in Human Corticospinal Excitability Reveal Value-Based Competition for Action during Decision Processing. *Journal of Neuroscience* 32: 8373–8382, 2012.
- Kornblum S, Hasbroucq T, Osman A.** Dimensional overlap: cognitive basis for stimulus-response compatibility--a model and taxonomy. *Psychol Rev* 97: 253–270, 1990.
- Ma WJ, Husain M, Bays PM.** Changing concepts of working memory. *Nat Neurosci* 17: 347–356, 2014.
- Medendorp WP.** Integration of Target and Effector Information in Human Posterior Parietal Cortex for the Planning of Action. *Journal of Neurophysiology* 93: 954–962, 2004.
- Messier J, Kalaska JF.** Comparison of variability of initial kinematics and endpoints of reaching movements. *Exp Brain Res* 125: 139–152, 1999.
- Moray N.** Where is capacity limited? A survey and a model. *Acta Psychol (Amst)* 27: 84–92, 1967.
- Nashed JY, Crevecoeur F, Scott SH.** Rapid Online Selection between Multiple Motor Plans. *Journal of Neuroscience* 34: 1769–1780, 2014.
- Norman DA, Shallice T.** Attention to action: Willed and automatic control of behavior. *CHIP Report* 99. .
- Oliveira FTP, Diedrichsen J, Verstynen T, Duque J, Ivry RB.** Transcranial magnetic stimulation of posterior parietal cortex affects decisions of hand choice. *Proceedings of the National Academy of Sciences* 107: 17751–17756, 2010.

- Oliveira FTP, Ivry RB.** The Representation of Action: Insights From Bimanual Coordination. *Curr Dir Psychol Sci* 17: 130–135, 2008.
- Oostwoud Wijdenes L, Brenner E, Smeets JBJ.** Fast and fine-tuned corrections when the target of a hand movement is displaced. *Experimental Brain Research* 214: 453–462, 2011.
- Palmer J.** Attentional limits on the perception and memory of visual information. *J Exp Psychol Hum Percept Perform* 16: 332–350, 1990.
- Pellizzer G, Hedges JH.** Motor planning: effect of directional uncertainty with discrete spatial cues. *Exp Brain Res* 150: 276–289, 2003.
- Prescott TJ, Redgrave P, Gurney K.** Layered Control Architectures in Robots and Vertebrates. *Adaptive Behavior* 7: 99–127, 1999.
- Roland PE, Larsen B, Lassen NA, Skinhøj E.** Supplementary motor area and other cortical areas in organization of voluntary movements in man. *J Neurophysiol* 43: 118–136, 1980.
- Rosenbaum DA.** Human movement initiation: specification of arm, direction, and extent. *J Exp Psychol Gen* 109: 444–474, 1980.
- Shadmehr R, Mussa-Ivaldi FA.** Adaptive representation of dynamics during learning of a motor task. *J Neurosci* 14: 3208–3224, 1994.
- Shaw ML.** A capacity allocation model for reaction time. *Journal of Experimental Psychology: Human Perception and Performance* 4: 586–598, 1978.
- Snyder LH, Batista AP, Andersen RA.** Coding of intention in the posterior parietal cortex. *Nature* 386: 167–170, 1997.
- Song J-H, Nakayama K.** Hidden cognitive states revealed in choice reaching tasks. *Trends in Cognitive Sciences* 13: 360–366, 2009.
- Sprague TC, Ester EF, Serences JT.** Reconstructions of Information in Visual Spatial Working Memory Degrade with Memory Load. *Current Biology* 24: 2174–2180, 2014.
- Stewart BM, Gallivan JP, Baugh LA, Flanagan JR.** Motor, not visual, encoding of potential reach targets. *Curr Biol* 24: R953–954, 2014.
- Van Sonderen JF, Denier van der Gon JJ.** Reaction-time-dependent differences in the initial movement direction of fast goal-directed arm movements. *Human Movement Science* 10: 713–726, 1991.
- Wilken P, Ma WJ.** A detection theory account of change detection. *J Vis* 4: 1120–1135, 2004.
- Wong AL, Haith AM, Krakauer JW.** Motor Planning. *Neuroscientist* (June 30, 2014). doi: 10.1177/1073858414541484.
- Zhang W, Luck SJ.** Discrete fixed-resolution representations in visual working memory. *Nature* 453: 233–235, 2008.

Figures




Fig. 1. Experimental configurations. Start point(s) and possible target locations (30° arcs) in (a) Exp. 1, in which auditory cues indicate which of two targets (red) to reach to, and (b) Exps. 2 and 3, in which cues indicated the response hand. Timing for the two cueing conditions (c). In these examples the cue indicates a movement to the left target or with the left hand respectively. In the *early cue* condition (top) this information is provided at the time of the first tone (and coincides with target onset in Exp.1, 2 and the target

presented early condition in 3). In the *late cue* condition (bottom), this information is only provided at the onset of the imperative. In the target presented late condition in Exp.3 the target was presented at the same time as the imperative.

Exp. 1 - target is cued




Fig. 2. Experiment 1: Effect of number of movement plans (within hand). Movement variability (a) and response times (b) for target cued early or late. Movement variability as a function of (c) response time and (d) peak speed for the different conditions. Error bars indicate ± 1 SE.




Fig. 3. Probability density distributions for direction error in Exp. 1 for *early cue – no jump* (a), *late cue – no jump* (b), *early cue – jump* (c) and *late cue – jump* (d) conditions. Direction errors to the right target are in dark grey and to the left target in light grey.

Exp. 2 - hand is cued




Fig. 4. Experiment 2: Effect of number of movement plans between hands. Movement variability (a) and response times (b) for hand cued early or late. Movement variability as a function of (c) response time and (d) peak speed for the different conditions.




Fig. 5. Probability density distributions for direction error in Exp. 2 for *early cue – no jump* (a), *late cue – no jump* (b), *early cue – jump* (c) and *late cue – jump* (d) conditions. Direction errors with the right hand are in dark grey and with the left hand in light grey.




Fig. 6. Experiment 3: Effect of general preparedness. Movement variability (a) and response time (b) for Exp. 3A, in which there was no hand uncertainty because the hand was cued early (purple), and Exp. 3B, in which movements with both hands had to be prepared because the hand was cued late (orange). Filled bars indicate conditions in which there was no target uncertainty because the target was presented at the first tone (*early*), as in the no-jump condition of Exp. 2. Hatched bars indicate conditions in which the target was presented at the time of the imperative (*late*).

Table

Table 1. Properties of the experiments.

	What is cued?	Timing cue	Timing target(s)	Target jumps
Exp. 1	target	early/late	early	20%
Exp. 2	hand	early/late	early	20%
Exp. 3A	hand	early	early/late	no
Exp. 3B	hand	late	early/late	no