

ANALYSIS & PDE

Volume 9

No. 8

2016

GABRIEL P. PATERNAIN AND HANMING ZHOU

**INVARIANT DISTRIBUTIONS AND THE GEODESIC RAY
TRANSFORM**

INVARIANT DISTRIBUTIONS AND THE GEODESIC RAY TRANSFORM

GABRIEL P. PATERNAIN AND HANMING ZHOU

We establish an equivalence principle between the solenoidal injectivity of the geodesic ray transform acting on symmetric m -tensors and the existence of invariant distributions or smooth first integrals with prescribed projection over the set of solenoidal m -tensors. We work with compact simple manifolds, but several of our results apply to nontrapping manifolds with strictly convex boundary.

1. Introduction

The present paper studies the geodesic ray transform of a compact simply connected Riemannian manifold with no conjugate points and strictly convex boundary. Our main objective is to establish an equivalence principle between injectivity of the ray transform acting on solenoidal symmetric m -tensors and the existence of solutions to the transport equation (associated with the geodesic vector field) with prescribed projection over the set of solenoidal m -tensors.

The Radon transform in the plane is the most fundamental example of the geodesic ray transform. It packs the integrals of a function f in \mathbb{R}^2 over straight lines:

$$Rf(s, \omega) = \int_{-\infty}^{\infty} f(s\omega + t\omega^\perp) dt, \quad s \in \mathbb{R}, \omega \in S^1.$$

Here ω^\perp is the rotation of ω by 90 degrees counterclockwise. The properties of this transform are well studied [Helgason 1999] and constitute the theoretical underpinnings for many medical imaging methods such as CT and PET. Generalizations of the Radon transform are often needed. In seismic and ultrasound imaging one finds ray transforms where the measurements are given by integrals over more general families of curves, often modeled as the geodesics of a Riemannian metric. Moreover, integrals of tensor fields over geodesics are ubiquitous in rigidity questions in differential geometry and dynamics.

In this paper we will relate the injectivity properties of the geodesic ray transform with a well-studied subject in classical mechanics: the existence of special first integrals of motion along geodesics. Some Riemannian metrics admit distinguished first integrals; e.g., the geodesic flow of an ellipsoid in \mathbb{R}^3 admits a nontrivial first integral which is quadratic in momenta. As recently shown in [Kruglikov and Matveev 2016], a generic metric does not admit a nontrivial first integral that is polynomial in momenta, but here we will show a complementary statement going in the opposite direction: from the injectivity of the geodesic ray transform on tensors, we will show that it is possible to construct a smooth first integral with *any* prescribed polynomial part. In other words, given a polynomial F of degree m in momenta

MSC2010: primary 53C65; secondary 58J40.

Keywords: geodesic ray transform, first integral, tensor tomography, invariant distribution.

satisfying a natural restriction condition (related with the transport equation, see Section 7), we will show that we can find a smooth function G whose dependence on momenta is of order $> m$ such that $F + G$ is a first integral of the geodesic flow. Generically G is nonvanishing and *not* polynomial in momenta.

Let us now explain our results in more detail. The geodesic ray transform acts on functions defined on the unit sphere bundle of a compact oriented n -dimensional Riemannian manifold (M, g) with boundary ∂M ($n \geq 2$). Let SM denote the unit sphere bundle on M ; i.e.,

$$SM := \{(x, \xi) \in TM : \|\xi\|_g = 1\}.$$

We define the volume form on SM by $d\Sigma^{2n-1}(x, \xi) = |dV^n(x) \wedge d\Omega_x(\xi)|$, where dV^n is the volume form on M and $d\Omega_x(\xi)$ is the volume form on the fiber S_xM . The boundary of SM is given by $\partial SM := \{(x, \xi) \in SM : x \in \partial M\}$. On ∂SM the natural volume form is $d\Sigma^{2n-2}(x, \xi) = |dV^{n-1}(x) \wedge d\Omega_x(\xi)|$, where dV^{n-1} is the volume form on ∂M . We define two subsets of ∂SM ,

$$\partial_{\pm} SM := \{(x, \xi) \in \partial SM : \pm \langle \xi, \nu(x) \rangle_g \leq 0\},$$

where $\nu(x)$ is the outward unit normal vector on ∂M at x . It is easy to see that

$$\partial_+ SM \cap \partial_- SM = S(\partial M).$$

Given $(x, \xi) \in SM$, we denote by $\gamma_{x,\xi}$ the unique geodesic with $\gamma_{x,\xi}(0) = x$ and $\dot{\gamma}_{x,\xi}(0) = \xi$ and let $\tau(x, \xi)$ be the first time when the geodesic $\gamma_{x,\xi}$ exits M .

We say that (M, g) is *nontrapping* if $\tau(x, \xi) < \infty$ for all $(x, \xi) \in SM$.

Definition 1.1. The *geodesic ray transform* of a function $f \in C^\infty(SM)$ is the function

$$If(x, \xi) = \int_0^{\tau(x,\xi)} f(\gamma_{x,\xi}(t), \dot{\gamma}_{x,\xi}(t)) dt, \quad (x, \xi) \in \partial_+ SM.$$

Note that if the manifold (M, g) is nontrapping and has strictly convex boundary, then $I : C^\infty(SM) \rightarrow C^\infty(\partial_+ SM)$, and Santaló’s formula (see Section 2) implies that I is also a bounded map $L^2(SM) \rightarrow L^2_\mu(\partial_+ SM)$, where $d\mu(x, \xi) = |\langle \nu(x), \xi \rangle| d\Sigma^{2n-2}(x, \xi)$ and $L^2_\mu(\partial_+ SM)$ is the space of functions on $\partial_+ SM$ with inner product

$$(u, v)_{L^2_\mu(\partial_+ SM)} = \int_{\partial_+ SM} u \bar{v} d\mu.$$

Given $f \in C^\infty(SM)$, what properties of f may be determined from the knowledge of If ? Clearly a general function f on SM is not determined by its geodesic ray transform alone, since f depends on more variables than If . In applications one often encounters the transform I acting on special functions on SM that arise from symmetric tensor fields, and we will now consider this case.

We denote by $C^\infty(S^m(T^*M))$ the space of smooth covariant symmetric tensor fields of rank m on M with L^2 inner product

$$(u, v) := \int_M u_{i_1 \dots i_m} \overline{v^{i_1 \dots i_m}} dV^n,$$

where $v^{i_1 \dots i_m} = g^{i_1 j_1} \dots g^{i_m j_m} v_{j_1 \dots j_m}$. There is a natural map

$$\ell_m : C^\infty(S^m(T^*M)) \rightarrow C^\infty(SM)$$

given by $\ell_m(f)(x, \xi) := f_x(\xi, \dots, \xi)$. We can now define the geodesic ray transform acting on symmetric m -tensors simply by setting $I_m := I \circ \ell_m$. Let $d = \sigma \nabla$ be the symmetric inner differentiation, where ∇ is the Levi-Civita connection associated with g , and σ denotes symmetrization. It is easy to check that if $v = dp$ for some $p \in C^\infty(S^{m-1}(T^*M))$ with $p|_{\partial M} = 0$, then $I_m v = 0$. The tensor tomography problem asks the following question: are such tensors the only obstructions for I_m to be injective? If this is the case, then we say I is *solenoidal injective* or *s-injective* for short. The problem is wide open for compact nontrapping manifolds with strictly convex boundary (but see [Uhlmann and Vasy 2016; Stefanov et al. 2014]). There are more results if one assumes the stronger condition of being *simple*, i.e., (M, g) is simply connected, has no conjugate points and strictly convex boundary. For simple surfaces, the tensor tomography problem has been completely solved [Paternain et al. 2013]. For simple manifolds of any dimension, solenoidal injectivity is known for I_0 and I_1 [Muhometov 1977; Anikonov and Romanov 1997]. For m -tensors, $m \geq 2$, the tensor tomography problem is still open, but some substantial partial results were established under additional assumptions; see, e.g., [Pestov and Sharafutdinov 1988; Sharafutdinov 1994; Stefanov and Uhlmann 2005; Paternain et al. 2015a; Stefanov et al. 2014].

Let us explain a bit further the term “solenoidal injective”. Consider the Sobolev space $H^k(S^m(T^*M))$ naturally associated with the L^2 inner product defined above. By [Sharafutdinov 1994; Sharafutdinov et al. 2005], there is an orthogonal decomposition of L^2 symmetric tensors fields. Given $v \in H^k(S^m(T^*M))$, $k \geq 0$, there exist uniquely determined $v^s \in H^k(S^m(T^*M))$ and $p \in H^{k+1}(S^{m-1}(T^*M))$ such that

$$v = v^s + dp, \quad \delta v^s = 0, \quad p|_{\partial M} = 0,$$

where δ is the divergence. We call v^s and dp the *solenoidal* part and *potential* part of v respectively. Moreover, we denote by $H^k(S_{\text{sol}}^m(T^*M))$ and $C^\infty(S_{\text{sol}}^m(T^*M))$ the subspaces of $H^k(S^m(T^*M))$ and $C^\infty(S^m(T^*M))$ respectively whose elements are solenoidal symmetric tensor fields. Solenoidal injectivity of I_m simply means that I_m is injective when restricted to $C^\infty(S_{\text{sol}}^m(T^*M))$.

Let I^* denote the adjoint of I using the L^2 inner products defined above; that is,

$$(Iu, \varphi) = (u, I^* \varphi)$$

for $u \in L^2(SM)$, $\varphi \in L^2_\mu(\partial_+ SM)$. A simple application of Santaló’s formula yields

$$I^* \varphi = \varphi^\sharp,$$

where $\varphi^\sharp(x, \xi) := \varphi(\gamma_{x, \xi}(-\tau(x, -\xi)), \dot{\gamma}_{x, \xi}(-\tau(x, -\xi)))$ (see Section 2 for details). Observe that by definition, φ^\sharp is constant along orbits of the geodesic flow. If we are now interested in I_m^* , we note that

$$I_m^* = \ell_m^* \circ I^*$$

and hence we just need to compute ℓ_m^* . This is easy (see Section 2) and one finds

$$L_m f := \ell_m^* f(x)_{i_1 \dots i_m} := g_{i_1 j_1} \dots g_{i_m j_m} \int_{S_x M} f(x, \xi) \xi^{j_1} \dots \xi^{j_m} d\Omega_x(\xi).$$

The fundamental microlocal property of the geodesic ray transform is that, for simple manifolds, $I_m^* I_m$ is a pseudodifferential operator of order -1 on a slightly larger open manifold engulfing M . Moreover, it has a suitable ellipticity property when acting on solenoidal tensors [Sharafutdinov et al. 2005]. This has been exploited to great effect to derive surjectivity of I_m^* knowing injectivity of I_m [Pestov and Uhlmann 2005; Dairbekov and Uhlmann 2010] for $m = 0, 1$. Since the range of I_m^* is contained in the space of solenoidal tensors, by saying I_m^* is *surjective* we mean that the range of I_m^* equals the latter. Surjectivity of I_m^* for tensors of order 0 and 1 has been the key for the recent success in the solution of several long standing questions in 2D [Salo and Uhlmann 2011; Pestov and Uhlmann 2005; Paternain et al. 2012; 2013; 2014; Guillarmou 2014]. However, very little is known about surjectivity for $m \geq 2$ and this largely motivates the present paper.

The surjectivity properties of the adjoint of the geodesic ray transform reveal themselves in the existence of solutions f to the transport equation $Xf = 0$ with prescribed values for $L_m f$ in the space of solenoidal tensors. Here X is the geodesic vector field acting on distributions by duality (recall that X preserves the volume form $d\Sigma^{2n-1}$). A distribution f on SM is said to be *invariant* if it satisfies $Xf = 0$. As we already mentioned, in this paper we mainly study the relation among the injectivity of I_m , the surjectivity of its adjoint I_m^* on solenoidal tensor fields and the existence of some invariant distributions or smooth first integrals associated with solenoidal tensor fields. On a compact nontrapping manifold with strictly convex boundary, the geodesic ray transform I_m is extendable to a bounded operator

$$I_m : H^k(S^m(T^*M)) \rightarrow H^k(\partial_+ SM)$$

for all $k \geq 0$ [Sharafutdinov 1994, Theorem 4.2.1]. Moreover, it can be easily checked that

$$I_m(H_0^k(S^m(T^*M))) \subset H_0^k(\partial_+ SM)$$

and hence we can define I_m^* by duality acting on negative Sobolev spaces to obtain a bounded operator:

$$I_m^* : H^{-k}(\partial_+ SM) \rightarrow H^{-k}(S^m(T^*M)).$$

In other words, for $\varphi \in H^{-k}(\partial_+ SM)$, we have $I_m^* \varphi$ is defined by $(I_m^* \varphi, u) = (\varphi, I_m u)$ for all $u \in H_0^k(S^m(T^*M))$. Let $C_\alpha^\infty(\partial_+ SM)$ denote the set of smooth functions φ for which φ^\sharp is also smooth.

Our main result is the following theorem:

Theorem 1.2. *Let M be a compact simple Riemannian manifold. Then the following are equivalent:*

- (1) I_m is s -injective on $C^\infty(S^m(T^*M))$.
- (2) For every $u \in L^2(S_{\text{sol}}^m(T^*M))$, there exists $\varphi \in H^{-1}(\partial_+ SM)$ such that $u = I_m^* \varphi$.
- (3) For every $u \in L^2(S_{\text{sol}}^m(T^*M))$, there exists $f \in H^{-1}(SM)$ satisfying $Xf = 0$ and $u = L_m f$.
- (4) For every $u \in C^\infty(S_{\text{sol}}^m(T^*M))$, there exists $\varphi \in C_\alpha^\infty(\partial_+ SM)$ such that $u = I_m^* \varphi$.
- (5) For every $u \in C^\infty(S_{\text{sol}}^m(T^*M))$, there exists $f \in C^\infty(SM)$ with $Xf = 0$ such that $L_m f = u$.

We observe that by [Sharafutdinov et al. 2005, Theorem 1.1], s -injectivity of I_m on $L^2(S^m(T^*M))$ is equivalent to s -injectivity of I_m on $C^\infty(S^m(T^*M))$.

Let us return to the subject of special first integrals associated with the geodesic flow. By considering the vertical Laplacian Δ on each fiber $S_x M$ of SM , we have a natural L^2 decomposition $L^2(SM) = \bigoplus_{m \geq 0} H_m(SM)$ into vertical spherical harmonics. We set $\Omega_m := H_m(SM) \cap C^\infty(SM)$. Then a function u belongs to Ω_m if and only if $\Delta u = m(m + n - 2)u$, where $n = \dim M$. The maps

$$\ell_m : C^\infty(S^m(T^*M)) \rightarrow \bigoplus_{k=0}^{[m/2]} \Omega_{m-2k}$$

and

$$L_m : \bigoplus_{k=0}^{[m/2]} \Omega_{m-2k} \rightarrow C^\infty(S^m(T^*M))$$

are isomorphisms. These maps give natural identification between functions in Ω_m and *trace-free* symmetric m -tensors (for details on this, see [Guillemin and Kazhdan 1980b; Dairbekov and Sharafutdinov 2010; Paternain et al. 2015a]). If (M, g) is a simple manifold with I_m s -injective, Theorem 1.2(5) says that given any $u \in C^\infty(S^m_{\text{sol}}(T^*M))$ there is a first integral of the geodesic flow f such that $L_m f = u$. In other words, if we let $F = L_m^{-1}u \in \bigoplus_{k=0}^{[m/2]} \Omega_{m-2k}$ and $G = f - F$, we see that F is polynomial of degree m in velocities and it can be completed by adding G to obtain a first integral. We also see that (taking the even or odd part of f if necessary) $G \in \bigoplus_{k \geq 1} \Omega_{m+2k}$. These were the functions mentioned earlier in the introduction. If G were to be zero, then there would be a first integral that is polynomial in velocities and generically these do not exist. We note that the paper [Paternain et al. 2015a] also constructs invariant distributions (they are not smooth in general) with prescribed m -th polynomial component using a different method (a Beurling transform), but it requires nonpositive curvature for it to work. As already mentioned, here we use instead the normal operator $I_m^* I_m$.

The results in [Pestov and Uhlmann 2005; Dairbekov and Uhlmann 2010] prove that (1) implies (4) or (5) in Theorem 1.2 for $m = 0, 1$, so the main contribution in the theorem is to cover the case $m \geq 2$ and also to provide additional invariant distributions associated with L^2 solenoidal tensors. The proof of Theorem 1.2 relies on a solenoidal extension of tensor fields. For $m = 0$ no extension is needed and for $m = 1$ the situation is considerably simpler and an extension result is already available in [Kato et al. 2000]. Paradoxically the need for a solenoidal extension does not arise in the more complicated setting of Anosov manifolds since there is no boundary. In this setting, an analogous result to Theorem 1.2 (in the L^2 setting) has been recently proved by C. Guillarmou [2014, Corollary 3.7] and these ideas gave rise to a full solution to the tensor tomography problem on an Anosov surface.

Since in 2D the tensor tomography problem has been fully solved [Paternain et al. 2013], we derive:

Corollary 1.3. *Let (M, g) be a compact simple surface. For every $u \in C^\infty(S^m_{\text{sol}}(T^*M))$, there exists $f \in C^\infty(SM)$ with $Xf = 0$ such that $L_m f = u$.*

We shall also give an alternative proof of the corollary using results from [Paternain et al. 2015b]. The alternative proof avoids the smooth solenoidal extension and sheds some light on the relationship between the transport equation and the solenoidal condition.

The rest of the paper is organized as follows. Section 2 contains some preliminaries. In Section 3 we establish the L^2 and C^∞ compactly supported solenoidal extension of tensor fields. This necessitates at some point the use of the generic nonexistence of nontrivial Killing tensor fields recently proved in [Kruglikov and Matveev 2016]. Section 4 uses the well-established microlocal analysis to prove a surjectivity result for $I_m^* I_m$ following the strategy in [Dairbekov and Uhlmann 2010]. Section 5 establishes various boundedness properties on Sobolev spaces that allow us to extend the relevant operators to negative Sobolev spaces (i.e., distributions). Section 6 bundles up everything together and proves Theorem 1.2. Section 7 gives an alternative proof of Corollary 1.3 and clarifies the connection between solenoidal tensors and the transport equation.

2. Preliminaries

In this section we provide details about the regularity properties of the operators introduced in the previous section. First we describe the basic notation we will use frequently in the rest of the paper. Given a compact Riemannian manifold M with boundary, we define

$$C_c^\infty(\mathcal{M}^{\text{int}}) := \{f \in C^\infty(M) : \text{supp } f \subset \mathcal{M}^{\text{int}}\},$$

$$H^k(\mathcal{M}^{\text{int}}) := \{f \in H^k(M) : \text{supp } f \subset \mathcal{M}^{\text{int}}\} \quad \text{for } k \in \mathbb{Z}.$$

Then for any $s > 0, s \in \mathbb{Z}$, we say $H_0^s(M)$ is the completion of $C_c^\infty(\mathcal{M}^{\text{int}})$ under the H^s norm.

Now let M be a compact manifold. Given $f \in C^\infty(SM)$ and $u \in C^\infty(S^m(T^*M))$, we have

$$(\ell_m u, f) = \int_{SM} u_{j_1 \dots j_m}(x) \xi^{j_1} \dots \xi^{j_m} f(x, \xi) d\Sigma^{2n-1}$$

$$= \int_M u_{j_1 \dots j_m}(x) \int_{S_x M} f(x, \xi) \xi^{j_1} \dots \xi^{j_m} d\Omega_x(\xi) dV^n(x).$$

This means that

$$L_m = \ell_m^* : C^\infty(SM) \rightarrow C^\infty(S^m(T^*M))$$

is given by

$$L_m f(x)_{i_1 \dots i_m} = g_{i_1 j_1} \dots g_{i_m j_m} \int_{S_x M} f(x, \xi) \xi^{j_1} \dots \xi^{j_m} d\Omega_x(\xi).$$

Since the metric tensor g is smooth, for the sake of simplicity, we identify $L_m f$ with its dual,

$$L_m f(x)^{j_1 \dots j_m} = \int_{S_x M} f(x, \xi) \xi^{j_1} \dots \xi^{j_m} d\Omega_x(\xi).$$

On the other hand, it is easy to see that the map ℓ_m can be extend to the bounded operator

$$\ell_m : H^k(S^m(T^*M)) \rightarrow H^k(SM)$$

for any integer $k \geq 0$. In particular $\ell_m(H_0^k(S^m(T^*M))) \subset H_0^k(SM)$. Therefore we can define

$$L_m : H^{-k}(SM) \rightarrow H^{-k}(S^m(T^*M)) \tag{1}$$

in the sense of distributions and it is bounded.

Next, if M is compact nontrapping with strictly convex boundary, we study the properties of I and its adjoint I^* . Recall a useful integral identity called Santaló's formula.

Lemma 2.1 [Sharafutdinov 1999, Lemma 3.3.2]. *Let M be a compact nontrapping Riemannian manifold with strictly convex boundary. For every function $f \in C(SM)$, the equality*

$$\int_{SM} f(x, \xi) d\Sigma^{2n-1}(x, \xi) = \int_{\partial_+ SM} d\mu(x, \xi) \int_0^{\tau(x, \xi)} f(\gamma_{x, \xi}(t), \dot{\gamma}_{x, \xi}(t)) dt$$

holds.

Notice that the definition of compact dissipative Riemannian manifold (CDRM) in [Sharafutdinov 1999] is equivalent to compact nontrapping manifolds with strictly convex boundary.

Now let $\varphi \in C_\alpha^\infty(\partial_+ SM)$ and $f \in C^\infty(SM)$. By Santaló's formula,

$$\begin{aligned} (If, \varphi) &= \int_{\partial_+ SM} \varphi(x, \xi) d\mu \int_0^{\tau(x, \xi)} f(\gamma_{x, \xi}(t), \dot{\gamma}_{x, \xi}(t)) dt \\ &= \int_{\partial_+ SM} d\mu \int_0^{\tau(x, \xi)} \varphi^\#(\gamma_{x, \xi}(t), \dot{\gamma}_{x, \xi}(t)) f(\gamma_{x, \xi}(t), \dot{\gamma}_{x, \xi}(t)) dt \\ &= \int_{SM} \varphi^\# f d\Sigma^{2n-1}. \end{aligned}$$

Thus $I^*\varphi = \varphi^\#$ with

$$I^* : C_\alpha^\infty(\partial_+ SM) \rightarrow C^\infty(SM)$$

bounded. By the proof of [Sharafutdinov 1994, Theorem 4.2.1], one can extend I to a bounded operator

$$I : H^k(SM) \rightarrow H^k(\partial_+ SM)$$

and $I(H_0^k(SM)) \subset H_0^k(\partial_+ SM)$ for any integer $k \geq 0$ (notice that $I(C_c^\infty((SM)^{\text{int}})) \subset C_c^\infty((\partial_+ SM)^{\text{int}})$). Thus we can define the bounded operator

$$I^* : H^{-k}(\partial_+ SM) \rightarrow H^{-k}(SM) \tag{2}$$

in the sense of distributions.

Given $u \in H_0^k(S^m(T^*M))$ and $\varphi \in H^{-k}(\partial_+ SM)$, we have $I_m^*\varphi$ is defined in the sense of distributions:

$$(I_m^*\varphi, u) := (I^*\varphi, \ell_m u) = (\varphi, I \circ \ell_m u) = (\varphi, I_m u).$$

Lemma 2.2. *Given a compact nontrapping Riemannian manifold M with strictly convex boundary,*

$$I_m^* = L_m \circ I^* : H^{-k}(\partial_+ SM) \rightarrow H^{-k}(S^m(T^*M))$$

is a bounded operator.

To conclude this section, we briefly discuss X , the generating vector field of the geodesic flow on the unit sphere bundle SM , acting on distributions. Since X is a differential operator on SM , it is obvious that

$$X : H^{k+1}(SM) \rightarrow H^k(SM), \quad k \geq 0.$$

For $f \in H^{-k}(SM)$ and $h \in H_0^{k+1}(SM)$ (so $Xh \in H_0^k(SM)$), we define $Xf \in H^{-k-1}(SM)$ in the sense of distributions (notice that the volume form $d\Sigma^{2n-1}$ is invariant under the geodesic flow):

$$(Xf, h) := (f, -Xh).$$

3. Solenoidal extensions

In the paper [Kato et al. 2000], the authors proved the existence of compactly supported solenoidal extensions of solenoidal 1-forms to some larger manifold in both L^2 and smooth cases.

Proposition 3.1. *Let Ω be a bounded simply connected domain, with smooth boundary, contained in some Riemannian manifold \mathcal{M} . Let U be an open neighborhood of Ω with ∂U smooth. Then there exists a bounded map $\mathcal{E} : L^2_{\text{sol}}(T^*\Omega) \rightarrow L^2_{U,\text{sol}}(T^*\mathcal{M})$ such that $\mathcal{E}|_{\Omega} = \text{Id}$. Moreover, $\mathcal{E}(C^\infty_{\text{sol}}(T^*\Omega)) \subset C^\infty_{U,\text{sol}}(T^*\mathcal{M})$.*

Here $L^2_{U,\text{sol}}(T^*\mathcal{M})$ and $C^\infty_{U,\text{sol}}(T^*\mathcal{M})$ denote the subspaces of $L^2_{\text{sol}}(T^*\mathcal{M})$ and $C^\infty_{\text{sol}}(T^*\mathcal{M})$ respectively consisting of elements supported in U .

Our goal is to extend this result to symmetric tensor fields of higher rank. However, for tensor fields of higher rank, new ideas are required and the argument is more involved.

L^2 solenoidal extensions. We first prove the extension in the L^2 category by solving a suitable elliptic system.

Proposition 3.2. *Let Ω be a bounded simply connected domain, with smooth boundary, contained in some Riemannian manifold (\mathcal{M}, g) . Let U be an open neighborhood of Ω with ∂U smooth. Then given $m \geq 2$, $K \geq 2$ and $\epsilon > 0$, there exist a Riemannian metric \tilde{g} and a bounded map $\mathcal{E} : L^2(S^m_{\text{sol}}(T^*_g\Omega)) \rightarrow L^2(S^m_{U,\text{sol}}(T^*_{\tilde{g}}\mathcal{M}))$ such that $\|\tilde{g} - g\|_{C^K} < \epsilon$, $\tilde{g}|_{\bar{\Omega}} = g$ and $\mathcal{E}|_{\Omega} = \text{Id}$.*

Proof. Suppose $u \in L^2(S^m_{\text{sol}}(T^*_g\Omega))$, i.e., $\delta u = 0$ in the sense of distributions. By the Green’s formula for symmetric tensor fields (see [Sharafutdinov 1994]) one can define the boundary contraction of u with the outward unit normal vector ν on $\partial\Omega$ in the sense of distributions; i.e., for $v \in H^1(S^{m-1}(T^*_g\Omega))$ we have

$$(u, dv)_{\Omega} = (j_{\nu}u, v)_{\partial\Omega}. \tag{3}$$

Since the trace operator $T : H^1(S^{m-1}(T^*_g\Omega)) \rightarrow H^{1/2}(S^{m-1}(\partial T^*_g\Omega))$, $Tv = v|_{\partial\Omega}$, is surjective, $j_{\nu}u \in H^{-1/2}(S^{m-1}(\partial T^*_g\Omega))$ is well-defined, and in local coordinates

$$(j_{\nu}u)_{i_1 i_2 \dots i_{m-1}} = u_{i_1 i_2 \dots i_{m-1} j} \nu^j.$$

By (3), for $v \in H^1(S^{m-1}(T^*_g\Omega))$ with $dv = 0$ (Killing tensor fields on Ω), we have $(j_{\nu}u, v)_{\partial\Omega} = 0$.

It is known that generic (in the C^K -topology for $K \geq 2$) metrics admit only trivial integrals polynomial in momenta [Kruglikov and Matveev 2016]; i.e., for a generic metric h , the only Killing tensor fields are of the form ch^k , where $c \in \mathbb{R}$ and

$$h^k = \sigma(\underbrace{h \otimes \cdots \otimes h}_k)$$

is the symmetric tensor product of k copies of h . Thus given any $\epsilon > 0$ and $K \geq 2$, there is a smooth metric \tilde{g} with $\|\tilde{g} - g\|_{C^K} < \epsilon$ and $\tilde{g}|_{\bar{\Omega}} = g$ so that $(U \setminus \bar{\Omega}, \tilde{g})$ (thus (U, \tilde{g})) does not have nontrivial Killing tensor fields.

Define

$$f = \begin{cases} -j_\nu u & \text{on } \partial\Omega, \\ 0 & \text{on } \partial U. \end{cases}$$

Let $D := U \setminus \bar{\Omega}$ and consider the following boundary value problem for systems of second-order partial differential equations:

$$\begin{cases} \delta dw = 0 & \text{in } D, \\ j_\mu dw = f \in H^{-1/2}(S^{m-1}(\partial T_{\tilde{g}}^* D)), \\ w \in H^1(S^{m-1}(T_{\tilde{g}}^* D)). \end{cases} \tag{4}$$

Here μ is the outward unit normal vector on ∂D for D ; notice $\mu|_{\partial\Omega} = -\nu$. We claim that the system (4) is a regular elliptic system (also called coercive in some texts). Assume that the claim is true for the moment and let us continue the proof.

Next, we study the solutions of the homogeneous problem. Let $\delta dv = 0$ and $j_\mu dv|_{\partial D} = 0$ for some $v \in H^1(S^{m-1}(T_{\tilde{g}}^* D))$; by ellipticity, v is smooth. Applying Green’s formula, one has

$$\int_D \langle dv, dv \rangle dV^n(x) = - \int_D \langle \delta dv, v \rangle dV^n(x) + \int_{\partial D} \langle j_\mu dv, v \rangle dV^n(x) = 0,$$

i.e., $dv \equiv 0$. So the solution set of the homogeneous problem is

$$\mathcal{K} = \{v \in C^\infty(S^{m-1}(T_{\tilde{g}}^* D)) : dv \equiv 0\},$$

the set of Killing tensor fields of rank $m - 1$ on D .

Now by [McLean 2000, Theorem 4.11], (4) is solvable in $H^1(S^{m-1}(T_{\tilde{g}}^* D))$ for the given boundary condition f if and only if $(v, f)_{\partial D} = 0$ for all $v \in \mathcal{K}$. Note that (D, \tilde{g}) does not have nontrivial Killing tensor fields. If m is even, the only Killing $(m-1)$ -tensor field is $v = 0$; then $(v, f)_{\partial D} = (0, f)_{\partial D} = 0$. If m is odd, the Killing $(m-1)$ -tensor fields in D are of the form $v = c\tilde{g}^{(m-1)/2}|_D$. Thus we can extend v to $v = c\tilde{g}^{(m-1)/2}|_U$, which is also a Killing tensor field in Ω . By the definition of f ,

$$(v, f)_{\partial D} = -(v, j_\nu u)_{\partial\Omega} = -(v, \delta u)_\Omega - (dv, u)_\Omega = 0,$$

since $\delta u = 0, dv = 0$ in Ω .

Thus the system (4) is solvable. Let $w \in H^1(S^{m-1}(T_{\tilde{g}}^* D))$ be a solution of (4) (the set of all solutions is $w + \mathcal{K}$) and define

$$\mathcal{E}u = \begin{cases} u & \text{in } \Omega, \\ dw & \text{in } D, \\ 0 & \text{in } \mathcal{M} \setminus \bar{U}. \end{cases}$$

It is easy to see that $\mathcal{E}u \in L^2(S^m(T_{\tilde{g}}^*\mathcal{M}))$ and $\text{supp } \mathcal{E}u \subset \bar{U}$. In particular, for $v \in H^1(S^{m-1}(T_{\tilde{g}}^*\mathcal{M}))$,

$$\begin{aligned} (\delta\mathcal{E}u, v)_{\mathcal{M}} &= -(\mathcal{E}u, dv)_{\mathcal{M}} = -(dw, dv)_D - (u, dv)_{\Omega} \\ &= -(j_{\mu}dw, v)_{\partial D} - (j_{\nu}u, v)_{\partial\Omega} \\ &= -(-j_{\nu}u, v)_{\partial\Omega} - (j_{\nu}u, v)_{\partial\Omega} \\ &= 0. \end{aligned}$$

Thus $\mathcal{E}u$ is solenoidal in the sense of distributions, and $\mathcal{E}u \in L^2(S_{U,\text{sol}}^m(T_{\tilde{g}}^*\mathcal{M}))$.

Moreover, by [McLean 2000, Theorem 4.11], we have the stability estimate

$$\|\mathcal{E}u\|_{L^2(\mathcal{M})}^2 = \|u\|_{L^2(\Omega)}^2 + \|dw\|_{L^2(D)}^2 \leq \|u\|_{L^2(\Omega)}^2 + C\|j_{\nu}u\|_{H^{-1/2}(\partial\Omega)}^2 \leq C'\|u\|_{L^2(\Omega)}^2,$$

i.e., \mathcal{E} is bounded. □

The only thing left to prove is the claim about ellipticity.

Lemma 3.3. *The system (4) above is a regular elliptic system.*

Proof. It is well known that δd is a self-adjoint elliptic operator; see, for example, [Sharafutdinov 1994]. We just need to show that the Neumann boundary value problem satisfies the Lopatinskii condition.

To check the Lopatinskii condition, we follow a similar procedure to that in the proof of [Sharafutdinov 1994, Theorem 3.3.2]. We choose local coordinates $(x^1, x^2, \dots, x^{n-1}, x^n = t \geq 0)$ in a neighborhood W of $x_0 = (x', 0) \in \partial D$ in D so that $\partial D \cap W = \{t = 0\}$ and $g_{ij}(x_0) = \delta_{ij}$. Define $d_0 = \sigma_p d$ and $\delta_0 = \sigma_p \delta$, the principal symbols of d and δ respectively. Then we need to show that the boundary value problem for systems of ordinary differential equations

$$\begin{cases} \delta_0(x', 0, \xi', D_t)d_0(x', 0, \xi', D_t)w(t) = 0, \\ j_{-\frac{\partial}{\partial t}}d_0(x', 0, \xi', D_t)w(t)|_{t=0} = f_0 \end{cases}$$

has a unique solution in \mathcal{N}_+ for all $\xi' \in \mathbb{R}^{n-1} \setminus \{0\}$ and $f_0 \in S^{m-1}(\mathbb{R}^n)$, symmetric $(m-1)$ -tensors on \mathbb{R}^n . Here $D_t = -id/dt$, and for the sake of simplicity, we drop the space variables $(x', 0)$ from the symbols so

$$\mathcal{N}_+ := \{w \in S^{m-1}(\mathbb{R}^n)|_{\{x'\} \times [0, \infty)} : \delta_0(\xi', D_t)d_0(\xi', D_t)w = 0 \text{ and } w \text{ decays rapidly together with all derivatives as } t \rightarrow +\infty\}.$$

Since the equation $\det(\delta_0(\xi', \zeta)d_0(\xi', \zeta)) = 0$ has real coefficients with no real root for $\xi' \neq 0$, it is not difficult to see that $\dim \mathcal{N}_+ = \dim S^{m-1}(\mathbb{R}^n)$. Thus it is sufficient to show that the homogeneous problem

$$\begin{cases} \delta_0(\xi', D_t)d_0(\xi', D_t)w(t) = 0, \\ j_{-\frac{\partial}{\partial t}}d_0(\xi', D_t)w(t)|_{t=0} = 0 \end{cases} \tag{5}$$

has only the zero solution in \mathcal{N}_+ .

By a similar computation to that in the proof of [Sharafutdinov 1994, Theorem 3.3.2], we have the following Green's formula. Let $v(t) \in C^\infty([0, \infty) \rightarrow S^m(\mathbb{R}^n))$ and $w(t) \in C^\infty([0, \infty) \rightarrow S^{m-1}(\mathbb{R}^n))$

such that both of them decay rapidly together with all derivatives as $t \rightarrow +\infty$. If $j_{-\frac{\partial}{\partial t}} v(0) = 0$ (notice that different from [Sharafutdinov 1994], here we use the Neumann boundary condition at $t = 0$) then

$$\int_0^\infty \langle \delta_0(\xi', D_t)v, w \rangle dt = - \int_0^\infty \langle v, d_0(\xi', D_t)w \rangle dt. \tag{6}$$

Now if $w(t) \in \mathcal{N}_+$ is a solution to (5), let $v(t) = d_0(\xi', D_t)w(t)$. By (6) we obtain

$$d_0(\xi', D_t)w(t) = 0.$$

Notice that

$$(d_0(\xi)w)_{i_1 \dots i_m} = \frac{i}{m} \sum_{k=1}^m \xi_{i_k} w_{i_1 \dots \widehat{i_k} \dots i_m},$$

where the $\widehat{}$ over i_k means this index is omitted. Let $i_m = n$ and $\xi = (\xi', D_t)$. We obtain the system of first-order ordinary differential equations

$$(d_0(\xi', D_t)w)_{ni_1 \dots i_{m-1}} = \frac{i}{m} \left\{ (\ell + 1)D_t w_{i_1 \dots i_{m-1}} + \sum_{i_k \neq n} \xi_{i_k} w_{ni_1 \dots \widehat{i_k} \dots i_{m-1}} \right\} = 0,$$

where $\ell = \ell(i_1, \dots, i_{m-1})$ is the number of occurrences of the index n in (i_1, \dots, i_{m-1}) . Since $\lim_{t \rightarrow +\infty} w(t) = 0$, by induction on ℓ , the only solution to the above first-order homogeneous system is $w \equiv 0$, and this shows that (4) satisfies the Lopatinskii condition. \square

Smooth solenoidal extensions. In this subsection we achieve C^∞ solenoidal extensions for tensors of arbitrary rank. Observe that the approach we use is quite different from the one of [Kato et al. 2000].

Proposition 3.4. *Let Ω be a bounded connected domain, with smooth boundary, contained in some Riemannian manifold (\mathcal{M}, g) . Let U be an open neighborhood of Ω with ∂U smooth. Then given $m \geq 2, K \geq 2$ and $\epsilon > 0$, there exist a Riemannian metric \tilde{g} and a bounded map $\mathcal{E} : H^k(S_{\text{sol}}^m(T_g^* \Omega)) \rightarrow L^2(S_{U, \text{sol}}^m(T_{\tilde{g}}^* \mathcal{M}))$ for some integer $k \geq 2$ such that $\|\tilde{g} - g\|_{C^K} < \epsilon, \tilde{g}|_{\bar{\Omega}} = g, \mathcal{E}|_{\Omega} = \text{Id}$ and*

$$\mathcal{E}(C^\infty(S_{\text{sol}}^m(T_g^* \Omega))) \subset C^\infty(S_{U, \text{sol}}^m(T_{\tilde{g}}^* \mathcal{M})).$$

To prove the proposition, we start with the following lemma on the existence of solenoidal extensions that might not be compactly supported.

Lemma 3.5. *Let Ω be a bounded connected domain, with smooth boundary, contained in some Riemannian manifold (\mathcal{M}, g) . There exists an open neighborhood U of Ω such that every $u \in C^\infty(S_{\text{sol}}^m(T^* \bar{\Omega}))$ can be extended to $\tilde{u} \in C^\infty(S_{\text{sol}}^m(T^* U))$ with $\tilde{u}|_{\bar{\Omega}} = u$.*

Proof. Let $u \in C^\infty(S_{\text{sol}}^m(T^* \bar{\Omega}))$, i.e., $\delta u = 0$, in local coordinates $u = u_{j_1 \dots j_m} dx^{j_1} \otimes \dots \otimes dx^{j_m}$ and

$$(\delta u)_{i_1 \dots i_{m-1}} = g^{jk} \nabla_j u_{ki_1 \dots i_{m-1}} = 0, \tag{7}$$

where

$$\nabla_j u_{ki_1 \dots i_{m-1}} = \partial_j u_{ki_1 \dots i_{m-1}} - \Gamma_{jk}^\ell u_{\ell i_1 \dots i_{m-1}} - \sum_{s=1}^{m-1} \Gamma_{j i_s}^\ell u_{\ell k i_1 \dots \widehat{i_s} \dots i_{m-1}}. \tag{8}$$

Pick $x_0 \in \partial\Omega$. We follow the idea of the proof in [Stefanov and Uhlmann 2005, Lemma 4.1] and choose semigeodesic coordinates $(x^1, \dots, x^{n-1}, x^n) = (x', x^n)$ near x_0 with $\partial\Omega = \{x^n = 0\}$ and $\partial_n = \nu$ the unit outward (with respect to Ω) vector normal to $\partial\Omega$; thus

$$g^{kn} = \delta_n^k, \quad \Gamma_{kn}^n = \Gamma_{nn}^k = 0 \quad \text{for all } k = 1, 2, \dots, n.$$

We extend the components $u_{j_1 \dots j_m}$, $j_s < n$ for all $1 \leq s \leq m$, smoothly to U (note that $U \setminus \bar{\Omega}$ is determined by the semigeodesic neighborhood of $\partial\Omega$), and denote the extensions by $v_{j_1 \dots j_m}$. We will construct the other components in $\{x^n > 0\}$ by induction on the number of appearances of n in $j_1 \dots j_m$. By equations (7) and (8), if $i_1, \dots, i_{m-1} < n$,

$$\begin{aligned} \partial_n v_{ni_1 \dots i_{m-1}} - \sum_{s=1}^{m-1} \sum_{\ell < n} \Gamma_{ni_s}^\ell v_{\ell ni_1 \dots \hat{i}_s \dots i_{m-1}} - \sum_{j,k < n} g^{jk} \left(\Gamma_{jk}^n v_{ni_1 \dots i_{m-1}} + \sum_{s=1}^{m-1} \Gamma_{ji_s}^n v_{nki_1 \dots \hat{i}_s \dots i_{m-1}} \right) \\ = - \sum_{j,k < n} g^{jk} \left(\partial_j v_{ki_1 \dots i_{m-1}} - \sum_{\ell < n} \Gamma_{jk}^\ell v_{\ell i_1 \dots i_{m-1}} - \sum_{s=1}^{m-1} \sum_{\ell < n} \Gamma_{ji_s}^\ell v_{\ell ki_1 \dots \hat{i}_s \dots i_{m-1}} \right). \end{aligned} \tag{9}$$

Notice that the right side of (9) is known, so it gives a system of first-order linear ODEs. Given the initial values $u_{ni_1 \dots i_{m-1}}(x', 0) = v_{ni_1 \dots i_{m-1}}(x', 0)$, there exists a unique solution to (9). Thus we obtain continuous $v_{ni_1 \dots i_{m-1}}$ with $i_1, \dots, i_{m-1} < n$ near ∂M . In particular, $v_{ni_1 \dots i_{m-1}}(x', x^n)$ depends smoothly on x' , the first $n - 1$ variables.

By differentiating (9) repeatedly with respect to x^n , we get that $\partial_n^s v_{ni_1 \dots i_{m-1}}(x', x^n)$, $s \geq 0$, are continuous in $x^n \geq 0$ and smooth with respect to x' . Moreover, by (9) and the fact that u is solenoidal we carry out an induction on s , so

$$\begin{aligned} \partial_n^s v_{ni_1 \dots i_{m-1}}(x', 0) \\ = G_{i_1 \dots i_{m-1}}^s \left(\partial_n^\ell v_{nj_1 \dots j_{m-1}}, \partial_n^\ell v_{j_1 \dots j_m}, \partial_n^\ell \partial_k v_{j_1 \dots j_m}; \ell < s; j_1, \dots, j_{m-1}, j_m, k < n \right)(x', 0) \\ = G_{i_1 \dots i_{m-1}}^s \left(\partial_n^\ell u_{nj_1 \dots j_{m-1}}, \partial_n^\ell u_{j_1 \dots j_m}, \partial_n^\ell \partial_k u_{j_1 \dots j_m}; \ell < s; j_1, \dots, j_{m-1}, j_m, k < n \right)(x', 0) \\ = \partial_n^s u_{ni_1 \dots i_{m-1}}(x', 0) \end{aligned}$$

for all $s \geq 0$; i.e., $\partial_n^s v_{ni_1 \dots i_{m-1}}$ are consistent with $\partial_n^s u_{ni_1 \dots i_{m-1}}$ at $(x', 0)$.

Next by induction on the number of appearances of n and repeatedly using equations (7) and (8), one can get unique

$$v_{ni_1 \dots i_{m-1}}, \quad v_{nni_1 \dots i_{m-2}}, \quad \dots, \quad v_{n \dots ni_1}, \quad v_{n \dots n},$$

which together with their normal derivatives with respect to x^n of all orders, are continuous (smooth with respect to x') and consistent with the corresponding $\partial_n^m u_{j_1 \dots j_m}$ at $(x', 0)$. Therefore we get a smooth solenoidal m -tensor

$$\tilde{u} = \begin{cases} u & \text{on } \bar{\Omega}, \\ v & \text{on } U \setminus \bar{\Omega}. \end{cases} \quad \square$$

Proof of Proposition 3.4. There exist two precompact open neighborhoods V, U of Ω which satisfy

$$\Omega \subset \bar{\Omega} \subset V \subset \bar{V} \subset U \subset \bar{U} \subset \mathcal{M}.$$

Given $u \in C^\infty(S^m_{\text{sol}}(T^*\Omega))$, by Lemma 3.5, we can extend u to get $u_V \in C^\infty(S^m_{\text{sol}}(T^*V))$ with $u_V|_{\bar{\Omega}} = u$. Then we extend u_V to a smooth m -tensor w on \mathcal{M} with $\text{supp } w \subset U$. Let $f = \delta w$ and $D = U \setminus \bar{\Omega}$ open, so $\text{supp } f \subset U \setminus V \subset D$.

Similar to the perturbation-of-metrics argument in the proof of Proposition 3.2, given any $\epsilon > 0$ and $K \geq 2$, there is a smooth metric \tilde{g} with $\|\tilde{g} - g\|_{C^K} < \epsilon$ and $\tilde{g}|_V = g$ so that (D, \tilde{g}) does not have nontrivial Killing tensor fields. Now if m is even, the only Killing $(m-1)$ -tensor field on (D, \tilde{g}) is $v = 0$. Then

$$(v, f)_D = (0, f)_D = 0.$$

If m is odd, Killing $(m-1)$ -tensor fields on (D, \tilde{g}) are of the form $v = c\tilde{g}^{(m-1)/2}|_D$. Thus we can extend v to $v = c\tilde{g}^{(m-1)/2}|_U$, which is also a Killing tensor field in Ω . By Green's formula,

$$(v, f)_D = (v, \delta w)_D = -(dv, w)_D + (v, j_\mu w)_{\partial D} = -(v, j_\nu u)_{\partial \Omega} = -(v, \delta u)_\Omega - (dv, u)_\Omega = 0.$$

since $\delta u = 0$ and $dv = 0$ in Ω . Here $\mu = -\nu$ is the unit outward normal vector on ∂D and

$$(j_\mu w)_{i_1 i_2 \dots i_{m-1}} = w_{i_1 i_2 \dots i_{m-1} j} \mu^j.$$

Now by [Delay 2012, Theorem 1.3], there exist $u_D \in C^\infty(S^m(T^*\mathcal{M}))$ with $\text{supp } u_D \subset U \setminus \Omega$ such that $\delta u_D = -f$. It is not difficult to check that the symmetric differentiation d satisfies the kernel restriction condition (KRC) and the asymptotic Poincaré inequality (API) of [Delay 2012]. We define $\mathcal{E}u = w + u_D$. Then $\delta \mathcal{E}u = \delta w + \delta u_D = f - f = 0$; i.e., $\mathcal{E}u \in C^\infty(S^m_{U, \text{sol}}(T^*_\tilde{g} \mathcal{M}))$. Moreover, $\mathcal{E}u|_\Omega = u$.

The argument above gives a construction for compactly supported smooth solenoidal extensions. One can further check that the extension can be constructed in a stable way. In view of the ODEs (9), the solution is controlled by the initial value and the nonhomogeneous term on the right side under Sobolev norms; see, e.g., [Han 2011]. By induction on the number of appearances of n and repeatedly differentiating (9), we have that

$$\|u_V\|_{H^1(V \setminus \bar{\Omega})} \leq C \left(\|j_\mu u\|_{H^{k_1}(\partial \Omega)} + \sum_{i_s < n} \|(u_V)_{i_1 \dots i_m}\|_{H^{k_2}(V \setminus \bar{\Omega})} \right)$$

for some $k_1, k_2 \geq 1$. Note that in boundary normal coordinates $\mu = -\partial_n$, and we have full freedom to control the elements $(u_V)_{i_1 \dots i_m}$, with $i_s < n$ for all $1 \leq s \leq m$, by $u|_\Omega$ due to the fact that δ is an underdetermined elliptic operator. Thus

$$\|u_V\|_{H^1(V \setminus \bar{\Omega})} \leq C \|u\|_{H^k(\Omega)}$$

for some integer $k \geq 2$. Then $\|w\|_{H^1(U)} \leq C \|u\|_{H^k(\Omega)}$ by extending u_V to w in a stable way.

Next we control the L^2 norm of u_D . Roughly speaking, u_D is the symmetric differentiation of some smooth $(m-1)$ -tensor p , multiplied by a smooth nonnegative weight which vanishes exponentially at the boundary of D ; concretely $u_D = \psi^2 \phi^2 dp$ with ϕ a boundary-defining function on D and ψ vanishes exponentially at the boundary ∂D . By [Delay 2012, Lemma 10.2], $\|p\|_{H^2_{\phi, \psi}(D)} \leq C \|\psi^{-2} \delta w\|_{L^2_\psi(D)}$,

where $H^2_{\phi,\psi}$ and L^2_{ψ} are some weighted Sobolev spaces; see [Delay 2012] for more details. Then one can check that the following inequality with unweighted Sobolev norms holds:

$$\|u_D\|_{L^2(D)} \leq C \|w\|_{H^1(U)}.$$

Now we combine the estimates above to obtain

$$\|\mathcal{E}u\|_{L^2(\mathcal{M})} \leq C_1(\|w\|_{L^2(U)} + \|u_D\|_{L^2(D)}) \leq C_2\|w\|_{H^1(U)} \leq C\|u\|_{H^k(\Omega)}$$

for some $C > 0$ independent of u . Since $C^\infty(S^m_{\text{sol}}(T^*\Omega))$ is dense in $H^k(S^m_{\text{sol}}(T^*\Omega))$ under the H^k norm, we can extend \mathcal{E} to a bounded map from H^k to L^2 with the same properties, which completes the proof. \square

Remark 3.6. We expect that the L^2 norm of $\mathcal{E}u$ can be bounded by the L^2 norm of $u|_\Omega$ through sharper estimates, similar to the result under the L^2 setting in the previous subsection. However, the H^k space is enough for carrying out the argument under the smooth setting in the next section; see Lemma 4.3.

4. Surjectivity of the normal operator $I_m^* I_m$

Since M is simple we can consider an extension \tilde{M} of M which is open ($\tilde{M} = \tilde{M}^{\text{int}}$) and whose compact closure is also simple. It is well known that the normal operator $N = I_m^* I_m$ is a pseudodifferential operator of order -1 on \tilde{M} ; see, for example, [Sharafutdinov 1994; Stefanov and Uhlmann 2004; 2008; Sharafutdinov et al. 2005]. Below is a lemma that, roughly speaking, gives a right parametrix for N on the space of solenoidal tensor fields. The proof is similar to [Sharafutdinov et al. 2005, Theorem 3.1].

Lemma 4.1. *Let S be a parametrix for the operator δd . There exists a pseudodifferential operator Q of order 1 on the bundle of symmetric m -tensor fields $S^m(T^*\tilde{M})$ such that*

$$E = NQ + dS\delta + K, \tag{10}$$

where E is the identity operator and K is a smoothing operator.

Proof. Let $\lambda(\xi)$ be the principal symbol of the pseudodifferential operator N and

$$S^m_\xi(T^*_x\tilde{M}) = \{u \in S^m(T^*_x\tilde{M}) : j_\xi u = 0\},$$

where $j_\xi = -i\sigma_p(\delta) : S^m(T^*_x\tilde{M}) \rightarrow S^{m-1}(T^*_x\tilde{M})$. By [Sharafutdinov 1994, Theorem 2.12.1],

$$\lambda(\xi) : S^m_\xi(T^*_x\tilde{M}) \rightarrow S^m_\xi(T^*_x\tilde{M})$$

is an isomorphism for $\xi \neq 0$. Thus there exists $p(\xi)$ such that $\lambda(\xi)p(\xi) = \text{Id}$ on $S^m_\xi(T^*_x\tilde{M})$. Namely, we can find some pseudodifferential operator P of order 1 such that on $S^m_\xi(T^*_x\tilde{M})$,

$$NP = E - B$$

for some operator B of order -1 . Now multiplying both sides by the ‘‘solenoidal projection’’ $E - dS\delta$, which is of order 0, one has

$$NP(E - dS\delta) = E - dS\delta - R \tag{11}$$

defined on $S^m(T^*\tilde{M})$.

Then we multiply both sides of (11) by δ to get $\delta R = R'$ with R' some smoothing operator. Let $C = \sum_{k=0}^{\infty} R^k$, which is a pseudodifferential operator of order 0 and a parametrix for $E - R$. Write (11) as

$$NP(E - dS\delta) + dS\delta = E - R,$$

and multiply both sides by C to get

$$NP(E - dS\delta)C + dS\delta + dS\delta \sum_{k=1}^{\infty} R^k = (E - R)C = E + R'',$$

with R'' a smoothing operator. Since δR is smoothing, $dS\delta \sum_{k=1}^{\infty} R^k$ is smoothing too. We arrive at the equation

$$NP(E - dS\delta)C + dS\delta + K = E,$$

where K is a smoothing operator. Denote $P(E - dS\delta)C$ by Q (note that one can make Q properly supported). Then we get (10), which finishes the proof. \square

Let U be a small open neighborhood of M in \tilde{M} . Denote the restriction operator from \tilde{M} to M by r_M . Then the following holds:

Lemma 4.2. *Suppose M is a compact simple Riemannian manifold, and assume I_m is s -injective on $C^\infty(S^m(T^*M))$. Then the operator*

$$r_M N : H_c^{-1}(S^m(T^*\tilde{M})) \rightarrow L^2(S_{\text{sol}}^m(T^*M))$$

is surjective.

Note that elements in $H_c^{-1}(S^m(T^*\tilde{M}))$ are defined in the sense of distributions, which are compactly supported in \tilde{M} .

Proof. We adopt the approach of [Dairbekov and Uhlmann 2010] for showing the surjectivity of N on 1-forms. By Lemma 4.1,

$$NQ u = u + K u$$

for all $u \in L_c^2(S_{\text{sol}}^m(T^*\tilde{M}))$ with K a smoothing operator on \tilde{M} . Since the simplicity is stable under small C^2 -perturbations of the metric g , by Proposition 3.2, we perturb the metric of $\tilde{M} \setminus \bar{M}$ a little bit (still denoted by g) so that under the new metric \tilde{M} is still simple and there exists a bounded operator $\mathcal{E} : L^2(S_{\text{sol}}^m(T^*M)) \rightarrow L^2(S_{U,\text{sol}}^m(T^*\tilde{M}))$ such that on $L^2(S_{\text{sol}}^m(T^*M))$,

$$r_M N Q \mathcal{E} = E + r_M K \mathcal{E}.$$

Since K is a smoothing operator, $r_M K \mathcal{E}$ is compact on $L^2(S_{\text{sol}}^m(T^*M))$, which implies that $E + r_M K \mathcal{E}$ has closed range and finite codimension. Thus we have $r_M N Q \mathcal{E} : L^2(S_{\text{sol}}^m(T^*M)) \rightarrow L^2(S_{\text{sol}}^m(T^*M))$ has closed range and finite codimension. By the inclusion relation

$$r_M N Q \mathcal{E}(L^2(S_{\text{sol}}^m(T^*M))) \subset r_M N(H_c^{-1}(S^m(T^*\tilde{M}))) \subset L^2(S_{\text{sol}}^m(T^*M)),$$

the intermediate space $r_M N(H_c^{-1}(S^m(T^* \tilde{M})))$ is also closed in $L^2(S_{\text{sol}}^m(T^* M))$. Thus it suffices to show that the adjoint $(r_M N)^*$ is injective, which will imply the surjectivity of $r_M N$.

For L^2 symmetric m -tensor fields, we have the decomposition

$$L^2(S^m(T^* M)) = L^2(S_{\text{sol}}^m(T^* M)) \oplus L^2(S_P^m(T^* M)), \tag{12}$$

where $L^2(S_P^m(T^* M))$ is the potential part. Thus the dual operator of $r_M N$ is

$$(r_M N)^* : L^2(S_{\text{sol}}^m(T^* M)) \rightarrow (H_c^{-1}(S^m(T^* \tilde{M})))^*.$$

For $u \in L^2(S_{\text{sol}}^m(T^* M))$ and $v \in H_c^{-1}(S^m(T^* \tilde{M}))$, if we denote by $\mathcal{E}_0 u$ the extension of u to \tilde{M} by zero (note that generally $\mathcal{E}_0 u$ is not solenoidal on \tilde{M}), we have

$$((r_M N)^* u, v) = (u, r_M N v) = (\mathcal{E}_0 u, N v) = (N \mathcal{E}_0 u, v),$$

i.e., $(r_M N)^* = N \mathcal{E}_0$.

Therefore given $u \in L^2(S_{\text{sol}}^m(T^* M))$, if $N \mathcal{E}_0 u = 0$, then

$$0 = (N \mathcal{E}_0 u, \mathcal{E}_0 u) = \|I_m \mathcal{E}_0 u\|_{L^2(\partial_+ S \tilde{M})}^2 \implies I_m \mathcal{E}_0 u = 0.$$

Since $\mathcal{E}_0 u = 0$ outside M and \tilde{M} is simple, this implies

$$I_m u = 0.$$

By [Sharafutdinov et al. 2005, Theorem 1.1], u is smooth and $\delta u = 0$. The s -injectivity assumption implies $u = 0$. This completes the proof of the lemma. \square

Next we prove the lemma in the smooth setting:

Lemma 4.3. *Suppose M is a compact simple Riemannian manifold, and assume I_m is s -injective on $C^\infty(S^m(T^* M))$. Then the operator*

$$r_M N : C_c^\infty(S^m(T^* \tilde{M})) \rightarrow C^\infty(S_{\text{sol}}^m(T^* M))$$

is surjective.

Proof. By Lemma 4.1,

$$N Q u = u + K u$$

for all $u \in C_c^\infty(S_{\text{sol}}^m(T^* \tilde{M}))$ with K a smoothing operator on \tilde{M} . Since the simplicity is stable under small C^2 -perturbations of the metric g , by Proposition 3.4, we perturb the metric of $\tilde{M} \setminus \bar{M}$ a little bit (still denoted by g) so that under the new metric \tilde{M} is still simple and there exists a bounded operator $\mathcal{E} : H^k(S_{\text{sol}}^m(T^* M)) \rightarrow L^2(S_{U, \text{sol}}^m(T^* \tilde{M}))$ for some integer $k \geq 2$ with $\mathcal{E}(C^\infty(S_{\text{sol}}^m(T^* M))) \subset C^\infty(S_{U, \text{sol}}^m(T^* \tilde{M}))$ such that on $H^k(S_{\text{sol}}^m(T^* M))$,

$$r_M N Q \mathcal{E} = E + r_M K \mathcal{E}.$$

Now the argument of [Dairbekov and Uhlmann 2010, Lemma 2.2] can be applied to tensors of any order to finish the proof. \square

Remark 4.4. One can actually prove Lemmas 4.2 and 4.3 just by applying Lemma 3.5. Given a smooth solenoidal tensor u on M , by Lemma 3.5 we first extend it to a smooth solenoidal tensor \tilde{u} on an arbitrarily small open neighborhood U ; then we extend \tilde{u} smoothly to \tilde{M} with compact support, denoted by $\mathcal{E}u$. Note that generally $\mathcal{E}u$ is not solenoidal. Since the Schwartz kernel of the parametrix S of δd is smooth away from the diagonal $\Delta_{\tilde{M} \times \tilde{M}}$, we can choose S to make the support of its Schwartz kernel sufficiently close to $\Delta_{\tilde{M} \times \tilde{M}}$ so that $dS \delta \mathcal{E}u = 0$ in an open neighborhood of M . This implies that $r_M dS \delta \mathcal{E}u = 0$, i.e., $r_M N Q \mathcal{E}u = u + r_M K \mathcal{E}u$. It also works for L^2 solenoidal tensors.

On the other hand, the original proof of [Dairbekov and Uhlmann 2010, Lemma 2.2] uses the existence of compactly supported solenoidal extensions of solenoidal 1-forms one more time at the very end to show that the adjoint $(r_M N)^*$ is injective. However, one can also avoid this. Notice that given a 1-form f in the kernel of $(r_M N)^*$, by [Dairbekov and Uhlmann 2010, equation (2.33)], $f = dp$ for some distribution p on \tilde{M} with $\text{sing supp } p \subset \partial \tilde{M}$ and $p|_{\partial \tilde{M}} = 0$. Moreover, since $\text{supp } f \subset M$, we have $dp = 0$ outside M . As p is smooth outside M and $p = 0$ on $\partial \tilde{M}$, strict convexity of ∂M implies $p \equiv 0$ in $\tilde{M} \setminus M$. Now given a smooth solenoidal 1-form u on M , by Lemma 3.5 let $\mathcal{E}u$ be the smooth compactly supported extension of u to \tilde{M} which is solenoidal in a small open neighborhood ($\neq \tilde{M}$) of M . Since the supports of $\delta \mathcal{E}u$ and p are disjoint, we have

$$(f, \mathcal{E}u) = (dp, \mathcal{E}u) = (p, \delta \mathcal{E}u) = 0,$$

which implies that $f = 0$, i.e., $(r_M N)^*$ has trivial kernel. The argument works for tensors of arbitrary rank.

At this point, we see that one can prove the surjectivity of $r_M N$ just using Lemma 3.5, without the need of knowing the generic absence of nontrivial Killing tensors [Kruglikov and Matveev 2016]. However, a perturbation of the metric seems still necessary so far for the proof of the existence of *compactly supported* solenoidal extensions, and Propositions 3.2 and 3.4 may find their applications in other areas.

5. Analysis of the adjoint I_m^*

Before proving the main result, we need to extend the definition of the geodesic ray transform I_m so that it acts on negative Sobolev spaces. To this end, we will study the regularity property of the adjoint of the geodesic ray transform, I_m^* .

As discussed in the Introduction, given M a compact nontrapping manifold with strictly convex boundary, the operator $I_m^* : C^\infty(\partial_+ SM) \rightarrow C^\infty(S^m(T^*M))$ is the product of two operators, i.e., $I_m^* = L_m \circ I^*$. We instead study the regularity properties of I^* and L_m . We start with the latter.

Lemma 5.1. *Given a compact Riemannian manifold M (with or without boundary), the operator*

$$L_m : H^k(SM) \rightarrow H^k(S^m(T^*M))$$

is bounded for every integer $k \geq 0$.

Proof. Our purpose is to show that there exists a constant $C > 0$ such that for any $w \in H^k(SM)$, the following holds:

$$\|L_m f\|_{H^k} \leq C \|f\|_{H^k}. \tag{13}$$

Since M is compact, by a partition of unit, it suffices to show the above inequality in local charts. Let U be a domain in SM with local coordinate system (z^1, \dots, z^{2n-1}) . We assume $\text{supp } f \subset U$. Let V be a domain in M with local coordinate system (x^1, \dots, x^n) , and ψ be a smooth function with support in V . We will show

$$\|\psi L_m f\|_{H^k(S^m(T^*V))} \leq C \|f\|_{H^k(U)}.$$

By the definition of the H^k norm of tensors, we only need to show the above inequality is true for each component of the tensor.

We start with $f \in C^\infty(SM)$ with support in U ; then $L_m f$ is also smooth. Let $J = (j_1 \cdots j_m)$ and $\xi^J := \xi^{j_1} \cdots \xi^{j_m}$. Then

$$\begin{aligned} & D_x^\alpha [\psi(x) L_m f(x)^J] \\ &= D_x^\alpha \left[\psi(x) \int_{S_x M} f(x, \xi) \xi^J d\Omega_x(\xi) \right] \\ &= D_x^\alpha \left[\psi(x) \int_{S^{n-1}} f(x, \xi(x, \eta)) \xi^J(x, \eta) P(x, \eta) d\Omega(\eta) \right] \\ &= \sum_{\alpha_1 + \alpha_2 + \alpha_3 = \alpha} D_x^{\alpha_1} \psi(x) \int_{S^{n-1}} D_x^{\alpha_2} f(x, \xi(x, \eta)) \cdot D_x^{\alpha_3} [\xi^J(x, \eta) P(x, \eta)] d\Omega(\eta) \\ &= \sum_{\alpha_1 + \alpha_2 + \alpha_3 = \alpha} D_x^{\alpha_1} \psi(x) \int_{S_x M} D_x^{\alpha_2} f(x, \xi) \cdot D_x^{\alpha_3} [\xi^J P(x, \eta(x, \xi))] \cdot P'(x, \xi) d\Omega_x(\xi). \end{aligned} \tag{14}$$

Here P and P' are corresponding Jacobians.

For $|\alpha| \leq k$, according to (14),

$$\begin{aligned} \|D_x^\alpha [\psi(x) L_m f(x)^J]\|_{L^2(V)}^2 &\leq \sum_{\beta \leq \alpha} C_{\beta, \alpha} \int_V \int_{S_x M} |D_x^\beta f(x, \xi)|^2 d\Omega_x(\xi) dx \\ &\leq \sum_{|\gamma| \leq |\alpha|} C_{\gamma, \alpha} \int_U |D_z^\gamma f(z)|^2 dz \leq C \|f\|_{H^k(U)}^2. \end{aligned}$$

Thus the estimate (13) is proved when $w \in C^\infty(SM)$.

For $f \in H^k(SM)$, since $C^\infty(SM)$ is dense in $H^k(SM)$, by an approximation argument, it is easy to show that $L_m f \in H^k(S^m(T^*M))$ and the estimate (13) holds too. This proves the lemma. \square

Now we turn to the analysis of the operator I^* , which basically is an invariant extension, along the geodesic flow, of functions on $\partial_+ SM$ to functions on SM . It is well known that given $\varphi \in C^\infty(\partial_+ SM)$, $\varphi^\# = I^*(\varphi)$ is not necessarily in $C^\infty(SM)$. The following subspace of $C^\infty(\partial_+ SM)$ has already been considered in the Introduction:

$$C_\alpha^\infty(\partial_+ SM) := \{\varphi \in C^\infty(\partial_+ SM) : \varphi^\# \in C^\infty(SM)\}.$$

In particular, by [Pestov and Uhlmann 2005, Lemma 1.1], if M is compact nontrapping with strictly convex boundary,

$$C_\alpha^\infty(\partial_+ SM) = \{\varphi \in C^\infty(\partial_+ SM) : A\varphi \in C^\infty(\partial SM)\}$$

where

$$A\varphi(x, \xi) = \begin{cases} \varphi(x, \xi), & (x, \xi) \in \partial_+ SM, \\ \varphi(\gamma_{x,\xi}(-\tau(x, -\xi)), \dot{\gamma}_{x,\xi}(-\tau(x, -\xi))), & (x, \xi) \in \partial_- SM. \end{cases}$$

Since $A\varphi$ is smooth in both $(\partial_+ SM)^{\text{int}}$ and $(\partial_- SM)^{\text{int}}$, the singularities can only come from $S(\partial M)$. We introduce the space $H_\alpha^k(\partial_+ SM)$, $k \geq 0$, to be the completion of $C_\alpha^\infty(\partial_+ SM)$ under the H^k norm. Obviously $H_\alpha^0(\partial_+ SM) = L^2(\partial_+ SM)$. It is easy to show that $C_c^\infty((\partial_+ SM)^{\text{int}}) \subset C_\alpha^\infty(\partial_+ SM)$ (this is from the fact that $\partial_+ SM$ is compact and the boundary ∂M is strictly convex), which implies that $H_0^k(\partial_+ SM) \subset H_\alpha^k(\partial_+ SM)$.

Lemma 5.2. *Given a compact nontrapping manifold M with strictly convex boundary, the operator*

$$I^* : H_\alpha^k(\partial_+ SM) \rightarrow H^k(SM)$$

is bounded for any integer $k \geq 0$.

Proof. The idea is similar to the proof of Lemma 5.1. First we consider the case $\varphi \in C_\alpha^\infty(\partial_+ SM)$; thus $\varphi^\# \in C^\infty(SM)$. Let U be a domain in $\partial_+ SM$ with local coordinate systems (y^1, \dots, y^{2n-2}) . We assume $\text{supp } \varphi \subset U$. Let V be a domain in SM with local coordinate systems (z^1, \dots, z^{2n-1}) , and ψ be a smooth function with support in V . Since M is compact, it suffices to show

$$\|\psi\varphi^\#\|_{H^k(V)} \leq C \|\varphi\|_{H^k(U)}.$$

Since

$$D_z^\alpha[\psi(z)\varphi^\#(z)] = \sum_{\beta+\gamma=\alpha} D_z^\gamma \psi(z) \cdot D_z^\beta \varphi^\#(z),$$

we obtain that for $|\alpha| \leq k$,

$$\|D_z^\alpha[\psi(z)\varphi^\#(z)]\|_{L^2(V)}^2 \leq \sum_{\beta \leq \alpha} C_{\beta,\alpha} \int_V |D_z^\beta \varphi^\#(z)|^2 dz.$$

Now let $D = \{(y, t) : y \in \partial_+ SM, 0 \leq t \leq \tau(y)\}$ be a closed domain in $\partial_+ SM \times \mathbb{R}$. Define the map $\Psi : D \rightarrow SM$ by $z = \Psi(y, t) = (\gamma_y(t), \dot{\gamma}_y(t))$. By [Sharafutdinov 1994, Lemma 4.2.2],

$$\begin{aligned} \int_V |D_z^\beta \varphi^\#(z)|^2 dz &\leq \sum_{|\sigma|+s=|\beta|} C_{\beta,\sigma,s} \int_U \int_0^{\tau(y)} |D_y^\sigma D_t^s \varphi^\#(z(y, t))|^2 |\xi(y), \nu(x(y))| dt dy \\ &= \sum_{|\sigma|=|\beta|} C_{\beta,\sigma} \int_U \int_0^{\tau(y)} |D_y^\sigma \varphi^\#(y, t)|^2 dt d\mu(y) \quad (\text{since } D_t^s D_y^\sigma \varphi^\# = D_y^\sigma D_t^s \varphi^\#) \\ &= \sum_{|\sigma|=|\beta|} C_{\beta,\sigma} \int_U \tau(y) |D_y^\sigma \varphi(y)|^2 d\mu(y) \\ &\leq \sum_{|\sigma|=|\beta|} C'_{\beta,\sigma} \int_U |D_y^\sigma \varphi(y)|^2 d\mu(y) \leq C \|\varphi\|_{H^k(U)}^2. \end{aligned}$$

Therefore, $\|\varphi^\#\|_{H^k(SM)} \leq C \|\varphi\|_{H^k(\partial_+ SM)}$ for $\varphi \in C_\alpha^\infty(\partial_+ SM)$.

If $\varphi \in H_\alpha^k(\partial_+ SM)$, since $C_\alpha^\infty(\partial_+ SM)$ is dense in $H_\alpha^k(\partial_+ SM)$, by an approximation argument, it is easy to show that $\varphi^\sharp \in H^k(SM)$ and the operator I^* is bounded, which proves the lemma. \square

Combining the two lemmas above, we obtain the desired regularity property of I_m^* .

Proposition 5.3. *Given a compact nontrapping Riemannian manifold M with strictly convex boundary, the adjoint operator of the geodesic ray transform on symmetric m -tensors*

$$I_m^* = L_m \circ I^* : H_\alpha^k(\partial_+ SM) \rightarrow H^k(S^m(T^*M))$$

is bounded for any integer $k \geq 0$.

Now we can extend the definition of the geodesic ray transform so that it acts on $(H^k(S^m(T^*M)))^*$ (the dual space is with respect to the L^2 inner product) for integers $k \geq 1$. Let $u \in (H^k(S^m(T^*M)))^*$ and $\varphi \in H_\alpha^k(\partial_+ SM)$. We define $I_m u$ in the sense of distributions:

$$(I_m u, \varphi) := (u, I_m^* \varphi). \tag{15}$$

By Proposition 5.3, the right-hand side of (15) is well-defined. We derive the following corollary:

Corollary 5.4. *Given M , a compact nontrapping manifold with strictly convex boundary, the operator*

$$I_m : (H^k(S^m(T^*M)))^* \rightarrow (H_\alpha^k(\partial_+ SM))^*$$

defined by (15) is bounded.

Here the dual space $(H_\alpha^k(\partial_+ SM))^*$ is also with respect to the L^2 inner product. Note $H_0^k(\partial_+ SM) \subset H_\alpha^k(\partial_+ SM)$; thus $(H_\alpha^k(\partial_+ SM))^* \subset H^{-k}(\partial_+ SM)$. On the other hand, since $C^\infty(S^m(T^*M))$ is dense in $H^k(S^m(T^*M))$ under the H^k -norm, it is clear that $H_c^{-k}(S^m(T^*M^{\text{int}})) \subset (H^k(S^m(T^*M)))^*$; we will use the weaker map in the next section:

$$I_m : H_c^{-k}(S^m(T^*M^{\text{int}})) \rightarrow H^{-k}(\partial_+ SM). \tag{16}$$

6. Proof of Theorem 1.2

Now we are in a position to prove our main theorem. We start by showing that (1), (2) and (3) are equivalent.

Proof. (1) \Rightarrow (2): Since M is simple, given $u \in L^2(S_{\text{sol}}^m(T^*M))$, by Lemma 4.2, there exists $v \in H_c^{-1}(S^m(T^*\tilde{M}))$ such that $r_M I_m^* I_m v = u$. Then (16) implies the existence of some $\tilde{\varphi} = I_m v \in H^{-1}(\partial_+ S\tilde{M})$ such that $u = r_M I_m^* \tilde{\varphi}$. For $w \in H_0^1(S^m(T^*M))$, we define the distribution φ acting on $I_m(H_0^1(S^m(T^*M)))$ by

$$(\varphi, I_m w) := (\tilde{\varphi}, I_m \tilde{w}) = (I_m^* \tilde{\varphi}, \tilde{w}),$$

where $\tilde{w} \in H_0^1(S^m(T^*\tilde{M}))$ is the extension of w which is zero outside M . We claim that there exists $C > 0$ such that

$$|(\varphi, I_m w)| \leq C \|I_m w\|_{H^1}$$

for all $w \in H_0^1(S^m(T^*M))$. Assuming the claim, note that $I_m w \in H_0^1(\partial_+ SM)$ and by the Hahn–Banach theorem, φ can be extended to a bounded linear functional on $H_0^1(\partial_+ SM)$, still denoted by φ , i.e., $\varphi \in H^{-1}(\partial_+ SM)$. By the definition of φ ,

$$|(\varphi, I_m w)| = |(\tilde{\varphi}, I_m \tilde{w})| \leq C \|I_m \tilde{w}\|_{H^1}.$$

Therefore to prove the claim, it suffices to show that

$$\|I_m \tilde{w}\|_{H^1(\partial_+ S\tilde{M})} \leq C \|I_m w\|_{H^1(\partial_+ SM)} \tag{17}$$

for some $C > 0$.

Assume at this point that inequality (17) holds and let us continue with the proof. Now $\varphi \in H^{-1}(\partial_+ SM)$ is well-defined. Let $w \in H_0^1(S^m(T^*M))$, and let \tilde{w} be the extension of w into \tilde{M} which is zero outside M , so $\tilde{w} \in H_0^1(S^m(T^*\tilde{M}))$. Then

$$(r_M I_m^* \tilde{\varphi}, w) = (I_m^* \tilde{\varphi}, \tilde{w}) = (\tilde{\varphi}, I_m \tilde{w}) = (\varphi, I_m w) = (I_m^* \varphi, w).$$

Thus $u = r_M I_m^* \tilde{\varphi} = I_m^* \varphi$. (The choice of φ is not unique.)

(2) \Rightarrow (3): Given $u \in L^2(S_{\text{sol}}^m(T^*M))$, by the assumption, there is $\varphi \in H^{-1}(\partial_+ SM)$ such that $u = I_m^* \varphi$. Since $I_m^* = L_m \circ I^*$, we define $f = I^* \varphi$; then $f \in H^{-1}(SM)$ and $u = L_m f$. Furthermore, given $h \in H_0^2(SM)$,

$$(Xf, h) = (f, -Xh) = (I^* \varphi, -Xh) = (\varphi, -I(Xh)) = 0,$$

i.e., $Xf = 0$.

(3) \Rightarrow (1): Assume $I_m u = 0$ for some $u \in C^\infty(S_{\text{sol}}^m(T^*M))$. Then it is well known that there exists $h \in C^\infty(SM)$ with $h|_{\partial SM} = 0$ such that

$$Xh = -\ell_m u.$$

Moreover, by [Sharafutdinov 2002, Lemma 2.3] there exists $p \in C^\infty(S^{m-1}(T^*M))$ with $p|_{\partial M} = 0$ such that $u|_{\partial M} = dp|_{\partial M}$. When $m = 0$, this just means $u|_{\partial M} = 0$. Calculations in local coordinates show that $X(\ell_{m-1} p) = \ell_m dp$. Thus we obtain

$$X(h + \ell_{m-1} p) = -\ell_m (u - dp),$$

with $(h + \ell_{m-1} p)|_{\partial SM} = 0$.

Under the projection $\pi : SM \rightarrow M$, the pullback of the unit normal vector ν to ∂M is the unit normal vector μ to ∂SM , and in local coordinates

$$X = \xi^i \frac{\partial}{\partial x^i} - \Gamma_{jk}^i \xi^j \xi^k \frac{\partial}{\partial \xi^i},$$

where Γ_{jk}^i are the Christoffel symbols. By taking the boundary normal coordinates (x', x^n) near $x \in \partial M$ (so $\nu(x) = \mu(x, \xi) = \partial/\partial x^n$), together with the fact that $(h + \ell_{m-1} p)|_{\partial SM} = 0$, we obtain that for $(x, \xi) \in \partial SM$,

$$0 = -\ell_m (u - dp)(x, \xi) = X(h + \ell_{m-1} p)(x, \xi) = \xi^n \partial_{x^n} (h + \ell_{m-1} p)(x, \xi).$$

The first equality comes from the fact $u - dp|_{\partial M} = 0$. Thus $\partial_\mu(h + \ell_{m-1}p)(x, \xi) = 0$ for all $\xi \notin S_x\partial M$. But since h and p are smooth, and the measure of $S_x\partial M$ is zero on S_xM , we get $\partial_\mu(h + \ell_{m-1}p)(x, \xi) = 0$ for all $\xi \in S_xM$, so $h + \ell_{m-1}p \in H_0^2(SM)$.

On the other hand, there exists $f \in H^{-1}(SM)$ with $Xf = 0$ such that $u = L_m f$. It follows that

$$0 = (Xf, h + \ell_{m-1}p) = (f, -X(h + \ell_{m-1}p)) = (f, \ell_m(u - dp)) = (L_m f, u - dp) = \|u\|^2,$$

where the last equality comes from the fact that u is orthogonal to dp . Thus $u = 0$, which implies the s-injectivity. □

Remark 6.1. By carrying out an argument similar to the one of [Stefanov and Uhlmann 2005, Lemma 4.1], one can actually show that there exists $p \in C^\infty(S^{m-1}(T^*M))$ with $p|_{\partial M} = 0$ such that $\partial_\nu^k u|_{\partial M} = \partial_\nu^k dp|_{\partial M}$ for all integers $k \geq 0$. When $m = 0$, this means the boundary jet of u is zero, i.e., $\partial_\nu^k u|_{\partial M} = 0$ for all $k \geq 0$. Note that [Stefanov and Uhlmann 2005] only considers the case that u is a symmetric 2-tensor field, but the proof works for tensors of any rank. On the other hand, given $\partial_\nu^k u|_{\partial M} = \partial_\nu^k dp|_{\partial M}$, one should be able to prove that $h + \ell_{m-1}p \in H_0^{k+2}(SM)$ for all $k \geq 0$, i.e., $h + \ell_{m-1}p$ also has zero boundary jet. However, for our purposes $k = 0$ is enough.

The thing left to prove is the inequality (17). Actually the H^k norms of $I_m w$ and $I_m \tilde{w}$ are equivalent for arbitrary $k \geq 0$, provided that w is in $H_0^k(S^m(T^*M))$. A simple calculation shows that $\|I_m \tilde{w}\|_{L^2}^2 = (\tilde{w}, I_m^* I_m \tilde{w}) = (w, r_M I_m^* I_m \tilde{w}) = (w, I_m^* I_m w) = \|I_m w\|_{L^2}^2$. We assume $\partial \tilde{M}$ and ∂M are sufficiently close.

Lemma 6.2. *Let M be a compact nontrapping manifold with strictly convex boundary. Given $w \in H_0^k(S^m(T^*M))$, $k \geq 1$, let $\tilde{w} \in H_0^k(S^m(T^*\tilde{M}))$ be the extension of w to \tilde{M} by zero. Then there exists $C > 1$ such that*

$$\frac{1}{C} \|I_m w\|_{H^k(\partial_+ SM)} \leq \|I_m \tilde{w}\|_{H^k(\partial_+ S\tilde{M})} \leq C \|I_m w\|_{H^k(\partial_+ SM)}. \tag{18}$$

Proof. We only need to show (17), which is half of (18). Since ∂M and $\partial \tilde{M}$ are close, we can assume the closure of \tilde{M} is still compact nontrapping with strictly convex boundary. Given a geodesic $\gamma_{x,\xi}$ on M determined by $(x, \xi) \in \partial_+ SM$, we can uniquely extend it to a geodesic $\gamma_{y,\eta}$ on \tilde{M} determined by $(y, \eta) \in \partial_+ S\tilde{M}$. It is not difficult to see that the map

$$T : \partial_+ SM \rightarrow \partial_+ S\tilde{M}, \quad \text{with } T(x, \xi) = (y, \eta),$$

is a diffeomorphism from $\partial_+ SM$ onto its image $T(\partial_+ SM)$. On the other hand, by the definition of \tilde{w} , $I_m w(x, \xi) = I_m \tilde{w}(T(x, \xi)) = I_m \tilde{w}(y, \eta)$ and $I_m \tilde{w}(y, \eta) = 0$ for $(y, \eta) \in \partial_+ S\tilde{M} \setminus T(\partial_+ SM)$.

Since $\partial_+ SM$ and $\partial_+ S\tilde{M}$ are compact, similar to the proofs of Lemmas 5.1 and 5.2, we will work in local charts. Let U be a domain in $\partial_+ S\tilde{M}$ with local coordinates $(\tilde{z}^1, \dots, \tilde{z}^{2n-2})$ and φ be a smooth function on $\partial_+ S\tilde{M}$ with $\text{supp } \varphi \subset U$. In the mean time, there is a domain V in $\partial_+ SM$ with local coordinates (z^1, \dots, z^{2n-2}) such that $T^{-1}(U \cap T(\partial_+ SM)) \subset V$, and ψ is a smooth function on $\partial_+ SM$ with $T^{-1}(U \cap T(\partial_+ SM)) \subset \text{supp } \psi \subset V$ and $\psi \equiv 1$ on $T^{-1}(U \cap T(\partial_+ SM))$. We first consider the case $w \in C_c^\infty(S^m(T^*M)^{\text{int}})$ and show that there exists $C > 0$ such that

$$\|\varphi \cdot I_m \tilde{w}\|_{H^k(U)} \leq C \|\psi \cdot I_m w\|_{H^k(V)}.$$

Notice that for $|\alpha| \leq k$,

$$D_{\bar{z}}^\alpha[\varphi \cdot I_m \tilde{w}] = \sum_{\beta+\gamma=\alpha} D_{\bar{z}}^\gamma \varphi \cdot D_{\bar{z}}^\beta I_m \tilde{w}.$$

Thus

$$\begin{aligned} \|D_{\bar{z}}^\alpha[\varphi \cdot I_m \tilde{w}]\|_{L^2(U)}^2 &\leq \sum_{\beta \leq \alpha} C_{\beta, \alpha} \int_U |D_{\bar{z}}^\beta I_m \tilde{w}|^2 d\bar{z} \\ &= \sum_{\beta \leq \alpha} C_{\beta, \alpha} \int_{U \cap T(\partial_+ SM)} |D_{\bar{z}}^\beta I_m \tilde{w}(\bar{z})|^2 d\bar{z} \\ &\leq \sum_{|\sigma| \leq |\alpha|} C_{\sigma, \alpha} \int_{T^{-1}(U \cap T(\partial_+ SM))} |D_z^\sigma I_m \tilde{w}(T(z))|^2 J dz \\ &\leq C' \sum_{|\sigma| \leq |\alpha|} \int_{T^{-1}(U \cap T(\partial_+ SM))} |D_z^\sigma(\psi \cdot I_m w)(z)|^2 dz \\ &\leq C' \sum_{|\sigma| \leq |\alpha|} \int_V |D_z^\sigma(\psi \cdot I_m w)(z)|^2 dz \leq C \|\psi \cdot I_m w\|_{H^k(V)}^2, \end{aligned}$$

where J is the Jacobian related to the diffeomorphism T . Therefore

$$\|I_m \tilde{w}\|_{H^k(\partial_+ S\tilde{M})} \leq C \|I_m w\|_{H^k(\partial_+ SM)}$$

for $w \in C_c^\infty(S^m(T^*M)^{\text{int}})$.

Now for $w \in H_0^k(S^m(T^*M))$, there is a sequence $w_k \in C_c^\infty(S^m(T^*M)^{\text{int}})$, $k = 1, 2, \dots$, which converges to w in the H^k norm. Then it is not difficult to see that the sequence $\tilde{w}_k \in C_c^\infty(S^m(T^*\tilde{M}))$ converges to $\tilde{w} \in H_0^k(S^m(T^*\tilde{M}))$. By the boundedness of the operator I_m , we know $I_m w_k$ and $I_m \tilde{w}_k$ converge to $I_m w$ and $I_m \tilde{w}$ respectively in the H^k norm. This implies that above estimates are valid for any $w \in H_0^k(S^m(T^*M))$. \square

The following proposition that holds on compact nontrapping manifolds with strictly convex boundary shows that items (4) and (5) in Theorem 1.2 are equivalent and any of them implies item (1).

Proposition 6.3. *Let M be a compact nontrapping Riemannian manifold with strictly convex boundary and let $u \in C^\infty(S_{\text{sol}}^m(T^*M))$. The following are equivalent:*

- (i) *There exists $\varphi \in C_\alpha^\infty(\partial_+ SM)$ such that $u = I_m^* \varphi$.*
- (ii) *There exists $f \in C^\infty(SM)$ satisfying $Xf = 0$ and $u = L_m f$.*

Either of these two conditions implies s -injectivity of I_m .

Proof. (i) \Rightarrow (ii): By the assumption, there is $\varphi \in C_\alpha^\infty(\partial_+ SM)$ such that $u = I_m^* \varphi = L_m \circ I^* \varphi$. Define $f = I^* \varphi = \varphi^\sharp \in C^\infty(SM)$ (since $\varphi \in C_\alpha^\infty(\partial_+ SM)$); then $u = L_m f$. Moreover, it is clear that $Xf = X\varphi^\sharp = 0$ by definition.

(ii) \Rightarrow (i): If there exists $f \in C^\infty(SM)$ with $Xf = 0$, this implies that $f = I^*(f|_{\partial_+ SM})$. We define $\varphi = f|_{\partial_+ SM} \in C^\infty(\partial_+ SM)$. However, since $\varphi^\sharp = f \in C^\infty(SM)$, we know φ actually sits in the space $C_\alpha^\infty(\partial_+ SM)$. By the assumption, $u = L_m f = L_m \circ I^* \varphi = I_m^* \varphi$.

The argument that shows that any of these conditions imply s-injectivity of I_m is even easier than the proof that (3) implies (1) in Theorem 1.2 since we do not have to worry about paring Xf with an element in $H_0^2(SM)$. Assuming (ii), integration by parts yields right away that

$$0 = (Xf, h) = (f, -Xh) = (f, \ell_m(u)) = (L_m f, u) = \|u\|^2. \quad \square$$

Finally we show that in Theorem 1.2, item (1) implies item (4):

Since M is simple, given $u \in C^\infty(S_{\text{sol}}^m(T^*M))$, by Lemma 4.3, there exists $v \in C_c^\infty(S^m(T^*\tilde{M}))$ such that $r_M I_m^* I_m v = u$. Then it is a standard argument that if we define $\varphi = I^*(I_m v)|_{\partial_+ SM}$, then $I_m^* \varphi = u$. Moreover, since $I^*(I_m v)$ is smooth in the interior of $S\tilde{M}$, we have $\varphi \in C_\alpha^\infty(\partial_+ SM)$.

The proof of Theorem 1.2 is now complete.

7. Alternative proof of Corollary 1.3

Before giving the alternative proof, we will explain how the solenoidal condition of a tensor manifests itself at the level of the transport equation. It seems that this basic relation has not appeared before in the literature, although we believe it was known to experts.

As we already pointed out in the Introduction, by considering the vertical Laplacian Δ on each fiber $S_x M$ of SM , we have a natural L^2 decomposition $L^2(SM) = \bigoplus_{m \geq 0} H_m(SM)$ into vertical spherical harmonics. We set $\Omega_m := H_m(SM) \cap C^\infty(SM)$. Then a function u belongs to Ω_m if and only if $\Delta u = m(m+n-2)u$, where $n = \dim M$. The maps

$$\ell_m : C^\infty(S^m(T^*M)) \rightarrow \bigoplus_{k=0}^{[m/2]} \Omega_{m-2k}$$

and

$$L_m : \bigoplus_{k=0}^{[m/2]} \Omega_{m-2k} \rightarrow C^\infty(S^m(T^*M))$$

are isomorphisms. These maps give natural identification between functions in Ω_m and *trace-free* symmetric m -tensors (for details on this, see [Guillemin and Kazhdan 1980b; Dairbekov and Sharafutdinov 2010; Paternain et al. 2015a]). The geodesic vector field X maps Ω_m to $\Omega_{m-1} \oplus \Omega_{m+1}$ and hence we can split it as $X = X_+ + X_-$, where $X_\pm : \Omega_m \rightarrow \Omega_{m\pm 1}$ and $X_+^* = -X_-$. Note that

$$X \ell_{m-1} = \ell_m d.$$

Given $f \in \bigoplus_{k=0}^{[m/2]} \Omega_{m-2k}$, in general $Xf \in \bigoplus_{k=0}^{[(m+1)/2]} \Omega_{m+1-2k}$. The next simple lemma characterizes the solenoidal condition in terms of Xf .

Lemma 7.1. *$Xf \in \Omega_{m+1}$ if and only if $L_m f$ is a solenoidal tensor.*

Proof. Note that $L_m f$ is solenoidal if and only if $(L_m f, dh) = 0$ for any $h \in C^\infty(S^{m-1}(T^*M))$ with $h|_{\partial M} = 0$. But

$$(L_m f, dh) = (f, \ell_m dh) = (f, X \ell_{m-1} h) = -(Xf, \ell_{m-1} h)$$

and the last term is zero if and only if $(Xf)_{m-2k-1} = 0$ for $0 \leq k \leq [(m-1)/2]$ since $\ell_{m-1}h \in \bigoplus_{k=0}^{[(m-1)/2]} \Omega_{m-1-2k}$. \square

Another way to look at the condition $Xf \in \Omega_{m+1}$ is that the following equations should hold:

$$X_- f_{m-2k} + X_+ f_{m-2k-2} = 0 \quad \text{for } 0 \leq k \leq [(m-1)/2].$$

Lemma 7.2. *The following are equivalent:*

- (1) *Given a nonnegative integer m and $a_m \in \Omega_m$ with $X_- a_m = 0$, there exists $w \in C^\infty(SM)$ such that $Xw = 0$ and $w_m = a_m$.*
- (2) *Given a nonnegative integer m and $f = \sum_{k=0}^m f_k$ such that $Xf \in \Omega_m \oplus \Omega_{m+1}$, there exists $w \in C^\infty(SM)$ such that $Xw = 0$ and $\sum_{k=0}^m w_k = f$.*

Proof. The fact that (2) implies (1) is quite obvious from the fact that $a_m \in \Omega_m$ with $X_- a_m = 0$ implies $Xa_m = X_+ a_m \in \Omega_{m+1}$.

To prove that (1) implies (2) we proceed by induction on m . The case $m = 0$ follows right away since $Xf_0 \in \Omega_1$ and $X_- f_0 = 0$.

Suppose the claim holds for m and let $f = \sum_{k=0}^{m+1} f_k$ be given with $Xf \in \Omega_{m+1} \oplus \Omega_{m+2}$. This is equivalent to saying that $X(\sum_{k=0}^m f_k) \in \Omega_m \oplus \Omega_{m+1}$ and $X_- f_{m+1} + X_+ f_{m-1} = 0$.

By the induction hypothesis, there exists $w \in C^\infty(SM)$ such that $Xw = 0$ and $w_k = f_k$ for all $k \leq m$. The equation $Xw = 0$ in degree m is

$$X_- w_{m+1} + X_+ f_{m-1} = 0$$

and thus

$$X_-(f_{m+1} - w_{m+1}) = 0.$$

Using item (1) in the lemma, there exists $w' = \sum_{k=0}^{m+1} w'_k \in C^\infty(SM)$ such that $Xw' = 0$ and $w'_{m+1} = f_{m+1} - w_{m+1}$. Then $X(w + w') = 0$ and $\sum_{k=0}^{m+1} (w + w')_k = f$ as desired. \square

Finally we show:

Proposition 7.3. *The following are equivalent:*

- (1) *Given a nonnegative integer m and $u \in C^\infty(S_{\text{sol}}^m(T^*M))$, there exists $f \in C^\infty(SM)$ with $Xf = 0$ such that $L_m f = u$.*
- (2) *Given a nonnegative integer m and $a_m \in \Omega_m$ with $X_- a_m = 0$, there exists $w \in C^\infty(SM)$ such that $Xw = 0$ and $w_m = a_m$.*

Proof. Assume (1) holds. Given $a_m \in \Omega_m$ with $X_- a_m = 0$, we see using Lemma 7.1 that $L_m a_m$ is a solenoidal tensor. Hence there is f such that $Xf = 0$ and $f_m = L_m^{-1} L_m f = a_m$ (note that $L_m f_k = 0$ for $k > m$). Thus (2) holds.

Conversely if (2) holds, then item (2) in Lemma 7.2 holds. Thus there exists $f \in C^\infty(SM)$ such that $Xf = 0$ and $\sum_{k=0}^{[m/2]} f_{m-2k} = L_m^{-1} a_m$ and (1) holds. \square

Proof of Corollary 1.3. On account of Proposition 7.3, it suffices to show that given $a_m \in \Omega_m$ with $X_-a_m = 0$, there exists $w \in C^\infty(SM)$ such that $Xw = 0$ and $w_m = a_m$. What makes this possible in dimension two is [Paternain et al. 2015b, Lemma 5.6], whose content we now explain.

If (M, g) is an oriented Riemannian surface, there is a global orthonormal frame $\{X, X_\perp, V\}$ of SM equipped with the Sasaki metric, where X is the geodesic vector field, V is the vertical vector field and $X_\perp = [X, V]$. We define the Guillemin–Kazhdan operators [1980a]

$$\eta_\pm = \frac{1}{2}(X \pm iX_\perp).$$

If $x = (x_1, x_2)$ are oriented isothermal coordinates near some point of M , we obtain local coordinates (x, θ) on SM , where θ is the angle between ξ and $\partial/\partial x_1$. In these coordinates $V = \partial/\partial\theta$ and η_+ and η_- are ∂ - and $\bar{\partial}$ -type operators; see [Paternain et al. 2015a, Appendix B].

For any $m \in \mathbb{Z}$ we define

$$\Lambda_m = \{u \in C^\infty(SM) : Vu = imu\}.$$

In the (x, θ) -coordinates elements of Λ_m look locally like $h(x)e^{im\theta}$. Spherical harmonics may be further decomposed as

$$\begin{aligned} \Omega_0 &= \Lambda_0, \\ \Omega_m &= \Lambda_m \oplus \Lambda_{-m} \quad \text{for } m \geq 1. \end{aligned}$$

Any $u \in C^\infty(SM)$ has a decomposition $u = \sum_{m=-\infty}^\infty u_m$, where $u_m \in \Lambda_m$. The geodesic vector field decomposes as

$$X = \eta_+ + \eta_-,$$

where $\eta_\pm : \Lambda_m \rightarrow \Lambda_{m\pm 1}$. If $m \geq 1$, the action of X_\pm on Ω_m is given by

$$X_\pm(e_m + e_{-m}) = \eta_\pm e_m + \eta_\mp e_{-m}, \quad e_j \in \Lambda_j,$$

and for $m = 0$, we have $X_+|_{\Omega_0} = \eta_+ + \eta_-$ and $X_-|_{\Omega_0} = 0$.

With these preliminaries out of the way, [Paternain et al. 2015b, Lemma 5.6] says that given $f \in \Lambda_m$, there is a smooth $w \in C^\infty(SM)$ with $Xw = 0$ and $w_m = f$. For $m = 0$, this gives the desired result right away.

Given $a_m \in \Omega_m$ with $X_-a_m = 0$ and $m \geq 1$, we write $a_m = e_m + e_{-m}$ with $e_j \in \Lambda_j$. Then $\eta_-e_m + \eta_+e_{-m} = 0$. Consider now smooth p, q with $Xp = Xq = 0$ and $p_m = e_m$ and $q_{-m} = e_{-m}$. Then

$$w = \sum_{k=-\infty}^{-m} q_k + \sum_{k=m}^{\infty} p_k$$

satisfies $Xw = 0$ and $w_m = a_m$. □

Acknowledgements

We are very grateful to Mikko Salo and Gunther Uhlmann for several discussions related to this paper; our main result is motivated by a question raised in [Paternain et al. 2015a, Remark 11.7]. We are also very

grateful to Boris Kruglikov and Vladimir Matveev for extensive discussions related to Killing tensors and for the input provided in [Kruglikov and Matveev 2016]. Finally we would like to thank Colin Guillarmou and Plamen Stefanov for invaluable comments on the first draft. In particular, Stefanov provided us with the proof of Lemma 3.5 and Guillarmou with the first part of Remark 4.4. This research was supported by EPSRC grant EP/M023842/1.

References

- [Anikonov and Romanov 1997] Y. E. Anikonov and V. G. Romanov, “On uniqueness of determination of a form of first degree by its integrals along geodesics”, *J. Inverse Ill-Posed Probl.* **5**:6 (1997), 487–490. MR Zbl
- [Dairbekov and Sharafutdinov 2010] N. S. Dairbekov and V. A. Sharafutdinov, “Conformal Killing symmetric tensor fields on Riemannian manifolds”, *Mat. Tr.* **13**:1 (2010), 85–145. In Russian; translated in *Siberian Advances in Mathematics* **21**:1 (2011), 1–41. MR Zbl
- [Dairbekov and Uhlmann 2010] N. Dairbekov and G. Uhlmann, “Reconstructing the metric and magnetic field from the scattering relation”, *Inverse Probl. Imaging* **4**:3 (2010), 397–409. MR Zbl
- [Delay 2012] E. Delay, “Smooth compactly supported solutions of some underdetermined elliptic PDE, with gluing applications”, *Comm. Partial Differential Equations* **37**:10 (2012), 1689–1716. MR Zbl
- [Guillarmou 2014] C. Guillarmou, “Invariant distributions and X-ray transform for Anosov flows”, preprint, 2014. To appear in *J. Differential Geom.* arXiv
- [Guillemin and Kazhdan 1980a] V. Guillemin and D. Kazhdan, “Some inverse spectral results for negatively curved 2-manifolds”, *Topology* **19**:3 (1980), 301–312. MR Zbl
- [Guillemin and Kazhdan 1980b] V. Guillemin and D. Kazhdan, “Some inverse spectral results for negatively curved n -manifolds”, pp. 153–180 in *Geometry of the Laplace operator* (Honolulu, HI, 1979), Proc. Sympos. Pure Math. **36**, Amer. Math. Soc., Providence, RI, 1980. MR Zbl
- [Han 2011] Q. Han, *A basic course in partial differential equations*, Graduate Studies in Mathematics **120**, American Mathematical Society, Providence, RI, 2011. MR Zbl
- [Helgason 1999] S. Helgason, *The Radon transform*, 2nd ed., Progress in Mathematics **5**, Birkhäuser, Boston, 1999. MR Zbl
- [Kato et al. 2000] T. Kato, M. Mitrea, G. Ponce, and M. Taylor, “Extension and representation of divergence-free vector fields on bounded domains”, *Math. Res. Lett.* **7**:5–6 (2000), 643–650. MR Zbl
- [Kruglikov and Matveev 2016] B. Kruglikov and V. S. Matveev, “The geodesic flow of a generic metric does not admit nontrivial integrals polynomial in momenta”, *Nonlinearity* **29**:6 (2016), 1755–1768. MR Zbl
- [McLean 2000] W. McLean, *Strongly elliptic systems and boundary integral equations*, Cambridge University Press, 2000. MR Zbl
- [Muhometov 1977] R. G. Muhometov, “The reconstruction problem of a two-dimensional Riemannian metric, and integral geometry”, *Dokl. Akad. Nauk SSSR* **232**:1 (1977), 32–35. In Russian. MR
- [Paternain et al. 2012] G. P. Paternain, M. Salo, and G. Uhlmann, “The attenuated ray transform for connections and Higgs fields”, *Geom. Funct. Anal.* **22**:5 (2012), 1460–1489. MR Zbl
- [Paternain et al. 2013] G. P. Paternain, M. Salo, and G. Uhlmann, “Tensor tomography on surfaces”, *Invent. Math.* **193**:1 (2013), 229–247. MR Zbl
- [Paternain et al. 2014] G. P. Paternain, M. Salo, and G. Uhlmann, “Spectral rigidity and invariant distributions on Anosov surfaces”, *J. Differential Geom.* **98**:1 (2014), 147–181. MR Zbl
- [Paternain et al. 2015a] G. P. Paternain, M. Salo, and G. Uhlmann, “Invariant distributions, Beurling transforms and tensor tomography in higher dimensions”, *Math. Ann.* **363**:1–2 (2015), 305–362. MR Zbl
- [Paternain et al. 2015b] G. P. Paternain, M. Salo, and G. Uhlmann, “On the range of the attenuated ray transform for unitary connections”, *Int. Math. Res. Not.* **2015**:4 (2015), 873–897. MR Zbl

- [Pestov and Sharafutdinov 1988] L. N. Pestov and V. A. Sharafutdinov, “Integral geometry of tensor fields on a manifold of negative curvature”, *Sibirsk. Mat. Zh.* **29**:3 (1988), 114–130. In Russian; translated in *Siberian Mathematical Journal* **29**:3 (1988), 427–441. MR Zbl
- [Pestov and Uhlmann 2005] L. Pestov and G. Uhlmann, “Two dimensional compact simple Riemannian manifolds are boundary distance rigid”, *Ann. of Math. (2)* **161**:2 (2005), 1093–1110. MR Zbl
- [Salo and Uhlmann 2011] M. Salo and G. Uhlmann, “The attenuated ray transform on simple surfaces”, *J. Differential Geom.* **88**:1 (2011), 161–187. MR Zbl
- [Sharafutdinov 1994] V. A. Sharafutdinov, *Integral geometry of tensor fields*, VSP, Utrecht, 1994. MR Zbl
- [Sharafutdinov 1999] V. Sharafutdinov, “Ray transform on Riemannian manifolds”, lecture notes, University of Oulu, 1999, available at <http://math.nsc.ru/~sharafutdinov/files/Lectures.pdf>.
- [Sharafutdinov 2002] V. A. Sharafutdinov, “An integral geometry problem in a nonconvex domain”, *Sibirsk. Mat. Zh.* **43**:6 (2002), 1430–1442. In Russian; translated in *Siberian Math. J.* **43**:6 (2002), 1159–1168. MR Zbl
- [Sharafutdinov et al. 2005] V. Sharafutdinov, M. Skokan, and G. Uhlmann, “Regularity of ghosts in tensor tomography”, *J. Geom. Anal.* **15**:3 (2005), 499–542. MR
- [Stefanov and Uhlmann 2004] P. Stefanov and G. Uhlmann, “Stability estimates for the X-ray transform of tensor fields and boundary rigidity”, *Duke Math. J.* **123**:3 (2004), 445–467. MR Zbl
- [Stefanov and Uhlmann 2005] P. Stefanov and G. Uhlmann, “Boundary rigidity and stability for generic simple metrics”, *J. Amer. Math. Soc.* **18**:4 (2005), 975–1003. MR Zbl
- [Stefanov and Uhlmann 2008] P. Stefanov and G. Uhlmann, “Integral geometry on tensor fields on a class of non-simple Riemannian manifolds”, *Amer. J. Math.* **130**:1 (2008), 239–268. MR Zbl
- [Stefanov et al. 2014] P. Stefanov, G. Uhlmann, and A. Vasy, “Inverting the local geodesic X-ray transform on tensors”, preprint, 2014. To appear in *J. Anal. Math.* arXiv
- [Uhlmann and Vasy 2016] G. Uhlmann and A. Vasy, “The inverse problem for the local geodesic ray transform”, *Invent. Math.* **205**:1 (2016), 83–120. With an appendix by H. Zhou. MR Zbl

Received 18 Nov 2015. Revised 28 Jul 2016. Accepted 12 Sep 2016.

GABRIEL P. PATERNAIN: g.p.paternain@dpms.cam.ac.uk

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Cambridge CB3 0WB, United Kingdom

HANMING ZHOU: h318@dpms.cam.ac.uk

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Cambridge CB3 0WB, United Kingdom

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 8 2016

Estimates for radial solutions of the homogeneous Landau equation with Coulomb potential	1773
MARIA PIA GUALDANI and NESTOR GUILLEN	
Forward self-similar solutions of the Navier–Stokes equations in the half space	1811
MIKHAIL KOROBKOV and TAI-PENG TSAI	
Decay of solutions of Maxwell–Klein–Gordon equations with arbitrary Maxwell field	1829
SHIWU YANG	
Invariant distributions and the geodesic ray transform	1903
GABRIEL P. PATERNAIN and HANMING ZHOU	
Multiple vector-valued inequalities via the helicoidal method	1931
CRISTINA BENEÀ and CAMIL MUSCALU	
Structure of modular invariant subalgebras in free Araki–Woods factors	1989
RÉMI BOUTONNET and CYRIL HOUDAYER	
Finite-time blowup for a supercritical defocusing nonlinear wave system	1999
TERENCE TAO	
A long \mathbb{C}^2 without holomorphic functions	2031
LUKA BOČ THALER and FRANČ FORSTNERIČ	



2157-5045(2016)9:8;1-4