

This is a repository copy of *Interacting effects of topography, vegetation, human activities and wildland-urban interfaces on wildfire ignition risk*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/115952/>

Version: Accepted Version

---

**Article:**

Calviño-Cancela, María, Chas-Amil, María Luisa, García-Martínez, Eduardo D. et al. (1 more author) (2017) Interacting effects of topography, vegetation, human activities and wildland-urban interfaces on wildfire ignition risk. *Forest Ecology and Management*. pp. 10-17. ISSN 0378-1127

<https://doi.org/10.1016/j.foreco.2017.04.033>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

**Interacting effects of topography, vegetation, human activities and  
wildland-urban interfaces on wildfire ignition risk**

María Calviño-Cancela • María L. Chas-Amil • Eduardo D. García-Martínez • Julia  
Touza

M. Calviño-Cancela (corresponding author)

Department of Ecology and Animal Biology, University of Vigo.

Experimental Sciences Building. University Campus. 36310 Vigo, Spain;

[maria@uvigo.es](mailto:maria@uvigo.es); Phone: +34 986 818 742

M. L. Chas-Amil

Departamento de Economía Cuantitativa, Universidade de Santiago de Compostela.

Baixada Burgo das Nacións s/n. 15782 Santiago de Compostela, Spain;

[marisa.chas@usc.es](mailto:marisa.chas@usc.es)

E. D. García-Martínez

Departamento de Geografía y Ordenación del Territorio, Universidad de Zaragoza.

Facultad de Filosofía y Letras. Pedro Cerbuna 12, 50009, Zaragoza, Spain;

[edgm\\_73@yahoo.es](mailto:edgm_73@yahoo.es)

J. Touza

Environment Department, University of York.

Wentworth Way, Heslington, YO104AD, York, UK; [julia.touza@york.ac.uk](mailto:julia.touza@york.ac.uk)

## **Abstract**

Effective fire prevention requires a better understanding of the patterns and causes of fire ignition. In this study, we focus on the interacting factors known to influence fire ignition risk, such as the type of vegetation, topographical features and the wildland-urban interface (WUI; i.e. where urban development meet or intermingle with wildland). We also analyze the human activities and motivations related to fires and whether they differ depending on the type of vegetation and the location within/outside WUI. There were significant interactions between topography, type of vegetation and location within/outside WUI. The risk of ignition was in general higher at lower elevations, and this tendency was more marked in forested land covers (all plantations and open woodlands), with the noticeable exception of native forests. North-facing sites had lower fire ignition risk outside the WUI, especially in native forests, while southern aspects showed higher fire ignition risk, especially in open shrublands. However, this effect of the aspect was only significant outside WUI areas. In relation to causes, there were also interactions between human activities/motivations related to fires, the type of vegetation and the location within/outside WUI. All forestry plantations appeared clustered in relation to fire causes, especially in the WUI, with high incidence of deliberately caused fires related to violent or mentally ill people and rekindle fires. In contrast, native forests, despite structural similarities with forestry plantations, showed more similarity with agricultural areas and open woodlands in relation to fire causes. In shrublands, there was a relatively high incidence of fires related to ranching, especially outside the WUI. This pattern of interactions depicts a complex scenario in relation to fire ignition risk and prompts to the importance of taking this complexity into account in order to adjust fire management measures for improved effectiveness.

48    **Keywords:** wildfire ignition risk, Galicia, forest fires, human-related causes,  
49    topography, Spain, WUI

50

## 51    **Abbreviations**

52    WUI: Wildland-urban interface

53    LULC: land use/land cover

54    Agr: Agriculture areas

55    OpShr: Open shrublands

56    Shr: Shrublands

57    OpWd: Open woodlands

58    AtlF: Atlantic forests

59    PiP: Pine plantations

60    EuP: Eucalypt plantations

61    MxAtl: Mixed Atlantic forests

62    MxEuPiP: Mixed plantations of pines and eucalypts

63    MxPiP: Mixed pine plantations

64    MxEuP: Mixed eucalypt plantations

65

## 1. Introduction

Fire is an important agent of change in natural ecosystems that has driven species adaptations and shaped landscapes over millions of years. As a consequence of human activities, current fire regimes have changed dramatically in many areas compared to natural regimes, causing impacts in both natural ecosystems as well as in the human society (Bowman et al. 2011). For the need to better understand fire patterns and improve fire prevention measures, there is an increasing interest on fire causes and risks. Fires occur as a consequence of both natural and human causes, with weather, topography, type of vegetation or proximity to human settlements being decisive factors in determining the likelihood of fire occurrence (e.g., Moreira et al. 2011).

The type of vegetation, as a land use/land cover (LULC) type, has been shown to be especially relevant for fire ignition risk (e.g., Bajocco and Ricotta 2008; Carmo et al. 2011; Cumming 2001; Nunes et al. 2005). Vegetation types differ in fuel loads and flammability as well as on fuel continuity, as determined by the structure of vegetation (Saura-Mas et al., 2010). For instance, in NW Spain, native forests and agricultural areas have the lowest fire ignition risk, whereas shrublands and mixed forestry plantations have the highest ignition risk (Calviño-Cancela et al. 2016). Knowledge on the fire ignition risk associated to different vegetation types can inform landscape management policy decisions, which can promote vegetation types with lower fire ignition risk.

LULCs have been shown to interact with other factors such as the proximity of human settlements (Calviño-Cancela et al. 2016). In relation to this, wildland-urban interfaces (WUIs) have been defined as areas where urban development meet or intermingle with wildland, and this interfaces are of special concern for fire risk management since fires are usually more frequent in these areas and the danger to

human lives and properties can be higher there (e.g., Cohen 2000). The only study, to our knowledge, addressing this interaction between LULC and the WUI revealed that the fire ignition risk associated to different LULC does differ between WUI and non-WUI areas, with forestry plantations showing the highest increase in the likelihood of fire occurrence in WUI compared to non-WUI areas (Calviño-Cancela et al. 2016). Topography can also interact with LULC to modify fire risk, since it affects the distribution of vegetation (e.g., agriculture fields are usually located in flat, low areas, while forest and plantations usually occupy steeper areas, less suitable for agriculture) and some abiotic factors such as temperature and moisture content of fuels (e.g., in North versus South facing slopes).

In addition to these elements, nowadays the human factor is essential to understand the patterns of fire risk. Human activities have altered fire regimes worldwide, modifying fire frequency, intensity, and size of wildfires (Bowman et al. 2011). Human-related causes, whether intentional or by accident, are the most frequent causes of fires (FAO 2007). In addition, certain human uses or activities are specifically associated to particular LULCs, being important drivers of fire risk in those LULCs. Common examples are agricultural burnings in farmlands or the periodical burnings in shrublands and grasslands to control woody encroachment and promote growth of new shoots, grasses and forbs for grazing (Ganteaume et al. 2013; Vélez 2002; Webb 1998). Similarly, socioeconomic factors, such as fragmentation of holdings, that limits the profit owners obtain from forestry products, urbanisation pressure, rural land abandonment or conflicts associated to forests' multiple uses have been shown to increase the probability of fire (e.g., Chas-Amil et al. 2015; Romero-Calcerrada et al. 2010; Yang et al. 2007). Moreover, since population density, human behaviour and activities differ markedly between WUI and non-WUI areas, human-related factors are

expected to modify the fire ignition risk associated to LULCs and topographical features depending on their location within or outside WUIs areas. Topography can also affect the risk of fire related to human causes, since human accessibility and activities can be markedly determined by topography (e.g., high and abrupt areas are less accessible).

In this study, we assess the fire ignition risk focusing on the interacting effects of LULC types, the WUI and topographical features. We also analyze the underlying causes related to fire occurrence, focusing on human activities and motivations, and how this is affected by location within or outside the WUI in different LULC types.

## **2. Materials and methods**

### **2.1. Study area**

This study was carried out in Galicia (NW of the Iberian Peninsula; Fig. 1), the most important forestry region in Spain (Manuel and Gil 2002), where c. 70% of the land is forested, mainly covered by tree plantations of *Pinus pinaster* and *Eucalyptus globulus*, in pure and mixed stands. Native forests dominated by *Quercus robur* have higher species diversity and more distinctive communities than tree plantations (Calviño-Cancela et al. 2012a, 2012b, Calviño-Cancela 2013), but are reduced to small, isolated patches (Díaz-Maroto and Vila-Lameiro 2008; Teixido et al., 2010). Eucalyptus plantations have expanded notably in the last decades, both by intentional planting and natural spread (Aguas et al. 2014, Calviño-Cancela and Rubido-Bará 2013, MAGRAMA, 2011). The frequency of wildfires is especially high in the study area: more than 30% of forest fires in Spain each year occur in this region, mainly associated with intentional behavior (75%) (MAGRAMA 2012).

### **2.2 Data**

A database of 26,838 wildfire reports for the period January 1<sup>st</sup>, 2006 to December 31<sup>st</sup>, 2011 obtained from the Spanish Forest Fire Statistics (EGIF) was used in this study, which includes the coordinates of each ignition point (see Calviño-Cancela et al. (2016) for details). Fire causes and motivations given in fire reports were grouped in 12 categories focusing, for deliberate and negligent fires, on human activities and behaviours to which the fire ignition was related: agriculture and vegetation management (including agricultural burnings and fires related to verge maintenance), ranching (fires related to pasture regeneration), forestry management, hunting, recreation, waste management (rubbish burning), profit gaining, conflicts, mentally ill or violent people, accidents, natural (lighting) and rekindle (Table 1). Fires caused by “other negligences”, “other deliberate causes” and with “unknown causes” were excluded from the study, due to the lack of definition of these categories, as they may include very different causes.

For each fire ignition point we determined the land use/land cover type (LULC), its topographic features (slope, elevation, and aspect) and the location inside or outside of the WUI. We determined the LULC type using information from the Fourth National Forest Inventory (IFN4, MAGRAMA 2011; see Calviño-Cancela et al. (2016) for further details). Areas with no or very scarce vegetation (e.g., water bodies, beaches, or artificial surfaces such as industrial or urban areas) were excluded, as well as the less frequent LULCs (grasslands, Mediterranean forests and Acacia woods), due to the low number of fires in WUI in these categories. WUI was defined as the area within a 50 m radius around buildings at a distance of up to 400 m from wildland vegetation (Law 3/2007 of April 9, 2007, addressing the issues of wildfire prevention and suppression, as modified by Law 7/2012 of June 28, 2012 of Galician Forestry). The mapping of WUIs in Galicia was obtained from Chas-Amil et al. (2013).

Topographic variables were calculated using the Spatial Analyst extension to ArcGIS® 10.2.2 by ESRI to derive the slope, elevation and aspect, based on a 10 m spatial resolution digital elevation model (DEM, 1:5,000 scale), developed by SITGA (Galician Territorial Information System). The slope was defined as a percentage and elevation in meters. Aspect was defined as the compass direction that the slope faces: N (315° to 360° and 0° to 45°), E (45° to 135°), S (135° to 225°) or W (225° to 315°) direction.

### **2.3. Data analyses**

In order to compare the patterns of distribution of ignition points with a random model, we selected 26,838 random locations in the region and characterized them in regard to WUI, LULCs and topography, as done for ignition points. To select random points we used the module Random Points Generation of Hawth's Analysis Tools, in ArcGIS. Then, we obtained 100 samples with 5,000 locations each, out of the 26,838 fire ignition and random points, using a Montecarlo method (bootstrapping; random resampling with replacement; Efron 1982, Manly 1998).

In relation to topography, we tested whether there were differences in elevation between ignition and random points, WUI/non-WUI areas and LULCs with ANOVA, with Random/Fire, WUI/non-WUI and LULCs as fixed factors and the elevation as the variate. For differences in slope, we followed the same approach but using a generalized linear model with the negative binomial distribution and logratio as the link function, because slope followed a negative binomial distribution instead of a normal distribution. To analyse the effect of the aspect (N, S, E and W, a categorical variable), we calculated the proportional differences between the number of fires recorded in each combination of topographic features x LULCs x within/outside WUI and that in the random set, which corresponds to the expected number according to a random probability. This is

equivalent to selection indexes used in other studies (e.g., Moreira et al. 2001; Bajocco and Ricotta 2008), since proportional differences are the observed minus the expected frequencies divided by the expected frequencies. We performed an ANOVA with LULC and within/outside WUI as fixed factors and the proportional differences between the fire and random sets in each compass aspect (N, S, E and W) as variates.

In relation to causes, we used again a Montecarlo method to resample from the original set of ignition points, obtaining 100 samples with 100 cases per each LULC category within and outside the WUI. Shrublands and Open shrublands were pooled together for this analysis. We used PERMANOVA to analyse differences in the causes of fires as affected by LULC and location within/outside the WUI (fixed factors). The proportion of fires for each cause in each combination of LULC and WUI/non-WUI was used as the variate. We used 9999 permutations for the analyses, with type III sums of squares, fixed effects sum set to zero and permutation of residuals under a reduced model. We used NMDS (non-metric multidimensional scaling) ordinations to represent graphically the difference between LULCs within and outside the WUI, showing the distance between LULCs in the fire-causes space. For the sake of clarity in figure 5, we used only 30 randomly chosen samples out of the 100 samples per LULC. To represent the main fire causes driving the patterns of distances (divergences) between LULCs in the NMDS ordinations, we calculated the spearman rank correlation of each fire cause with the axes and represented those with  $r > 0.5$ . Both PERMANOVA and NMDS ordinations were based on Bray-Curtis similarities of square-root transformed data. We used PRIMER 6.1.12 (Clarke and Gorley 2006) with the PERMANOVA + 1.0.2 add-on (Anderson et al. 2008) for these analyses.

### **3. Results**

### 3.1. Topography

We found 2<sup>nd</sup> and 3<sup>rd</sup> order interactions of Ignition/Random with WUI/non-WUI and LULCs in relation to elevation (Table 2) and slope (Table 3), which means that the divergence in elevation and slope between random and ignition points differed between WUI and non-WUI areas and depending on the LULC. For elevation, ignition points had in general lower elevation than random points (Fig. 2). This pattern was noticeable in tree plantations, with 14-39% lower elevation in ignition points, whereas differences in the rest of LULCs were lower than 10%. In addition to tree plantations, this pattern was also noticeable in mixed Atlantic forest and open woodlands within the WUI (17% and 15% lower, respectively), whereas ignitions in open shrublands had in the WUI higher elevations than expected under the random model (35% higher), in contrast with the similar elevation between random and ignition points outside the WUI (Fig. 2). Regarding slope (Fig. 3), agricultural areas had the lowest slope (c. 10%), that was similar in ignition (10.4%) and random points (10.1%) in non-WUI areas, with higher slopes for ignition points in the WUI (12%). Slopes were similar in the rest of LULCs, varying between 16% and 29% in random points and 16-24% in ignition points. Despite higher slopes outside the WUI in random points (18% on average), fires occurred at similar slopes within and outside the WUI, thus at flatter areas than average outside the WUI but steeper than average within the WUI, except for MxAtl, with ignition points tending to be in flatter areas in the WUI (Fig. 3).

There was a clear contrast between WUI and non-WUI areas in the risk of fire ignition associated with aspect (Table 4; Fig. 4). Outside the WUI, the percentage of fires occurring in sites facing North was lower than expected by random chance, especially in Atlantic forests (AtlF and MxAtl; Fig. 4). In contrast, southern aspects

showed the opposite pattern, especially in open shrublands (Fig. 4). Within the WUI, however, there was not a clear pattern in regard to aspect (Fig. 4).

### **3.2. Fire causes**

The PERMANOVA analysis of differences in the causes of fires revealed a significant effect of both locations within/outside the WUI and the LULC, as well as a significant interaction between these two factors (WUI:LULC, Table 5). Despite this interaction, the general pattern was similar outside and within the WUI, as showed in the distribution of LULCs in the causes space (NMDS ordinations; Fig. 5). All plantations appeared clustered in this ordination (EuP, MxEuP, PiP, MxPiP, and MxEuPiP; on the right in Fig. 5), especially outside the WUI, which reveal similarities in the causes associated with the fires occurring in these LULCs. The difference between plantations and other LULCs (shrublands, Atlantic forests, agricultural areas and open woodlands) was mostly due to a higher frequency of fires in plantations caused by violent or mentally ill people and, in a lesser degree, of rekindle fires outside the WUI, as well as a lower incidence of fires related to ranching, and to agriculture and vegetation management outside the WUI (Table 6). Shrublands appear as the most distant to plantations (Fig. 5), with Atlantic forests, agricultural areas and open woodlands occupying intermediate positions. Shrublands differ mainly because of the relatively high incidence of fires related to ranching, especially outside WUIs, and the highest frequency of fires related to hunting, although this activity caused a low number of fires (1.6%). Agricultural areas and open woodlands appear very close in the fire causes space, especially in the WUI, where they intermingle (Fig. 5). The relative importance of the different fire causes is very similar in these LULCs, especially in relation to rekindle fires, fires caused by mentally ill or violent people, and related to

agricultural and vegetation management (Table 6). Recreation was mainly related to Mixed Atlantic forests, particularly in the WUI.

## **4. Discussion**

### **4.1. Topography**

As shown in previous studies, topography had a significant effect on the risk of fire (e.g., Carmo et al. 2011, Guglietta et al. 2015, Oliveira et al. 2013) but, interestingly, this effect differed depending on the LULC and the location within or outside WUI areas. These interaction effects have not been previously explored in detail, despite their interest for management. There was a general tendency of higher fire ignition risk at lower elevations. However, this tendency was not consistent for all LULCs within/outside the WUI. It was more marked in forested land covers (all plantations and open woodlands), with the noticeable exception of native forests (AtlF). A higher fire ignition risk at lower elevations has been related to better accessibility (more and better roads at low elevation), which increases the risk of human-related fires both within and outside the WUI (Chas-Amil et al. 2015; Ganteaume et al. 2013). The impact of this increasing accessibility might have been especially important on deliberate fires, and the high incidence of fires caused by arsonists in plantations might explain the notable effect of low elevation in these land covers, in contrast with native forests, where these fires are relatively infrequent. Shrublands and agricultural areas showed also a contrasting pattern, with a striking higher ignition risk at higher elevations observed in open shrublands in the WUI. Shublands had the highest average elevations of all vegetation types considered (c. 780 m outside the WUI and c. 530 m in the WUI in contrast with an average of c. 410 m and 330 m, respectively, in the rest of LULCs), and suffer the highest ignition risk in the region (Calviño-Cancela et al. 2016).

The most common causes of ignition in this vegetation type are related to the use of fire as a tool, for vegetation management and in relation to ranching. Limited accessibility does not probably discourage ranchers and farmers in the same way as arsonists, who need a quick escape. On the other hand, at the high elevations typical of shrublands, the microclimate may play an important role, with higher elevations having dryer and windier conditions, which increase fire hazard. In addition, limited accessibility could increase the risk of spread of these fires, for it complicates fire-fighting operations.

Accessibility can also explain the higher ignition risk of flatter areas outside the WUI, where steeper areas are difficult to access. Within the WUI, fires occurred at similar steepness than outside the WUI but, since the terrain is flatter in general, these areas were steeper than average. The flattest areas within the WUI are occupied by the more valuable uses (e.g., residences or crops), more protected against fire.

In relation to aspect, lower ignition risk in northern slopes is a common pattern in temperate zones in the northern hemisphere (see e.g., González et al. 2005, Mermoz et al. 2005; but see Carmo et al. 2011). North facing slopes receive less solar radiation, which translates into lower temperatures, higher moisture contents and thus reduced flammability. The shade effect is more pronounced at lower sun elevation angles (i.e., at higher latitudes and closer to the winter solstice) and at steeper slopes. This explains the interaction with the WUI: the terrain is flatter within WUIs, which reduces the shade in north facing slopes. The effect of reducing fire ignition risk in northern slopes outside the WUI was more marked in tree covered land covers and, especially, in native forests (AtlF and MxAtl), where the dominant broadleaved trees (e.g. *Quercus robur*, *Castanea sativa*) contribute to maintain the typical fresh and humid microclimate of northern slopes and to reduce fire risk.

#### **4.2. Fire causes**

Human activities have been shown as important determinants of fire occurrence in the region. Increased fire ignition risk in WUIs is the result of the proximity of human settlements that affects the kind of activities performed in the surrounding landscape (Bar-Massada et al. 2014). Fire ignitions were most frequently related to agriculture and vegetation management, despite regulations devised to limit fire hazards (e.g., banning of agricultural burnings in summer) (Moreira et al. 2011). More awareness among citizens regarding the danger involved in this activity is thus necessary.

The distribution of LULCs in the causes space, as depicted in the ordination (Fig. 5), was very intuitive, with LULCs that seem *a priori* similar (for instance in terms of habitat structure, species composition or uses) appearing close, for the accompanying similarity in the causes of their fires. This is very revealing of the close relationship between causes and LULCs. For instance, all forestry plantations appeared clustered, especially outside the WUI, and at a certain distance from native forests (AtlF), which are very similar in structure. Note the higher incidence in forestry plantations of deliberately caused fires related to violent or mentally ill people. The economic value of these plantations may make them the target for individuals willing to cause damage to land owners. However, most fires in this category (68.6%) were assigned to pyromaniacs, which are supposed to have no conscious motivation to set fires. But the incidence of fires related to this mental disorder are often overestimated, due to the poor understanding of this condition by fire reporters and officials (Doley 2003 and references therein), which may be hiding the true conscious motivations of arsonists. AtlF appeared relatively close to agriculture (Agr), with open woodlands (OpWd) occupying intermediate positions. AtlF are expanding in some areas as a result of natural regeneration after land abandonment by farmers in rural areas (Calvo-Iglesias et

al. 2009, Corbelle-Rico et al. 2012). Thus, their proximity to active agricultural areas may explain their similarity in fire causes. This would also explain the intermediate position of OpWd, which are often transitional stages of colonization of abandoned fields towards forests or mixed formations (Calvo-Iglesias et al. 2009; Escribano-Avila et al. 2014). The relatively high incidence of fires related to ranching in shrublands, especially outside the WUI, is probably related to their use for extensive livestock grazing, since deliberate periodical burnings have been traditionally practiced in these areas to provide a flush of new growth more nutritious for grazers (Webb 1998). Shrublands are also especially important for hunting in Galicia, where hunting is centred on small game and particularly on rabbits (*Oryctolagus cuniculus*), which are most abundant in this type of habitat (Gálvez-Bravo 2011, Tapia et al. 2014).

#### **4.3. Implications for management**

Our results highlight the importance of considering the interactions between factors known to influence fire ignition risk, such as the WUI, LULCs and topography. The pattern of interactions found depicts a complex scenario in relation to fire ignition risk and prompts to the importance of taking this complexity into account in order to adjust fire management measures for improved effectiveness. A better understanding of the fire ignition risk associated with different landscape features, such as vegetation, topography and proximity to urban areas, together with the underlying human-causes of fire ignitions increases the efficiency in the allocation of fire prevention measures such as surveillance or vegetation management, and facilitates the devising of regulations or education campaigns focused on increasing citizen awareness on the fire hazards related to particular activities or behaviours in certain environments (e.g. vegetation management practices in agricultural land and native vegetation, and arsonists in forestry plantations). As commented previously, knowledge on the effect of vegetation

on the risk of fire is especially interesting for fire prevention since vegetation can be subject to active management. Our results show that other factors such as topography and location within or outside the WUI, and differences in fire causes may affect the fire proneness of vegetation types. Certain vegetation types show more fire resistance in certain contexts (e.g. Atlantic forests in northern slopes in non-WUI areas), so that they can be used, or be promoted, to reduce fire hazard at the landscape scale. On the other hand, land covers that are particularly fire-prone in certain circumstances (e.g. open shrublands in Southern slopes outside the WUI or in higher altitudes in the WUI), require increased efforts in preventing wildfire occurrence.

#### **Acknowledgments**

This research was funded in part by the Spanish Ministry of Economy and Competitiveness (Grant ECO2012-39098-C06- 05). The Spanish Ministry of Agriculture, Food and Environment (MAGRAMA) provided the wildfire database.

## References

- Aguas A, Ferreira A, Maia P, Fernandes PM, Roxo L, Keizer J, Silva JS, Rego FC, Moreira F. (2014). Natural establishment of *Eucalyptus globulus* Labill. in burnt stands in Portugal. *Forest Ecology and Management* 323, 47-56.
- Anderson MJ, Gorley RN, Clarke KR (2008) *PERMANOVA + for PRIMER: Guide to Software and Statistical Methods*. PRIMER-E, Plymouth
- Bajocco S, Ricotta C. (2008). Evidence of selective burning in Sardinia (Italy): which land-cover classes do wildfires prefer? *Landscape Ecology* 23 (2):241-248
- Bar-Massada A, Radeloff VC, Stewart SI. (2014). Biotic and abiotic effects of human settlements in the wildland-urban interface. *BioScience* 64(5):429-437
- Bowman, D. M. J. S., Balch, J., Artaxo, P., Bond, W. J., Cochrane, M. A., D'Antonio, C. M., DeFries, R., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Mack, M., Moritz, M. A., Pyne, S., Roos, C. I., Scott, A. C., Sodhi, N. S. and Swetnam, T. W. (2011). The human dimension of fire regimes on Earth. *Journal of Biogeography*, 38: 2223–2236.
- Calviño-Cancela M, Rubido-Bará M, van Etten EJB. (2012a). Do eucalypt plantations provide habitat for native forest biodiversity? *Forest Ecology and Management* 270, 153-162.
- Calviño-Cancela M, Eugenia López de Silanes M, Rubido-Bará M, Uribarri J. 2012b. The potential role of tree plantations in providing habitat for lichen epiphytes. *Forest Ecology and Management* 291, 386-395.
- Calviño-Cancela M. (2013). Effectiveness of eucalypt plantations as a surrogate habitat for birds. *Forest Ecology and Management* 310, 692-699.

402 Calviño-Cancela M, Rubido-Bará M. (2013). Invasive potential of *Eucalyptus globulus*:  
403 Seed dispersal, seedling recruitment and survival in habitats surrounding  
404 plantations. *Forest Ecology and Management* 305, 129-137.

405 Calviño-Cancela M, Chas-Amil ML, García-Martínez E, Touza J. (2016). Wildfire risk  
406 associated with different land covers within and outside wildland-urban interfaces.  
407 *Forest Ecology and Management*, 372: 1-9.

408 Calvo-Iglesias MS, Fra-Paleo U, Diaz-Varela RA. (2009). Changes in farming system  
409 and population as drivers of land cover and landscape dynamics: The case of  
410 enclosed and semi-openfield systems in Northern Galicia (Spain). *Landscape and*  
411 *Urban Planning* 90:168-177

412 Carmo M, Moreira F, Casimiro P, Vaz P. (2011). Land use and topography influences  
413 on wildfire occurrence in northern Portugal. *Landscape and Urban Planning*  
414 100:169–176

415 Chas-Amil ML, Prestemon JP, McClean C, Touza J. (2015). Human-ignited wildfire  
416 patterns and responses to policy shifts. *Applied Geography* 5:164-176

417 Chas-Amil ML, Touza J, García-Martínez E. (2013). Forest fires in the wildland-urban  
418 interface: a spatial analysis of forest fragmentation and human impacts. *Applied*  
419 *Geography* 43:127-137

420 Clarke KR, Gorley RN. (2006). PRIMER v6: User Manual/Tutorial. PRIMER-E,  
421 Plymouth, 192pp.

422 Cohen, J.D. (2000). Preventing disaster: home ignitability in the wildland-urban  
423 interface. *Journal of Forestry* 98, 15-21.

424 Corbelle-Rico E, Crecente-Maseda R, Santé-Riveira I. (2012). Multi-scale assessment  
425 and spatial modelling of agricultural land abandonment in a European peripheral  
426 region: Galicia (Spain), 1956-2004. *Land Use Policy* 29: 493-501.

427 Cumming SG. (2001). Forest type and wildfire in the Alberta boreal mixedwood: what  
 428 do fires burn? *Ecol Appl* 11:97-110  
 429 Díaz-Maroto IJ, Vila-Lameiro P. (2008). Historical evolution and land-use changes in  
 430 natural broadleaved forests in the north-west Iberian Peninsula. *Scandinavian*  
 431 *Journal of Forest Research* 23:371-379  
 432 Doley, R. 2003. Pyromania - Fact or fiction? *Brithish Journal of Criminology* 43: 797-  
 433 807.  
 434 Efron, B. 1982. The jackknife, the bootstrap, and other resampling plans. Society for  
 435 Industrial and Applied Mathematics, Philadelphia, Pennsylvania, USA.  
 436 Escribano-Ávila G, Calviño-Cancela M, Pías B, Virgós E, Valladares F, Escudero A.  
 437 (2014). Diverse guilds provide complementary dispersal services in a woodland  
 438 expansion process after land abandonment. *Journal of Applied Ecology* 51:1701-  
 439 1711  
 440 FAO. (2007). Fire management-Global assessment 2006. FAO Forestry Paper 151.  
 441 Rome  
 442 Finney, MA. (2005). The challenge of quantitative risk analysis for wildland fire. *Forest*  
 443 *Ecology and Management* 211(1): 97-108  
 444 Gálvez-Bravo L. (2011). Conejo – *Oryctolagus cuniculus*. In: Salvador A, Cassinello J  
 445 (Eds) Enciclopedia Virtual de los Vertebrados Españoles. Museo Nacional de  
 446 Ciencias Naturales. [In Spanish] <http://www.vertebradosibericos.org/>  
 447 Ganteaume A, Jappiot M, Lampin C, Guijarro M, Hernando C. (2013). Flammability of  
 448 some ornamental species in wildland-urban interfaces in Southeaster France:  
 449 laboratory assessment at particle level. *Environmental Management* 52:467-480

450 González J, Palahi M, Pukkala T. (2005). Integrating fire risk considerations in forest  
 451 management planning in Spain - a landscape level perspective. *Landscape Ecology*  
 452 20(8):957-970

453 Guglietta D, Migliozi A, Ricotta C. (2015). A Multivariate Approach for Mapping  
 454 Fire Ignition Risk: The Example of the National Park of Cilento (Southern Italy).  
 455 *Environmental Management* 56: 157-164

456 MAGRAMA. (2011). *Cuarto Inventario Forestal Nacional. Galicia*. [DVD]. Ministerio  
 457 de Agricultura, Alimentación y Medio Ambiente

458 MAGRAMA. (2012). *Los incendios forestales en España*. Decenio 2001-2010.

459 Manuel C, Gil L. (2002). *La transformación histórica del paisaje forestal en Galicia*.  
 460 Tercer Inventario Forestal Nacional. Ministerio de Medio Ambiente, Madrid, Spain

461 Manly BFJ. 1998. Randomization, bootstrap, and Monte Carlo methods in biology.  
 462 Chapman and Hall, London, UK.

463 Mermoz M, Kitzberger T, Veblen TT. (2005). Landscape influences on occurrence and  
 464 spread of wildfires in Patagonian forests and shrublands. *Ecology* 86 (10):2705–  
 465 2715

466 Moreira F, Rego FC, Ferreira PG. (2001). Temporal (1958–1995) pattern of change in a  
 467 cultural landscape of northwestern Portugal: implications for fire occurrence.  
 468 *Landscape Ecology* 16:557–567

469 Moreira F, Viedma O, Arianoutsou, M, Curt T, Koutsias N, Rigolot E, Barbati A,  
 470 Corona P, Vaz P, Xanthopoulos G, Mouillot F, Bilgili E. (2011). Landscape--  
 471 wildfire interactions in southern Europe: implications for landscape management.  
 472 *Journal of Environmental Management* 92(10):2389-2402

473 Nunes MCS, Vasconcelos MJ, Pereira JMC, Dasgupta N, Alldredge RJ, Rego FC. 2005.  
 474 Land-cover type and fire in Portugal: do fires burn land cover selectively?  
 475 *Landscape Ecology* 20:661–673

476 Oliveira S, Moreira F, Boca R, San-Miguel-Ayanz J, Pereira JMC. (2013). Assessment  
 477 of fire selectivity in relation to land cover and topography: a comparison between  
 478 Southern European countries. *International Journal of Wildland Fire* 23(5):620-630

479 Romero-Calcerrada R, Barrio-Parra F, Millington JDA, Novillo CJ. (2010). Spatial  
 480 modelling of socioeconomic data to understand patterns of human-caused wildfire  
 481 ignition risk in the SW of Madrid (central Spain). *Ecological Modelling* 221:34–45

482 Saura-Mas S, Paula J, Pausas JG, Lloret F. (2010). Fuel loading and flammability in the  
 483 Mediterranean Basin woody species with different post-fire regenerative strategies.  
 484 *International Journal of Wildland Fire* 1:783–794

485 Tapia, L., Domínguez, J., Regos, A., Vidal, M. (2014). Using remote sensing data to  
 486 model European wild rabbit (*Oryctolagus cuniculus*) occurrence in a highly  
 487 fragmented landscape in northwestern Spain. *Acta Theriol* 59:289–298.

488 Teixido AL, Quintanilla LG, Carreno F, Gutiérrez D. (2010). Impacts of changes in  
 489 land use and fragmentation patterns on Atlantic coastal forests in northern Spain.  
 490 *Journal of Environmental Management* 91(4):879-886

491 Vélez R. (2002). Causes of forest fires in the Mediterranean basin. In Arbez, M., Birot,  
 492 Y. Carnus, J. M. Risk Management and Sustainable Forestry. EFI Proceedings  
 493 2002 No. 45 pp 35-42

494 Webb NR. (1998). The traditional management of European heathlands. *Journal of*  
 495 *Applied Ecology* 35:987-990

496 Yang J, He HS, Shifley SR, Gustafson EJ. (2007). Spatial patterns of modern period  
497 human-caused fire occurrence in the Missouri Ozark Highlands. *Forest Science*  
498 53(1):1-15.

499 Table 1: Fire causes categories used in this study.

Category	Definition
Agriculture and vegetation management	Fires caused by farmers in agricultural burnings, verge maintenance, bush clearing, control of animals considered harmful for crops or livestock and those related to beekeeping.
Ranching	Fires set to promote forage production for grazers.
Forestry management	Fires related to forest works.
Hunting	Fires caused by hunter to facilitate hunting or to protest against hunting restrictions.
Recreation	Fires caused by campfires, fireworks, cigarettes, hot air balloons or children.
Waste management	Rubbish burning.
Profit gaining	Fires set to create job opportunities in fire fighting brigades or restoration activities, to affect wood prices, to force land-use changes or to increase forest productivity.
Conflicts	Fires related to revenges, disagreements related to land ownership, protests against reductions in public investment, expropriations or the establishment of Natural Protected Areas, or caused by political groups to cause social unrest.
Mentally ill or violent people	Fires caused by arsonists, for excitement, in pseudo-religious or satanic rites or by vandals.
Accidents	Fires caused by accidents, related to railroads, electric power, vehicles, engines or machinery or by army manoeuvres.
Natural	Fires caused by lighting.
Rekindle	Restart of fires.

500  
501

502 Table 2: Topography. Analysis of differences in elevation between ignition and random  
503 points, WUI and non-WUI areas and LULCs using ANOVA.

Source of variation	d.f.	<i>SS</i>	<i>F</i>	<i>P</i> value
Ignition/Random	1	$1.015 \cdot 10^7$	151.83	<0.001
WUI	1	$1.235 \cdot 10^8$	1847.08	<0.001
LULC	10	$1.440 \cdot 10^9$	2154.22	<0.001
Ign/Rand : WUI	1	$2.244 \cdot 10^7$	335.69	<0.001
Ign/Rand : LULC	10	$3.297 \cdot 10^7$	49.32	<0.001
WUI: LULC	10	$4.690 \cdot 10^6$	7.02	<0.001
Ign/Rand : WUI: LULC	10	$1.339 \cdot 10^6$	2.00	<0.001
Residual	50423	$3.370 \cdot 10^9$		
Total	50466	$5.005 \cdot 10^9$		

504

Table 3: Topography. Analysis of differences in slope between ignition and random points, WUI and Non-WUI areas and LULCs using a Generalized Linear Model with negative binomial distribution and logratio as link function.

Source of variation	d.f.	Deviance ( $\chi^2$ )	<i>P</i> value
Ignition/Random	1	301	<0.001
WUI	1	351	<0.001
LULC	10	7562	<0.001
Ign/Rand : WUI	1	103	<0.001
Ign/Rand : LULC	10	149	<0.001
WUI: LULC	10	211	<0.001
Ign/Rand : WUI: LULC	10	359	<0.001
Residual	50423	41898	
Total	50466	50173	

Table 4: Topography. Analysis of departures between the frequency of ignition points in each compass aspect (N, S, W and E) and that expected by random chance (i.e., obtained in random points) as affected by location within/outside the WUI and LULCs using ANOVA.

Source of variation	d.f.	<i>SS</i> (N; S; E; W)	<i>F</i> (N; S; E; W)	<i>P</i> value (N; S; E; W)
WUI	1	479995; 38934; 518093; 50304	1572.5; 106.6; 1300.8; 122.9	<0.001; <0.001; <0.001; <0.001
LULC	10	9716668; 12741357; 4339542; 2950043	3183.2; 3489.2; 1089.6; 720.9	<0.001; <0.001; <0.001; <0.001
WUI: LULC	10	7267035; 8260603; 386331; 34599694	2380.7; 2262.1; 970.0; 845.5	<0.001; <0.001; <0.001; <0.001
Residual	2178	664828; 795318; 867433; 891248		
Total	2199	18128527; 21836212; 9588383; 7351565		


515 Table 5: Results of the PERMANOVA analysis on differences in fire causes as affected  
 516 by location within and outside the WUI and LULC.

Source of variation	d.f.	SS	PseudoF	P value
WUI	1	58370	771.1	0.001
LULC	9	$3.81 \cdot 10^5$	559.2	0.001
WUI: LULC	9	$1.50 \cdot 10^5$	219.4	0.001
Residual	1980	$1.50 \cdot 10^5$		
Total	1999	$7.39 \cdot 10^5$		

517

Table 6: Percentage of fires occurring in each LULC that were associated to different causes, as detailed in Table 1, outside the WUI (upper value) and within the WUI (bottom value).

Causes	Land uses / covers									
	Agr	Shr	OpWd	AtlF	MxAtl	PiP	EuP	MxPiP	MxEuP	MxEuPiP
Agr. & Veg. Management	63.25 55.83	59.16 64.56	57.03 54.37	63.76 61.43	54.74 27.78	36.01 44.44	35.52 47.76	39.80 53.33	36.28 60.00	37.94 50.00
Ranching	7.73 6.95	17.67 10.13	6.40 3.88	9.80 10.00	2.63 11.11	2.31 0.00	3.00 1.49	0.76 0.00	3.98 0.00	2.41 0.00
Forestry Management	0.93 1.64	0.30 0.00	0.61 0.00	0.62 1.43	0.53 0.00	2.31 0.85	3.00 2.99	1.76 2.22	2.65 4.00	1.74 0.00
Hunting	0.96 1.02	5.75 5.06	3.14 1.94	2.40 0.00	2.63 0.00	0.74 0.00	1.43 0.00	1.01 8.89	1.77 0.00	1.07 0.00
Recreation	1.54 3.48	0.48 0.00	1.05 4.85	1.42 0.00	2.11 11.11	2.03 1.71	3.99 7.46	2.02 4.44	2.65 0.00	1.61 2.00
Waste Management	0.80 0.61	0.30 1.27	0.77 0.00	0.53 0.00	1.58 0.00	0.65 0.00	1.14 1.49	0.25 2.22	2.21 0.00	0.67 0.00
Profit gaining	1.03 0.51	0.14 0.00	0.72 0.00	0.18 0.00	0.00 0.00	1.94 1.71	1.00 0.00	1.51 0.00	2.21 4.00	1.47 1.00
Conflicts	1.31 1.43	1.43 2.53	2.10 0.00	2.58 0.00	2.11 5.56	3.32 1.71	3.85 1.49	3.78 2.22	3.10 0.00	2.01 3.00
Mentally ill or violent people	12.06 17.59	5.91 12.66	15.28 17.48	6.06 7.14	17.37 16.67	31.02 33.33	32.67 31.34	30.23 15.56	26.99 24.00	36.73 29.00
Accidents	2.31 4.09	1.23 3.80	2.98 8.74	2.85 10.00	3.68 16.67	3.14 4.27	2.57 1.49	3.02 2.22	4.42 4.00	1.88 2.00
Natural	1.48 1.64	2.73 0.00	3.14 2.91	2.94 4.29	4.74 0.00	6.28 1.71	1.28 0.00	2.77 0.00	1.77 0.00	1.34 1.00
Rekindle	6.61 5.21	4.91 0.00	6.78 5.83	6.86 5.71	7.89 11.11	10.25 10.26	10.56 4.48	13.10 8.89	11.95 4.00	11.13 12.00




522

523 Figure 1: Study area location map.

524

525



526

527 Fig. 2: Differences in elevation (in percentage) between ignition and random points in  
 528 the LULC types, outside the WUI (light grey) and within the WUI (dark grey).






Fig. 3: Differences in slope (in percentage) between ignition and random points in the LULC types, outside the WUI (light grey) and within the WUI (dark grey).

533



534



535

536

537 Fig. 4: Fire risk depending on site aspect in non-WUI (a) and WUI areas (b). Departures  
538 from 0 show percentage increases or decreases in fire risk compared to that expected by  
539 random in each aspect (N, S, W, E) for each LULCs.

540




Fig. 5: Fire causes in LULCs. NMDs ordinations of LULC types based on Bray-Curtis similarities on square root transformed data of fire causes in Non-WUI (a) and WUI areas (b), showing distances between LULCs in the fire causes space. See the key for symbols of each type of LULC. Superimposed vectors show the fire causes driving the patterns of distance between LULCs.