

This is a repository copy of *Urban eddy covariance measurements reveal significant missing NOx emissions in Central Europe*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/117076/>

Version: Published Version

Article:

Karl, T., Graus, M., Striednig, M. et al. (10 more authors) (2017) Urban eddy covariance measurements reveal significant missing NOx emissions in Central Europe. Scientific Reports. 2536. ISSN 2045-2322

<https://doi.org/10.1038/s41598-017-02699-9>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

SCIENTIFIC REPORTS



OPEN

Urban eddy covariance measurements reveal significant missing NO_x emissions in Central Europe

T. Karl¹, M. Graus¹, M. Striednig¹, C. Lamprecht¹, A. Hammerle², G. Wohlfahrt², A. Held³, L. von der Heyden³, M. J. Deventer⁴, A. Krismer⁵, C. Haun⁶, R. Feichter⁷ & J. Lee⁸

Nitrogen oxide (NO_x) pollution is emerging as a primary environmental concern across Europe. While some large European metropolitan areas are already in breach of EU safety limits for NO₂, this phenomenon does not seem to be only restricted to large industrialized areas anymore. Many smaller scale populated agglomerations including their surrounding rural areas are seeing frequent NO₂ concentration violations. The question of a quantitative understanding of different NO_x emission sources is therefore of immanent relevance for climate and air chemistry models as well as air pollution management and health. Here we report simultaneous eddy covariance flux measurements of NO_x, CO₂, CO and non methane volatile organic compound tracers in a city that might be considered representative for Central Europe and the greater Alpine region. Our data show that NO_x fluxes are largely at variance with modelled emission projections, suggesting an appreciable underestimation of the traffic related atmospheric NO_x input in Europe, comparable to the weekend-weekday effect, which locally changes ozone production rates by 40%.

The nitrogen cycle¹ is essential for maintaining the oxidizing capacity of the atmosphere and regulating ozone in the lower atmosphere². Perturbations due to rapid industrialization and agricultural activities have led to a significant increase of atmospheric nitrogen oxides (NO_x) during the 20th century³. A regionally intense buildup of photochemical smog due to the presence of nitrogen oxides, CO and non-methane volatile organic compounds (NMVOC) was first identified in the US and attributed as the main cause of severe ozone pollution in many areas⁴. Decades of subsequent research activities ranging from detailed laboratory^{5,6} and smog chamber⁷⁻⁹ studies to large scale field campaigns¹⁰⁻¹² have led to a reasonably good mechanistic understanding of the formation of tropospheric ozone, which is characterized by a complex nonlinear relationship between NO_x and reactive carbon species¹³. This interdependency gives regulators two key strategies to mitigate ozone pollution. The effectiveness to control ozone thereby very much depends on the ratio between ambient OH reactivity and NO_x concentrations¹⁴, which can be described by relatively simple analytical relationships¹⁵. The development of mechanistic regional¹⁶ and global air chemistry models¹⁷ has further given regulators and scientists powerful tools to study tropospheric ozone formation¹⁶, where the mitigation of NO_x emissions has emerged as one of the key air pollution control strategies for ozone¹⁸⁻²⁰ and more recently also for particulate matter with a diameter of 1 μm or less (PM1)²¹. Due to the toxicity, nitrogen dioxide (NO₂) is also regulated as a hazardous air pollutant itself²². For example, in Europe regulatory action under the EU Thematic Strategy on Air Pollution is in place to limit urban street canyon NO₂ concentrations to 40 μg/m³ per year (or 200 μg/m³/h on less than 18 days/year)²³. Current trends across European air quality networks show that regulatory thresholds of NO₂ are violated at many stations, which does not seem to be limited to large population centers anymore (ref. 24, SI). In fact many rural

¹Institute of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria. ²Institute of Ecology, University of Innsbruck, Innsbruck, Austria. ³Atmospheric Chemistry, University of Bayreuth, Innsbruck, Germany. ⁴Department of Geography, University of California, Berkeley, USA. ⁵Abteilung Waldschutz, Amt der Tiroler Landesregierung, Innsbruck, Austria. ⁶Abteilung Geoinformation, Amt der Tiroler Landesregierung, Innsbruck, Austria. ⁷Amt für Verkehrsplanung, Umwelt, Magistrat III Stadt Innsbruck, Innsbruck, Austria. ⁸National Centre for Atmospheric Science and Department of Chemistry, University of York, York, UK. Correspondence and requests for materials should be addressed to T.K. (email: thomas.karl@uibk.ac.at)

areas and smaller towns see NO₂ concentration levels rivaling those of large metropolitan areas. Owing to the spatiotemporal variability and uncertainty of different anthropogenic NO_x sources, it is difficult to attribute emission uncertainties to specific sectors in complex bottom-up emission inventories²⁵ or top-down remote sensing assessments²⁶. Recently evidence has accumulated that rapid shifts in transportation fuels can have significant impacts on air quality^{27,28}. In Europe for example the question about the increasing penetration of Diesel cars raises concerns as to what extent such a technological change has been counterproductive to mitigating atmospheric NO₂ pollution under new emission regulation standards^{19,29}. The United States environmental protection agency's (US EPA) notice of violation of the Clean Air Act to a German automaker regarding Diesel engines has sparked a number of new real world driving (RDE) emission tests across Europe, which show significant manufacturer and vehicle specific variability^{30,31}. These new data suggest that the impact on up-scaled average fleet emissions needed for accurate air quality predictions remains unclear³².

A number of urban flux measurement sites for energy and CO₂ have been established highlighting their potential for surface-atmosphere exchange studies^{33,34}. In contrast, similar measurements for reactive gases are often still quite limited, owing to the complexity of the required measurement systems. A set of recently conducted urban ground based and airborne NMVOC flux measurements revealed the usefulness to test bottom-up emission inventories and revealed significant discrepancies for some species^{35–39}. Urban flux measurements for NO_x are even more scarce^{32,40,41} indicating that constraints on emission sources in urban areas can be quite uncertain. Here we improve upon existing work, by simultaneously measuring NO_x, selected tracer NMVOCs, CO and CO₂ leading to a well constrained flux dataset, that allows testing our understanding of prominent NO_x emission sources.

Results

Obtaining ensemble average statistics on fleet emissions by eddy covariance flux measurements.

A comprehensive set of eddy covariance measurements for NO_x, marker NMVOC, CO and CO₂ at an urban location allows a direct comparison of relative flux ratios with bottom-up emission sources. The study site located in Innsbruck (N 47°15'51.50", E 11°23'6.77") at the center of the Inn valley, represents one of the most strategically important Alpine crossing points for the transport of goods between Northern and Southern Europe. Each year approximately 6 million vehicles⁴² pass through the east-west facing valley, which is about 10 km wide surrounded by mountain ridges about 2.5 km high. The valley topography leads to a very predictable and pronounced wind system characterized by a topographic amplification factor (TAF) of about 3⁴³. Due to the combination of significant traffic induced NO_x emissions and increasingly stringent NO₂ limit values, the area is in non-attainment. Local authorities are facing legal proceedings by the European Commission for their failure to control excessive levels of nitrogen oxides (Fig. S2), similar to many areas across Europe²³. Tracer flux relationships allow investigating to what extent urban emissions are caused by (a) traffic, (b) urban residential and (c) biomass burning/biofuel activities. Figure 1 shows the diurnal evolution of Weekday (Tuesday–Thursday) and Sunday NO_x fluxes and concentrations along with mean traffic count data at the site during the measurement campaign (July – October, 2015). Median measured midday NO_x mixing ratios in Innsbruck are comparable to values reported for central London (10–14 ppbv), while corresponding observed fluxes are about a factor of 3–4 lower (i.e. 3000–4000 ng/m²/s vs 700–1440 ng/m²/s)³². These observations are consistent with the idea of an intensification of air pollution proportional to TAF, and a corresponding effective lower air volume that pollutants are being mixed into in steep valleys (Fig. S2, ref. 43). This comparison likely indicates that a much stronger reduction of NO_x emissions from the transport sector would be required in the Alps than for example in London in order to achieve current air pollution standards along one of the busiest EU transport corridors across the Alps. Several lines of evidence exclude significant presence of biomass burning during the present study. The average ratio between benzene and toluene fluxes exhibited a typical value (3.3 ± 0.7 ; $R^2 = 0.85$) characteristic for urban emission sources, dominated by fossil fuel combustion and evaporative/cold-start emissions. The correlation between acetonitrile and benzene, toluene, NO_x or CO₂ fluxes was low with an R^2 of 0.07, 0.08, 0.02 and 0.06 respectively. We also did not observe significant excursions of other species recently suggested as additional biomass burning markers⁴⁴ such as furfural and furan showing a correlation coefficient of $R^2 < 0.2$ above their background fluxes. We observed an excellent correlation between CO₂, benzene and NO_x fluxes (CO_2/NO_x ; $R^2 = 0.86$; benzene/NO_x; $R^2 = 0.75$). The covariance between between NO_x and CO₂ (benzene) fluxes yielded values of 0.91 (0.86). We interpret these observations such that benzene, NO_x and CO₂ emissions are dominated by road traffic with contributions from residential combustion sources.

Benchmarking urban source emission ratios and inventories.

To gain a more quantitative insight, we investigated flux ratios between NO_x and CO₂ ($F_{\text{NO}_x}/F_{\text{CO}_2}$; Fig. 2). The advantage of this approach is that it allows determining the actual ensemble average of different emission sources based on measured flux ratios, similar to an end-member un-mixing regression analysis. This allows us to compare our measurements to relative emission strengths reported in emission models and inventories. For large scale emission inventories (e.g. grid cells > 1 km²) this approach also circumvents uncertainties related to assumptions of various downscaling approaches. The observed $F_{\text{NO}_x}/F_{\text{CO}_2}$ ratios follow a diurnal cycle showing a ~40–50% variation throughout a day, which reflects the pronounced fluctuation of traffic activity across the city (e.g. ranging from about 97 vehicles/h at night to 890 vehicles/h during daytime at a traffic count station within the flux footprint). Since we can exclude significant industrial emissions within the flux footprint as well as biomass burning, the variation of $F_{\text{NO}_x}/F_{\text{CO}_2}$ should exhibit the characteristic behavior of city scale sources comprised of (1) a combination of vehicular emissions and (2) residential/domestic combustion sources (e.g. oil and gas heating units). A minimization routine (SI) allowed un-mixing these two end-members of the compositional data, reflecting the actual emission ratios for traffic and urban residential combustion sources (Fig. 2). The fitted model (SI) can reproduce the diurnal cycle and activity factors reasonably well, leading to a NO_x/CO₂ emission ratio for traffic of $4.2(\pm 0.3) \times 10^{-3}$ ([mg/




Figure 1. Statistical plot of measured NO_x fluxes, mixing ratios and traffic count data. The center dot shows the ensemble median, where the box around it represents one standard deviation and whiskers the 25 and 75% percentile. Individual extreme values are plotted as open circles. Panels A, B and C represent weekdays (i.e. TUE-THU; composite of 609 individual data points) and panels D, E and F depict Sundays (193 individual data points).




Figure 2. Diurnal cycle of median traffic count data (black line - left axis), measured (blue circles) and model fitted (dashed blue line) mass flux ratio of NO_x/CO_2 (right axis). The corresponding calculated end members for traffic and residential combustion ratios are indicated by blue and green horizontal lines. The shading reflects one standard deviation. In addition colored dashed lines show predictions using fixed emission ratios from COPERT (magenta) and ACCMIP (red).

m^2/h $\text{NO}_x/[\text{mg}/\text{m}^2/\text{h}] \text{CO}_2$), and $0.20(\pm 0.05) \times 10^{-3}$ ($[\text{mg}/\text{m}^2/\text{h}] \text{NO}_x/[\text{mg}/\text{m}^2/\text{h}] \text{CO}_2$) for residential combustion sources. These calculated ratios are also depicted in Fig. 2 by horizontal shaded blue and green lines. The activity factors suggest that the ratio is dominated by traffic, comprising about 85% of the activity averaged over the entire day (and >95% during peak traffic). We also investigated $\text{NO}_x/\text{benzene}$ flux source ratios revealing

Measurement/inventory ratio or Measurement/emission standard ratio	NO _x /CO ₂ NO _x /CO	NO _x /benzene
INNAQS/COPERT ^{1,‡}	1.7 ^{1.8} _{1.6}	3.9 ^{4.7} _{3.1}
INNAQS/HBFA3.2 ²	1.5 ^{1.6} _{1.4}	N/A
INNAQS/ACCMIP (traffic) ³	4.0 ^{5.3*} _{2.8*}	2.1 ^{2.5} _{1.7}
INNAQS/EMEP (traffic) [‡]	2.9 ^{3.3*} _{2.6*}	N/A
INNAQS/US Tier II ⁴	36.0 ^{38.6} _{33.4}	N/A
	NO _x /CO ₂ PC and LCV < 1305 kg	NO _x /CO ₂ LCV 1305 kg–3500 kg
INNAQS/Euro6 ⁵	Diesel	4.7 ^{5.0} _{4.3}
	Petrol	12.6 ^{13.5} _{11.7}
INNAQS/Euro5 ⁵	Diesel	2.1 ^{2.3} _{1.9}
	Petrol	12.6 ^{13.5} _{11.7}
INNAQS/Euro4 ⁵	Diesel	1.5 ^{1.6} _{1.4}
	Petrol	4.7 ^{5.0} _{4.4}
INNAQS/Euro3 ⁵	Diesel	0.75 ^{0.8} _{0.7}
	Petrol	2.5 ^{2.7} _{2.3}

Table 1. Measurement – inventory comparison (i.e. measured/modelled flux (emission) ratio). The uncertainty range is given by sub- and superscripts. PC: passenger cars; LCV: light commercial vehicle. ¹COPERT emission model (SI). ²HBFA 3.2 (SI). Comparison with HBFA 3.3, that was published during the copy editing phase, yielded an average bias of 1.2. ³ACCMIP – (SI). ⁴US EPA²². ⁵EEA²⁴. *For inventories that do not explicitly report CO₂, we converted data using the measured midday range of CO to CO₂ flux ratios (3.6 to 4.6 ppbv/ppmv), which fall close to a recent evaluation based on a road tunnel study⁴⁵. ‡Data are trend adjusted for 2015 according to GAINS (<http://gains.iiasa.ac.at/models/>)²³.

comparable differences as observed for $F_{\text{NO}_x}/F_{\text{CO}_2}$. The CO₂ flux weekend-weekday effect and CO/CO₂ ratios close to a recent road tunnel study⁴⁵, all imply that a biogenic influence on CO₂ fluxes due to photosynthetic uptake or respiration can be considered negligible at this site. The obtained NO_x/CO₂ flux ratio for traffic is significantly larger than predicted by a number of state of the art emission inventories and emission standards (Table 1). Generally, we observe 50–70% higher NO_x emissions relative to CO₂ from road traffic than what is calculated with the most recent traffic emission models. In these detailed bottom – up models, mobile source emissions are treated for different engine sizes and fuels, that, in the past, relied on standardized protocols obtained in test facilities, but were recently updated based on a number of RDE tests^{30,31}. Exhaust from modern gasoline powered engines, despite higher ignition temperatures than those powered by Diesel, can be effectively treated for NO along with NMVOC (and CO) using a three way catalytic (TWC) converter. TWC treatment can lead to a 10 fold reduction of NO. This has been hard to achieve for Diesel powered cars, which nowadays mostly rely on selective catalytic reduction due to the high air to fuel ratio during combustion. Generally, different combustion and exhaust treatment characteristics result in significantly higher NO_x/CO₂ emission ratios for Diesel powered cars than for gasoline. The modelled fleet average contribution suggests that at least 90% of urban NO_x emissions should originate from Diesel driven vehicles at the present location. 85% is modelled to be emitted by the passenger car fleet based on the COPERT model and TRACCS database (ref. 46, SI). The current Austrian passenger car fleet comprises about 50% Diesel cars and the percentage across Europe grew at a substantially faster rate compared to the US⁴⁷. Based on our measurements the current average Austrian car fleet emits about 36 times more NO_x per CO₂ molecule compared to the US TIER II emission standard and a factor of 8–10 more than Euro 6 emission standards³¹. Factoring in differences in fuel economy between the European and US car fleet, this would equate to about an order of magnitude more NO_x emissions per travelled distance compared to newly introduced emission standards. How comparable are these results to other European countries? Diesel engines dominate the European passenger car market: 55% of all newly registered vehicles in the EU were powered by Diesel-fuel in 2012⁴²; the penetration of Diesel cars of the two largest European economies bordering the Alps ranges from 30% (Germany) to 70% (France)⁴², more than an order of magnitude higher than in the US⁴⁷. When comparing measured NO_x/CO₂ flux ratios with current inventories used for IPCC atmospheric chemistry/climate⁴⁸ and air quality models⁴⁹ we obtain a discrepancy up to about a factor of 3–4 (Table 1). Incidentally, a recent comprehensive model evaluation has suggested significant discrepancies between regionally modelled and observed surface NO_x concentrations that seemed worse (e.g. by a factor of 2 during summers) over Europe than over the US²⁵. While a number of uncertainties (agricultural emissions, vertical mixing, biomass burning emissions, deposition) can potentially result in modelled concentration biases in these models, our measurements suggest that road transport related biases likely contribute significantly to these discrepancies as they account for about half of the total NO_x emissions across Europe⁵⁰.

What is the impact on atmospheric chemistry? The weekend – weekday effect (Fig. 1) allows to gain insight into changing NO_x and NMVOC fluxes on ozone production in more detail. In Austria, heavy duty vehicle traffic (trucks heavier than 7.5t and all road trains) is banned between Saturday 15:00 and Sunday 22:00 and on public holidays between midnight and 22:00. Traffic count data generally show a pronounced difference in

driving habits resulting in a factor of 1.9 ± 0.2 lower vehicle counts on Sunday than on weekdays. We calculated typical ozone production rates for midday-afternoon conditions (11–16 h LT), when photochemistry peaks. The corresponding NO_x fluxes are a factor of 2.1 ± 0.2 lower on Sundays, closely matching observations of vehicle activity. Benzene and toluene fluxes, representing the variation of anthropogenic NMVOC emissions, were lower by a factor of 1.8 ± 0.3 and 2.0 ± 0.3 respectively. CO_2 fluxes changed by a similar factor of 2.3 ± 0.5 . The weekend-weekday comparison provides an independent confirmation that these pollutant emissions are dominated by traffic activity during the day. Average NO_x concentrations are a factor of 2.5 ± 0.2 lower during Sundays. Incidentally, the observed weekend – weekday reduction is comparable to the observed measurement-inventory discrepancies or the effect if an entire car fleet was converted from a Euro 5 (0.18 g/km) to a Euro 6 (0.08 g/km) NO_x emission standard. Our measurements therefore allows us to benchmark such a hypothetical regulatory action in the real atmosphere. Changes in local ozone production are calculated following procedures outlined before^{15,51}, where the sensitivity of local ozone production can be approximated by the ratio of radical termination (L_N) processes (e.g. $\text{NO}_2 + \text{OH}$) and photochemical radical production (Q):

$$\frac{L_N}{Q} \approx \frac{2\delta O_x - 2\delta \text{NO}_x - \delta J}{\delta O_x + \delta \text{NMVOC} - 3\delta \text{NO}_x} \quad (1)$$

Here the δ symbol indicates the relative change between weekday and Sunday, J represents the photolysis rates, and $O_x = \text{O}_3 + \text{NO}_2$. All terms on the right side can be inferred from measurements of the weekend effect, where the anthropogenic change of NMVOCs is assumed to follow benzene. There is evidence of a non-negligible biogenic NMVOC (BVOC) presence at the site (e.g. 20–50% of the NMVOC reactivity) and these BVOCs do not exhibit any anthropogenically related variation between weekdays and weekend. We apportioned the change of NMVOC reactivity therefore into a biogenic and anthropogenic part using data from the PTR-QiTOF-MS instrument. To achieve this, we estimated the total anthropogenic NMVOC reactivity from known urban concentration ratios^{52,53} scaled to benzene and compared this to the measured reactivity of BVOC, in particular the reactive biogenic marker species isoprene and monoterpenes. This allowed to obtain upper ($\delta \text{NMVOC} = \delta \text{benzene}$) and lower ($\delta \text{NMVOC} = \delta \text{benzene} \times [\text{reactivity anthropogenic NMVOC}] / [\text{reactivity BVOC}]$) bounds for L_N/Q . It can further be shown¹⁵ that the sensitivity of local ozone production can be related to L_N/Q according to:

$$\frac{d \ln P(\text{O}_3)}{d \ln(\text{NO}_x)} = \frac{1 - \frac{3}{2} \frac{L_N}{Q}}{1 - \frac{1}{2} \frac{L_N}{Q}} \quad (2a)$$

$$\frac{d \ln P(\text{O}_3)}{d \ln(\text{NMVOC})} = \frac{\frac{1}{2} \frac{L_N}{Q}}{1 - \frac{1}{2} \frac{L_N}{Q}} \quad (2b)$$

The calculated range for L_N/Q delimits a ratio between 0.82 and 0.92 (theoretical maximum = 1) and places the site in a NO_x inhibited chemical regime for ozone production during weekdays^{15,51}. Incidentally these afternoon values are systematically higher than reported for a rural site in Northern Italy with comparable NO_x concentrations, and more closely follow conditions reported for Milan in 1998⁵⁴. Our flux data clearly demonstrate that the weekend effect of ozone production found in Innsbruck is largely a direct result of lower weekend NO_x emissions. As a consequence of the dominating nitrogen chemistry for radical loss calculated via eqs 1 and 2, local ozone production will therefore increase by 39% to 70% proportional to a reduction of NO_x mixing ratios in Innsbruck. It will decrease between 70% to 85% proportional to a reduction of NMVOC or CO. The impact on gross ozone production ($P(\text{O}_3)$) was also investigated independently using the Leeds Master Chemical Mechanism (MCM) as a photochemical box model (refs 13 and 55, SI, Fig. 3). The model simulates an increase of $P(\text{O}_3)$ by 40% (from 2.5 to 3.5 ppbv/h) on Sundays due to a 2 fold reduction of NO_x , giving an explanation for the observed weekend effect, which results in an increase of local afternoon ozone concentrations up to 24% (Fig. 4). The model simulation suggests a maximum increase of $P(\text{O}_3)$ up to 3 times (2.5 ppbv/h to 7.8 ppbv/h), if solely NO_x concentrations were decreased from current levels to about 2 ppbv (Fig. 3). Below 2–4 ppbv of NO_x , ozone production would enter the NO_x sensitive regime and gradually lead to decreasing $P(\text{O}_3)$. Next, we setup the MCM as a diurnally constrained 0-dimensional diluting box model (SI) to study the sensitivity with respect to changes in modelled ozone concentrations between weekdays and weekends. The model is thereby fully constrained by methane, CO, measured NMVOC, NO, photolysis rates, and PBL height along with ancillary meteorological variables (e.g. temperature, humidity). NO_2 and ozone are initialized by their measured initial concentrations, but are then allowed to freely adjust during the model run. We chose a spin-up time of 3 days and used model results from the last day of simulation to compare with measured ozone concentrations. Figure 4 shows the diurnal patterns of modelled and observed ozone concentrations. Observed (modelled) peak ozone concentrations are 49 ± 2 (52) ppbv and 43 ± 2 (44) ppbv on weekends and weekdays respectively. Modelled and observed weekend-weekday differences averaged over daytime hours correspond to 7.2 ± 1 and 7.6 ± 1.2 . While the model reproduces the general diurnal ozone cycle and midday peak reasonably well, it underestimates absolute nighttime concentrations exhibiting a stronger diurnal cycle. We attribute this shortcoming mainly to poor knowledge and constraints on entrainment and advection processes that dominate observed nighttime distributions of ozone. While this approach has its limitations, we believe it does a sufficient job for investigating relative daytime changes of ozone concentrations caused by the weekend effect. On average, daytime weekday concentrations are about 7–8 ppbv lower than on weekends. These results can be compared to two quite contrasting areas. In Mexico City⁵¹, ozone concentrations change very little despite similar relative variations of NO_x mixing ratios between weekday and weekend, which




Figure 3. Calculated ozone production rates (solid lines) and derivatives (dashed lines) as a function of NO_x . Blue and cyan represent base cases for the study site, taking into account changes in O_3 , NO_x and NMVOCs concentrations. The solid black line follows the observed weekend-weekday trajectory.

can be attributed to a completely NO_x saturated environment. On the other hand, a quite pronounced variation is found in the southern California Air basin⁵⁶, where average ozone concentrations are observed to change almost twice as much compared to the present study. We interpret these observations such that they reflect the change in ozone production efficiency (Fig. 4), when going from environments with NO_x saturated conditions to more VOC limited conditions. In this context a number of explanations of the weekend effect on ozone mixing ratios are often discussed⁵¹: (1) different timing of NO_x emissions on weekends and associated chemical repartitioning, (2) carryover of previous day pollutants at the surface and aloft, (3) higher weekend VOC emissions, (4) higher weekend photolysis frequencies due to lower aerosol loadings, (5) changes in biogenic emissions due to a different radiation regime caused by lower aerosol loadings and (6) lower overall weekend NO_x emissions. We show for the first time, that, for Innsbruck, at least 85% of the NO_x concentration change, leading to a 7–8 ppbv ozone increase on weekends, can be directly associated with a change of the overall local atmospheric NO_x flux.

Discussion

Continued urbanization in conjunction with rapid technological changes in the mobility sector poses a challenge for accurate up to date predictions of pollutant emissions. By constraining the actual fluxes of nitrogen oxides, NMVOC, CO_2 and CO into the atmosphere our measurements provide an observationally based explanation why NO_x concentrations have hardly declined since the introduction of EURO 3 emission standards in Central Europe. While the technological shift towards Diesel passenger cars might have helped curb CO_2 emissions through better fuel economy compared to gasoline powered cars in the past, it created a widespread problem of NO_2 pollution across Europe²³ that does not seem to be exclusively limited to the largest metropolitan and industrialized areas. The presented flux measurements indicate that traffic related NO_x emissions in current operational air quality models can be significantly underestimated by up to a factor of 4 across countries exhibiting a sizeable fraction of Diesel powered cars in their fleet. As Diesel fuels (including bio-diesel) could account for 70% of the growth in transportation fuels by 2040, with significant demand in Asian markets according to industry projections⁵⁷, a better understanding of the uncertainty in associated changes of NO_x fluxes and ozone chemistry will therefore be important for future environmental impact studies. Our measurements show that projected significant decreases in European NO_x emissions from the mobile transport sector will lead to conditions improving NO_2 exposure limits, but could locally increase ozone levels on the short term. Concomitant reduction measures for NO_x and NMVOC (CO) might therefore still prove most effective to avoid parallel increases of local ozone levels due to new NO_x emission standards. This might be particularly important in areas where topographic amplification can lead to a stronger accumulation of air pollutants than over flat land and be a relevant consideration for mountainous mega-cities⁵⁸. Using the observed weekend effect as proxy for underestimated NO_x emissions (i.e. a factor 2–4 difference), models would overestimate $P(\text{O}_3)$ by 30–40% under the observed NO_x inhibited – VOC limited regime and underestimate $P(\text{O}_3)$ downwind, once NO_x concentrations fall below 2–4 ppbv. Here we demonstrate that parallel flux measurements of a wide range of chemical species can be used to benchmark urban emission sources, complement traditional approaches and significantly improve uncertainties inherent to bottom-up scaling in atmospheric chemistry models.




Figure 4. Observed and modelled changes in ozone concentrations on weekdays (Tuesday – Thursday) and weekends (Sunday). Panel A depicts a statistical boxplot showing weekend-weekday differences during daytime, which are represented by the colored sections in panels B and C.

Methods

Instrumentation. *NM VOC.* A PTR-QiTOF instrument (Ionicon, Austria) was operated in hydronium mode at standard conditions in the drift tube of 112 Townsend. The instrument was set up to sample ambient air from a turbulently purged “3/8” Teflon line. Every seven hours, zero calibrations were performed for 30 minutes providing VOC free air from a continuously purged catalytical converter through a setup of software controlled solenoid valves. In addition, known quantities of a suite of VOC from a 1 ppm calibration gas standard (Apel & Riemer, USA) were periodically added to the VOC free air and dynamically diluted into low ppbv mixing ratios. *NO_x.* A dual channel chemiluminescence instrument (CLD 899 Y; Ecophysics) was used for NO and NO_x measurements. The instrument was equipped with a metal oxide converter operated at 375°C. The instrument was operated in flux mode acquiring data at about 5 Hz, similar to measurements performed over a pasture⁵⁹ and forest⁶⁰. A NO standard was periodically introduced for calibration. Zeroing was performed once a day close to midnight. *CO₂, H₂O.* A closed path eddy covariance system (CPEC 200; short inlet, enclosed IRGA design; Campbell Scientific) measured three dimensional winds along with CO₂ and H₂O. An additional 3D sonic anemometer (CSAT3; Campbell Scientific) was available for turbulence measurements at an alternative height level (Fig. S3). Calibration for CO₂ was performed once a day. *O₃.* Ozone concentrations were obtained from a closed-path UV photometric analyser (APOA-370, Horiba, Japan); *CO.* CO measurements were available for a limited amount of time in August 2015. Ambient mole fractions of carbon monoxide (CO) were measured with a quantum cascade laser spectrometer (CWQC-TILDAS-76-D, Aerodyne, USA) with a 76 m path length optical cell at a wavenumber of ca. 2190 cm⁻¹. The QCL was operated at a pressure of ca. 4 kPa using a built-in pressure controller and temperature of the optical bench and housing controlled to 35 °C. Fitting of absorption spectra at 2 Hz, storing of calculated dry mole fractions, switching of zero/calibration valves, control of pressure lock, correction for band broadening and other system controls were realized by the TDLWintel software (Aerodyne, USA) run on a PC synchronized in time with the system collecting the anemometer data using the NTP software (Meinberg, Germany).

Eddy covariance data analysis. The eddy covariance method is derived from the scalar budget equation after Reynolds decomposition, and in its simplest form for horizontally homogeneous flows normal to the surface, where the mean vertical motion of wind (\overline{w}) can be considered 0, relates the measured surface-atmosphere exchange flux (F) to the covariance between vertical wind and concentration fluctuation according to:

$$F = \langle w'c' \rangle,$$

where w' represents the vertical fluctuation of wind speed, and c' the concentration fluctuation. Brackets denote the averaging interval. The ensemble average used here is 30 minutes. Fluxes were selected according to standard quality control criteria, such as raw data despiking, correcting for high and low pass filtering biases, applying a stationarity test and test on developed turbulent conditions (e.g. u^* filtering)⁶¹. In addition we parsed the data to make sure that the flux footprint would reflect a representative urban area (SI).

References

- Fowler, D. *et al.* Effects of global change during the 21st century on the nitrogen cycle. *Atmos. Chem. Phys.* **15**, 13849–13893 (2015).
- Monks, P. S. *et al.* Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer. *Atmos. Chem. Phys.* **15**, 8889–8973 (2015).
- Intergovernmental Panel on Climate Change (IPCC) “Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change”, (IPCC, 2014, Geneva, <https://www.ipcc.ch/report/ar5/syr/>).
- Haagen-Smit, A. J., Bradley, C. E. & Fox, M. M. Ozone formation in photochemical oxidation of organic substances. *Indust. and Engineering Chemistry* **45**, 2086–2089 (1953).
- Calvert, J. G. Gaseous reactions. *Annual Review of Physical Chemistry* **11**, 41–64 (1960).
- Atkinson, R. Kinetics and mechanism of the gasphase reactions of the hydroxylradical with organic compounds under atmospheric conditions. *Chem. Rev.* **86**, 69–201 (1985).
- Leone, J. A., Flagan, R. C., Grosjean, D. & Seinfeld, J. H. An outdoor smog chamber and modeling study of toluene-NO_x photooxidation. *Int. J. of Chem. Kinetics* **17**, 177–216 (1985).
- Martin, M., Pons, M., Wirtz, K., Etzkorn, T. & Senzig, J. An outdoor smog chamber study of the gas-phase chemistry of toluene and xylenes in NO_x/air systems, in *The oxidizing capacity of the troposphere* (Proceedings of the seventh European symposium on physicochemical behavior of atmospheric pollutants, European Commission, Luxembourg, 1996) pp. 350–354.
- Karl, M. *et al.* Kinetic Study of the OH-isoprene and O₃-isoprene reaction in the atmosphere simulation chamber SAPHIR. *GRL*. **31**, doi:10.1029/2003GL019189 (2004).
- Meagher, J. F., Cowling, E. B., Fehsenfeld, F. C. & Parkhurst, W. J. Ozone formation and transport in the southeastern United States: Overview of the SOS Nashville Middle Tennessee Ozone Study. *J. Geophys. Res.* **103**, 22213–22223 (1998).
- Ridley, B. A. & Robinson, E. The Mauna-Loa-Observatory Photochemistry Experiment. *J. Geophys. Res.* **97**, 10285–10290 (1992).
- Jacob, D. J. *et al.* Transport and Chemical Evolution over the Pacific (TRACE-P) aircraft mission: Design, execution, and first results. *J. Geophys. Res.* **108**, 1–19 (2003).
- Jenkin, M. E., Saunders, S. M. & Pilling, M. J. The tropospheric degradation of volatile organic compounds: A protocol for mechanism development. *Atmos. Environ.* **31**, 81–104 (1997).
- Brune, W. H. *et al.* Ozone production chemistry in the presence of urban plumes, *Faraday Discussions*, doi:10.1039/c5fd00204d (2015).
- Kleinman, L. I. *et al.* Dependence of ozone production on NO and hydrocarbons in the troposphere. *GRL* **24**, 2299–2302 (1997).
- Im, U. *et al.* Evaluation of operational on-line-coupled regional air quality models over Europe and North America in the context of AQMEII phase 2. Part I: Ozone. *Atmos. Environm.* **115**, 404–420 (2015).
- Ehhalt, D. & Prather, M. “Atmospheric Chemistry and Greenhouse Gases” (Intergovernmental Panel on Climate Change, 3rd assessment report, Chapter 4, 2001; http://www.grida.no/publications/other/ipcc_tar/).
- de Gouw, J. A., Parrish, D. D., Frost, G. J. & Trainer, M. Reduced emissions of CO₂, NO_x, and SO₂ from US power plants owing to switch from coal to natural gas with combined cycle technology. *Earths Future* **2**, 75–82 (2014).
- Ehlers, C. *et al.* Twenty years of ambient observations of nitrogen oxides and specified hydrocarbons in air masses dominated by traffic emissions in Germany, *Faraday Discussions*, doi:10.1039/c5fd00180c (2015).
- Carlsaw, D. C. & Beevers, S. D. New Directions: Should road vehicle emissions legislation consider primary NO₂? *Atmos. Environ.* **38**, 1233–1234 (2004).
- Kiendler-Scharr, A. *et al.* Ubiquity of organic nitrates from nighttime chemistry in the European submicron aerosol. *GRL* **43**, 7735–7744 (2016).
- U. S. Environmental Protection Agency, (EPA), “NAAQS Scientific and Technical Information”, <https://www.epa.gov/naaqs/nitrogen-dioxide-no2-primary-air-quality-standards> retrieved on 07-01-2016, (2016).
- Kiesewetter, G. *et al.* Modelling NO₂ concentrations at the street level in the GAINS integrated assessment model: projections under current legislation. *Atmos. Chem. Phys.* **14**, 813–829 (2014).
- European Environment Agency, “Air quality in Europe – 2015” (Tech. Rep. 5/2015, ISBN 978-92-9213-702-1, doi:10.2800/62459, 2015).
- Giordano, L. *et al.* Assessment of the MACC reanalysis and its influence as chemical boundary conditions for regional air quality modeling in AQMEII-2. *Atmos. Environ.* **115**, 371–388 (2015).
- Heckel, A. *et al.* Influence of low spatial resolution a priori data on tropospheric NO₂ satellite retrievals. *Atmos. Meas. Technol.* **4**, 1805–1820 (2011).
- Salvo, A. & Geiger, F. M. Reduction in local ozone levels in urban Sao Paulo due to a shift from ethanol to gasoline use. *Nature Geoscience*. **7**, 450–458 (2014).
- Madronich, S. Atmospheric chemistry: Ethanol and ozone. *Nature Geoscience* **7**, 395–397 (2014).
- Carlsaw, D. C. Evidence of an increasing NO₂/NO_x emissions ratio from road traffic emissions. *Atmos. Environm* **39**, 4793–4802 (2005).
- Bundesministerium für Verkehr und digitale Infrastruktur (BMVI), “Bericht der Untersuchungskommission “Volkswagen” 2016” (<https://www.bmvi.de/SharedDocs/DE/Publikationen/LA/bericht-untersuchungskommission-volkswagen.html>, retrieved 2016) [the easiest access to this source is via the URL].
- Franco, V., Posada Sanches, F., German, J. & Mock, P. “Real-world exhaust emissions from modern diesel cars, Part 2” (Tech. Report: The International Council on Clean Transportation, www.theicct.org, Washington, DC, 2014).
- Lee, J. *et al.* Measurement of NO_x fluxes from a tall tower in central London, UK and comparison with emission inventories. *Environ. Sci. & Tech* **49**, 1025–1034 (2015).
- Christen, A. Atmospheric measurement techniques to quantify greenhouse gas emissions from cities. *Urban Climate* **10**, 241–260 (2014).
- Nemitz, E., Hargreaves, K. J., McDonald, A. G., Dorsey, J. R. & Fowler, D. Meteorological measurements of the urban heat budget and CO₂ emissions on a city scale. *Environ. Sci. Tech.* **36**, 3139–3146 (2002).
- Velasco, H. E. *et al.* Eddy covariance flux measurements of pollutant gases in urban Mexico City. *Atmos. Chem. Phys.* **9**, 7325–7342 (2009).
- Valach, A. C., Langford, B., Nemitz, E., MacKenzie, A. R. & Hewitt, C. N. Seasonal and diurnal trends in concentrations and fluxes of volatile organic compounds in central London. *Atmos. Chem. Phys.* **15**, 7777–7796 (2015).
- Karl, T. *et al.* Emissions of volatile organic compounds inferred from airborne flux measurements over a megacity. *Atmos. Chem. Phys.* **9**, 271–285 (2009).
- Langford, B., Davison, B., Nemitz, E. & Hewitt, C. N. Mixing ratios and eddy covariance flux measurements of volatile organic compounds from an urban canopy (Manchester, UK). *Atm. Chem. Phys* **9**, 1971–1987 (2009).
- Rantala, P. *et al.* Anthropogenic and biogenic influence on VOC flux at an urban background site in Helsinki, Finland. *Atmos. Chem. Phys.* **16**, 7981–8007 (2016).
- Marr, L. C., Moore, T. O., Klappmeyer, M. E. & Killar, M. B. Comparison on NO_x fluxes measured by eddy covariance to emission inventories and land use. *Environ. Sci. & Techn* **47**, 1800–1808 (2013).
- Vaughan, A. R. *et al.* Spatially resolved flux measurements of NO_x from London suggest significantly higher emissions than predicted by inventories. *Faraday Discussions* **189**, 455–472 (2016).

42. EUROSTAT, “Energy, transport and environment indicators” (Tech. Report 201, *European Commission*, ec.europa.eu, ISBN 978-92-79-49471-0, doi:10.2785/547816, 2015).
43. Vergreiner, I. & Dreiseitl, E. Valley winds and slope winds – observations and elementary thoughts. *Meteorol. Atmos. Phys.* **36**, 24–286 (1987).
44. Coggon, M. M. *et al.* Emissions of nitrogen-containing organic compounds from the burning of herbaceous and arboraceous biomass: Fuel composition dependence and the variability of commonly used nitrile tracers. *GRL*, doi:10.1002/2016GL070562 (2016).
45. Popa, M. E. *et al.* Vehicle emissions of greenhouse gases and related tracers from a tunnel study: CO, CO₂, N₂O, CO₂, CH₄, CO₂, O₂:CO₂ ratios, and the stable isotopes ¹³C, ¹⁸O in CO₂ and CO. *Atmos. Chem. Phys.* **14**, 210–2123 (2014).
46. Papadimitriou, G. *et al.* “TRACCS - Transport data collection supporting the quantitative analysis of measures relating to transport and climate change” (Tech. Rep. EMISIA No. 13.RE.025.V1, 2013).
47. Harley, R. A., Marr, L. C., Lehner, J. K. & Giddings, S. N. Changes in motor vehicle emissions on diurnal to decadal time scales and effects on atmospheric composition. *Environ. Sci. Tech.* **39**, 5356–5362 (2005).
48. Lamarque, J. F. *et al.* Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application. *Atmos. Chem. Phys.* **10**, 7017–7039 (2010).
49. Vestreng, V. *et al.* Evolution of NO_x emissions in Europe with focus on road transport control measures. *Atmos. Chem. Phys.* **9**, 1503–1520 (2009).
50. Reis, S. *et al.* A. Road traffic emissions –predictions of future contributions to regional ozone levels in Europe. *Atmos. Environ.* **34**, 4701–4710 (2000).
51. Stephens, S. *et al.* Weekly patterns of Mexico City’s surface concentrations of CO, NO_x, PM₁₀ and O₃ during 1986–2007. *Atmos. Chem. Phys.* **8**, 5313–5325 (2008).
52. Hellen, H., Hakola, H. & Laurila, T. Determination of source contributions of NMHCs in Helsinki (60 degrees N, 25 degrees E) using chemical mass balance and the Unmix multivariate receptor models. *Atmos. Environ.* **37**, 1413–1424 (2003).
53. Borbon, A. *et al.* Emission ratios of anthropogenic volatile organic compounds in northern mid-latitude megacities: Observations versus emission inventories in Los Angeles and Paris. *J. Geophys. Res.* **118**, 2041–2057 (2013).
54. Spirig, C., Neftel, A., Kleinman, L. I. & Hjorth, J. NO_x versus VOC limitation of O₃ production in the Po valley: Local and integrated view based on observations, *J. Geophys. Res.* **107**, doi:10.1029/2001JD000561 (2002)
55. Wolfe, G. & Thornton, J. The Chemistry of atmosphere-forest exchange (CAFÉ) model – Part 1: Model description and characterization. *Atmos. Chem. Phys.* **11**, 77–101 (2011).
56. Kim, S. W. *et al.* Modeling the weekly cycle of NO_x and CO emissions and their impacts on O₃ in the Los Angeles – South Coast air Basin during the CalNex 2010 field campaign. *J. Geophys. Res. Atmos.* **121**, 1340–1360 (2016).
57. ExxonMobil, “The Outlook for Energy: A view to 2040” (Techn. Report, <http://corporate.exxonmobil.com/en/energy/energy-outlook/introduction/a-view-to-2040> retrieved on 1st of August 2016).
58. Molina, L. T. *et al.* An overview of the MILAGRO 2006 Campaign: Mexico City emissions and their transport and transformation. *Atmos. Chem. Phys.* **10**, 8697–8760 (2010).
59. Marx, O., Brümmer, C., Ammann, C., Wolff, V. & Freibauer, A. TRANC – a novel fast-response converter to measure total reactive atmospheric nitrogen. *Atmos. Meas. Tech.* **5**, 1045–1057 (2012).
60. Rummel, U., Ammann, C., Gut, A., Meixner, F. X. & Andreae, M. O. Eddy covariance measurements of nitric oxide flux within an Amazonian rainforest, *J. Geophys. Res.* **107**, doi:10.1029/2001JD000520 (2002).
61. Foken, T., Leuning, R., Oncley, S. P., Mauder, M. & Aubinet, M. Corrections and data quality, Eddy Covariance: A Practical Guide to Measurement and Data Analysis (Kluwer Academic Publishers, Dordrecht, ed. 1, 2010), pp. 85–131.

Acknowledgements

T.K. received support from the EC Seventh Framework Program (Marie Curie Reintegration Program, “ALP-AIR”, grant no. 334084).

Author Contributions

Karl, T. and Graus, M. conceived the experiment. Karl, T., Graus, M., Striednig, M., Lamprecht C., Hammerle, A., Wohlfahrt, G., Held, A., von der Heyden, L., and Deventer J. designed and performed the INNAQS field experiment, interpreted the data and wrote the manuscript with inputs from all co-authors; Lee, J. provided input on the interpretation of NO_x flux data; Krismer A. provided air quality data, Feichter R. performed and analyzed traffic flow data; Haun C. provided the bottom up stationary emission source data; Karl T. performed the model simulations.

Additional Information

Supplementary information accompanies this paper at doi:10.1038/s41598-017-02699-9

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017