

# 1 Space invaders; biological invasions in marine conservation planning

2 **Authors:** Sylvaine Giakoumi<sup>1,2</sup>, François Guilhaumon<sup>3</sup>, Salit Kark<sup>2</sup>, Antonio Terlizzi<sup>4,5</sup>, Joachim Claudet<sup>6,7</sup>,  
3 Serena Fellingine<sup>4</sup>, Carlo Cerrano<sup>8</sup>, Marta Coll<sup>9,10</sup>, Roberto Danovaro<sup>5,8</sup>, Simonetta Fraschetti<sup>4</sup>, Drosos  
4 Koutsoubas<sup>11,12</sup>, Jean-Batiste Ledoux<sup>10,13</sup>, Tessa Mazor<sup>14</sup>, Bastien Mérigot<sup>15</sup>, Fiorenza Micheli<sup>16</sup> and Stelios  
5 Katsanevakis<sup>11</sup>

6 <sup>1</sup>Université Nice Sophia Antipolis, CNRS, FRE 3729 ECOMERS, Parc Valrose, 28 Avenue Valrose, 06108  
7 Nice, France; s.giakoumi@uq.edu.au

8 <sup>2</sup> The Biodiversity Research Group, ARC Centre of Excellence for Environmental Decisions and NESP  
9 Threatened Species Recovery hub, School of Biological Sciences, The University of Queensland, Brisbane,  
10 Queensland, Australia; salit.kark@gmail.com

11 <sup>3</sup>Institut de Recherche pour le Développement (IRD), MARBEC - Biodiversité Marine et ses usages, UMR  
12 9190 - University of Montpellier, Montpellier, France; francois.guilhaumon@ird.fr

13 <sup>4</sup>Dipartimento di Scienze e Tecnologie Biologiche ed Ambientali, Università del Salento, CoNISMa, 73100  
14 Lecce, Italy; antonio.terlizzi@unisalento.it; serena.fellingine@unisalento.it; simona.fraschetti@unisalento.it

15 <sup>5</sup> Stazione Zoologica Anton Dohrn, Villa Comunale I, Napoli, Italia

16 <sup>6</sup> National Center for Scientific Research, CRIOBE, USR 3278 CNRS-EPHE-UPVD, Perpignan, France;  
17 joachim.claudet@gmail.com

18 <sup>7</sup> Laboratoire d'Excellence CORAIL, 58 Avenue Paul Alduy, 66860, Perpignan cedex, France

19 <sup>8</sup> Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, UO CoNISMa,  
20 via Breccie Bianche, I-60131, Ancona, Italy; c.cerrano@univpm.it; r.danovaro@univpm.it

21 <sup>9</sup> Institut de Recherche pour le Développement (IRD), UMR MARBEC & LMI ICEMASA, University of  
22 Cape Town, Private Bag X3, Rondebosch, Cape Town 7701, South Africa; marta.coll.work@gmail.com

23 <sup>10</sup> Institut de Ciències del Mar CSIC, Passeig Marítim de la Barceloneta 37-49, E-08003, Barcelona, Spain;  
24 jbbaptiste.ledoux@gmail.com

25 <sup>11</sup> University of the Aegean, Department of Marine Sciences, University Hill, Mytilene 81100, Greece;  
26 drosos@aegean.gr; stelios@katsanevakis.com

27 <sup>12</sup> National Marine Park of Zakynthos, Zakynthos 29100, Greece

28 <sup>13</sup> CIIMAR/CIMAR, Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto,  
29 Porto, Portugal

30 <sup>14</sup> CSIRO Oceans and Atmosphere Flagship, EcoSciences Precinct 41, Brisbane, QLD, Australia;  
31 tessa.mazor@csiro.au

32 <sup>15</sup> University of Montpellier, UMR 9190 MARBEC, Station Ifremer, Avenue Jean Monnet, BP 171, 34203  
33 Sète Cedex, France; bastien.merigot@univ-montp2.fr

34 <sup>16</sup> Hopkins Marine Station, Stanford University, Pacific Grove CA 93950, USA; micheli@stanford.edu

35

36 **Keywords:** alien species, biological invasions, conservation planning, impacts, marine biogeographic  
37 regions, marine protected areas, management actions, Mediterranean Sea.

38 **Running title:** Biological invasions in marine conservation planning

39 **Article Type:** Biodiversity Research

40 **Corresponding author:** Sylvaine Giakoumi

41 **Word count**

42 **Main body: 4884**

43 **Abstract: 298**

44 **References: 59**

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60 **ABSTRACT**

61 **Aim** Biological invasions are major contributors to global change and native biodiversity decline. However,  
62 they are overlooked in marine conservation plans. Here, we examine for the first time the extent to which  
63 marine conservation planning research has addressed (or ignored) biological invasions. Furthermore, we  
64 explore the change of spatial priorities in conservation plans when different approaches are used to  
65 incorporate the presence and impacts of invasive species.

66 **Location** Global analysis with a focus on the Mediterranean Sea region.

67 **Methods** We conducted a systematic literature review consisting of three steps: 1) article selection using a  
68 search engine, 2) abstract screening, and 3) review of pertinent articles, which were identified in the second  
69 step. The information extracted included the scale and geographic location of each case study as well as the  
70 approach followed regarding invasive species. We also applied the software Marxan to produce and compare  
71 conservation plans for the Mediterranean Sea that either protect, or avoid areas impacted by invasives, or  
72 ignore the issue. One case study focused on the protection of critical habitats, and the other on endemic fish  
73 species.

74 **Results** We found that of 119 papers on marine spatial plans in specific biogeographic regions only three  
75 (2.5%) explicitly took into account invasive species. When comparing the different conservation plans for  
76 each case study, we found that the majority of selected sites for protection (ca. 80%) changed in the critical  
77 habitat case study, while this proportion was lower but substantial (27%) in the endemic fish species case  
78 study.

79 **Main conclusions** Biological invasions are being widely disregarded when planning for conservation in the  
80 marine environment across local to global scales. More explicit consideration of biological invasions can  
81 significantly alter spatial conservation priorities. Future conservation plans should explicitly account for  
82 biological invasions to optimize the selection of marine protected areas.

83

84

85

## 86 INTRODUCTION

87 Biological invasions are amongst the major components of current global change and drivers of native  
88 biodiversity loss in terrestrial, freshwater, and marine ecosystems (Pyšek & Richardson, 2010; Simberloff *et*  
89 *al.*, 2013). Alien species (i.e. organisms introduced outside their natural range) can become invasive and  
90 substantially change species composition and the functioning of native ecosystems by a range of processes:  
91 competition, predation, overgrazing, release of toxins, hybridization, disease transmission, and habitat  
92 alteration (Levine, 2008; Vilà *et al.*, 2011). In the marine environment, ecological impacts including the loss  
93 of native genotypes, degradation of habitats, changes in trophic interactions, and displacement of native  
94 species have been documented (Albins, 2012; Katsanevakis *et al.*, 2014; Verges *et al.*, 2014). Invasives can  
95 also impact the provision of ecosystem services with negative socio-economic consequences for coastal  
96 communities, for instance causing the decline of commercial fish and shellfish stocks or decreasing the  
97 potential for recreational activities (Bax *et al.*, 2003; Katsanevakis *et al.*, 2014). Moreover, some marine  
98 invasives are venomous or toxic and can have negative impacts on human health (Streftaris & Zenetos,  
99 2006). The multi-dimensional consequences of invasives render their distribution and impacts major topics  
100 of scientific interest with crucial conservation implications (Molnar *et al.*, 2008; Katsanevakis *et al.*, 2016).

101 Globally, there is an urgent need to adopt management strategies for the control of invasive populations and  
102 the mitigation of their impacts. The Aichi Target 9 of the Convention on Biological Diversity (CBD) states  
103 that by 2020: i) invasive alien species and pathways are identified and prioritized, ii) priority species are  
104 controlled or eradicated, and iii) measures are in place to manage pathways to prevent their introduction and  
105 establishment (Convention on Biological Diversity, 2015). Regional policies have also focused on the uptake  
106 of management actions for the mitigation of invasives' impacts. For instance, under the European Union  
107 Marine Strategy Framework Directive (EU, 2008), member states are committed to developing strategies to  
108 achieve Good Environmental Status (GES) by 2020. One of the GES descriptors dictates that alien species  
109 should be at density levels that do not adversely alter ecosystems. Nevertheless, comprehensive strategies to  
110 mitigate impacts of alien species on marine biodiversity and ecosystem services have not yet been developed  
111 in the EU.

112 Despite the increasing number of studies addressing the assessment of invasion pathways (e.g., Seebens *et*  
113 *al.*, 2013; Essl *et al.*, 2015) and impacts of biological invasions on marine ecosystems (e.g., Katsanevakis *et*

114 *al.*, 2014; Katsanevakis *et al.*, 2016), there is still a gap in our understanding of how to use such information  
115 to guide conservation planning. Should conservation plans target areas that are highly invaded by alien  
116 species and invest resources in mitigating negative impacts of invasives? Alternatively, should plans avoid  
117 highly invaded areas and invest resources in non-invaded or less invaded areas? In marine conservation  
118 planning, the first hypothesis would favour an approach to *protect* areas highly impacted by invasives in  
119 order to restore them by taking additional management actions, e.g., eradication, within those areas. The  
120 second hypothesis would lead planners to *avoid* such areas and protect areas less vulnerable to invasions. In  
121 the absence of a good knowledge base on which hypothesis is valid under which conditions, the easy  
122 approach is to just *ignore* the issue.

123 Here, we examine whether marine conservation plans have directly addressed biological invasions by either  
124 protecting or avoiding impacted areas, or not (thus they have ignored the issue deliberately or not).

125 Furthermore, we use two case studies (one habitat-based and one species-based) to explore how spatial  
126 priorities change when areas with high alien species density and impacts are protected, avoided, or ignored  
127 (i.e. information about biological invasions was not considered). We base our case studies in the  
128 Mediterranean Sea, one of the major hotspots of marine biological invasions (Edelist *et al.*, 2013).

129 Approximately 1,000 alien species have been reported in the Mediterranean Sea (Zenetos *et al.*, 2012), and  
130 this number is expected to grow after the enlargement of the Suez Canal (Galil *et al.*, 2014). Simultaneously,  
131 the identification of priority areas for conservation is ongoing in the region, as Mediterranean countries aim  
132 to achieve Aichi Target 11 of the CBD by protecting 10% of the sea under their jurisdiction. Invasive species  
133 may nullify or in some cases benefit (Schlaepfer *et al.*, 2011) the effects of protection, such as ecosystem  
134 recovery. Thus, the presence of such species and their impacts should be explicitly considered when  
135 selecting marine protected areas (MPAs). Synthesizing our findings we identify gaps in knowledge that need  
136 to be filled in order to optimize MPA site-selection under global changes, specifically when accounting for  
137 invasive species, in the Mediterranean region and beyond.

## 138 **METHODS**

### 139 **Literature review and synthesis**

140 We performed a bibliographic search using the Elsevier's Scopus database ([www.scopus.com](http://www.scopus.com)). Eligibility  
141 criteria included any paper or review published between 1950 and the cut-off date 18 April 2015 with the

142 terms ‘conservation planning’ and ‘marine’ or ‘sea’ in the title, keywords or abstract. Grey literature and  
143 non-English publications were not considered in this review.

144 The results summed up to 793 peer-reviewed papers. Our review started with a screening of these 793 paper  
145 abstracts. Articles were excluded if they: 1) were unrelated to conservation planning, 2) did not include a  
146 specific case study for which a conservation plan was developed, 3) took into account only terrestrial or  
147 freshwater species, habitats, or ecosystems and not marine, or 4) mentioned the term “conservation planning”  
148 only for justification or discussion of results but did not produce a conservation plan. As a result, 214  
149 abstracts (27%) qualified for the next round of reviews. These were papers that presented conservation plans  
150 in marine environments, or included content that was potentially relevant after reading the abstract alone, and  
151 were thus retained for the second step of the analysis.

152 In the second selection process, the entire 214 articles were read, using the same exclusion criteria listed  
153 above. Finally, 119 studies were suitable for the qualitative and quantitative synthesis (see Appendix S1 in  
154 Supporting Information for final list of articles).

155 The following information was extracted from each article (Table S1): 1) year of publication; 2) scale of case  
156 study (local < national < regional < global); 3) geographic location of the case study; 4) the relevant marine  
157 biogeographic region ("realm" according to Spalding *et al.* (2007)); 5) the features (species, habitats,  
158 ecosystems) that were targeted for conservation; 6) the conservation planning method/tool that was used  
159 (e.g., Marxan, Zonation); 7) the approach the study followed regarding biological invasions, i.e. whether  
160 biological invasions were taken into account in the planning process by ‘protecting’ or ‘avoiding’ areas  
161 impacted by invasive species or the issue was ‘ignored’; and 8) the method that was used if the ‘avoid’ or  
162 ‘protect’ approach was followed.

### 163 **Conservation plans: applying the ‘protect’, ‘avoid’, or ‘ignore’ approaches in two Mediterranean case** 164 **studies**

165 In addition to the literature review exploring how biological invasions have been treated in past conservation  
166 plans, we examined whether and how spatial priorities change when biological invasions are explicitly  
167 accounted in conservation planning. Here, we used two case studies to compare systematic conservation  
168 plans that followed three different approaches for dealing with invasive species: protect, avoid, or ignore

169 areas impacted by invasives. One case study aimed to account for impacts of invasives on two critical marine  
170 habitats, the seagrass *Posidonia oceanica* meadows and coralligenous formations. The second case study  
171 aimed to assess changes in priority conservation areas for endemic fish species when accounting (or not) for  
172 invasives.

173 To identify conservation priority areas for our features of interest (habitats and species), we used the  
174 conservation planning software Marxan (Ball *et al.*, 2009). This software uses a simulated annealing  
175 algorithm to find a suite of good near-optimal systems of priority areas that meet conservation targets while  
176 minimizing socio-economic costs. In Marxan, the user sets a target for every feature to be protected, which  
177 in our case was expressed as the percentage of the feature's overall distribution range (see below case studies  
178 1 & 2). The study area was the entire Mediterranean Sea excluding areas deeper than 1,000 m, where the  
179 habitats and species included in these analyses do not occur (Giakoumi *et al.*, 2013; Guilhaumon *et al.*,  
180 2015). The study area was divided into a grid of 12,828 cells (hereafter planning units) each of 10 x10 km.  
181 Marxan was run 1,000 times and consisted of 1,000,000 iterations per run. We defined areas of greater  
182 irreplaceability by using the selection frequency of each planning unit, which is the proportion of runs in  
183 which a planning unit is selected amongst the 1,000 runs. These areas were considered higher priority for  
184 protection. The Boundary Length Modifier (BLM, measure of trade-off between cost and compactness of the  
185 solution) was set to 0, as our aim was to examine differences in the selection of priority areas among the  
186 scenarios and not to design an MPA network with a desirable level of compactness.

#### 187 *Case study 1: Critical habitats*

188 Data (presence/absence) on the distribution of seagrass *P. oceanica* meadows and coralligenous formations  
189 were obtained from Giakoumi *et al.* (2013). We set a 60% target of the current distribution of the *P.*  
190 *oceanica* meadows and 40% of the distribution of coralligenous formations as per Giakoumi *et al.* (2013)  
191 following guidelines by the EU (ETC/BD, 2010). Although these targets are policy-based and are not  
192 supported by solid ecological evidence, they represent the current practice in EU and it is thus a pragmatic  
193 approach to follow. In the 'protect' scenario we targeted the proportion of seagrass meadows and  
194 coralligenous formations impacted by alien species in each planning unit. The impacted habitat feature  
195 within each site was estimated based on the CIMPAL index (Cumulative IMPacts of invasive ALien species)  
196 developed by Katsanevakis *et al.* (2016). For the CIMPAL index, cumulative impact scores were estimated

197 on the basis of the distributions of habitats and alien species, the reported magnitude of ecological impacts,  
198 and the strength of such evidence. Evidence for most of the reported impacts of marine aliens in the literature  
199 is weak, mostly based on expert judgement or dubious correlations (Katsanevakis *et al.*, 2014). Hence, in the  
200 estimation of the CIMPAL index the weights of impacts with low supporting evidence are downweighted, in  
201 comparison to impacts documented through manipulative or descriptive experiments (Katsanevakis *et al.*,  
202 2016). The index was normalized as follows to obtain values between 0 and 1:  $I_i = \frac{x_i - \min(x)}{\max(x) - \min(x)}$ ,

203 where  $I_i$  is the normalized index value and  $x_i$  is the initial index value for the planning unit  $i$ .

204 Then, to estimate an index ( $E$ ) of the magnitude of impacts on each planning unit  $i$  in which a specific feature  
205 is present, the presence or absence of the feature ( $F$ ) was multiplied by the index value ( $I$ ):

$$206 \quad E_i = F_i * I_i$$

207 In the ‘avoid’ scenario, we only set targets for the features in good condition (i.e. not impacted by alien  
208 species). An index of the condition ( $H$ ) of a specific feature in each planning unit  $i$  was estimated as:

209

210 In the ‘ignore’ scenario we did not consider the information about impacts from invasives on the critical  
211 habitats as per Giakoumi *et al.* (2013).

212 The most commonly accounted for and significant cost in marine planning is opportunity cost, e.g., fishing  
213 profits that are forgone when an area is made a no-take zone (Ban & Klein, 2009). The socio-economic cost  
214 used herein represents the spatial distribution of the combined opportunity cost for three marine sectors:  
215 commercial (small and large-scale) fishing, non-commercial fishing (recreational and subsistence), and  
216 aquaculture. Data were obtained from Mazor *et al.* (2014).

### 217 *Case study 2: Endemic fish species*

218 Data on the distribution of 80 endemic fish species were obtained from Guilhaumon *et al.* (2015). Among the  
219 80 species, 54 were benthic, 18 demersal, and 8 pelagic (Appendix S2). We used area-based species-specific  
220 representation targets following the methods in Guilhaumon *et al.* (2015). A representation target of 100%  
221 was set for endemic species with restricted-ranges (geographic range of <1,000 km<sup>2</sup>) and a target of 10% was



222 used for widespread endemics (those endemic species with a geographic range  $> 35,860 \text{ km}^2$ , corresponding  
223 to one third of the species). For endemics with intermediate range sizes, the target was interpolated as a  
224 linear function of log-transformed area of occupancy. Additionally, we modified the area-based targets  
225 according to the species level of threat as determined by the IUCN Red List categories (Abdul Malak *et al.*,  
226 2011). Following Kark *et al.* (2009) the representation target of critically endangered species ( $n=1$ ) was set  
227 to 100% irrespective of their geographic range; the targets for species that are vulnerable ( $n=1$ ) or  
228 endangered ( $n=3$ ) were defined as the maximum between the 30% of their geographic range and their  
229 linearly interpolated target. Data deficient species ( $n=1$ ) and species not evaluated by IUCN ( $n=71$ ) were  
230 attributed the “least-concern” IUCN category (Appendix S2).

231 We accounted for impacts of alien species by combining the values of the relative Functional Nearest  
232 Neighbour index (FNNr; see Elleouet *et al.*, 2014) with the socio-economic cost (Mazor *et al.*, 2014). The  
233 FNNr index arises from a trait-based approach and expresses the magnitude of functional similarity (or niche  
234 overlap) between endemic and alien species as a proportion of the total number of endemic species per  
235 planning unit. The FNNr index assumes that co-occurring native and alien species are more likely to interact  
236 if they have greater similarity in their ecological (e.g. habitat use) and biological (e.g. diet) attributes, that is,  
237 greater similarity in their ecological niches (*sensu* Violle & Jiang, 2009).

238 In the ‘avoid’ scenario, we summed the values of FNNr index (ranging from 0 to 1) and the socio-economic  
239 cost in each planning unit. In order to give the same weight to the two components, the FNNr index and the  
240 socio-economic cost were rescaled to range in the same magnitude. High FNNr index values increased the  
241 cost of planning units in the ‘avoid’ scenario, and thus the optimization algorithm avoided the selection of  
242 these areas. This scheme was reversed in the ‘protect’ scenario, where  $1-\text{FNNr}$  values were added to the  
243 socio-economic cost. Planning units with high FNNr values contributed less to the cost of the planning units,  
244 and these areas were more likely to be selected for protection. In the ‘ignore’ scenario, we did not consider  
245 the information about potential ecological interactions between endemics and aliens and ran Marxan  
246 considering only the socio-economic cost.

## 247 **RESULTS**

### 248 **Biological invasions in past marine conservation plans**

249 Since 2000, there has been a progressive increase in the number of publications on marine conservation  
250 plans, resulting in a total of 119 publications (Appendix S1; Fig. S1A). Most of these publications (57%)  
251 referred to local scales (Fig. S1B). The reviewed conservation plans covered all marine realms, with a higher  
252 concentration in the Temperate Northern Atlantic and the Central Indo-Pacific realms (Fig. 1). The majority  
253 of conservation plans (58%) included habitats or ecosystems as features to conserve (Fig. 2). A large  
254 percentage of studies also set fish species distributions as conservation features (33%). Charismatic marine  
255 animals, particularly mammals and birds, were also commonly targeted for protection (23% and 22%  
256 respectively). For the identification of priority areas for conservation of these features, half the studies used  
257 conservation planning software. Of those, the vast majority (88%) used some version of the software  
258 Marxan, whereas the rest of them used C-Plan (Pressey *et al.*, 2009) and Zonation (Moilanen *et al.*, 2009).  
259 The other half of the studies used a variety of tools: geospatial analyses (e.g., ArcGIS), species distribution  
260 and habitat suitability models, complementarity analyses, hotspot analyses, food-web models, univariate and  
261 multivariate statistical methods, GLM models, tracking methods, scoring methods, vulnerability assessments,  
262 and combinations of those.

263 Out of the 119 papers included in our analyses we found only three papers (Tallis *et al.*, 2008; Giakoumi *et*  
264 *al.*, 2011; Klein *et al.*, 2013) that explicitly took into account invasive species in their conservation plans  
265 (Table S1). All other papers ignored invasives' presence and/or impacts (Table S1; Fig. S1A). All three  
266 studies used Marxan software. Tallis *et al.* (2008) incorporated threats in a site-prioritization exercise for the  
267 Pacific Northwest coast ecoregion (U.S.A.), including invasive species, into Marxan's cost function. Areas  
268 with higher threat had higher cost, thus, highly invaded areas were avoided. Similarly, Klein *et al.* (2013) in  
269 a conservation plan for California incorporated threats, including invasives, into Marxan by adding an  
270 additional constraint: minimize the chance that the reserved features are in poor condition. The algorithm,  
271 therefore, favoured the selection of priority conservation areas less impacted by threats, one of which was  
272 vulnerability to invasives. In contrast, Giakoumi *et al.* (2011) set conservation targets for all fish species of  
273 the shallow sublittoral of the Cyclades Archipelago (Greece), including the invasive herbivore species  
274 *Siganus luridus*; following, thus, the 'protect' approach.

## 275 **Comparing the consequences of 'protect', 'avoid', or 'ignore' strategies for conservation plans**

### 276 *Critical habitats case study*

277 We found that the selection frequency of the great majority of planning units changed depending on the  
278 approach that was followed (protect, avoid or ignore). Only ~13% of the planning units containing a  
279 conservation feature had maximum irreplaceability (i.e., a selection frequency of 1,000) across all three  
280 scenarios (green-bordered planning units in Fig. 3). In all pairwise scenario comparisons ('protect' *versus*  
281 'ignore', 'avoid' *versus* 'ignore', and 'protect' *versus* 'avoid'), the selection of ~80% of planning units  
282 differed (Table 1; Fig. 3). Areas highly impacted by invasive species, such as the Balearic Islands (Eastern  
283 Spain), Sicily (South Italy), and the Greek Ionian coastal waters (Western Greece) presented higher selection  
284 frequency in the 'protect' rather than the 'ignore' scenario. These same areas presented higher selection in  
285 the 'ignore' scenario than in the 'avoid'. When comparing the 'protect' and 'avoid' scenarios, the highly  
286 impacted areas presented higher selection in the 'protect' than in the 'avoid' scenario.

#### 287 *Endemic fish species case study*

288 In all pairwise scenario comparisons, the selection of nearly one third (27%) of planning units differed  
289 (Table 1; Fig. 4). Only ~3% of planning units presented maximum irreplaceability across all three scenarios  
290 (green-bordered planning units in Fig. 4). When comparing the 'protect' and 'ignore' scenarios, no clear  
291 geographical pattern arose. Planning units showing greater irreplaceability in the 'protect' approach were  
292 spread across the Mediterranean Sea. However, some patches of markedly higher irreplaceability could be  
293 identified in the Gulf of Lions (France) and in the Adriatic Sea (eastern Italian coast). These areas presented  
294 higher irreplaceability in the 'avoid' scenario compared to the 'ignore' scenario. Finally, in the pairwise  
295 comparison 'protect' *versus* 'avoid' scenario, irreplaceability substantially increased in the 'avoid' scenario  
296 along the coastal waters of Italy in the Adriatic Sea and moderately increased in patchy locations along all  
297 Mediterranean coasts. Planning units exhibiting higher irreplaceability in the 'protect' scenario were mainly  
298 located along the Greek coast and remaining Adriatic Sea.

## 299 **DISCUSSION**

300 Our literature review demonstrates that the role of biological invasions has been widely overlooked when  
301 planning for conservation in the marine environment, at all spatial scales. Yet, the explicit consideration of  
302 biological invasions can significantly change spatial conservation priorities. This is clearly shown by the  
303 comparison we made of conservation plans following three different approaches: 'avoid', 'protect' or  
304 'ignore' areas with high presence and/or impacts of invasives. Our findings have important implications on

305 the placement of new MPAs in order for countries to achieve the 10% goal set by Aichi Target 11 of the  
306 Convention on Biological Diversity (2015).

307 In the Mediterranean Sea, invasive species are considered one of the most severe threats to species and  
308 ecosystems (Coll *et al.*, 2012; Micheli *et al.*, 2013a). When making decisions about the establishment of new  
309 MPAs, this threat should be explicitly taken into account for an effective allocation of conservation funds.  
310 Particular attention should be given to areas where changes in the priority selection among scenarios are  
311 more pronounced: the Balearic Islands in Spain, the Gulf of Lions in France, Sicily in Italy, the Adriatic Sea,  
312 and the Greek coasts (especially in the west). The importance of biological invasions in these areas differed  
313 depending on which features were targeted for protection (habitats or fish species). To make informed  
314 decisions about the placement of new MPAs, a holistic approach targeting numerous species and habitats  
315 would be desirable.

316 We propose that in order to effectively incorporate biological invasions into marine conservation planning in  
317 the future, the scientific community should urgently fill information gaps regarding: 1) the spatial  
318 distribution of invasive species both at present and in the future; 2) the ecological and socio-economic  
319 impacts of biological invasions; and 3) the role of MPAs in controlling invasive populations and mitigating  
320 their impacts.

321 Extensive mapping efforts of invasive species distributions should urgently be applied. Whether the planning  
322 approach is 'avoid' or 'protect', accurate information about the distribution of alien species is a prerequisite  
323 for effective planning as we demonstrated in our case studies. Several governmental and intergovernmental  
324 bodies have already invested important resources in the creation of georeferenced databases of the current  
325 distribution of alien species (e.g. Katsanevakis *et al.*, 2015). Nevertheless, biological invasions are a dynamic  
326 threat (Strayer *et al.*, 2006), and predictions of their future distributions is crucial for effective management  
327 plans and selection of new MPA sites. Areas that are currently unaffected by biological invasions may be  
328 severely affected in the future, therefore a dynamic conservation is required. At present, accurate projections  
329 of future distributions of marine alien species are limited. Species distribution models forecasting the spread  
330 of aliens are currently based on climate predictions and may underestimate the potential spread of aliens  
331 (Parravicini *et al.*, 2015), and interactions with other sources of disturbance (Bulleri *et al.*, 2011). Studies  
332 comparing source and front populations across a species new range could prove useful for better

333 understanding and predicting populations dynamics of marine aliens and thus, providing guidance for  
334 potential mitigation actions and for new MPA siting.

335 Further research is required to better understand the ecosystem changes biological invasions may cause to  
336 native ecosystems and their impacts on socio-economic activities. To date, evidence shows that most alien  
337 species have negative impacts on native biodiversity and human wellbeing (e.g. Katsanevakis *et al.*, 2014).  
338 However, in some cases, alien species can provide conservation benefits and contribute to conservation  
339 objectives; for instance, they can provide habitat or food resources to rare species, serve as functional  
340 substitutes for extinct taxa, and facilitate the recovery of degraded ecosystems (Schlaepfer *et al.*, 2011). For  
341 example, in New England, USA, the invasion of green crabs, *Carcinus maenas*, into heavily burrowed salt  
342 marshes partially promoted cordgrass recovery by reversing trophic cascades that were triggered by  
343 overfishing of salt marsh predators (Bertness & Coverdale, 2013). Invasives can also provide new economic  
344 opportunities. For instance, Mollo *et al.* (2014) showed how targeted exploitation of invasives can lead to  
345 new biotechnological and pharmacological applications. In the Levantine Sea, the world's most invaded sea,  
346 a large percentage of fisheries is now composed of invasive fish species (Edelist *et al.*, 2013). The  
347 commercial exploitation of such species has created new opportunities for local fisheries. Schlaepfer *et al.*  
348 (2011) speculate that alien species might contribute to achieving conservation goals in the future because  
349 they may be more likely than native species to persist and provide ecosystem services in areas where climate  
350 and land use are changing rapidly. Nevertheless, the contribution of alien species to achieving conservation  
351 and economic goals is likely species-specific, as is their response to the alternative planning strategies  
352 ('protect' or 'avoid'). Therefore, additional information on the impacts (negative or positive) alien species  
353 have on ecosystems and human activities is crucial for the formulation of conservation targets for specific  
354 species or habitats during the planning process.

355 Lastly, more information is required on whether MPAs are a useful conservation strategy for the  
356 management of alien populations. The 'biotic resistance hypothesis' states that ecosystems with high species  
357 richness are more resistant to invaders than those with low biodiversity (Levine & D'Antonio, 1999; Jeschke,  
358 2014). Hence, the expected recovery of native species richness within MPAs could prevent the penetration  
359 and settlement of alien species. Furthermore, the restoration of top-down regulation processes (e.g.,  
360 restoration of top predators' populations) in MPAs could help control the spreading of some alien species

361 inside MPAs (Mumby *et al.*, 2011). Nonetheless, numerous studies have reported the opposite pattern, i.e.  
362 positive relationships between the numbers of native and alien species (McKinney, 2002). These  
363 observations led to the ‘biotic acceptance hypothesis’ – which supports the notion that ecosystems can  
364 accommodate the establishment of aliens and their coexistence with native species –and to a rich-get-richer  
365 pattern where areas with high native species richness support high numbers of alien species (Stohlgren *et al.*,  
366 2006). Moreover, the populations of some alien species could be enhanced in MPAs mainly because they  
367 would benefit from non-harvesting (Burfeind *et al.*, 2013). Therefore, further empirical studies are necessary  
368 to assess the potential role of MPAs in controlling alien species and mitigating their impacts. If MPAs prove  
369 to have no effect or even favour invasive species then their establishment in impacted areas should either be  
370 avoided (Boudouresque & Verlaque, 2005), or complemented with other management measures for  
371 successful invasion control and mitigation of invasives’ impacts (Thresher & Kuris, 2004).

372 Based on current evidence and until the effects of MPAs on alien and particularly invasive species are clearly  
373 demonstrated the ‘protect’ or ‘avoid’ planning approaches should be selected. This selection will depend on  
374 the specificities of the study area, the expected response of invasive populations to protection, and their  
375 negative or positive impacts on ecosystem functioning and services. A ‘protect’ approach could be followed  
376 for the restoration of some habitats and the protection of specific populations impacted by invasives, or for  
377 the protection of alien species that have proven to be beneficial for ecosystems or human wellbeing.  
378 Conversely, an ‘avoid’ approach may be developed for harmful alien species that cannot be controlled at a  
379 reasonable cost as well as for habitats on which no substantial effect of protection is anticipated. An  
380 alternative would be to prioritize for conservation areas that are always selected as priorities regardless of the  
381 approach, and are thus less susceptible to biological invasions. In our case studies, these areas are those  
382 highlighted in green in Figs 3 and 4, and interestingly most of them coincide with ‘consensus areas’  
383 proposed by Micheli *et al.* (2013b).

384 Despite the potential effectiveness of MPAs in mitigating the impacts of invasive species locally, MPAs  
385 alone are unlikely to be sufficient for managing the impact of invasives. Additional management actions  
386 aimed at prevention as well as mitigation of invasives’ impacts are required both inside and outside MPAs.  
387 For instance, eradication of recent alien introductions (Myers *et al.*, 2000; Anderson, 2005) and actions to  
388 control well-established invasive populations, such as harvesting by divers (Green *et al.*, 2014), the use of

389 selective fishing gear (Archdale *et al.*, 2010) and the controlled development of targeted fisheries may be  
390 examined as management actions to assist the recovery of highly impacted areas under a ‘protect’ approach.  
391 Suppressing invasives below population densities that cause environmental harm can have a similar effect to  
392 complete eradication, in terms of protecting the native biodiversity on a local scale (Green *et al.*, 2014). Such  
393 management actions should be incorporated into spatial plans and be prioritized on the basis of their cost-  
394 effectiveness, accounting for the cost of actions and their expected benefits on ecosystems (Giakoumi *et al.*,  
395 2015).

396

## 397 **CONCLUSION**

398 Our review reveals that explicit consideration of biological invasions is lacking in marine conservation plans.  
399 At the same time, our case studies highlight that the approach taken to include this issue (protect or avoid  
400 invasive species) or not (ignore the relevant information) can lead to different recommendations regarding  
401 conservation priorities. The lack of explicit consideration of biological invasion in conservation planning  
402 might be partly driven by the large remaining uncertainty regarding how invasive species respond to  
403 conservation actions, and how they may influence the outcomes of such actions. Other reasons might be: the  
404 limited data availability and scientific understanding of biological invasions; the limited awareness and  
405 concern by policy makers; and consequently, the limited funding directed to the control of alien populations  
406 and mitigation of their impacts. More research is clearly needed to determine the more effective strategy for  
407 incorporating biological invasions in marine conservation planning. Research priorities should involve  
408 multidisciplinary approaches and include: 1) extensive mapping efforts of invasive species distributions and  
409 development of accurate models for the prediction of their future distributions; 2) assessment of invasive  
410 species ecological and socio-economic impacts in host ecosystems; and 3) assessment of the role MPAs have  
411 in controlling invasive populations and mitigating their impacts. Ultimately, the management of invasives  
412 and their potential integration into conservation plans depend on how conservation goals are set in the future.  
413 A shift from a species-based towards a function-based approach, focusing on invasives’ functional role and  
414 their interactions with native communities (see Brown and Mumby (2014) would provide better guidance on  
415 the appropriate strategies for managing invasive species.

## 416 **ACKNOWLEDGMENTS**

417 We thank all participants of the 3<sup>rd</sup> International Workshop “Advancing Conservation Planning in the  
418 Mediterranean Sea”, (Lecce, Italy) for discussions that inspired and shaped this paper. We also thank L.  
419 Hawkes, S. Maxwell and two anonymous reviewers for their helpful suggestions. S.G. was supported by  
420 ARC CEED (University of Queensland) funding and the ANR project PAVIS; S.Kark by the Australian  
421 Research Council; J.C. by ERa-Net BiodivERsA (BUFFER project); R.D. by the programme DEVOTES  
422 (7FP); S.F. by the EU Project COCONET (7FP, Grant Agreement No. 287844); M.C. by a Marie Curie  
423 Career Integration Grant Fellowship (PCIG10-GA-2011-303534); and J.B.L by a Postdoctoral grant  
424 (SFRH/BPD/74400/2010) from the Portuguese Foundation for Science and Technology. F.M. acknowledges  
425 the support of the Pew Charitable Trust.

## 426 REFERENCES

- 427 Abdul Malak, D., Livingstone, S.R., Pollard, D., Polidoro, B.A., Cuttelod, A., Bariche, M., Bilecenoglu, M.,  
428 Carpenter, K.E., Collette, B.B., Francour, P., Goren, M., Kara, M.H., Massutí, E., Papaconstantinou,  
429 C. & Tunesi, L. (2011) Overview Of The Conservation Status Of The Marine Fishes Of The  
430 Mediterranean Sea. In, p. 61pp. IUCN, Gland, Switzerland and Malaga, Spain.
- 431 Albins, M.A. (2012) Effects of invasive Pacific red lionfish *Pterois volitans* versus a native predator on  
432 Bahamian coral-reef fish communities. *Biological Invasions*, **15**, 29-43.
- 433 Anderson, L.W.J. (2005) California’s Reaction to *Caulerpa taxifolia*: A Model for Invasive Species Rapid  
434 Response\*. *Biological Invasions*, **7**, 1003-1016.
- 435 Archdale, M.V., Anasco, C.P. & Nakagawa, A. (2010) Liftnets Compare Favorably with Pots as Harvesting  
436 Fishing Gear for Invasive Swimming Crabs. *Journal of Fisheries and Aquatic Science*, **5**, 510-516.
- 437 Ball, I.R., Possingham, H.P. & Watts, M. (2009) Marxan and relatives: Software for spatial conservation  
438 prioritisation. *Spatial conservation prioritisation: Quantitative methods and computational tools* (ed.  
439 by A. Moilanen, K.A. Wilson and H.P. Possingham), pp. 185-195. Oxford University Press, Oxford,  
440 UK.
- 441 Ban, N.C. & Klein, C.J. (2009) Spatial socioeconomic data as a cost in systematic marine conservation  
442 planning. *Conservation Letters*, **2**, 206-215.
- 443 Bax, N., Williamson, A., Aguero, M., Gonzalez, E. & Geeves, W. (2003) Marine invasive alien species: a  
444 threat to global biodiversity. *Marine Policy*, **27**, 313-323.



- 445 Bertness, M.D. & Coverdale, T.C. (2013) An invasive species facilitates the recovery of salt marsh  
446 ecosystems on Cape Cod. *Ecology*, **94**, 1937-43.
- 447 Boudouresque, C.F. & Verlaque, M. (2005) Nature conservation, marine protected areas, sustainable  
448 development and the flow of invasive species to the Mediterranean Sea. *Travaux Scientifiques du*  
449 *Parc National de Port-Cros*, **21**, 29-54.
- 450 Brown, C.J. & Mumby, P.J. (2014) Trade-offs between fisheries and the conservation of ecosystem function  
451 are defined by management strategy. *Frontiers in Ecology and the Environment*, **12**, 324-329.
- 452 Bulleri, F., Alestra, T., Ceccherelli, G., Tamburello, L., Pinna, S., Sechi, N. & Benedetti-Cecchi, L. (2011)  
453 Determinants of *Caulerpa racemosa* distribution in the north-western Mediterranean. *Marine*  
454 *Ecology Progress Series*, **431**, 55-67.
- 455 Burfeind, D.D., Pitt, K.A., Connolly, R.M. & Byers, J.E. (2013) Performance of non-native species within  
456 marine reserves. *Biological Invasions*, **15**, 17-28.
- 457 Coll, M., Piroddi, C., Albouy, C., Ben Rais Lasram, F., Cheung, W.W.L., Christensen, V., Karpouzi, V.S.,  
458 Guilhaumon, F., Mouillot, D., Paleczny, M., Palomares, M.L., Steenbeek, J., Trujillo, P., Watson, R.  
459 & Pauly, D. (2012) The Mediterranean Sea under siege: spatial overlap between marine biodiversity,  
460 cumulative threats and marine reserves. *Global Ecology and Biogeography*, **21**, 465-480.
- 461 Convention on Biological Diversity (2015) *Aichi biodiversity targets*. Available at:  
462 <https://www.cbd.int/sp/targets/> (accessed 6 November 2015).
- 463 Edelist, D., Rilov, G., Golani, D., Carlton, J.T. & Spanier, E. (2013) Restructuring the Sea: profound shifts in  
464 the world's most invaded marine ecosystem. *Diversity and Distributions*, **19**, 69-77.
- 465 Elleouet, J., Albouy, C., Ben Rais Lasram, F., Mouillot, D. & Leprieur, F. (2014) A trait-based approach for  
466 assessing and mapping niche overlap between native and exotic species: the Mediterranean coastal  
467 fish fauna as a case study. *Diversity and Distributions*, **20**, 1333-1344.
- 468 Essl, F., Bacher, S., Blackburn, T.M., Booy, O., Brundu, G., Brunel, S., Cardoso, A.-C., Eschen, R.,  
469 Gallardo, B., Galil, B., García-Berthou, E., Genovesi, P., Groom, Q., Harrower, C., Hulme, P.E.,  
470 Katsanevakis, S., Kenis, M., Kühn, I., Kumschick, S., Martinou, A.F., Nentwig, W., O'Flynn, C.,  
471 Pagad, S., Pergl, J., Pyšek, P., Rabitsch, W., Richardson, D.M., Roques, A., Roy, H.E., Scalera, R.,  
472 Schindler, S., Seebens, H., Vanderhoeven, S., Vilà, M., Wilson, J.R.U., Zenetos, A. & Jeschke, J.M.  
473 (2015) Crossing Frontiers in Tackling Pathways of Biological Invasions. *BioScience*, **65**, 769-782.

474 ETC/BD (European Topic Centre on Biological Diversity) (2010).  
475 [http://bd.eionet.europa.eu/activities/Natura\\_2000/pdfs/Additional\\_marine\\_guidelines.pdf](http://bd.eionet.europa.eu/activities/Natura_2000/pdfs/Additional_marine_guidelines.pdf) (accessed  
476 14 May 2013)

477 EU (2008) Directive of the European Parliament and the Council Establishing a Framework for Community  
478 Action in the Field of Marine Environmental Policy (Marine Strategy Framework Directive),  
479 European Commission Directive 2008/56/EC, OJ L 164. In:

480 Galil, B.S., Boero, F., Campbell, M.L., Carlton, J.T., Cook, E., Fraschetti, S., Gollasch, S., Hewitt, C.L.,  
481 Jelmert, A., Macpherson, E., Marchini, A., McKenzie, C., Minchin, D., Occhipinti-Ambrogi, A.,  
482 Ojaveer, H., Olenin, S., Piraino, S. & Ruiz, G.M. (2014) 'Double trouble': the expansion of the Suez  
483 Canal and marine bioinvasions in the Mediterranean Sea. *Biological Invasions*, **17**, 973-976.

484 Giakoumi, S., Grantham, H.S., Kokkoris, G.D. & Possingham, H.P. (2011) Designing a network of marine  
485 reserves in the Mediterranean Sea with limited socio-economic data. *Biological Conservation*, **144**,  
486 753-763.

487 Giakoumi, S., Brown, C.J., Katsanevakis, S., Saunders, M.I. & Possingham, H.P. (2015) Using threat maps  
488 for cost-effective prioritization of actions to conserve coastal habitats. *Marine Policy*, **61**, 95-102.

489 Giakoumi, S., Sini, M., Gerovasileiou, V., Mazor, T., Beher, J., Possingham, H.P., Abdulla, A., Çinar, M.E.,  
490 Dendrinou, P., Gucu, A.C., Karamanlidis, A.A., Rodic, P., Panayotidis, P., Taskin, E., Jaklin, A.,  
491 Voultsiadou, E., Webster, C., Zenetos, A. & Katsanevakis, S. (2013) Ecoregion-Based Conservation  
492 Planning in the Mediterranean: Dealing with Large-Scale Heterogeneity. *PLoS ONE*, **8**, e76449.

493 Green, S.J., Dulvy, N.K., Brooks, A.M.L., Akins, J.L., Cooper, A.B., Miller, S. & Côté, I.M. (2014) Linking  
494 removal targets to the ecological effects of invaders: a predictive model and field test. *Ecological  
495 Applications*, **24**, 1311-1322.

496 Guilhaumon, F., Albouy, C., Claudet, J., Velez, L., Ben Rais Lasram, F., Tomasini, J.-A., Douzery, E.J.P.,  
497 Meynard, C.N., Mouquet, N., Troussellier, M., Araújo, M.B. & Mouillot, D. (2015) Representing  
498 taxonomic, phylogenetic and functional diversity: new challenges for Mediterranean marine-  
499 protected areas. *Diversity and Distributions*, **21**, 175-187.

500 Jeschke, J.M. (2014) General hypotheses in invasion ecology. *Diversity and Distributions*, **20**, 1229-1234.

501 Kark, S., Levin, N., Grantham, H.S. & Possingham, H.P. (2009) Between-country collaboration and  
502 consideration of costs increase conservation planning efficiency in the Mediterranean Basin.  
503 *Proceedings of the National Academy of Sciences*, **106**, 15368-15373.

504 Katsanevakis, S., Deriu, I., D'Amico, F., Nunes, A.L., Pelaez Sanchez, S., Crocetta, F., Arianoutsou, M.,  
505 Bazos, I., Christopoulou, A., Curto, G., Delipetrou, P., Kokkoris, Y., Panov, V., Rabitsch, W.,  
506 Roques, A., Scalera, R., Shirley, S.M., Tricarico, E., Vannini, A., Zenetos, A., Zervou, S., Zikos, A.,  
507 Cardoso, A.C.C. (2015) European Alien Species Information Network (EASIN): supporting  
508 European policies and scientific research. *Management of Biological Invasions*, **6**, 147–157.

509 Katsanevakis, S., Tempera, F. & Teixeira, H. (2016) Mapping the impact of alien species on marine  
510 ecosystems: the Mediterranean Sea case study. *Diversity and Distributions*, n/a-n/a.

511 Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çinar, M.E., Oztürk, B., Grabowski, M.,  
512 Golani, D. & Cardoso, A.C. (2014) Impacts of invasive alien marine species on ecosystem services  
513 and biodiversity: a pan-European review. *Aquatic Invasions*, **9**, 391–423.

514 Klein, C.J., Tulloch, V.J., Halpern, B.S., Selkoe, K.A., Watts, M.E., Steinback, C., Scholz, A. &  
515 Possingham, H.P. (2013) Tradeoffs in marine reserve design: habitat condition, representation, and  
516 socioeconomic costs. *Conservation Letters*, **6**, 324-332.

517 Levine, J.M. (2008) Biological invasions. *Current Biology*, **18**, R57-R60.

518 Levine, J.M. & D'Antonio, C.M. (1999) Elton revisited: a review of evidence linking diversity and  
519 invasibility. *Oikos*, **87**, 15-26.

520 Mazor, T., Giakoumi, S., Kark, S. & Possingham, H.P. (2014) Large-scale conservation planning in a  
521 multinational marine environment: cost matters. *Ecological Applications*, **24**, 1115-1130.

522 McKinney, M.L. (2002) Influence of settlement time, human population, park shape and age, visitation and  
523 roads on the number of alien plant species in protected areas in the USA. *Diversity and*  
524 *Distributions*, **8**, 311-318.

525 Micheli, F., Halpern, B.S., Walbridge, S., Ciriaco, S., Ferretti, F., Fraschetti, S., Lewison, R., Nykjaer, L. &  
526 Rosenberg, A.A. (2013a) Cumulative Human Impacts on Mediterranean and Black Sea Marine  
527 Ecosystems: Assessing Current Pressures and Opportunities. *PLoS ONE*, **8**, e79889.

528 Micheli, F., Levin, N., Giakoumi, S., Katsanevakis, S., Abdulla, A., Coll, M., Frascetti, S., Kark, S.,  
529 Koutsoubas, D., Mackelworth, P., Maiorano, L. & Possingham, H.P. (2013b) Setting Priorities for  
530 Regional Conservation Planning in the Mediterranean Sea. *PLoS ONE*, **8**, e59038.

531 Moilanen, A., Kujala, H. & Leathwick, J.R. (2009) The Zonation framework and software for conservation  
532 prioritization. *Spatial Conservation Prioritization: Quantitative Methods and Computational Tools*  
533 (ed. by A. Moilanen, K.A. Wilson and H.P. Possingham), pp. 196-210. Oxford University Press  
534 Oxford, UK.

535 Mollo, E., Cimino, G. & Ghiselin, M.T. (2014) Alien biomolecules: a new challenge for natural product  
536 chemists. *Biological Invasions*, **17**, 941-950.

537 Molnar, J.L., Gamboa, R.L., Revenga, C. & Spalding, M.D. (2008) Assessing the global threat of invasive  
538 species to marine biodiversity. *Frontiers in Ecology and the Environment*, **6**, 485-492.

539 Mumby, P.J., Harborne, A.R. & Brumbaugh, D.R. (2011) Grouper as a Natural Biocontrol of Invasive  
540 Lionfish. *PLoS ONE*, **6**, e21510.

541 Myers, J.H., Simberloff, D., Kuris, A.M. & Carey, J.R. (2000) Eradication revisited: dealing with exotic  
542 species. *Trends in Ecology & Evolution*, **15**, 316-320.

543 Parravicini, V., Mangialajo, L., Mousseau, L., Peirano, A., Morri, C., Montefalcone, M., Francour, P.,  
544 Kulbicki, M. & Bianchi, C.N. (2015) Climate change and warm-water species at the north-western  
545 boundary of the Mediterranean Sea. *Marine Ecology*, **36**, 897-909.

546 Pressey, R.L., Watts, M.E., Barrett, T.W. & Ridges, M.J. (2009) The C-plan conservation planning system:  
547 origins, applications, and possible futures. *Spatial Conservation Prioritization: quantitative methods*  
548 *and computational tools* (ed. by A. Moilanen, W. K.A and H.P. Possingham), pp. 211-234. Oxford  
549 University Press, Oxford, UK.

550 Pyšek, P. & Richardson, D. (2010) Invasive species, environmental change and management, and health. .  
551 *Annual Review of Environment and Resources*, **35**, 25–55.

552 Schlaepfer, M.A., Sax, D.F. & Olden, J.D. (2011) The potential conservation value of non-native species.  
553 *Conserv Biol*, **25**, 428-37.

554 Seebens, H., Gastner, M.T. & Blasius, B. (2013) The risk of marine bioinvasion caused by global shipping.  
555 *Ecology Letters*, **16**, 782-790.

556 Simberloff, D., Martin, J.L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., Courchamp, F., Galil, B.,  
557 Garcia-Berthou, E., Pascal, M., Pysek, P., Sousa, R., Tabacchi, E. & Vila, M. (2013) Impacts of  
558 biological invasions: what's what and the way forward. *Trends Ecol Evol*, **28**, 58-66.

559 Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M., Halpern, B.S., Jorge,  
560 M.A., Lombana, A., Lourie, S.A., Martin, K.D., McManus, E., Molnar, J., Recchia, C.A. &  
561 Robertson, J. (2007) Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf  
562 Areas. *BioScience*, **57**, 573-583.

563 Stohlgren, T.J., Barnett, D., Flather, C., Fuller, P., Peterjohn, B., Kartesz, J. & Master, L.L. (2006) Species  
564 richness and patterns of invasion in plants, birds, and fishes in the United States. *Biological*  
565 *Invasions*, **8**, 427-447.

566 Strayer, D.L., Eviner, V.T., Jeschke, J.M. & Pace, M.L. (2006) Understanding the long-term effects of  
567 species invasions. *Trends in Ecology & Evolution*, **21**, 645-651.

568 Streftaris, N. & Zenetos, A. (2006) *Alien Marine Species in the Mediterranean - the 100 'Worst Invasives'*  
569 *and their Impact*.

570 Tallis, H., Ferdana, Z. & Gray, E. (2008) Linking terrestrial and marine conservation planning and threats  
571 analysis. *Conserv Biol*, **22**, 120-30.

572 Thresher, R.E. & Kuris, A.M. (2004) Options for Managing Invasive Marine Species. *Biological Invasions*,  
573 **6**, 295-300.

574 Verges, A., Steinberg, P.D., Hay, M.E., Poore, A.G., Campbell, A.H., Ballesteros, E., Heck, K.L., Jr., Booth,  
575 D.J., Coleman, M.A., Feary, D.A., Figueira, W., Langlois, T., Marzinelli, E.M., Mizerek, T.,  
576 Mumby, P.J., Nakamura, Y., Roughan, M., van Sebille, E., Gupta, A.S., Smale, D.A., Tomas, F.,  
577 Wernberg, T. & Wilson, S.K. (2014) The tropicalization of temperate marine ecosystems: climate-  
578 mediated changes in herbivory and community phase shifts. *Proc Biol Sci*, **281**, 20140846.

579 Vilà, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J.L., Pergl, J., Schaffner, U., Sun, Y. &  
580 Pyšek, P. (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on  
581 species, communities and ecosystems. *Ecology Letters*, **14**, 702-708.

582 Violle, C. & Jiang, L. (2009) Towards a trait-based quantification of species niche. *Journal of Plant Ecology*,  
583 **2**, 87-93.

584 Zenetos, A., Gofas, S., Morri, C., Rosso, A., Violanti, D., Garcia Raso, J.E., Cinar, M.E., Almogi-labin, A.,  
585 Ates, A.S., Azzurro, E., Ballesteros, E., Bianchi, C.N., Bilecenoglu, M., Gambi, M.C., Giangrande,  
586 A., Gravili, C., Hyams-kaphzan, O., Karachle, P.K., Katsanevakis, S., Lipej, I., Mastrototaro, F.,  
587 Mineur, F., Pancucci-papadopoulou, M.A., Ramos espla, A., Salas, C., San martin, G., Sfriso, A.,  
588 Streftaris, N. & Verlaque, M. (2012) *Alien species in the Mediterranean Sea by 2012. A contribution*  
589 *to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2.*  
590 *Introduction trends and pathways.*

591

## 592 **SUPPORTING INFORMATION**

593 Additional Supporting Information may be found in the online version of this article:

594 **Appendix S1.** List of 119 articles included in the synthesis.

595 **Table S1.** Articles' attributes included in the analyses.

596 **Figure S1.** Time trend in marine conservation plans and their scale.

597 **Appendix S2.** List of endemic fish included in the case study with their functional traits and IUCN category.

598

## 599 **DATA ACCESSIBILITY**

600 Critical habitats GIS layers (distribution of seagrass meadows *Posidonia oceanica* and coralligenous

601 formations) used in this paper are available on MedOBIS database: <http://lifewww->

602 [00.her.hcmr.gr:8080/medobis/resource.do?r=posidonia](http://lifewww-00.her.hcmr.gr:8080/medobis/resource.do?r=posidonia), <http://lifewww->

603 [00.her.hcmr.gr:8080/medobis/resource.do?r=coralligenous](http://lifewww-00.her.hcmr.gr:8080/medobis/resource.do?r=coralligenous). Endemic fish GIS layers are available on

604 Ecological Archives: <http://www.esapubs.org/archive/ecol/E096/203/#data>.

605

## 606 **BIOSKETCH**

607 The authors belong to a larger group of scientists focusing on Advancing Marine Conservation Planning in

608 the Mediterranean Sea. This group was formed in 2012 (<http://link.springer.com/article/10.1007/s11160-012->

609 [9272-8#](http://link.springer.com/article/10.1007/s11160-012-9272-8#/)/page-1 focusing) and its research interests include: marine conservation planning, integrated

610 conservation planning across realms (land-freshwater-sea), impact assessment on food webs and implications

611 for conservation planning, transboundary conservation, governance of marine protected areas, cost-effective  
612 action prioritization accounting for climate change and biological invasions.

613 S.G., S.K., S.Kark, F.G., J.C., and A.T. conceived the ideas; S.G. led the writing and all aspects of the  
614 project; all authors conducted the literature review and revised the text; S.G. and F.G. analysed the data;  
615 S.G., F.G., S.K., and S.F. produced the figures.

616  
617 **FIGURE LEGENDS**

618 **Figure 1.** Distribution of marine conservation plans across realms. The different realms (biogeographic  
619 regions) are presented with different colours, whereas conservation plans following: the ‘ignore’ approach is  
620 presented in red, the ‘protect’ in yellow, and the ‘avoid’ in blue. Realms are defined according to Spalding et  
621 al. (2007).

622 **Figure 2.** Conservation features accounted for in the conservation plans (frequency computed over a total of  
623 119 publications).

624 **Figure 3.** Critical habitats case study (data from Giakoumi et al. 2013). Difference in planning unit (12,828  
625 cells, 10 x 10 km) selection frequency, from Marxan outputs, when following the different approaches: a)  
626 ‘ignore’ vs ‘protect’, b) ‘ignore’ vs ‘avoid’, and c) ‘avoid’ vs ‘protect’. Planning units in red are those  
627 presenting higher selection frequency in the ‘ignore’ scenario, in orange those with higher selection in the  
628 ‘protect’ scenario, and in blue those with higher selection in the ‘avoid’ scenario. Planning units are black if  
629 they had maximum selection frequency (1000) in all three scenarios. Scatter plots show the selection  
630 frequency for the planning units under the different scenarios. For the maps we used ETRS89 Lambert  
631 Azimuthal Equal-Area projection.

632 **Figure 4.** Fish species case study (data from Guilhaumon et al. 2015). Difference in planning unit (12,828  
633 cells, 10 x 10 km) selection frequency, from Marxan outputs, when following the different approaches: a)  
634 ‘ignore’ vs ‘protect’, b) ‘ignore’ vs ‘avoid’, and c) ‘avoid’ vs ‘protect’. Planning units in red are those with a  
635 higher selection frequency in the ‘ignore’ scenario, in orange those with higher selection in the ‘protect’  
636 scenario, and in blue those with higher selection in the ‘avoid’ scenario. Planning units are black if they had  
637 maximum selection frequency (1000) in all three scenarios. Scatter plots show the selection frequency for the

638 planning units under the different scenarios. For the maps we used ETRS89 Lambert Azimuthal Equal-  
639 Area projection.