

1 **For:** *Biological Conservation* as a research report

2 **Recent increases in human pressure and forest loss threaten many**

3 **Natural World Heritage Sites**

4 **Author List:** James R. Allan^{ab}, Oscar Venter^c, Sean Maxwell^{ab}, Bastian Bertzky^{de}, Kendall Jones^{ab},

5 Yichuan Shi^{df} James E.M. Watson^{bg}

6 **Affiliations:**

7 ^aCentre of Excellence for Environmental Decisions, School of Biological Sciences, The University
8 of Queensland, St Lucia QLD 4072, Australia.

9 ^bSchool of Geography, Planning and Environmental Management, University of Queensland, St
10 Lucia QLD 4072, Australia

11 ^cEcosystem Science and Management Program, University of Northern British Columbia, Prince
12 George, v2n4z9, Canada

13 ^dInternational Union for Conservation of Nature (IUCN) Rue Mauverney 28, 1196 Gland,
14 Switzerland

15 ^eEuropean Commission, Joint Research Centre (JRC), Via Enrico Fermi 2749, 21027 Ispra (VA),
16 Italy

17 ^fUnited Nations Environment Programme World Conservation Monitoring Centre (UNEP-
18 WCMC), Cambridge UK

19 ^gWildlife Conservation Society, Global Conservation Program, Bronx, NY, USA, 10460

20

21 *Corresponding Author: James R. Allan. E-mail: James.allan@ugconnect.edu.au, phone +64 424

22 982 651. Address 1/201 Gladstone road, Highgate Hill, QLD, Australia.

23 **Highlights**

- 24 • Natural World Heritage Sites (NWHS) are designated because of their global
25 significance, yet there has been no systematic quantitative assessment of how humanity
26 is negatively affecting them.
- 27 • Increases in human pressure and forest loss are occurring across the vast majority of
28 forested NWHS.
- 29 • NWHS are becoming isolated by substantial increases in human pressure and forest loss
30 in the landscapes surrounding them.
- 31 • We demonstrate how globally comparable quantitative metrics can be used to help
32 monitor NWHS and provide crucial baseline information necessary for their long-term
33 preservation.

34 **Key Words**

35 World Heritage, Habitat loss, Habitat fragmentation, Human Footprint, Forest loss, Monitoring,
36 Cumulative threat mapping, Biodiversity conservation

37

38 **Abstract**

39 Natural World Heritage Sites (NWHS), via their formal designation through the United Nations,
40 are globally recognized as containing some of the Earth's most valuable natural assets.
41 Understanding changes in their ecological condition is essential for their ongoing preservation.
42 Here we use two newly available globally consistent data sets that assess changes in human
43 pressure (Human Footprint) and forest loss (Global Forest Watch) over time across the global
44 network of terrestrial NWHS. We show that human pressure has increased in 63% of NWHS
45 since 1993 and across all continents except Europe. The largest increases in pressure occurred
46 in Asian NWHS, many of which were substantially damaged such as *Manas Wildlife Sanctuary*
47 and *Simien National Park*. Forest loss occurred in 91% of NWHS that contain forests, with a
48 global mean loss of 1.5% per site since 2000, with the largest areas of forest lost occurring in
49 the Americas. For example *Wood Buffalo National Park* and *Río Plátano Biosphere Reserve* lost
50 2581km² (11.7%) and 365km² (8.5%) of their forest respectively. We found that on average
51 human pressure increased faster and more forest loss occurred in areas surrounding NWHS,
52 suggesting they are becoming increasingly isolated and are under threat from processes
53 occurring outside their borders. While some NWHS such as the *Sinharaja Forest Reserve* and
54 *Mana Pools National Park* showed minimal change in forest loss or human pressure, they are in
55 the minority and our results also suggest many NWHS are rapidly deteriorating and are more
56 threatened than previously thought.

57

58

59 **1. Introduction**

60 The World Heritage Convention was adopted in 1972 to ensure the world’s most valuable
61 natural and cultural resources could be conserved in perpetuity (UNESCO 1972). The
62 Convention aims to protect places with Outstanding Universal Value that transcend national
63 boundaries, and are worth conserving for humanity as a whole. These places are granted World
64 Heritage Status, the highest level of recognition afforded globally (UNESCO 2015). A unique
65 aspect of The Convention is that host nations are held accountable for the preservation of their
66 World Heritage Sites by the international community, and must report on their progress to the
67 United Nations Educational, Scientific and Cultural Organisation (UNESCO). Over 190 countries
68 are signatories to The Convention, committing to conserving the 1031 World Heritage Sites
69 listed at the time of this study (UNESCO 2015). Of these, 229 are Natural World Heritage Sites
70 (NWHS), inscribed for their unique natural beauty and biological importance, including many of
71 the world’s most important places for biodiversity conservation such as the *Pantanal*
72 *Conservation Area* in Brazil (UNESCO 2016a) and the iconic *Serengeti National Park* in Tanzania
73 (UNESCO 2016b).

74 As the number of NWHS has increased over the last few decades, so have the pressures
75 humanity is exerting on the natural environment (Rockstrom et al. 2009, Steffen et al. 2015,
76 Venter et al. 2016b). Anthropogenic habitat conversion due to human activities such as
77 agriculture and urbanisation are driving biodiversity extinction rates well above background
78 levels, and the condition of many ecosystems is in decline worldwide (Barnosky et al. 2012,
79 Hansen et al. 2013, Pimm et al. 2014, Watson et al. 2016). If significant human activity occurs

80 inside a NWHS it could potentially damage the ecological condition of that site and compromise
81 its Outstanding Universal Value, and is therefore incompatible with the objectives of the World
82 Heritage Convention (UNESCO 2015). If a site's condition and values are compromised it could
83 be placed on the list of World Heritage in Danger and, ultimately, its World Heritage Status can
84 be revoked if the ecological condition inside a site continues to decline to the extent it loses the
85 values that are the basis for its listing. The consequences for a host nation could be substantial,
86 since they would be denied access to the World Heritage Fund and other financial mechanisms,
87 technical support provided by UNESCO and the Advisory Bodies, and lose the sustainable
88 development opportunities a World Heritage Site creates (Conradin et al. 2014). Accurate and
89 transparent monitoring and reporting of both the human pressures facing NWHS, and the
90 ecological condition within NWHS is therefore essential for both host nations and UNESCO.

91 Current monitoring of NWHS is summarised in site-level reports and surveys. This
92 includes periodic reporting on progress and condition by States Parties on a 6-year regional
93 cycle, reactive monitoring led by UNESCO and the Advisory Bodies in response to current issues,
94 and site-level monitoring and evaluation systems (Hockings et al. 2006, Hockings et al. 2008,
95 Stolton et al. 2012). The IUCN's World Heritage Outlook initiative and its expert-driven
96 evaluations also provide important information on the conservation outlook for all NWHS
97 (Osipova et al. 2014). These monitoring approaches are important and capture diverse site-level
98 data, but do not include monitoring based on globally comparable quantitative datasets. We
99 argue that these current monitoring approaches could be further strengthened by additionally
100 using globally comparable datasets to assess increases in human pressure or changes in
101 ecological state such as forest loss (Leverington et al. 2010). Thanks to recent advances in

102 remote sensing technology, globally comparable data on human pressure and ecological state is
103 now available, allowing trends to be analysed across the entire network of NWHS for the first
104 time. This important baseline information allows States Parties to assess their progress in
105 preserving their NWHS and enables rapid reporting of their progress to the World Heritage
106 Committee.

107 In this study we quantify changes in spatial and temporal patterns of human pressure
108 and ecological state across the entire global network of NWHS and their surrounding
109 landscapes for the first time. We examine human pressure in NWHS in 1993 and 2009 using the
110 most comprehensive cumulative threat map available, the recently updated Human Footprint
111 (Venter et al. 2016b, Venter et al. 2016a) which is a temporally explicit map of eight
112 anthropogenic pressures on the terrestrial environment. An increasingly popular approach for
113 monitoring ecological state is to monitor forest cover, which responds to anthropogenic
114 pressures (Nagendra et al. 2013, Tracewski et al. 2016). Therefore we also examine patterns of
115 forest cover loss in NWHS between 2000 and 2012 using high resolution maps of global forest
116 cover (Hansen et al. 2013). We identify which NWHS have suffered the greatest forest loss, and
117 largest increases in human pressure, as well as sites which are performing well at limiting these
118 negative changes and maintaining their ecological integrity.

119 **2. Methods**

120 **2.1 World Heritage Site Data**

121 Data on NWHS location, boundary and year of inscription was obtained from the 2015 World
122 Database on Protected Areas (UNEP-WCMC 2015). We applied filtering criteria to identify

123 which NWHS qualified for our analysis. Out of all natural sites, sites inscribed only under
124 criterion (viii), which covers sites of geological importance including fossil sites and caves
125 (UNESCO 1972), were excluded from this analysis, with the exception of *Vredefort Dome* in
126 South Africa, *Phong Nha-Ke Bhang National Park* in Vietnam, *Lena Pillars Nature Park* in Russia
127 and *Ischigualasto/Talampaya Natural Parks* in Argentina, because they are part of larger
128 conservation areas. In addition, we constrained our analysis to terrestrial NWHS, and the
129 terrestrial component of marine NWHS. Due to the 1km² resolution of the Human Footprint
130 data, we chose to exclude NWHS smaller than 5km². Initially 190 NWHS qualified for our
131 analysis.

132 **2.2 Analyzing Human Pressure**

133 To measure human pressure on the natural environment we used the recently updated Human
134 Footprint (Venter et al. 2016a, Venter et al. 2016b), which is a globally-standardised measure of
135 cumulative human pressure on the terrestrial environment. The updated Human Footprint is
136 based on the original methodology developed by (Sanderson et al. 2002); however, the update
137 is temporally explicit, quantifying changes in human pressure over the period 1993 to 2009. At
138 a 1km² resolution, the Human Footprint includes global data on: built environments, crop lands,
139 pasture lands, population density, night lights, railways, major roadways and navigable
140 waterways. This makes the Human Footprint the most comprehensive cumulative threat map
141 available (McGowan 2016). Still, it is important to note that it does not include data on all the
142 possible threats and pressures facing NWHS. Other threats, including invasive species
143 (Bradshaw et al. 2007), overabundant species (Ndoro et al. 2015), wildlife poaching (Plumptre

144 et al. 2007, Wittemyer et al. 2014), tourism pressure (Li et al. 2008), and rapid climate change
145 (Scheffer et al. 2015), are not directly accounted for in the Human Footprint data. Although in
146 some cases the included pressure data, including population density, night lights, railways,
147 major roadways and navigable waterways, can contribute to these threats (e.g. invasive species
148 and some forms of poaching), we acknowledge that some threats are not well covered, which
149 makes this a conservative assessment of threats.

150 In the Human Footprint, individual pressures were placed within a 0 - 10 scale and
151 summed, giving a cumulative score of human pressure ranging from 0 - 50. A Human Footprint
152 score below 3 indicates land which is predominantly free of permanent infrastructure, but may
153 hold sparse human populations. A Human Footprint score of 4 is equal to pasture lands, and is a
154 reasonable threshold of when land can be considered “human dominated” and species are
155 likely to be threatened by habitat conversion (Watson et al. 2016). A Human Footprint score of
156 7 is equal to agriculture, above which a landscape will contain multiple pressures, for example
157 agriculture with roads and other associated infrastructure, and is therefore highly modified by
158 humans.

159 To compare mean changes in Human Footprint between NWHS and their surroundings,
160 we calculated the mean change in Human Footprint between 1993 and 2009 in NWHS and a
161 surrounding 10 km buffer zone. Calculating the Human Footprint in surrounding buffer zones
162 allows us to infer how much pressure a NWHS is under from developments surrounding the
163 protected area. Buffer zones were defined as a 10km buffer of land directly adjacent to and
164 surrounding each NWHS, and were created using the Geographic Information System ArcMap

165 version 10.2.1. Because NWHS inscribed post 1993 could potentially have been impacted
166 before their inscription as a NWHS, we included only sites inscribed during or before 1993 when
167 calculating the change in Human Footprint (n = 94).

168 **2.3 Analysing Forest Loss**

169 To assess forest loss, we followed Hansen et al. (2013), and defined forest cover as vegetation
170 taller than 5m and forest loss as the complete removal of tree canopy at a 30m resolution
171 (Hansen et al. 2013). Hansen forest-cover change data was extracted and processed in the
172 Google Earth Engine (<http://earthengine.google.org/>), a cloud platform for earth-observation
173 data analysis. Sites which had zero percent forest cover in 2000 were excluded from the
174 analysis. Only NWHS inscribed during or before 2000 were included in the forest loss analysis (n
175 = 134), since NWHS inscribed post 2000 could potentially have been impacted before
176 inscription. We then calculated total forest loss between the years 2000 and 2012 as a
177 percentage of forest extent in 2000 for all NWHS and buffer zones. We adapted JavaScript code
178 developed by Tracewski (2016) for analysing Hansen forest-cover data within specified spatial
179 zones, which is freely available online (<https://github.com/RSPB/IBA>). Gain in forest cover was
180 not included in this analysis for two reasons: young forests are unlikely to support forest-
181 dependant species, and much of the gain can be attributed to monoculture plantations of oil
182 palm or rubber which are major threats to tropical forests (Tropek et al. 2014). There are
183 limitations of satellite-derived estimates of global forest change, such as an inability to
184 differentiate between ecologically valuable forest and agro-forests, such as oil palm, and lower
185 accuracy in more arid environments (Hansen et al. 2013, Achard et al. 2014, Tropek et al. 2014).

186 Likewise, ground truthing is required to infer the causes of forest loss since the dataset does
187 not differentiate between ecologically harmful clearing, and purposeful clearing for example of
188 invasive species, which has a conservation benefit. But even with these limitations, the Hansen
189 et al. (2013) forest data product is considered the most accurate global representation of
190 temporal loss of forest available (McRoberts et al. 2016).

191 **3. Results**

192 **3.1 Human Pressure**

193 **3.1.1 Human Pressure in NWHS**

194 The average Human Footprint per NWHS in 2009 is 6.4, which is higher than the global average
195 Human Footprint of 5.6, and there was considerable variation between regions and individual
196 sites. Out of 94 NWHS considered in this analysis, the majority of them (63%, n=59) had an
197 average Human Footprint ≥ 4 , and many NWHS (38%, n=36) had a Human Footprint ≥ 7
198 meaning they are highly modified by humans. *Keoladeo National Park* in India was subject to
199 the highest levels of human pressure of any NWHS, with a 2009 Human Footprint of 23.
200 *Göreme National Park* in Turkey, *Mount Taishan* in China, and *Manas Wildlife Sanctuary* in India
201 were also subject to some of the highest levels of human pressure, with a Human Footprint of
202 19, 17 and 17 respectively. European and Asian NWHS were under the highest levels of human
203 pressure of all the continents, whereas NWHS in North America and Oceania are under the
204 lowest (Table 1.). *Nahanni National Park* in Canada had the lowest 2009 Human Footprint of
205 0.08, along with *Kluane/ Wrangel-St. Elias/ Glacier Bay/ Tatshenshini-Alsek* in Canada/USA (0.3)
206 and *Aïr and Ténéré Natural Reserves* in Niger (0.4). These three NWHS are essentially free of

207 human pressure but no NWHS had a Human Footprint of zero (see supplementary Table A1 for
208 a full list of NWHS and their Human Footprint scores).

209 **3.1.2 Changes in Human Pressure in NWHS over time**

210 The Human Footprint in NWHS increased far more slowly than the global average, rising 1.7%
211 between 1993 and 2009, compared to the global increase of 9%. However, human pressure did
212 increase in the majority of NWHS (63% n = 58) and across all continents except Europe (Figure
213 1.). In most cases the increases were small; however, 14 sites (15%) were subject to substantial
214 increases in human pressure (average Human Footprint increase > 1) (Table 2.). The *Manas*
215 *Wildlife Sanctuary* in India underwent the largest increase in human pressure of any NWHS,
216 with its Human Footprint rising by 5 to a score of 17 and is now one of the most highly modified
217 by humans. *Komodo National Park* in Indonesia also underwent one of the largest increases in
218 human pressure with its Human Footprint rising by 4.

219 The largest increases in human pressure occurred in Asian NWHS, where the regional mean
220 Human Footprint increased by 8% between 1993 and 2009 (Figure 2.). NWHS in Oceania and
221 South America also underwent relatively large increases in human pressure, with their mean
222 Human Footprints rising by 6.8% and 4.3% respectively. The Human Footprint in European
223 NWHS decreased by 10% during the time period, however they were highly modified NWHS to
224 begin with and thus still face the highest levels of human pressure of all continents. Some
225 notable decreases occurred in the *Sinharaja Forest Reserve* in Sri Lanka, *Hierapolis-Pamukkale*
226 and *Göreme National Park* in Turkey, whose Human Footprint decreased by 7, 6.5 and 4
227 respectively.

228 3.1.3 Comparison with Buffer Zones

229 The 2009 average Human Footprint per buffer zone is 7.8, which is slightly higher than the
230 average Human Footprint per NWHS of 6.4. The trend of human pressure being higher in the
231 landscapes surrounding NWHS held across all continents and for the majority of NWHS (78%
232 n=70). European and Asian NWHS had the greatest levels of human pressure in their buffer
233 zones, which were considerably higher than the global average. The *Danube Delta* in Romania
234 had the greatest difference in human pressure compared to its buffer zone, with the relatively
235 low 2009 average Human Footprint of 4.5 inside the NWHS compared to a relatively high 13.9
236 in its buffer zone. Interestingly, some NWHS such as *Sagarmatha National Park* in Nepal had
237 very high levels of human pressure inside their boundaries compared to their buffer zones, with
238 2009 average Human Footprint scores of 6.5 and 3.7 respectively.

239 Globally, the average Human Footprint in buffer zones increased much faster than inside
240 NWHS, rising by 4.5% compared to 1.7% between 1993 and 2009. These increases were largest
241 in buffer zones in South America and Australia where the Human Footprint increased by 16%
242 and 11% respectively. Many NWHS performed well at limiting increases in human pressure
243 relative to the amount of pressure they are under from the surrounding landscape. For example
244 in *Iguaçu National Park* in Brazil the Human Footprint stayed almost constant within the NWHS
245 between 1993 and 2009, increasing by 0.2 compared to a large increase of 4.5 in its buffer
246 zone. Likewise in *Mount Taishan* in China the Human Footprint only increased by 1.1 inside the
247 NWHS but by 3.3 in its buffer zone. Conversely, some NWHS underwent larger increases in
248 human pressure within their borders than in their buffer zones. These include *Manas Wildlife*

249 *Sanctuary* in India where the Human Footprint inside the NWHS increased by 5.3 compared to
250 2.2 in the buffer zone, and *Simien National Park* in Ethiopia where the Human Footprint inside
251 the NWHS increased by 2.9, compared to 2.2 in its buffer zone.

252 **3.2 Forest Cover Loss**

253 **3.2.1 Forest Loss in NWHS**

254 Forest loss occurred in the majority of forested NWHS (91%, n=122) with a mean percentage
255 loss of 1.48% per NWHS (Figure 4.). In the year 2000 there was 433,173 km² of forest cover
256 inside all NWHS and by the end of 2012 the total area of forest cover lost was 7,271 km²
257 (1.67%). The majority of NWHS suffered low levels of forest loss, with 72% (n=97) of NWHS
258 losing < 1%. However, 8% (n=11) of NWHS suffered substantial forest loss (>5%), the majority of
259 which are North American NWHS (Figure 5.). North American NWHS accounted for 57% of all
260 the forest lost in NWHS globally (Table 3.). *Waterton Glacier International Peace Park* that
261 crosses the Canadian and USA border lost almost one quarter of its forested area (23%,
262 540km²), *Wood Buffalo National Park* in Canada lost 12% (2,582km²) of forest cover, and
263 *Yellowstone National Park* in the USA lost 6% (217km²)(Table 4.). *Río Plátano Biosphere Reserve*
264 in Honduras and *Lake Baikal* in Russia also lost large proportions of forest cover, 8% (365km²)
265 and 5% (1332km²) respectively (see supplementary Table A2 for a full list of NWHS and forest
266 loss statistics). After North America, Asian and South American NWHS lost the largest areas of
267 forest within their NWHS. NWHS in Oceania lost an above average percentage of their forested
268 area.

269 **3.2.2 Forest Loss in Buffer Zones**

270 Forest loss was higher in the buffer zones surrounding NWHS than in the sites themselves with
271 a mean percentage loss of 2.9% per NWHS buffer zone. This trend held for all continents except
272 for North America, where forest loss in the buffer zones was at very similar levels to inside
273 NWHS. NWHS in Oceania lost the highest percentage of forest cover in their buffer zones and
274 European NWHS the least. There was a clear increase in the number of NWHS suffering
275 substantial forest losses of > 5% in their buffer zones (19% n=25), compared to within their
276 boundaries. Forest loss was low (<1%) in only half of the NWHS buffer zones (48% n=58), while
277 72% of NWHS (n = 97) had low rates within their borders. Some notable NWHS which lost large
278 proportions of forest in their buffer zones are the *Australian Fossil Mammal Sites (Riversleigh /*
279 *Naracoorte)* which lost 33% (9km²), *The Discovery Coast Atlantic Forests* in Brazil which lost 11%
280 (192 km²), and *Kinabalu Park* in Malaysia which lost 10% (150 km²). Many NWHS performed
281 well at limiting forest loss within their borders, despite considerable losses in their buffer zones
282 (Figure 6). *Mount Wuyi* in China, for example lost only 1% (7km²) within its borders compared
283 to 9% (122 km²) in its buffer zone. And *Iguazu National Park* in Argentina lost almost no forest
284 inside its borders (0.02% <1km²) compared to extensive loss in its buffer zone (13% 110km²).

285 **4. Discussion**

286 Our analysis is the first globally comparable quantitative assessment of changes in human
287 pressure and ecological state across the entire network of NWHS, which is important baseline
288 information for the UNESCO World Heritage Convention, the IUCN as the advisory body to
289 UNESCO for Natural World Heritage, and the States Parties to monitor their progress at
290 conserving NWHS. We found that human pressure is increasing and forest loss is occurring in
291 the majority of forested NWHS worldwide, threatening to undermine their Outstanding

292 Universal Value. Our most concerning finding is that a number of NWHS are severely
293 threatened by large increases in human footprint (>1) (14 NWHS = 15% of the 94 NWHS
294 analyzed), and extensive forest loss (>5%) (11 NWHS = 8% of the 134 NWHS analyzed). The
295 negative impact occurring in these sites requires large scale conservation interventions to
296 ensure their value remains protected and sustained in the future. Our findings support
297 qualitative assessments from case-by-case reports, which corroborates that NWHS are
298 becoming increasingly threatened globally, and that the condition of a third of NWHS is now of
299 significant concern (Osipova et al. 2014, Wang et al. 2014). Our results also support other
300 studies showing that habitat extent and condition are declining in many protected areas across
301 the globe (Laurance et al. 2012, Geldmann et al. 2014). However our findings are particularly
302 concerning since NWHS are flagship protected areas afforded the highest level of international
303 protection.

304 There have been alarming rates of forest loss in the buffer zones surrounding nationally
305 designated protected areas over the last three decades (DeFries et al. 2005, Bailey et al. 2016,
306 Lui and Coomes 2016), and our results confirm this is also the case for many NWHS. We found
307 that forest loss and increases in human pressure were considerably higher in the buffer zones
308 surrounding the vast majority of NWHS. This suggests that NWHS may be performing well at
309 limiting negative changes within their boundaries (Bruner et al. 2001). However our findings
310 clearly show that NWHS are becoming increasingly isolated which is concerning since the
311 ecological integrity of many NWHS depend on links with the broader landscape (Naughton-
312 Treves et al. 2005, Kormos et al. 2015). Environmental degradation around NWHS could
313 decrease their area and increase edge effects, which are important determinants of biodiversity

314 persistence (Woodroffe and Ginsberg 1998, Hansen and DeFries 2007, Newmark 2008).
315 Furthermore, Laurance et al. (2012) found that degradation occurring around a protected area
316 strongly predisposes it to similar degradation within its borders, including trends in forest loss
317 and human pressure. To avert further damage to NWHS the World Heritage Committee should
318 consider directing more resources to conservation in the landscapes surrounding NWHS, and
319 continue designating and strengthening official buffer zones around NWHS, where communities
320 are engaged and low impact land uses promoted (Laurance et al. 2012, Kormos et al. 2015,
321 UNESCO 2015, Weisse and Naughton-Treves 2016).

322 We found that North American NWHS suffered such high levels of forest loss, despite
323 their protection and management being considered highly effective (Osipova et al. 2014). This
324 forest loss is almost certainly due to the largest pine beetle outbreaks on record, which are
325 causing widespread forest mortality, leaving dead trees prone to fires across large areas of
326 North America and causing substantial ecological damage (Westerling et al. 2006, MacFarlane
327 et al. 2013, True et al. 2014). This process is semi-natural; however, pine beetle outbreaks are
328 being assisted by anthropogenic climate change, because winters are no longer cold enough or
329 long enough to kill the beetles and reduce their numbers substantially (Westerling et al. 2006,
330 Raffa et al. 2008). Pine beetle outbreaks are proving incredibly difficult to manage, making
331 North American NWHS some of the most threatened worldwide with regard to forest loss.
332 While pine beetle outbreaks may explain forest loss in North America, globally the drivers and
333 mechanisms of forest loss in NWHS are diverse. For example, NWHS in Central America also lost
334 some of the largest areas of forest, which can be directly attributed to direct deforestation
335 activities undertaken by humans. Illegal drug trafficking in the *Río Plátano Biosphere Reserve* in

336 Honduras led to insecurity and instability, allowing widespread illegal deforestation and illegal
337 settlement to occur. Our findings show that *Río Plátano* lost 8% (365km²) of its forested area
338 since 1993 and had an above average increase in Human Footprint, supporting the World
339 Heritage Committee's decision in 2011 to inscribe it on the List of World Heritage in Danger.

340 We found that one third of NWHS underwent a decrease in human pressure, which is a
341 good result for conservation and a benchmark for other NWHS and protected areas to strive
342 towards. The Human Footprint decreased on average across European NWHS, which is also
343 encouraging, however we suggest that decreases in the Human Footprint should be interpreted
344 with care. Although the Human Footprint is the most comprehensive cumulative threat map
345 available, it does not include data on all the possible threats and pressures facing NWHS,
346 suggesting our results are conservative, and that NWHS may be even more threatened than we
347 have demonstrated. For example in *Aïr and Ténéré National Park* in Niger we found that
348 changes in the Human Footprint were minimal (0.1) but understand that political instability and
349 civil strife, along with poaching are the main pressures threatening the park (UNESCO 2016c).
350 These limitations can be largely overcome by combining our data with site level case-by-case
351 reports and therefore our study complements statutory monitoring mechanisms under the
352 World Heritage Convention (UNESCO 2015) and IUCN's World Heritage Outlook initiative
353 (Osipova et al. 2014). As discussed in the methods section, there are also limitations with
354 satellite derived estimates of global forest change, for example it is impossible to infer the
355 causes of forest loss without the use of site-level data, and not all forest loss in NWHS is
356 necessarily negative. For example, *iSimangaliso Wetland Park* in South Africa lost 18% (161km²)
357 of the forest in its buffer zone, but this is due to the purposeful clearing of pine and eucalyptus

358 plantations for restoration (Zaloumis and Bond 2011), so clearly serves a positive conservation
359 purpose. However, given the impacts of habitat loss on biodiversity (Maxwell et al. 2016) and
360 the prevalence of forest loss in protected areas globally (Heino et al. 2015), we do assume in
361 the majority of cases that forest loss is detrimental to the ecological state of NWHS. We also
362 note that forest loss is also just one indicator of ecological state, and a measure of intact forest
363 cover does not necessarily guarantee a NWHS is in good condition. For example the *Dja Faunal*
364 *Reserve* in Cameroon lost almost no forest during the time period; however, it has suffered
365 intense poaching in recent times threatening wildlife populations within its borders (UNESCO
366 2016d). The limitations of remotely sensed data are widely recognized and need to be
367 acknowledged, yet it remains an increasingly important tool for conservation monitoring, and
368 its overall utility is broadly acknowledged (Turner et al. 2003, Buchanan et al. 2009, Tracewski
369 et al. 2016).

370 **5. Conclusion**

371 The World Heritage Convention should be one of the world's most effective conservation
372 instruments globally, identifying and protecting the Earth's most valuable natural landscapes.
373 Our aim is to highlight growing challenges which are undermining its success. New globally
374 comparable data sets such as the Human Footprint and the Global Forest Change data have
375 provided an urgently needed opportunity to measure how well NWHS are maintaining their
376 ecological integrity (Watson et al. 2015). We used these metrics to analyse spatial and temporal
377 trends in human pressure for 94 NWHS, and forest loss in 134 NWHS, presenting baseline data
378 for the World Heritage Committee and the States Parties. There is a clear opportunity for the

379 World Heritage Committee to establish thresholds and targets with regard to human pressure
380 and forest loss in NWHS, and measure the effectiveness of management interventions across
381 sites. We urge the World Heritage Committee to assess the status of the NWHS which our
382 analysis suggests are highly threatened, since urgent conservation intervention is now clearly
383 needed to save many of these NWHS and their outstanding and unique values in perpetuity.

384 **6. Acknowledgements**

385 We thank Joseph Maina, Moreno Di Marco, Marc Hockings, Hugh Possingham and Cyril Kormos
386 for their valuable discussions about this idea over the past two years, and Tim Badman, Cyril
387 Kormos and Remco van Merm for valuable comments on the manuscript. We also thank two
388 anonymous reviewers for their helpful comments and feedback. The contents of this article do
389 not necessarily reflect the views or policies of IUCN or other contributory organizations.

390 **References**

- 391 Achard, F., R. Beuchle, P. Mayaux, H.-J. Stibig, C. Bodart, A. Brink, S. Carboni, B. Desclée, F. Donnay, H. D.
392 Eva, A. Lupi, R. Raši, R. Seliger, and D. Simonetti. 2014. Determination of tropical deforestation
393 rates and related carbon losses from 1990 to 2010. *Global Change Biology* **20**:2540-2554.
- 394 Bailey, K. M., R. A. McCleery, M. W. Binford, and C. Zweig. 2016. Land-cover change within and around
395 protected areas in a biodiversity hotspot. *Journal of Land Use Science* **11**:154-176.
- 396 Barnosky, A. D., E. A. Hadly, J. Bascompte, E. L. Berlow, J. H. Brown, M. Fortelius, W. M. Getz, J. Harte, A.
397 Hastings, P. A. Marquet, N. D. Martinez, A. Mooers, P. Roopnarine, G. Vermeij, J. W. Williams, R.
398 Gillespie, J. Kitzes, C. Marshall, N. Matzke, D. P. Mindell, E. Revilla, and A. B. Smith. 2012.
399 Approaching a state shift in Earth/'s biosphere. *Nature* **486**:52-58.
- 400 Bradshaw, C. J. A., I. C. Field, D. M. J. S. Bowman, C. Haynes, and B. W. Brook. 2007. Current and future
401 threats from non-indigenous animal species in northern Australia: a spotlight on World Heritage
402 Area Kakadu National Park. *Wildlife Research* **34**:419 - 436.
- 403 Bruner, A. G., R. E. Gullison, R. E. Rice, and G. A. B. da Fonseca. 2001. Effectiveness of parks in protecting
404 tropical biodiversity. *Science* **291**:125-128.
- 405 Buchanan, G. M., A. Nelson, P. Mayaux, A. Hartley, and P. F. Donald. 2009. Delivering a Global,
406 Terrestrial, Biodiversity Observation System through Remote Sensing. *Conservation Biology*
407 **23**:499-502.
- 408 Conradin, K., M. Engesser, and U. Wiesmann. 2014. Four decades of World Natural Heritage - how
409 changing protected area values influence the UNESCO label. *DIE ERDE* **146**:34 - 46.

410 DeFries, R., A. Hansen, A. C. Newton, and M. C. Hansen. 2005. Increasing Isolation of Protected Areas in
411 Tropical Forests over the Past Twenty Years. *Ecological Applications* **15**:19-26.

412 Geldmann, J., L. N. Joppa, and N. D. Burgess. 2014. Mapping Change in Human Pressure Globally on Land
413 and within Protected Areas. *Conservation Biology* **28**:1604-1616.

414 Hansen, A. J. and R. DeFries. 2007. Ecological Mechanisms Linking Protected Areas To Surrounding
415 Lands. *Ecological Applications* **17**:974-988.

416 Hansen, M. C., P. V. Potapov, R. Moore, M. Hancher, S. A. Turubanova, A. Tyukavina, D. Thau, S. V.
417 Stehman, S. J. Goetz, T. R. Loveland, A. Kommareddy, A. Egorov, L. Chini, C. O. Justice, and J. R.
418 G. Townshend. 2013. High-Resolution Maps of 21st-Century Forest Cover Change. *Science*
419 **342**:850 - 853.

420 Heino, M., M. Kumm, M. Makkonen, M. Mulligan, P. H. Verburg, M. Jalava, and T. A. Räsänen. 2015.
421 Forest Loss in Protected Areas and Intact Forest Landscapes: A Global Analysis. *PLoS ONE*
422 **10**:e0138918.

423 Hockings, M., R. James, S. Stolton, N. Dudley, V. Mathur, J. Makombo, J. Courrau, and J. Parrish. 2008.
424 Enhancing Our Heritage Toolkit. Assessing Management Effectiveness of Natural World Heritage
425 Sites., UNESCO, Paris, France.

426 Hockings, M., S. Stolton, F. Leverington, N. Dudley, and J. Courrau. 2006. Evaluating Effectiveness. A
427 framework for assessing management of Protected Areas. Best Practice Protected Area
428 Guidelines. IUCN, Gland, Switzerland and Cambridge, UK.

429 Kormos, C. F., B. Bertzky, T. Jaeger, Y. Shi, T. Badman, J. A. Hilty, B. G. Mackey, R. A. Mittermeier, H.
430 Locke, E. Osipova, and J. E. M. Watson. 2015. A Wilderness Approach under the World Heritage
431 Convention. *Conservation Letters* **9**:228-235.

432 Laurance, W. F. and D. C. Useche and J. Rendeiro and M. Kalka and C. J. Bradshaw and S. P. Sloan and S.
433 G. Laurance and M. Campbell and K. Abernethy and P. Alvarez and V. Arroyo-Rodriguez and P.
434 Ashton and J. Benitez-Malvido and A. Blom and K. S. Bobo and C. H. Cannon and M. Cao and R.
435 Carroll and C. Chapman and R. Coates and M. Cords and F. Danielsen and B. De Dijn and E.
436 Dinerstein and M. A. Donnelly and D. Edwards and F. Edwards and N. Farwig and P. Fashing and
437 P. M. Forget and M. Foster and G. Gale and D. Harris and R. Harrison and J. Hart and S. Karpanty
438 and W. J. Kress and J. Krishnaswamy and W. Logsdon and J. Lovett and W. Magnusson and F.
439 Maisels and A. R. Marshall and D. McClearn and D. Mudappa and M. R. Nielsen and R. Pearson
440 and N. Pitman and J. van der Ploeg and A. Plumptre and J. Poulsen and M. Quesada and H.
441 Rainey and D. Robinson and C. Roetgers and F. Rovero and F. Scatena and C. Schulze and D. Sheil
442 and T. Struhsaker and J. Terborgh and D. Thomas and R. Timm and J. N. Urbina-Cardona and K.
443 Vasudevan and S. J. Wright and G. J. Arias and L. Arroyo and M. Ashton and P. Auzel and D.
444 Babaasa and F. Babweteera and P. Baker and O. Banki and M. Bass and I. Bila-Isia and S. Blake
445 and W. Brockelman and N. Brokaw and C. A. Bruhl and S. Bunyavejchewin and J. T. Chao and J.
446 Chave and R. Chellam and C. J. Clark and J. Clavijo and R. Congdon and R. Corlett and H. S.
447 Dattaraja and C. Dave and G. Davies and M. Beisiegel Bde and N. da Silva Rde and A. Di Fiore and
448 A. Diesmos and R. Dirzo and D. Doran-Sheehy and M. Eaton and L. Emmons and A. Estrada and
449 C. Ewango and L. Fedigan and F. Feer and B. Fruth and J. G. Willis and U. Goodale and S.
450 Goodman and J. C. Guix and P. Guthiga and W. Haber and K. Hamer and I. Herbinger and J. Hill
451 and Z. Huang and I. F. Sun and K. Ickes and A. Itoh and N. Ivanauskas and B. Jackes and J. Janovec
452 and D. Janzen and M. Jiangming and C. Jin and T. Jones and H. Justiniano and E. Kalko and A.
453 Kasangaki and T. Killeen and H. B. King and E. Klop and C. Knott and I. Kone and E. Kudavidanage
454 and J. L. Ribeiro and J. Lattke and R. Laval and R. Lawton and M. Leal and M. Leighton and M.
455 Lentino and C. Leonel and J. Lindsell and L. Ling-Ling and K. E. Linsenmair and E. Losos and A.
456 Lugo and J. Lwanga and A. L. Mack and M. Martins and W. S. McGraw and R. McNab and L.
457 Montag and J. M. Thompson and J. Nabe-Nielsen and M. Nakagawa and S. Nepal and M.

458 Norconk and V. Novotny and S. O'Donnell and M. Opiang and P. Ouboter and K. Parker and N.
459 Parthasarathy and K. Pisciotta and D. Prawiradilaga and C. Pringle and S. Rajathurai and U.
460 Reichard and G. Reinartz and K. Renton and G. Reynolds and V. Reynolds and E. Riley and M. O.
461 Rodel and J. Rothman and P. Round and S. Sakai and T. Sanaiotti and T. Savini and G. Schaab and
462 J. Seidensticker and A. Siaka and M. R. Silman and T. B. Smith and S. S. de Almeida and N. Sodhi
463 and C. Stanford and K. Stewart and E. Stokes and K. E. Stoner and R. Sukumar and M. Surbeck
464 and M. Tobler and T. Tschardt and A. Turkalo and G. Umapathy and M. van Weerd and J. V.
465 Rivera and M. Venkataraman and L. Venn and C. Vereza and C. V. de Castilho and M. Waltert and
466 B. Wang and D. Watts and W. Weber and P. West and D. Whitacre and K. Whitney and D. Wilkie
467 and S. Williams and D. D. Wright and P. Wright and L. Xiankai and P. Yonzon and F. Zamzani.
468 2012. Averting biodiversity collapse in tropical forest protected areas. *Nature* **489**:290-294.
469 Leverington, F., K. L. Costa, H. Pavese, A. Lisle, and M. Hockings. 2010. A Global Analysis of Protected
470 Area Management Effectiveness. *Environmental Management* **46**:685-698.
471 Li, M., B. Wu, and L. Cai. 2008. Tourism development of World Heritage Sites in China: A geographic
472 perspective. *Tourism Management* **29**:308-319.
473 Lui, G. V. and D. A. Coomes. 2016. Tropical nature reserves are losing their buffer zones, but leakage is
474 not to blame. *Environmental Research* **147**:580-589.
475 MacFarlane, W. W., J. A. Logan, and W. R. Kern. 2013. An innovative aerial assessment of Greater
476 Yellowstone Ecosystem mountain pine beetle-caused whitebark pine mortality. *Ecological*
477 *Applications* **23**:421-437.
478 Maxwell, S. L., R. F. Fuller, T. M. Brooks, and J. E. M. Watson. 2016. Biodiversity: The ravages of guns
479 nets and bulldozers. *Nature* **536**:143 - 145.
480 McGowan, P. J. K. 2016. Conservation: Mapping the terrestrial human footprint. *Nature* **537**:172-173.
481 McRoberts, R. E., A. C. Vibrans, C. Sannier, E. Naesset, M. C. Hansen, B. F. Walters, and D. V. Lingner.
482 2016. Methods for evaluating the utilities of local and global maps for increasing the precision of
483 estimates of subtropical forest area. *Canadian Journal of Forest Research* **46**:924-932.
484 Nagendra, H., R. Lucas, J. P. Honrado, R. H. G. Jongman, C. Tarantino, M. Adamo, and P. Mairota. 2013.
485 Remote sensing for conservation monitoring: Assessing protected areas, habitat extent, habitat
486 condition, species diversity, and threats. *Ecological Indicators* **33**:45-59.
487 Naughton-Treves, L., M. B. Holland, and K. Brandon. 2005. The Role of Protected Areas in Conserving
488 Biodiversity and Sustaining Local Livelihoods. *Annual Review of Environment and Resources*
489 **30**:219-252.
490 Ndoró, O., C. Mashapa, S. Kativu, and E. Gandiwa. 2015. A comparative assessment of baobab density in
491 northern Mana Pools National Park, Zimbabwe. *African Journal of Ecology* **53**:109-111.
492 Newmark, W. D. 2008. Isolation of African protected areas. *Frontiers in Ecology and the Environment*
493 **6**:321-328.
494 Osipova, E., Y. Shi, C. Kormos, P. Shadie, C. Zwahlen, and T. Badman. 2014. IUCN World Heritage Outlook
495 2014: A conservation assessment of all natural world heritage sites. IUCN, Gland, Switzerland.
496 Pimm, S. L., C. N. Jenkins, R. Abell, T. M. Brooks, J. L. Gittleman, L. N. Joppa, P. H. Raven, C. M. Roberts,
497 and J. O. Sexton. 2014. The biodiversity of species and their rates of extinction, distribution, and
498 protection. *Science* **344**: DOI: 10.1126/science.1246752
499 Plumptre, A., D. Kujirakwinja, A. Treves, I. Owiunji, and H. Rainer. 2007. Transboundary conservation in
500 the greater Virunga landscape: Its importance for landscape species. *Biological Conservation*
501 **134**:279-287.
502 Raffa, K. F., B. H. Aukema, B. J. Bentz, A. L. Carroll, J. A. Hicke, M. G. Turner, and W. H. Romme. 2008.
503 Cross-scale Drivers of Natural Disturbances Prone to Anthropogenic Amplification: The Dynamics
504 of Bark Beetle Eruptions. *BioScience* **58**:501 - 517.

505 Rockstrom, J., W. Steffen, K. Noone, A. Persson, F. S. Chapin, E. F. Lambin, T. M. Lenton, M. Scheffer, C.
506 Folke, H. J. Schellnhuber, B. Nykvist, C. A. de Wit, T. Hughes, S. van der Leeuw, H. Rodhe, S.
507 Sorlin, P. K. Snyder, R. Costanza, U. Svedin, M. Falkenmark, L. Karlberg, R. W. Corell, V. J. Fabry, J.
508 Hansen, B. Walker, D. Liverman, K. Richardson, P. Crutzen, and J. A. Foley. 2009. A safe operating
509 space for humanity. *Nature* **461**:472-475.

510 Sanderson, E. W., M. Jaith, M. A. Levy, K. H. Redford, A. V. Wannebo, and G. Woolmer. 2002. The human
511 footprint and the last of the wild. *BioScience* **52**:891 - 903.

512 Scheffer, M., S. Barrett, S. R. Carpenter, C. Folke, A. J. Green, M. Holmgren, T. P. Hughes, S. Kosten, I. A.
513 van de Leemput, D. C. Nepstad, E. H. van Nes, E. T. H. M. Peeters, and B. Walker. 2015. Creating
514 a safe operating space for iconic ecosystems. *Science* **347**:1317-1319.

515 Steffen, W., K. Richardson, J. Rockstrom, S. E. Cornell, I. Fetzer, E. M. Bennett, R. Biggs, S. R. Carpenter,
516 W. de Vries, C. A. de Wit, C. Folke, D. Gerten, J. Heinke, G. M. Mace, L. M. Persson, V.
517 Ramanathan, B. Meyers, and S. Sorlin. 2015. Planetary boundaries: Guiding human development
518 on a changing planet. *Science* **347**:1259855-1259855.

519 Stolton, S., N. Dudley, and P. Shadie. 2012. *Managing Natural World Heritage*. UNESCO, Paris, France.

520 Tracewski, Ł., S. H. M. Butchart, P. F. Donald, M. Evans, L. D. C. Fishpool, and B. Graeme M. 2016.
521 Patterns of twenty-first century forest loss across a global network of important sites for
522 biodiversity. *Remote Sensing in Ecology and Conservation* **2**:37-44

523 Tropek, R., O. Sedláček, J. Beck, P. Keil, Z. Musilová, I. Šimová, and D. Storch. 2014. Comment on “High-
524 resolution global maps of 21st-century forest cover change”. *Science* **344**:981-981.

525 True, R., J. Karst, and M. Randall. 2014. Decline of ectomycorrhizal fungi following a mountain pine
526 beetle epidemic. *Ecology* **95**:1096-1103.

527 Turner, W., S. Spector, N. Gardiner, M. Fladeland, E. Sterling, and M. Steininger. 2003. Remote sensing
528 for biodiversity science and conservation. *Trends in Ecology & Evolution* **18**:306-314.

529 UNEP-WCMC. 2015. World Database on Protected Areas (WDPA) www.protectedplanet.net.

530 UNESCO. 1972. *Convention Concerning the Protection of the World Cultural and Natural Heritage*.

531 UNESCO. 2015. *Operational guidelines for the implementation of the World Heritage Convention*.
532 UNESCO.

533 UNESCO. 2016a. Pantanal Conservation Area. World Heritage List, <http://whc.unesco.org/en/list/999>.

534 UNESCO. 2016b. Serengeti National Park. World Heritage List, <http://whc.unesco.org/en/list/156>.

535 UNESCO. 2016c. State of Conservation Report: Air and Tenere National Park, Niger.
536 <http://whc.unesco.org/en/soc/3372>.

537 UNESCO. 2016d. State of Conservation Report: Dja Faunal Reserve, Cameroon.
538 <http://whc.unesco.org/en/soc/3454>.

539 Venter, O., E. W. Sanderson, A. Magrath, J. R. Allan, J. Beher, K. R. Jones, H. P. Possingham, W. F.
540 Laurance, P. Wood, B. M. Fekete, M. A. Levy, and J. E. M. Watson. 2016a. Global Human
541 Footprint maps for 1993 and 2009. *Scientific Data* **3**:10067.

542 Venter, O., E. W. Sanderson, A. Magrath, J. R. Allan, J. Beher, K. R. Jones, H. P. Possingham, W. F.
543 Laurance, P. Wood, B. M. Fekete, M. A. Levy, and J. E. M. Watson. 2016b. Sixteen years of
544 change in the global terrestrial human footprint and implications for biodiversity conservation.
545 *Nat Commun* **7**:12558.

546 Wang, Z., Z. Yang, and X. Du. 2014. Analysis on the threats and spatiotemporal distribution pattern of
547 security in World Natural Heritage Sites. *Environmental Monitoring and Assessment* **187**.

548 Watson, J. E. M., E. S. Darling, O. Venter, M. Maron, J. Walston, H. P. Possingham, N. Dudley, M.
549 Hockings, M. Barnes, and T. M. Brooks. 2015. Bolder science needed now for protected areas.
550 *Conservation Biology* **30**:243-248.

551 Watson, J. E. M., K. R. Jones, R. A. Fuller, M. Di Marco, D. B. Segan, S. H. Butchart, J. R. Allan, E.
552 McDonald-Madden, and O. Venter. 2016. Persistent disparities between recent rates of habitat

553 conversion and protection and implications for future global conservation targets. Conservation
554 Letters DOI: 10.1111/conl.12295
555 Weisse, M. J. and L. C. Naughton-Treves. 2016. Conservation Beyond Park Boundaries: The Impact of
556 Buffer Zones on Deforestation and Mining Concessions in the Peruvian Amazon. Environmental
557 Management **58**:297-311.
558 Westerling, A. L., H. G. Hidalgo, D. R. Cayan, and T. W. Swetnam. 2006. Warming and Earlier Spring
559 Increase Western U.S. Forest Fire Activity. Science **313**:940 - 943.
560 Wittemyer, G., J. M. Northrup, J. Blanc, I. Douglas-Hamilton, P. Omondi, and K. P. Burnham. 2014. Illegal
561 killing for ivory drives global decline in African elephants. Proceedings of the National Academy
562 of Sciences **111**:13117-13121.
563 Woodroffe, R. and J. R. Ginsberg. 1998. Edge Effects and the Extinction of Populations Inside Protected
564 Areas. Science **280**:2126-2128.
565 Zaloumis, N. P. and W. J. Bond. 2011. Grassland restoration after afforestation: No direction home?
566 Austral Ecology **36**:357-366.

567

568 **Figure and Table Headings**

569 **Table 1.** Global and continental mean Human Footprint score per Natural World Heritage Site
570 (NWHS) and percentage change 1993 - 2009. Scores exceeding the global mean are shown in
571 bold.

572 **Table 2.** Natural World Heritage Sites (NWHS) with the greatest increases and decreases in
573 Human Footprint between 1993 and 2009.

574 **Table 3.** Global and continental mean percentage forest loss per Natural World Heritage Site
575 (NWHS), and total area of forest lost between 2001 and 2012. Percentages exceeding the global
576 average are shown in bold.

577 **Table 4.** Natural World Heritage Sites (NWHS) with high percentage forest loss between 2001
578 and 2012. The total area of forest lost over the time period is also shown.

579 **Figure 1.** Frequency distribution of changes in Human Footprint between 1993 and 2009 in
580 Natural World Heritage Sites (NWHS). * indicates the median change in HF and the arrow
581 indicates the mean change in HF. Colors specify the continent in which the NWHS is situated.

582 **Figure 2.** Change in mean Human Footprint between 1993 and 2009 across Natural World
583 Heritage Sites (NWHS) inscribed prior to 1993. NWHS which experienced an increase (which
584 may threaten their unique values) are shown in red, whilst NWHS which experienced a
585 decrease are shown in green. Site boundaries are not to scale, and have been enlarged for
586 clarity.

587 **Figure 3.** (a) Change in Human Footprint between 1993 and 2009 inside Natural World Heritage
588 Sites (NWHS) versus buffer zones. NWHS are coloured according to continent. (b) NWHS below
589 the identity line have undergone less change than their surrounding buffers indicating good
590 relative performance. (c) NWHS below the x-axis have undergone a mean decrease in Human
591 Footprint indicating good overall performance. (d) We can visualise sites performing well on
592 both the absolute and relative scales (green), or poorly on both (red).

593 **Figure 4.** Frequency distribution of percent forest loss between 2000 and 2012 in Natural World
594 Heritage Sites (NWHS). * indicates the median % loss and the arrow indicates the mean % loss.
595 Colours specify the continent in which the NWHS is situated.

596 **Figure 5.** Percent forest loss between 2000 and 2012 in Natural World Heritage Sites inscribed
597 prior to 2000. Sites experiencing substantial forest loss (>5%) are shown in red. Site boundaries
598 are not to scale, and have been enlarged for clarity.

599 **Figure 6.** Percent forest loss between 2000 and 2012 in Natural World Heritage Sites (NWHS)
600 versus buffer zones. NWHS are coloured according to continent. NWHS below the identity line
601 have suffered higher forest loss in the buffer zone compared to within the NWHS boundaries.

602

Tables and Figures:

Table 1. Global and continental mean Human Footprint score per Natural World Heritage Site (NWHS) and percentage change 1993 - 2009. Scores exceeding the global mean are shown in bold.

Continent	Human Footprint 1993		Human Footprint 2009		% Change 1993 - 2009		# sites
	NWHS	Buffer	NWHS	Buffer	NWHS	Buffer	
Africa	6.0	6.9	6.2	7.1	2.9	2.8	25
Asia	9.3	11.4	10.0	12.0	8.1	4.6	18
Australia	3.3	4.2	3.6	4.6	6.8	10.5	10
Europe	11.2	12.5	10.2	12.4	-9.6	0.0	13
North America	2.8	3.9	2.9	4.0	2.9	2.6	16
South America	4.2	5.4	4.5	6.3	4.8	15.8	12
Global	6.3	7.4	6.4	7.8	1.7	4.5	94

Table 2. Natural World Heritage Sites (NWHS) with the greatest increases and decreases in Human Footprint between 1993 and 2009.

	Human Footprint 1993		Human Footprint 2009		Change 1993 - 2009	
	NWHS	Buffer	NWHS	Buffer	NWHS	Buffer
Increases						
Manas Wildlife Sanctuary	11.8	12.0	17.0	14.2	5.3	2.2
Komodo National Park	6.2	n/a	10.6	n/a	4.3	n/a
St Kilda	4.9	n/a	8.4	n/a	3.5	n/a
Chitwan National Park	11.5	13.9	14.5	17.5	3.0	3.5
Simien National Park	5.7	8.2	8.6	10.1	2.9	2.2
Decreases						
Sinharaja Forest Reserve	16.7	17.7	9.7	11.5	-7.0	-6.3
Hierapolis-Pamukkale	23.5	14.6	17.0	14.3	-6.5	-0.2
Bialowieża Forest	12.6	9.7	8.5	10.8	-4.1	1.2
Göreme National Park and the Rock Sites of Cappadocia	22.0	13.2	18.8	12.9	-3.3	0.0
Mana Pools National Park, Sapi and Chewore Safari Areas	9.0	8.9	6.2	6.7	-2.9	-2.2

603

604

Table 3. Global and continental mean percentage forest loss per Natural World Heritage Site (NWHS), and total area of forest lost between 2001 and 2012. Percentages exceeding the global average are shown in bold.

Continent	Mean % forest loss per NWHS		Summed forest loss (km ²)		# sites
	NWHS	Buffer	NWHS	Buffer	
Africa	0.6	2.4	523.4	1220.4	32
Asia	1.2	2.3	1599.2	1628.9	31
Australia	1.6	6.2	237.8	524.6	12
Europe	1.5	1.9	51.1	89.0	16
North America	3.9	3.8	4131.8	1814.3	21
South America	0.7	2.7	728.0	1479.3	22
Global	1.5	2.9	7271.2	6756.6	134


605

Table 4. Natural World Heritage Sites (NWHS) with high percentage forest loss between 2001 and 2012. The total area of forest lost over the time period is also shown.

	% forest loss		Summed forest loss (km ²)	
	NWHS	Buffer	NWHS	Buffer
Waterton Glacier International Peace Park	23.1	14.9	540.7	317.1
Shark Bay	12.4	14.3	5.8	2.7
Wood Buffalo National Park	11.7	8.9	2581.5	513.4
Grand Canyon National Park	9.8	1.1	38.2	5.1
Río Plátano Biosphere Reserve	8.5	10.1	365.6	252.0
Doñana National Park	7.3	0.8	2.1	1.0
Yellowstone National Park	6.3	3.1	217.0	59.4
Mount Athos	5.8	6.1	13.1	0.7
Canadian Rocky Mountain Parks	5.3	3.7	424.5	176.4
Lake Baikal	4.8	10.9	1332.6	1044.7

606

607



608

609 **Figure 1.** Frequency distribution of changes in Human Footprint between 1993 and 2009 in
 610 Natural World Heritage Sites (NWHS). * indicates the median change in HF and the arrow
 611 indicates the mean change in HF. Colors specify the continent in which the NWHS is situated.

612




Figure 2. Change in mean Human Footprint between 1993 and 2009 across Natural World Heritage Sites (NWHS) inscribed prior to 1993. NWHS which experienced an increase (which may threaten their unique values) are shown in red, whilst NWHS which experienced a decrease are shown in green. Site boundaries are not to scale, and have been enlarged for clarity.




Figure 3. (a) Change in Human Footprint between 1993 and 2009 inside Natural World Heritage Sites (NWHS) versus buffer zones. NWHS are coloured according to continent. (b) NWHS below the identity line have undergone less change than their surrounding buffers indicating good relative performance. (c) NWHS below the x-axis have undergone a mean decrease in Human Footprint indicating good overall performance. (d) We can visualise sites performing well on both the absolute and relative scales (green), or poorly on both (red).




Figure 4. Frequency distribution of percent forest loss between 2000 and 2012 in Natural World Heritage Sites (NWHS). * indicates the median % loss and the arrow indicates the mean % loss.

Colours specify the continent in which the NWHS is situated.




Figure 5. Percent forest loss between 2000 and 2012 in Natural World Heritage Sites inscribed prior to 2000. Sites experiencing substantial forest loss (>5%) are shown in red. Site boundaries are not to scale, and have been enlarged for clarity.




Figure 6. Percent forest loss between 2000 and 2012 in Natural World Heritage Sites (NWHS) versus buffer zones. NWHS are coloured according to continent. NWHS below the identity line have suffered higher forest loss in the buffer zone compared to within the NWHS boundaries.