## Technical University of Denmark



### Drivers of flood damage on event level

Kreibich, H.; Aerts, J. C. J. H.; Apel, H.; Arnbjerg-Nielsen, Karsten; Baldassarre, G. Di; Bouwer, L. M.; Bubeck, Ph.; Caloiero, T.; Cortes, M.; Do, Ch.; Gain, A. K.; Giampa, V.; Kuhlicke, Ch.; Kundewicz, Z. W.; Llasat, M. C.; Mård, J.; Matczak, P.; Mazzoleni, M.; Molinari, D.; Nguyen, V. D.; Petrucci, O.; Schröter, K.; Slager, K.; Thieken, A. H.; Vorogushyn, S.; Merz, B.

Published in: Geophysical Research Abstracts

Publication date: 2016

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

*Citation (APA):* Kreibich, H., Aerts, J. C. J. H., Apel, H., Arnbjerg-Nielsen, K., Baldassarre, G. D., Bouwer, L. M., ... Merz, B. (2016). Drivers of flood damage on event level. Geophysical Research Abstracts, 18.

## DTU Library Technical Information Center of Denmark

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Geophysical Research Abstracts Vol. 18, EGU2016-**PREVIEW**, 2016 EGU General Assembly 2016 © Author(s) 2016. CC Attribution 3.0 License.



# Drivers of flood damage on event level

Heidi Kreibich and the IAHS Panta Rhei Working group "Changes in Flood Risk" Team German Research Centre for Geosciences GFZ, Section 5.4 Hydrology, Potsdam, Germany (kreib@gfz-potsdam.de)

Flood risk is dynamic and influenced by many processes related to hazard, exposure and vulnerability. Flood damage increased significantly over the past decades, however, resulting overall economic loss per event is an aggregated indicator and it is difficult to attribute causes to this increasing trend. Much has been learned about damaging processes during floods at the micro-scale, e.g. building level. However, little is known about the main factors determining the amount of flood damage on event level. Thus, we analyse and compare paired flood events, i.e. consecutive, similar damaging floods that occurred in the same area. In analogy to 'Paired catchment studies' - a well-established method in hydrology to understand how changes in land use affect streamflow – we will investigate how and why resulting flood damage in a region differed between the first and second consecutive flood events.

One example are the 2002 and 2013 floods in the Elbe and Danube catchments in Germany. The 2002 flood caused the highest economic damage (EUR 11600 million) due to a natural hazard event in Germany. Damage was so high due to extreme flood hazard triggered by extreme precipitation and a high number of resulting dyke breaches. Additionally, exposure hotspots like the city of Dresden at the Elbe river as well as some smaller municipalities at the river Mulde (e.g. Grimma, Eilenburg, Bitterfeld, Dessau) were severely impacted. However, affected parties and authorities learned from the extreme flood in 2002, and many governmental flood risk programs and initiatives were launched. Considerable improvements since 2002 occurred on many levels that deal with flood risk reduction and disaster response, in particular in 1) increased flood prevention by improved spatial planning, 2) an increased number of property-level mitigation measures, 3) more effective early warning and improved coordination of disaster response and 4) a more targeted maintenance of flood defence systems and their deliberate relocation. Thus, despite higher hydrological severity damage due to the 2013 flood was significantly lower than in 2002. In our international comparative paired event study we investigate under which circumstances similar or contrasting processes occurred and hope to identify common key processes which determine flood damage on event level.