

Technical University of Denmark



Characterization of membrane foulants at ambient temperature anaerobic membrane bioreactor treating low-strength industrial wastewater

Zarebska, Agata; Kjerstadius, Hamse; Petrinic, Irena; Buksek, Hermina; Korenak, Jasmina; Jansen, Jes La Cour; Hélix-Nielsen, Claus

Published in:
16th Nordic Filtration Symposium 2016

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Zarebska, A., Kjerstadius, H., Petrinic, I., Buksek, H., Korenak, J., Jansen, J. L. C., & Hélix-Nielsen, C. (2016). Characterization of membrane foulants at ambient temperature anaerobic membrane bioreactor treating low-strength industrial wastewater. In 16th Nordic Filtration Symposium 2016: Book of abstract Lappeenranta, Finland: Lappeenranta University of Technology Press .

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Characterization of membrane foulants at ambient temperature anaerobic membrane bioreactor treating low-strength industrial wastewater

Agata Zarebska ^a, Hamse Kjerstadius ^b, Irena Petrinic ^c, Hermina Buksek ^c, Jasmina Korenak ^c, Jes la Cour Jansen ^b, Claus Helix-Nielsen ^{a,c}

^a Department of Environmental Engineering, Technical University of Denmark, Bygningstorvet Building: 115, 139, 2800 Kgs. Lyngby, Denmark

^b Department of Chemical Engineering, Faculty of Engineering, Lund University, Getingevägen 60, 221 00 Lund, Sweden

^c Faculty of Chemistry and Chemical Engineering, University of Maribor, Smetanova ulica 17, 2000 Maribor, Slovenia

KEYWORDS

Anaerobic membrane bioreactor, wastewater, fouling

ABSTRACT

The large volume of industrial low-strength wastewaters has a potential for biogas production through conventional anaerobic digestion (AD), limited though by the need of heating and concentrating of the wastewaters. The use of anaerobic membrane bioreactor (AnMBR) combining membrane filtration with anaerobic biological treatment at low temperature could not only reduce the operational cost of AD, but also alleviate environmental problems. However, at low temperature the AnMBR may suffer more fouling due to the increased extracellular polymeric substances production excreted by bacteria hampering the application of the process for the industrial wastewater treatment.

In order to solve or reduce the fouling problem it is necessary to have a good insight into the processes that take place both on and in the membrane pores during filtration. Therefore, the objective of this study is to contribute to a better understanding of organic and biofouling in AnMBR.

An AnMBR consisting of external PVDF membrane was operated at 25°C and fed with synthetic dairy wastewater. Intensity, morphology and composition of foulants were determined using Scanning Electron Microscopy coupled with X-ray Energy Dispersive Spectrometry (EDS), Fourier Transform Infrared Spectrometry (ATR-FTIR), Inductively Coupled Plasma-Optical Emission Spectrometry (ICP-OES), Ion chromatography (IC), zeta potential, and adenosine triphosphate measurements.

Based on membrane autopsies, it can be concluded that prevailing fouling is mainly of biological and organic origin. SEM observations demonstrated presence of numerous bacteria incorporated with the fouling layer composed of mainly proteins, carbohydrates and lipids as revealed by ATR-FTIR measurements. Furthermore the amounts of ions found by EDS & ICP-OES analysis do not support scaling layer formation.