

Technical University of Denmark



Vancomycin gene selection in the microbiome of urban *Rattus norvegicus* from hospital environment

Arn Hansen, Thomas; Joshi, Tejal; Larsen, Anders Rhod; Andersen, Paal Skytt; Harms, Klaus; Mollerup, Sarah; Willerslev, Eske; Fursted, Kurt; Nielsen, Lars Peter; Hansen, Anders Johannes

Published in:

Evolution, Medicine, and Public Health

Link to article, DOI:

[10.1093/emph/eow021](https://doi.org/10.1093/emph/eow021)

Publication date:

2016

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Arn Hansen, T., Joshi, T., Larsen, A. R., Andersen, P. S., Harms, K., Mollerup, S., ... Hansen, A. J. (2016). Vancomycin gene selection in the microbiome of urban *Rattus norvegicus* from hospital environment. *Evolution, Medicine, and Public Health*, 2016(1), 219-226. DOI: 10.1093/emph/eow021

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Vancomycin gene selection in the microbiome of urban *Rattus norvegicus* from hospital environment

Thomas Arn Hansen^{1,†}, Tejal Joshi^{2,†}, Anders Rhod Larsen³, Paal Skytt Andersen³, Klaus Harms¹, Sarah Mollerup¹, Eske Willerslev¹, Kurt Fuursted³, Lars Peter Nielsen⁴ and Anders Johannes Hansen^{*,1}

¹Centre for GeoGenetics, Natural History Museum of Denmark, University of Copenhagen, Copenhagen, DK-1350, Denmark; ²Center for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, Kemitorvet, 2800, Kgs. Lyngby, Denmark; ³Department of Microbiology & Infection Control, Statens Serum Institut; and ⁴Department of Autoimmunology and Biomarkers, Statens Serum Institut, Copenhagen S, DK-2300, Denmark
*Corresponding author. Centre for GeoGenetics, Natural History Museum of Denmark, University of Copenhagen, Copenhagen, DK-1350, Denmark; E-mail: ajhansen@snm.ku.dk

[†]These authors contributed equally to this work

Received 15 January 2016; revised version accepted 29 June 2016

ABSTRACT

Background and objectives: Widespread use of antibiotics has resulted in selection pressure on genes that make bacteria non-responsive to antibiotics. These antibiotic-resistant bacteria are currently a major threat to global health. There are various possibilities for the transfer of antibiotic resistance genes. It has been argued that animal vectors such as *Rattus norvegicus* (*R. norvegicus*) living in hospital sewage systems are ideal for carrying pathogens responsible for fatal diseases in humans. **Methodology:** Using a metagenomic sequencing approach, we investigated faecal samples of *R. norvegicus* from three major cities for the presence of antibiotic resistance genes. **Results:** We show that despite the shared resistome within samples from the same geographic locations, samples from hospital area carry significantly abundant vancomycin resistance genes. **Conclusions and implications:** The observed pattern is consistent with a selection for vancomycin genes in the *R. norvegicus* microbiome, potentially driven by the outflow of antibiotics and antibiotic-resistant bacteria into the wastewater systems. Carriage of vancomycin resistance may suggest that *R. norvegicus* is acting as a reservoir for possible transmission to the human population.

KEYWORDS: *vanb*; vancomycin; rats; antibiotics; *Rattus*; wild; metagenomics; selection

BACKGROUND AND OBJECTIVES

The clinical burden of infections caused by antibiotic-resistant pathogenic bacteria is an increasing challenge worldwide. Especially the nosocomial infections with multi-resistant bacteria are problematic because of the rising difficulties in targeted treatment that result in increased morbidity and mortality particularly in immunocompromised patients [1,2]. For four decades, the preferred choice of antibiotic drug to target methicillin-resistant *Staphylococcus aureus* (MRSA) has been vancomycin [3]; this has likely contributed to an increase in the number of vancomycin-resistant *Enterococcus* sp. (VRE) as well as vancomycin-intermediate-resistant *S. aureus* and vancomycin-resistant *S. aureus* strains identified in patients [4].

Resistance to vancomycin was first reported in *Enterococci* sp. in 1988 [5, 6] and is conferred by *van* operons, where *vanA* and *vanB* are clinically relevant and most investigated *van* operons [7]. Both *vanA* and *vanB* operons are mobilizable. Although *vanA* is typically located in Tn1546-like transposable elements that frequently reside on plasmids [8], *vanB* is often a part of a larger conjugable chromosomal element (Tn1549) [9, 10] but has also been observed on plasmids [11]. The pathogenesis of VRE ranges from infections of the urinary tract, biliary tract and wounds to severe bacteraemia and endocarditis, frequently linked to fatal outcome [12, 13]. VRE are more difficult to treat than the antibiotic susceptible species and therefore VRE infections are more often correlated with poorer prognosis and increased number of hospitalization days [14–17]. The resistance rate of clinical VRE in the period from 1995 to 2002 has increased from ~47 to ~70% in the USA [1], likely accounted by an increased use of vancomycin and other antibiotics in hospitals.

The use of antibiotics in the hospitals can be traced in the wastewater system of the hospitals, where both antibiotics and antibiotic-resistant bacteria are readily detected [18–22]. There is a risk that the antibiotic-resistant bacteria in the wastewater systems can interfere with the local ecosystem. First, antibiotics in the environment can select for resistant bacteria. Second, resistance traits can spread horizontally to the locally adapted sensitive bacteria when resistance functions are mobilizable. The gut of *Rattus norvegicus* (*R. norvegicus*) has been hypothesized to act as an incubator for antibiotic-resistant bacteria from the hospitals [23]. In this case, *R. norvegicus* could migrate and carry the

antibiotic resistance genes or possibly carry mobile genetic elements acquired from the hospital wastewater systems.

In this study, we compare *R. norvegicus* faecal samples from hospital and non-hospital environments using a metagenomics DNA sequencing approach. We show that the faeces of *R. norvegicus* from hospital environment have elevated levels of vancomycin resistance genes.

METHODOLOGY

Faecal samples were collected from urban areas of Malaysia, Hong Kong and Denmark. All Danish samples from wild rats ($n=20$) were collected from four locations within the Copenhagen area: Egedal municipality (EM) ($n=3$), Copenhagen University Hospital (CUH) ($n=6$), Botanical Garden of Copenhagen (BGC) ($n=2$) and Amager East (AE) ($n=9$). In addition, five samples were collected in Kuala Lumpur (KLU), Malaysia, one in Kuala Langat, Malaysia and two samples were obtained from Hong Kong, China. Freshness of the samples was assessed by visual and tactile inspection. Rat faecal matter is easy to recognize visually. However, we performed a metabarcoding study on six of the samples, data published elsewhere [24], to confirm that they had *R. norvegicus* origins. The samples collected in Asia were shipped at ambient temperature in Falcon tubes and immediately frozen upon arrival. The samples from Denmark were frozen at -20°C within 24 h of collection.

The frozen faecal samples were vortexed vigorously in 800 μl of PBS for 1 min and incubated at room temperature for 30 min. Following the incubation, the samples were re-vortexed vigorously for a minute and then centrifuged at 12 000 g for 5 min. The supernatant was split into three aliquots of 160 μl and subsequently passed through 0.22 μm sterile filters at 6000 \times g for 5 min. Each of the three filtrates were nuclease treated using 14 μl Turbo DNase (2 U/ μl) (Ambion), 6 μl Baseline ZERO DNase (1 U/ μl) (Epicentre), 6 μl RNase Cocktail (Ambion), 8.5 μl sterile water and 20.5 μl 10 \times Turbo buffer in a total volume of 205 μl and incubated at 37°C for 2 h. The three aliquots were pooled and nucleic acid extracted using the QIAamp Viral RNA Mini Kit (Qiagen), followed by the addition of 1 μl RNase Out (Invitrogen) to the extract. Indexed DNA libraries were subsequently prepared using Nextera XT DNA Sample Preparation kit (Illumina), according to the manufacturers' guidelines. All subsequent sequencing

was performed by 100 bp paired-end sequencing on an Illumina HiSeq 2000.

Raw reads from the HiSeq platform were demultiplexed using Novobarcode (<http://www.novocraft.com>, vBeta-0.8). For each sample, AdapterRemoval (v1.1) [25] was used to trim low quality bases, to remove adapter sequences from paired-end reads and to merge paired-end reads overlapping with more than 11 nucleotides.

The sequences were aligned to the ARG-ANNOT [26] nucleotide database using Bowtie2 [27] (`-no-unal -end-to-end -q`) and only perfect matches were used for further analyses. All matches were masked using Dustmasker (v1.0.0) [28] and read matches with less than 20% low-complexity nucleotide sequence and longer than 75 nucleotides were considered a real match and counted. Genes with less than ten reads mapping to them ($n = 297$) from all samples were excluded.

A global mean normalization was applied to the remaining 240 313 reads that mapped to the nucleotide sequences in the ARG-ANNOT database [26]. Empirical Bayes moderated t-tests were obtained using limma [29]. *P*-values were adjusted using the Bonferroni method of multiple-testing correction. To further characterize genes differentially distributed between hospital and non-hospital samples, we performed two-dimensional hierarchical clustering using Euclidean distance measure.

To identify genes shared among sample locations, a gene was considered present in a location if at least one read mapped to the gene sequence. Assembly of the reads was performed using Ray Meta (v2.2.0 default settings) [30] and contigs were mapped to the ARG-ANNOT database using bowtie2 and the contigs with a match were aligned using BLASTn (v2.2.29+ default settings) to verify their origin. To establish the microbial composition of the samples we applied MetaPhlAn (v1.7.7 default settings) [31] on the reads of the samples.

RESULTS

Rat faecal samples were collected in the vicinity of Copenhagen University Hospital (CUH), and several urban, non-hospital locations in Copenhagen, Malaysia and Hong Kong. DNA from the samples was isolated and sequenced using a metagenomic approach and a total 240 313 reads were mapped onto the ARG-ANNOT [26] database that comprises 1689 curated antibiotic resistance genes of which 89 are vancomycin resistance genes. Read mapping to

the antibiotic resistance genes showed that 63 of these 1689 genes had at least 10 mapped reads across samples, and they were considered hits (Supplementary Fig. S1). There were 2697 unique reads mapping to vancomycin resistance genes. A student's *t*-test demonstrated a significant difference between the samples from the hospital area and all other locations in terms of the total uniquely mapped reads. Out of the 63 genes with at least 10 mapped reads, 13 genes had significantly higher amounts of mapped reads in hospital compared to non-hospital samples. Nine of these significantly abundant genes were *vanB* genes. Two-dimensional hierarchical clustering of these genes with all samples demonstrated high levels of vancomycin genes in the CUH samples (Fig. 1a). Hierarchical clustering of all normalized reads mapping to ARG-ANNOT genes showed a clear clustering among CUH samples (Fig. 1b). To establish the common resistome and shared antibiotic resistance genes, we grouped the samples based on locations and compared the read counts of antibiotic resistance genes. If a gene had at least one read match, it was considered to be present. A majority of genes (15) were shared among all locations in the Copenhagen area (Fig. 1c), whereas 13 genes were specifically present in CUH samples. The urban Copenhagen area (AE, CUH, BCG) had a shared resistome (18 genes) that was not shared with the rural EM samples. In summary, there clearly appeared to be a shared resistome among samples in the Copenhagen area, but the samples from CUH carried a specific set of genes (13), that were not present in other samples from Copenhagen (Fig. 1c).

To obtain more information on the underlying genomic sequences represented by all reads, we *de novo* assembled the reads into contigs. The contigs resembling vancomycin-resistance genes were short (100–750 nucleotides), but the majority of them mapped with high similarity to the *vanB* operon SAU16 (Acc:KF823968) or SAU28 (Acc:KF823969), which is typically located on the transposon Tn1549. We therefore mapped all reads from the hospital samples to the Tn1549 transposon (accession number: NG_035288) to see the combined read distribution (Supplementary Fig. S2). The consensus sequence of the mapped reads had 99% identity to Tn1549, an average read depth of 54 and the entire Tn1549 sequence had a read depth of at least one. Furthermore, when mapping to MetaPhlAn marker genes in samples across the geographical locations, we observed low quantities of *Enterococcus faecalis*




Figure 1. Summary of resistome comparison. **(a)** Two-dimensional hierarchical clustering of significantly differentially abundant genes between hospital and non-hospital environments. Hospital samples (labelled CUH) show higher levels of vancomycin resistance genes compared to the non-hospital samples. Abbreviations are EM, Denmark, CUH, Denmark, BGC, Denmark, AE, Denmark, KLU, Malaysia, Kuala Langat, Malaysia (KLA) and Hong Kong, (HK). **(b)** Hierarchical clustering of normalized read counts of the ARG-ANNOT mapped genes, using Euclidean distance method. **(c)** A Venn diagram showing the number of genes shared among sample locations in Copenhagen area

and *Enterococcus faecium* (<0.5% in samples from AE, EM, HK, KLU and CUH) and *S. aureus* (< 0.08% in samples from AE, EM, BGC, KLU and CUH).

CONCLUSIONS AND IMPLICATIONS

Recent investigations of antibiotic resistance genes in environmental, human gut and faecal samples from wildlife have revealed the presence of a variety of antibiotic resistance genes, collectively called a resistome [32–34]. The resistome from urban *R. norvegicus* faecal samples from seven locations in three major cities around the world revealed the presence of a variety of genes associated with

antibiotic resistance (Supplementary Fig. S1). The resistomes from the hospital locations clearly cluster together, whereas the other samples show no tendency of geographical clustering (Fig. 1a and b). A *t*-test analysis showed significant differences between the resistome of hospital and non-hospital samples. Highly abundant vancomycin resistance genes present in the hospital samples primarily accounted for these differences (Fig. 1a and c).

Contigs mapping to the vancomycin resistance genes of the *vanB* genotype (Fig. 1a) had high similarity to the SAU28 operon and the nearly identical SAU16 operon in clinical *E. faecium* strains. One of

the most common means of the transfer of antibiotic resistance genes between bacterial populations is horizontal gene transfer (HGT) via mobile genetic elements. The conjugative transposon Tn1549 carries the *vanB* operon as well as elements for HGT of the transposon by conjugation [10], allowing itself to spread to bacterial populations independent of other mobile elements. We found that a consensus sequence derived from all reads from the hospital-associated samples mapped to the complete transposon Tn1549 with 99% identity. Although normally found in the chromosome, Tn1549 has also been found on plasmids in clinical isolates of *E. faecalis* [11, 35], suggesting multiple ways of transfer of the *vanB* resistance traits.

Rodents' intestines may constitute an important reservoir of microbes including human pathogens such as MRSA and VRE [36, 37] and antibiotics present in their guts may favour colonization of intestines by opportunistic, antibiotic-resistant pathogens [38]. Additionally, exposure to low levels of antibiotics has been shown to drive the selection for antibiotic resistance genes in bacterial communities [39]. Hence, long-term exposure of rats to antibiotics could drive a positive selection for resistance genes in their guts. The hierarchical clustering of all normalized reads showed a clear clustering among CUH samples, indicating that *vanB* genes have been selected for in the guts of *R. norvegicus* near hospitals (Fig. 1a and b). Interestingly, high levels of vancomycin have been detected from the CUH wastewater [22] (median vancomycin concentration of 9.1 µg/l), which could be the main driver of the selection for *vanB* genes. The detection of *vanB* in hospital samples suggests that antibiotics used in hospitals and their subsequent spread to the environment through sewage might select for VREs in the rats' intestines.

As mentioned earlier, *Rattus* spp. are well-known carriers of bacterial pathogens like MRSA, *Leptospira*, *Streptobacillus moniliformis* etc [40, 41], and our results indicate presence of *Enterococcus* spp. as well. In addition, individual rats residing close to hospital environments have been shown to carry hundreds of plasmids in their gut microbiome [23]. It could therefore be of interest to explore if *R. norvegicus* can act as a vector of plasmids and transposons carrying vancomycin resistance genes to humans around hospitals.

Recently, risks associated with antibiotic resistance in the environment were ranked, and highest

risk was attributed to antibiotic resistance genes on mobile elements that can be acquired or hosted by known human pathogens [42]. In this study, we highlight this risk by demonstrating significantly higher abundance of resistance genes exclusively in hospital samples and showing that these abundant genes are associated with mobile genetic elements. Furthermore, the risk situation can be assessed with an ecological perspective in which the ecological connectivity is deemed to be a major determinant of the horizontal transfer of antibiotic resistance [43]. Essentially, a donor and a recipient need to be in close contact, and still a subsequent transfer event between habitats would be low unless the recipient is under positive selection, which is the case in the presence of antibiotics [42]. We have discovered a high-risk situation that might be driven by the elevated number of antibiotic resistance genes and influx of antibiotics, as seen in the sewage from the hospital use, in combination with an environmental connectivity between wastewater, rats and humans. If humans are infected with vancomycin-resistant pathogenic bacteria spread by *R. norvegicus* as carrier, there could be a complete environmental loop from rodents to humans requiring hospitalization, and thus spreading the resistance genes further to other rodents or patients (Fig. 2).

A transmission route has been proposed for the spread of VRE between animals and humans, but it does not include the spread from sewage to wildlife and subsequently to humans [44]. In this study, we demonstrate that the transfer of antibiotic-resistant bacteria from hospital environment to rodents is a strong possibility. Several studies demonstrate antibiotics and antibiotic-resistant bacteria in sewage around hospital environments [18–22, 45]. One study showed a higher prevalence of extended-spectrum beta-lactamase-producing *Escherichia coli* in rats around hospital wastewater [46]. If elevated levels of antibiotics in the hospital wastewater are driving the selection of antibiotic resistance in an important vector like *R. norvegicus*, it is imperative to investigate the full extent of this phenomenon on a global scale. Besides hospitals, where the same antibiotic selection-driven ecosystem probably resides, farms with livestock also have an extensive use of antibiotics. Assessment of the spread of antibiotic resistance genes into important vector species in these areas will also be highly relevant for risk assessment. It would also be of outmost importance to determine what relevant pathways exist for spread of




Figure 2. Proposed ecosystem. A proposed system of antibiotic resistance from hospitals acquired by bacteria that are carried further by rodents and passed on to humans that return to the hospital for treatment

antibiotic resistance genes from *R. norvegicus* to the human microbiome, so that appropriate timely measures can be taken to limit the spread of fatal infections to humans.

SUPPLEMENTARY DATA

Supplementary data is available at *EMPH* online.

ACKNOWLEDGEMENTS

We would like to acknowledge Claus Schultz, Andy Brigham and Picky Chan from Rentokil and Claus Christensen from HOFOR for arranging and assisting with sample collection and the Danish National Hospital for delivering samples. Also we would like to thank The National High-throughput DNA-Sequencing Centre for excellent execution of sequencing.

FUNDING

Villum Fonden and Innovationsfonden funded this study.

Conflict of interest: None declared.

REFERENCES

1. Wisplinghoff H, Bischoff T, Tallent SM *et al.* Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective nationwide surveillance study. *Clin Infect Dis* 2004;**39**:309–17.
2. Hidron AI, Edwards JR, Patel J *et al.* NHSN annual update: antimicrobial-resistant pathogens associated with healthcare-associated infections: annual summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2006–2007. *Infect Control Hosp Epidemiol* 2008;**29**:996–1011.
3. van Hal SJ, Fowler VG. Is it time to replace vancomycin in the treatment of methicillin-resistant *Staphylococcus aureus* infections? *Clin Infect Dis* 2013;**56**:1779–88.

4. Gardete S, Tomasz A. Mechanisms of vancomycin resistance in *Staphylococcus aureus*. *J Clin Invest* 2014; **124**:2836–40.
5. Leclercq R, Derlot E, Duval J *et al*. Plasmid-mediated resistance to vancomycin and teicoplanin in *Enterococcus faecium*. *N Engl J Med* 1988; **319**:157–61.
6. Uttley AH, George RC, Naidoo J *et al*. High-level vancomycin-resistant enterococci causing hospital infections. *Epidemiol Infect* 1989; **103**:173–81.
7. Courvalin P. Vancomycin resistance in gram-positive cocci. *Clin Infect Dis* 2006; **42**:S25–34.
8. Arthur M, Molinas C, Depardieu F *et al*. Characterization of Tn1546, a Tn3-related transposon conferring glycopeptide resistance by synthesis of depsipeptide peptidoglycan precursors in *Enterococcus faecium* BM4147. *J Bacteriol* 1993; **175**:117–27.
9. Quintiliani R, Courvalin P. Conjugal transfer of the vancomycin resistance determinant *vanB* between enterococci involves the movement of large genetic elements from chromosome to chromosome. *FEMS Microbiol Lett* 1994; **119**:359–63.
10. Tsvetkova K, Marvaud J-C, Lambert T. Analysis of the mobilization functions of the vancomycin resistance transposon Tn1549, a member of a new family of conjugative elements. *J Bacteriol* 2010; **192**:702–13.
11. Rice LB, Carias LL, Donskey CL *et al*. Transferable, plasmid-mediated vanB-type glycopeptide resistance in *Enterococcus faecium*. *Antimicrob Agents Chemother* 1998; **42**:963–4.
12. Murray BE. The life and times of the Enterococcus. *Clin Microbiol Rev* 1990; **3**:46–65.
13. Rubinstein E, Keynan Y. Vancomycin-resistant enterococci. *Crit Care Clin* 2013; **29**:841–52.
14. Linden PK, Pasculle AW, Manez R *et al*. Differences in outcomes for patients with bacteremia due to vancomycin-resistant *Enterococcus faecium* or vancomycin-susceptible *E. faecium*. *Clin Infect Dis* 1996; **22**:663–70.
15. Cetinkaya Y, Falk P, Mayhall CG. Vancomycin-resistant enterococci. *Clin Microbiol Rev* 2000; **13**:686–707.
16. DiazGranados CA, Zimmer SM, Klein M *et al*. Comparison of mortality associated with vancomycin-resistant and vancomycin-susceptible enterococcal bloodstream infections: a meta-analysis. *Clin Infect Dis* 2005; **41**:327–33.
17. DiazGranados CA, Jernigan JA. Impact of vancomycin resistance on mortality among patients with neutropenia and enterococcal bloodstream infection. *J Infect Dis* 2005; **191**:588–95.
18. Caplin JL, Hanlon GW, Taylor HD. Presence of vancomycin and ampicillin-resistant *Enterococcus faecium* of epidemic clonal complex-17 in wastewaters from the south coast of England. *Environ Microbiol* 2008; **10**:885–92.
19. Kjerulf A, Pallesen L, Westh H. Vancomycin-resistant enterococci at a large university hospital in Denmark. *Apmis* 1996; **104**:475–9.
20. Novais C, Coque TM, Ferreira H *et al*. Environmental contamination with vancomycin-resistant enterococci from hospital sewage in Portugal. *Appl Environ Microbiol* 2005; **71**:3364–8.
21. Sadowy E, Luczkiewicz A. Drug-resistant and hospital-associated *Enterococcus faecium* from wastewater, riverine estuary and anthropogenically impacted marine catchment basin. *BMC Microbiol* 2014; **14**:66.
22. Fredskilde JW, Nielsen U. Måleprogram på Rigshospitalet. *dhi* 2008; 1–59.
23. Jørgensen TS, Xu Z, Hansen MA *et al*. Hundreds of circular novel plasmids and DNA elements identified in a rat cecum metagenome. *PLoS One* 2014; **9**:e87924.
24. Hansen TA, Møllerup S, Nguyen N-P *et al*. High diversity of picornaviruses in rats from different continents revealed by deep sequencing. *Emerg Microbes Infect* 2016; **5**:e90.
25. Lindgreen S. AdapterRemoval: easy cleaning of next-generation sequencing reads. *BMC Res Notes* 2012; **5**:337.
26. Gupta SK, Padmanabhan BR, Diene SM *et al*. ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrob Agents Chemother* 2014; **58**:212–20.
27. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012; **9**:357–9.
28. Morgulis A, Gertz EM, Schäffer AA *et al*. A fast and symmetric DUST implementation to mask low-complexity DNA sequences. *J Comput Biol* 2006; **13**:1028–40.
29. Smyth GK. limma: Linear Models for Microarray Data. *Bioinformatics and Computational Biology Solutions Using R and Bioconductor*. New York: Springer-Verlag, 2005, 397–420.
30. Boisvert S, Raymond F, Godzaridis E *et al*. Ray Meta: scalable de novo metagenome assembly and profiling. *Genome Biol* 2012; **13**:R122.
31. Segata N, Waldron L, Ballarini A *et al*. Metagenomic microbial community profiling using unique clade-specific marker genes. *Nat Methods* 2012; **9**:811–4.
32. Forsberg KJ, Reyes A, Wang B *et al*. The shared antibiotic resistome of soil bacteria and human pathogens. *Science* 2012; **337**:1107–11.
33. Hu Y, Yang X, Qin J *et al*. Metagenome-wide analysis of antibiotic resistance genes in a large cohort of human gut microbiota. *Nat Commun* 2013; **4**:2151.
34. Gilliver MA, Bennett M, Begon M *et al*. Antibiotic resistance found in wild rodents. *Nature* 1999; **401**:233–4.
35. Zheng B, Tomita H, Inoue T *et al*. Isolation of VanB-type *Enterococcus faecalis* strains from nosocomial infections: first report of the isolation and identification of the pheromone-responsive plasmids pMG2200, encoding VanB-type vancomycin resistance and a Bac41-type bacteriocin, and pMG2201, encoding erythromycin resistance and cytolysin (Hly/Bac). *Antimicrob Agents Chemother* 2009; **53**:735–47.
36. Gries DM, Pultz NJ, Donskey CJ. Growth in cecal mucus facilitates colonization of the mouse intestinal tract by methicillin-resistant *Staphylococcus aureus*. *J Infect Dis* 2005; **192**:1621–7.

37. Lozano C, González-Barrio D, García JT et al. Detection of vancomycin-resistant *Enterococcus faecalis* ST6-*vanB2* and *E. faecium* ST915-*vanA* in faecal samples of wild *Rattus rattus* in Spain. *Vet Microbiol* 2015;**177**: 168–74.
38. Kesteman A-S, Perrin-Guyomard A, Laurentie M et al. Emergence of resistant *Klebsiella pneumoniae* in the intestinal tract during successful treatment of *Klebsiella pneumoniae* lung infection in rats. *Antimicrob Agents Chemother* 2010;**54**:2960–4.
39. Gullberg E, Albrecht LM, Karlsson C et al. Selection of a multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals. *MBio* 2014;**5**:e01918–14–14.
40. Himsworth CG, Miller RR, Montoya V et al. Carriage of methicillin-resistant *Staphylococcus aureus* by wild urban Norway rats (*Rattus norvegicus*). de Lencastre H (ed.). *PLoS One* 2014;**9**:e87983.
41. Himsworth CG, Parsons KL, Jardine C et al. Rats, cities, people, and pathogens: a systematic review and narrative synthesis of literature regarding the ecology of rat-associated zoonoses in urban centers. *Vector Borne Zoonotic Dis* 2013;**13**:349–59.
42. Martínez JL, Coque TM, Baquero F. What is a resistance gene? Ranking risk in resistomes. *Nat Rev Microbiol* 2015;**13**:116–23.
43. Cohen O, Gophna U, Pupko T. The complexity hypothesis revisited: connectivity rather than function constitutes a barrier to horizontal gene transfer. *Mol Biol Evol* 2011;**28**:1481–9.
44. Nilsson O. Vancomycin resistant enterococci in farm animals - occurrence and importance. *Infect Ecol Epidemiol* 2012;**2**:606.
45. Moges F, Endris M, Belyhun Y et al. Isolation and characterization of multiple drug resistance bacterial pathogens from waste water in hospital and non-hospital environments, Northwest Ethiopia. *BMC Res Notes* 2014;**7**:215.
46. Guenther S, Wuttke J, Bethe A et al. Is fecal carriage of extended-spectrum- β -lactamase-producing *Escherichia coli* in urban rats a risk for public health? *Antimicrob Agents Chemother* 2013;**57**:2424–5..