

## **Aeromonas salmonicida - Epidemiology, whole genome sequencing, detection and in vivo imaging**

**Bartkova, Simona; Dalsgaard, Inger; Kokotovic, Branko**

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Bartkova, S., Dalsgaard, I., & Kokotovic, B. (2016). *Aeromonas salmonicida - Epidemiology, whole genome sequencing, detection and in vivo imaging*. Frederiksberg C: National Veterinary Institute, Technical University of Denmark.

## **DTU Library** Technical Information Center of Denmark

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# *Aeromonas salmonicida*

**Epidemiology, whole genome sequencing, detection and *in vivo* imaging**

Simona Bartkova  
PhD Thesis  
2016

National Veterinary Institute  
Section for Bacteriology and Pathology  
Technical University of Denmark, Frederiksberg C

 **DTU Vet**  
National Veterinary Institute

**Supervisors:**

Associate professor Inger Dalsgaard  
Technical University of Denmark

Senior researcher Branko Kokotovic  
Technical University of Denmark

**Assessment committee:**

Professor Karl Pedersen  
Technical University of Denmark

Professor Anders Miki Bojesen  
University of Copenhagen

Senior researcher Duncan John Colquhoun  
Norwegian Veterinary Institute

## Table of contents

Acknowledgements	3
List of manuscripts	4
List of Abbreviations	5
Summary	6
Sammendrag (summary in Danish)	8
Introduction	10
Chapter 1: <i>Aeromonas salmonicida</i> subsp. <i>salmonicida</i>	12
1.1 Background and taxonomy	12
1.2 Biochemical and morphological characteristics	12
1.3 Cultivation	14
1.4 Virulence	14
Chapter 2: Furunculosis	18
2.1 Historical background	18
2.2 Clinical signs of disease	18
2.3 Antibiotic treatment	19
2.4 Vaccination	20
2.5 Transmission	21
2.6 Route of entry and colonization site(s)	23
2.7 Susceptibility in fish species	24
2.8 <i>In vivo</i> imaging	25
2.9 Manuscript I	26
Chapter 3: Detection	28
3.1 Detection of carrier fish	28
3.2 PCR	29
3.3 Real-time PCR	30
3.4 Manuscript II	31
Chapter 4: Epidemiology	33
4.1 Epidemiology and genetic variation	33
4.2 MLST-v	34
4.3 Whole genome sequencing	37
4.4 Manuscript III	39
Chapter 5: Methodological considerations	40
5.1 Materials and methods for MLST-v	40
5.2 <i>In vivo</i> imaging (Manuscript I)	41
5.3 Real-time PCR (Manuscript II)	42
5.4 Whole genome sequencing (Manuscript III)	43
Chapter 6: Discussion and future perspectives	44
6.1 Discussion	44
6.2 Conclusion and future perspective	46
References	48
Accompanying manuscripts	68

## **Acknowledgements**

This research was financed by the Danish Council for Strategic Research under the ProFish project (Grant no. DSF: 11-116252) and the National Veterinary Institute (DTU). I would like to acknowledge all the ProFish project partners for their work and commitment to the ProFish project and for all the informative discussions we have had. It has been a pleasure collaborating with all of you.

First and foremost, I am truly thankful for all the guidance I have received throughout my PhD from my main supervisor Inger Dalsgaard and co-supervisor Branko Kokotovic. Over the last three years (or a little more to be correct) I have gained a lot of valuable knowledge from you both, which I will take with me wherever my future career takes me. I know it has not always been easy with me as a student, but you have somehow managed to keep me grounded and helped me whenever needed. Although especially the last couple of months were intense and stressful for all of us (not to mention full of unexpected events), in the end we managed to put it all together. Finally, I would like to say that I will always be grateful to you Inger Dalsgaard for giving me the opportunity to do a PhD at the National Veterinary Institute with you as my supervisor and supporting me all the way to the end.

I would also like to give a special thank you to the laboratory technicians Lisbeth Schade Hansen, Lene Gertman, Katja Ann Kristensen and Margrethe Carlsen as well as other technicians that have helped me in the laboratory along the way. This project would not have been possible without you. Especially Lene Gertman who patiently introduced me to the laboratory world at the National Veterinary Institute and Lisbeth Schade Hansen who took over the difficult task of keeping an eye on me and helping me with the project.

This thesis would also have not been accomplished without the help of all the manuscript co-authors and especially without the whole genome sequencing analysis and guidance of “Shinny” Pimlapas Leekitcharoenphon.

Through my time as a PhD student there have also been several other PhD students at this section, whom I have had the pleasure of getting to know. We have all had our ups and downs during our PhDs, though luckily we always had each other for support and most importantly entertainment. Some of you have already finished your PhD and have moved on in the science world, while others still have some time before finishing and I wish everyone all the best!

Finally, I wish to express my gratitude to my family, friends and basketball teammates and coaches. You have supported me the whole way and especially during the last couple of stressful months when I needed it the most; particularly my parents who I could always turn to for extra guidance, inspiration, support and of course comfort and homemade food.

## List of manuscripts

This thesis includes the following original manuscripts that have either been submitted, accepted or are published online.

### **Manuscript I** (accepted in Journal of Fish Diseases)

*Title:* Infection routes of *Aeromonas salmonicida* in rainbow trout monitored *in vivo* by real-time bioluminescence imaging

*Authors:* Simona Bartkova, Branko Kokotovic, Inger Dalsgaard

### **Manuscript II** (published online in Journal of Fish Diseases)

*Title:* Detection and quantification of *Aeromonas salmonicida* in fish tissue by real-time PCR

*Authors:* Simona Bartkova, Branko Kokotovic, Helle Frank Skall, Niels Lorenzen, Inger Dalsgaard

### **Manuscript III** (submitted to Frontiers in Microbiology)

*Title:* Epidemiology and genetics of *Aeromonas salmonicida* using whole genome sequencing

*Authors:* Simona Bartkova, Pimlapas Leekitcharoenphon, Frank Møller Aarestrup, Inger Dalsgaard

## List of Abbreviations

AFLP	Amplified fragment length polymorphism
A-layer	Virulence associated surface protein array
ARGs	Antibiotic resistance genes
<i>A. salmonicida</i>	<i>Aeromonas salmonicida</i> subsp. <i>salmonicida</i>
BHI	Brain heart infusion
BLI	Bioluminescence imaging
bp	Base pairs
β	Beta
CBB	Coomassie brilliant blue
CC	Clonal complex
CFU	Colony-forming units
ECPs	Extracellular products
ELISA	Enzyme-linked immunosorbent assay
FMNH <sub>2</sub>	Reduced riboflavin phosphate
GCAT	Glycerophospholipid:cholesterol acyltransferase
GFP	Green fluorescence protein
IROMPs	Iron-regulated outer membrane proteins
LB	Luria Bertani
LPS	Lipopolysaccharide
LUX	Light upon eXtension
M-CGH	Microarray-based comparative genomic hybridization
MLST	Multilocus sequence typing
MLST-v	MLST with housekeeping genes and virulence associated genes
NGS	Next-generation sequencing
PCR	Polymerase chain reaction
PFGE	Pulsed-field gel electrophoresis
RAPD	Randomly amplified DNA polymorphism fingerprinting analysis
Real-time PCR	Quantitative real-time polymerase chain reaction
R plasmids	Plasmids carrying antibiotic resistance genes
SIF	Stress-inducible furunculosis
SLVs	Single-locus variants
SNP	Single nucleotide polymorphism
ST	Sequence type
TSA	Tryptic soy agar
T2SS	Type II secretion system
T3SS	Type III secretion system
T6SS	Type VI secretion system
VBNC	Viable but non-culturable cells
VIB	Veal infusion broth
WGS	Whole genome sequencing

## Summary

*Aeromonas salmonicida* subsp. *salmonicida* is a bacterial fish pathogen that is the causative agent of furunculosis, a septicemic infection responsible for great losses in aquaculture around the world. In Denmark furunculosis was first seen in freshwater in the 1950s, though currently the infection causes problems in sea reared rainbow trout (*Oncorhynchus mykiss*) production. Outbreaks occur repeatedly during stressful conditions such as elevated temperatures, in spite of commercial vaccines being applied. Besides seemingly lacking adequate protection, the vaccines also produce undesirable side effects. Antibiotics are therefore used as treatment, which due to the possibility of developing resistance is neither a favorable nor sustainable solution. To complicate things further, it is possible that fish can be carriers of *A. salmonicida* and transfer the bacterium from freshwater to the sea where they develop septicemia when exposed to stressful sea-rearing conditions and high temperatures. By use of traditional bacteriological methods, continuous investigation of bacterial diagnostics on samples from different rainbow trout farms in Denmark was done, while studying the following three aspects of the concerns regarding *A. salmonicida*.

First, we focused on investigation of the route of entry and initial dissemination of *A. salmonicida* in fish. This was done by tracing the bacterium using *in vivo* bioluminescence imaging. A Danish strain was transformed with a plasmid vector containing a green fluorescence protein gene and bacterial luciferase genes that served as fluorescent and bioluminescent reporters respectively. The transformed *A. salmonicida* was used in a series of immersion experiments where fish were followed over a 24-hour period. Results showed that probable main colonization sites of *A. salmonicida* were the gills and the dorsal and pectoral fins. This was followed by dissemination through internal organs. Although optimization and further immersion experiments are needed, our results indicated that this tool could be a valuable approach for visualizing *A. salmonicida* in fish.

Focus was subsequently turned to finding a sensitive method for detecting *A. salmonicida* in infected and possible carrier fish. For this, a previously developed quantitative real-time polymerase chain reaction (real-time PCR) targeting the *aopP* gene located on *A. salmonicida* plasmid pAsal1 was assessed. The real-time PCR and bacterial culturing were employed for preliminary screening of *A. salmonicida* in 40 fish from Danish fresh- and seawater farms. *A. salmonicida* was detected by real-time PCR in freshwater farm fish showing no sign of disease, indicating possible presence of carrier fish. Out of five examined organs: spleen, kidney, intestine, gills and brain in each fish, *A. salmonicida* was most frequently detected in the spleen, brain and intestine, indicating that these three organs could play an important role in *A. salmonicida* infection. The real-time PCR exhibited highly sensitive detection of *A. salmonicida* as well as a high reproducibility and efficiency, though due to the fact that not all *A. salmonicida* seem to possess the target plasmid pAsal1, another sensitive detection method with a different and/or complementary target would need to be employed to be certain of avoiding false negatives.

The final focal point of this thesis revolved around obtaining knowledge on genetic and virulence variation as well as epidemiology of the disease causing Danish *A. salmonicida*. Due to high homogeneity among the *A. salmonicida* subspecies population, standard molecular methods for



bacterial typing cannot distinguish among *A. salmonicida* isolates. Whole genome sequencing was therefore applied on 99 Danish *A. salmonicida* isolated between years 1980 and 2014 from different geographical regions, one Scottish strain and the type strain NCIMB 1102. Sequences of the *A. salmonicida* were *de novo* assembled and then examined for presence of plasmids, virulence and iron acquisition proteins, and antibiotic resistance genes. The chromosome was also examined for single nucleotide polymorphisms that were aligned and subjected to Bayesian temporal tree reconstruction using the published genome of *A. salmonicida* A449 as reference. Main results revealed that there have been four major introductions of *A. salmonicida* into Denmark, *A. salmonicida* are highly homogenous with the exception of certain plasmids and virulence factors encoded on these plasmids, and nine *A. salmonicida* harbored several worldwide known genes encoding resistance against antibiotics. This study provided valuable information regarding the Danish disease causing *A. salmonicida*.

## Sammendrag (summary in Danish)

*Aeromonas salmonicida* subsp. *salmonicida* er en fiskepatogen bakterie, som forårsager sygdommen furunkulose, der medfører store tab i akvakultur over hele verden. I Danmark blev furunkulose første gang beskrevet i ferskvand i 1950'erne, men i øjeblikket forårsager sygdommen problemer i regnbueørred (*Oncorhynchus mykiss*) produceret i havbrug. Udbrud forekommer gentagne gange under stressende forhold, såsom forhøjede vandtemperaturer, på trods af at kommercielle vacciner anvendes. Udover tilsyneladende at mangle tilstrækkelig beskyttelse, medfører vaccinerne også uønskede bivirkninger. Antibiotika anvendes derfor som behandling, men på grund af muligheden for udvikling af resistens er dette hverken en positiv eller holdbar løsning. For at komplicere tingene yderligere, kan fisk være såkaldte skjulte bærere af *A. salmonicida* og overføre bakterien fra ferskvand til havet, hvor sygdom udvikles når fiskene udsættes for stressende forhold. Ved brug af traditionelle bakteriologiske metoder, blev der udført kontinuerlig undersøgelse på fisk fra forskellige dam- og havbrug i Danmark, samtidig blev det forsøgt at klarlægge nogle af de ovennævnte problemer vedrørende *A. salmonicida*.

Først har vi fokuseret på at undersøge, hvordan *A. salmonicida* kommer ind i fiskene, samt bakteriens indledende udbredelse inde i fiskene. Dette blev gjort ved at spore bakterien med *in vivo* bioluminescens imaging. En dansk *A. salmonicida* stamme blev transformeret med en plasmid vektor, der indeholder et gen kodende for et grønt fluorescerende protein og bakterielle luciferase gener kodende for bioluminescence. Den transformerede *A. salmonicida* blev brugt i et foreløbigt bad eksperiment, hvor fiskene blev fulgt over en 24-timers periode. Resultaterne viste, at sandsynlige fasthæftnings steder af *A. salmonicida* var gællerne og ryg- og brystfinner. Dette blev efterfulgt af udbredelse til de indre organer. Selvom der er behov for optimering og yderligere bad eksperimenter, viste foreløbige resultater, at denne metode kunne være et værdifuldt redskab til at visualisere *A. salmonicida* i fisk.

Fokus blev derefter vendt imod at finde en følsom metode til påvisning af *A. salmonicida* i inficerede og mulige bærerfisk. Til dette blev en tidligere udviklet kvantitativ real-time polymerasekædereaktion (real-time PCR) rettet mod *aopP* genet, der er lokaliseret på *A. salmonicida* plasmidet pAsal1 vurderet. Real-time PCR og dyrkning af bakterien blev brugt til en foreløbig undersøgelse af *A. salmonicida* i 40 fisk fra danske dam- og havbrug. *A. salmonicida* blev påvist med real-time PCR i dambrugs fisk uden tegn på sygdom, hvilket tyder på tilstedeværelse af bærerfisk. Følgende fem organer: milt, nyre, tarm, gæller og hjerne fra hver fisk blev undersøgt, hvorfra *A. salmonicida* hyppigst blev påvist i milten, hjernen og tarmen, hvilket viser at disse tre organer kan spille en vigtig rolle i *A. salmonicida* infektionen. Real-time PCR udviste høj følsomhed for påvisning af *A. salmonicida* samt en høj reproducerbarhed og effektivitet, men på grund af at ikke alle *A. salmonicida* har plasmidet pAsal1, er det nødvendigt at anvende en anden følsom påvisnings metode rettet imod et andet gen, for at være sikker på at undgå falske negative resultater.

Det sidste omdrejningspunkt for denne afhandling drejer sig om at skaffe viden om den genetiske variation, samt oprindelse og spredning af danske *A. salmonicida* som forårsager sygdom. På grund af høj homogenitet i *A. salmonicida* bakterierne kan standard molekylære metoder til bakteriel

typning ikke skelne mellem forskellige *A. salmonicida* isolater. Helgenomsekventering blev derfor anvendt på 99 danske *A. salmonicida* stammer der blev isoleret i perioden 1980 til 2014 fra forskellige geografiske områder. Yderligere blev en skotsk bakterie og type bakterien NCIMB 1102 undersøgt. Sekvenser af bakterierne blev *de novo* samlet og derefter undersøgt for tilstedeværelse af plasmider, virulens og jern protein sekvenser og gener for antibiotika resistens. Kromosomet blev også undersøgt for nukleotid polymorfenheder (SNPs), der blev brugt til konstruktion af et Bayesian fylogenetisk træ ved hjælp af det publicerede genom af *A. salmonicida* A449 som reference. De vigtigste resultater viste, at der har været fire store introduktioner af *A. salmonicida* i Danmark, at *A. salmonicida* er meget homogene med undtagelse af visse plasmider og virulensfaktorer kodet på disse plasmider, og at ni *A. salmonicida* havde flere globalt kendte gener, der koder for antibiotika resistens. Studiet resulterede i værdifuld viden om *A. salmonicida* bakterier, der forårsager sygdom i akvakultur i Danmark.

## **Introduction**

### **Development in aquaculture**

According to the Food and Agriculture Organization of the United Nations (2016), aquaculture is the fastest growing food-producing sector. This intensified fish-farming combined with elevated water temperature due to global warming, creates epidemiological opportunities for pathogens and thereby causing problems in aquaculture (Tan et al., 2002; Ganusov and Antia, 2003; De Silva and Soto, 2009; Pulkkinen et al., 2010). One of the problems is that the fish immune system becomes adversely affected by physiological stress arising during high stocking density and abnormal climate conditions such as prolonged rise in temperature (Harvell et al., 1999; Vargas-Chacoffa et al., 2014). The above mentioned conditions also create a favorable environment for activity of pathogens that otherwise would typically remain dormant (Lafferty, 2009) and general increase of virulence and/or transmission rate of pathogens (Marcogliese, 2008; Marcos-López et al., 2010).

In contrast to the rest of the world's aquaculture industry, the Danish industry has in the recent years remained stagnant. However, fish consumption in Denmark has increased during the last few years (Miljø- og Fødevareministeriet, 2015), leading to the Danish authorities' proposal for expanding aquaculture in order to increase production (Miljø- og Fødevareministeriet, 2014). Use of antibiotics in aquaculture has decreased due to implementation of vaccines. However, in comparison with other animal productions in Denmark, marine aquaculture lies at the top in use of antibiotics along with pig production when calculated in DAPD (Defined animal daily dose per 1,000 animals per day), which includes changes in live biomass and thus enables comparison of different animals (DANMAP, 2012). An increase in production and a possible rise in water temperature will not alleviate this problem. On the contrary, it will increase the potential threat of highly virulent pathogens emerging.

### **Problem of furunculosis**

Furunculosis is a septicemic infection caused by the highly homogenous Gram-negative bacterium *Aeromonas salmonicida* subsp. *salmonicida* (Bernoth et al., 1997; Garcia et al., 2000). Though *A. salmonicida* can be present in fish without inducing signs of disease where the infection is said to be in a 'covert' stage (Hiney et al., 1997). It is believed that these carrier fish transfer the bacterium from freshwater farms out to seawater farms, where the fish develop septicemia when exposed to stressful sea-rearing conditions and high temperatures (Dalsgaard and Madsen, 2000; Pedersen *et al.*, 2008). *A. salmonicida* can be diagnosed via traditional bacteriological methods (Dalsgaard et al., 1994; Austin and Austin, 2007), however, detection of the bacterium in carrier fish based on these methods has thus far been unsuccessful (Dalsgaard and Madsen, 2000). New and more sensitive methods need to be developed in order to detect *A. salmonicida* in carrier fish.

Furunculosis was first described from freshwater farms in Denmark in the 1950s (Rasmussen, 1964) and now causes great problems in seawater rainbow trout farms. There is only one vaccine against *A. salmonicida* that is licensed for Danish rainbow trout, which is the Norwegian commercial vaccine

AlphaJect 3000, initially developed for Atlantic salmon (*Salmo salar*) against vibriosis and furunculosis (Pharmaq, 2016). Although the vaccine is being implemented, outbreaks of furunculosis still occur (Dalsgaard and Madsen, 2000; Pedersen *et al.*, 2008), causing substantial economic losses in aquaculture. Side effects, likely caused by oil adjuvants, have also been reported (Haugarvoll *et al.*, 2010; Mutoloki *et al.*, 2010). In order to develop an effective strategy for preventing furunculosis, a more effective vaccine against Danish *A. salmonicida* needs to be developed and knowledge about the epidemiology, genetic and virulence variation of the Danish disease causing *A. salmonicida* isolates needs to be obtained.

### **Objective of this thesis**

The objective of this PhD project was to contribute to ongoing research on resolving the current concerns of furunculosis in Danish rainbow trout production by: 1) investigating route of entry and dissemination of *A. salmonicida* in fish in order to study the host-pathogen relationship that could provide new knowledge for improvement of detection and sampling strategies of the bacterium, 2) developing a highly sensitive method for detection of *A. salmonicida* in possible carriers and fish showing signs of disease, and 3) determining the epidemiology, genetic and virulence variability of the Danish *A. salmonicida* isolates that could aid in the development of an effective strategy for preventing furunculosis.


## Chapter 1: *Aeromonas salmonicida* subsp. *salmonicida*

### 1.1 Background and taxonomy

*Aeromonas salmonicida* subsp. *salmonicida* is an important bacterial fish pathogen, which was originally isolated at a German freshwater farm by Emmerich and Weibel (1894) and was given the name *Bacterium salmonicida*. Subsequently it was proposed by Griffin et al. (1953) to place the bacterium in the genus *Aeromonas* and re-classify the name of the species as *Aeromonas salmonicida* (Snieszko, 1957). The genus of *Aeromonas* has also gone through many taxonomic re-classifications and was eventually placed in the family *Aeromonadaceae* by Colwell et al. (1986). Although the species of *Aeromonas salmonicida* was first thought to be homogenous, by use of biochemical and molecular methods it has thus far been divided into five subspecies: *salmonicida*, *masoucida*, *achromogenes*, *smithia*, and *pectinolytica* (Austin, 1993; Wiklund and Dalsgaard, 1998; Kozinska et al., 2002; Beaz-Hidalgo et al., 2008; Studer et al., 2013). The four latter subspecies all belong to the so called “atypical” group, while subspecies *salmonicida* is the only *Aeromonas salmonicida* known as “typical” and is the causative agent of furunculosis. Subspecies *salmonicida* is also the focal point of this PhD project and is in this thesis referred to as *A. salmonicida*.

### 1.2 Biochemical and morphological characteristics

The bacterium *A. salmonicida* is Gram-negative, facultative anaerobic, non-motile, psychrophilic and consists of coccoide cells with the measurement of 0.5-6.0 x 1-2  $\mu\text{m}$  (Marsh, 1902; Griffin et al., 1953; Cipriano and Austin, 2011). One of the most basic characteristics of the subspecies is that colonies produce a brown water-soluble pigment after growth on agar in the presence of 0.1% tyrosine or phenylalanine for two to four days (Fig. 1) (Marsh, 1902; Griffin et al., 1953; Boone et al., 2001; Cipriano and Austin, 2011). The subspecies must, however, not be identified solely based on this characteristic, since some *A. salmonicida* strains do not produce this pigment (Wiklund et al., 1993; Koppang et al., 2000) while some other bacteria like *A. hydrophila* also produce diffusible brown pigment (Austin and Austin, 2012). There are also numerous well-known biochemical characteristics, which are frequently used for identification. This includes production of catalase and cytochrome oxidase and gelatin liquefaction, although exceptions have been found (Böhm et al., 1986; Wichardt et al., 1989; Chapman et al., 1991). Generally the bacterium has also been reported of being positive for the following carbohydrates and glycosides: glycerol, glucose, fructose, galactose, mannitol, mannose, maltose, dextrin, glycogen, starch, aesculin and salicin, as well as being positive in L-arabinose but negative in D-arabinose and not being able to convert tryptophan into indole (Dalsgaard et al., 1994). Though, gas production by *A. salmonicida* from fermented glucose might be weak in some strains or as seen with a strain from Canada, gas might not be produced (Dalsgaard et al., 1994). *A. salmonicida* negative for acid production from mannitol and hydrolysis in aesculin have, however, also been found (Austin et al., 1989).



**Figure 1.** *A. salmonicida* colonies grown on tryptic soy agar (left) and on Coomassie brilliant blue agar (right, from Austin and Austin, 2012)

After incubation on nutrient agar for about 24 hours at 18 - 20°C, *A. salmonicida* colonies are flat and punctiform with a diameter of less than 1 mm (Griffin et al., 1953; Boone et al., 2001). It is not until three to four days of growth (Griffin et al., 1953; Boone et al., 2001) that the colonies become circular, convex and entire with a diameter of 1 - 2 mm, which is the characteristic morphology of *A. salmonicida* colonies on solid media. Colonies also become friable at this point, such that if pushed, they can slide on the agar surface without being damaged (Duff and Stewart, 1933; Munro and Hastings, 1993; Boone et al., 2001). When the bacterium is grown on blood agar for 2 - 4 days the colonies also become grayish in color and form a zone of haemolysis around them due to production of  $\beta$ -haemolysis (Griffin et al., 1953; Dalsgaard et al., 1994).

Additionally, *A. salmonicida* has the ability to autoagglutinate in static liquid media and form “smooth” and “rough” colonies on solid media (Arkwright, 1912; Williamson, 1928; Udey, 1978; Kay and Trust, 1997). These characteristics are generally based on the presence or absence of the virulence associated surface protein array called the A-layer, with A-layer positive *A. salmonicida* strains autoagglutinating and forming “rough” colonies (Johnson et al., 1985; Dalsgaard et al., 1994; Kay and Trust, 1997; Austin and Austin, 2012). A-layer negative strains that autoagglutinate have, however, also been found (Johnson et al., 1985) and single bacterial isolates can form morphologically different colonies on solid media (Anderson, 1972; Dalsgaard et al., 1994; Austin and Austin, 2012). This is supported by our findings, where all except one of 101 sequenced *A. salmonicida* possessed the A-layer protein sequence (Manuscript III), though only 76 of the isolates seemed to autoagglutinate in static liquid media (unpublished results).

Another property of *A. salmonicida* that has been used in order to distinguish A-layer positive from A-layer negative strains is the ability to bind the dyes Coomassie brilliant blue (CBB) (Wilson and Horne, 1986; Cipriano and Bertolini, 1988; Dalsgaard et al., 1994; Austin and Austin, 2012) and

Congo red (Ishiguro et al., 1985; Dalsgaard et al., 1994). The A-layer positive *A. salmonicida* bind the dye and grow as dark blue and red colonies on tryptic soy agar (TSA) with added CBB and Congo red respectively, while the A-layer negative colonies grow as whitish colonies. Markwardt et al. (1989) also reported that CBB agar could be used to distinguish A-layer positive *A. salmonicida* from other bacteria in mixed populations isolated from clinical samples. However, Teska and Cipriano (1993) reported that bacteria such as *Aeromonas hydrophila*, *Pasteurella multocida* and various *Pseudomonas* species can also grow with dark blue colonies on CBB agar and dark blue colonies must therefore be subcultivated and tested for other characteristics before any confirmations can be made regarding their identification.

### 1.3 Cultivation

The recommended medium for isolation and cultivation of *A. salmonicida* by diagnostic manuals has usually been brain heart infusion (BHI) agar or TSA and both have been used in several studies (e.g. Beaz-Hidalgo et al., 2008b; Beaz-Hidalgo et al., 2013; Austin and Austin, 2012). TSA has the advantage of the possibility of supplementation with CBB or Congo red for selection of A-layer positive *A. salmonicida* as mentioned in the previous section, however, selection using this medium is not consistent and Austin and Austin (2012) reported that *A. salmonicida* “rough” colonies that are associated with A-layer and virulence are recovered better on BHI agar than TSA. *A. salmonicida* can also grow on common laboratory media such as Luria Bertani (LB) and blood agar and the latter has especially been used frequently (e.g. Böhm et al., 1986; Dalsgaard et al., 1994; Pedersen et al., 2008) due to the usual *A. salmonicida* production of  $\beta$ -haemolysis on this medium. The choice of liquid medium for cultivation of *A. salmonicida*, which enables the possibility for observing autoagglutination of A-layer positive isolates, has also varied between studies e.g. veal infusion (VIB) broth (Dalsgaard et al., 1994), Trypticase soy broth (Ishiguro et al., 1981) and BHI broth (Johnson et al., 1985). All *A. salmonicida* used in the present thesis were always cultivated in VIB broth and isolated on blood agar, with the exception of one study (Manuscript I) where LB agar with added ampicillin and BHI broth were also employed due to special circumstances.

Ideal growth conditions for *A. salmonicida* include aerobic conditions and pH from 5.3 to 9.0, although this can vary depending on the composition of the culture medium (Griffin et al., 1953). For many years the optimal growth temperature for *A. salmonicida* has been reported as being 22 - 25°C (Griffin et al., 1953; Brenner et al., 2005) and the maximum growth temperature 34.5°C (Griffin et al., 1953). However, it has now been reported by several studies that when *A. salmonicida* is grown at 25°C, or in some studies even 22°C, some of the plasmid encoded virulence genes can become inactivated or lost due to plasmid rearrangement or loss of the plasmids (Ishiguro et al., 1981; Stuber et al., 2003; Daher et al., 2011). This has led to Daher et al. (2011) suggesting that *A. salmonicida* should be grown at a maximal temperature of 20°C. All *A. salmonicida* used in this thesis were grown at 20°C.

### 1.4 Virulence

There are many virulence factors that *A. salmonicida* possesses, which can be used against the defense mechanisms of the host in order to establish an infection (Reith et al., 2008; Beaz-Hidalgo



and Figueras, 2013; Dallaire-Dufresne et al., 2014). Though, not all virulence factors are equally important and some are even non-functional (Reith et al., 2008). In fact, the virulence mechanism of *A. salmonicida* has proven to be a complex and intertwined system (Reith et al., 2008; Beaz-Hidalgo and Figueras, 2013; Dallaire-Dufresne et al., 2014).

One of the known virulence components of *A. salmonicida* that are important for attachment of host cells and entry into the host are adhesins e.g. the surface layer and pili (Austin and Austin, 2007; Reith et al., 2008). Though, it is only the A-layer, a tetragonal protein array that is associated with lipopolysaccharides (LPSs) on the cell surface, which has been described as being a major virulence factor of *A. salmonicida* (Udey and Fryer, 1978; Ishiguro et al., 1981; Phipps et al., 1983; Trust et al., 1983; Chart et al., 1984). This protein, encoded by the *vapA* gene (Chart et al., 1984), also displayed high sequence homogeneity in 101 sequenced *A. salmonicida*, of which 99 isolates were from Denmark and only one Danish isolate did not harbor the protein (Manuscript III). The A-layer also has a high proportion of hydrophobic amino acids; a property that increases hydrophobicity of the bacterial surface (Phipps et al., 1983). In agreement, Dalsgaard et al. (1994) reported that the “rough” colony forming strains of *A. salmonicida* believed to be A-layer positive had a more hydrophobic outer cell surface than the strains forming “smooth” colonies. When studying *in vitro* cultured macrophages Trust et al. (1983) found that A-layer positive *A. salmonicida* had an enhanced ability to associate with the macrophages, which was also enabled by the increase in hydrophobicity. The increased hydrophobicity is also responsible for A-layer positive *A. salmonicida* strains’ ability to autoagglutinate and to adhere to host tissue, while the A-layer negative strains have been reported to be avirulent (Udey and Fryer, 1978). The association of hydrophobicity with aggregation and sedimentation is also supported by Enger and Thorsen (1992), who added that the property could also play a role in the ecology of the bacterium outside its host, since *A. salmonicida* was detected by use of immunofluorescence in sediment beneath net pens, seawater, and surface film from samples at fish farms with furunculosis outbreaks. Nevertheless, A-layer negative virulent as well as A-layer positive non-virulent *A. salmonicida* have been observed, exemplifying that *in vitro* tests do not necessarily give an accurate result when assessing virulence of *A. salmonicida* (Johnson et al., 1985; Ellis, 1997), which as suggested by Olivier (1990) should be assessed through *in vivo* challenges instead.

Other virulence factors that have been reported as being important for virulence of *A. salmonicida* are extracellular products (ECPs) such as haemolysins, aerolysins, lipopolysaccharides, proteases and various toxins, which have been the subject of many *A. salmonicida* studies over the years (Ellis et al., 1981; Ellis et al., 1988; Ellis, 1997; Austin and Austin, 2007; Beaz-Hidalgo and Figueras, 2013). Already in 1953, Griffin et al. believed that the observed  $\beta$ -haemolysis and gelatin liquefaction on blood and gelatin plates respectively could be related to the characteristic tissue lesions of furunculosis and thus caused by production of protease enzymes. Years later in a study by Ellis et al. (1981), it was possible to reproduce the furunculosis associated lesions when ECPs were injected into rainbow trout. In another study, Ellis et al. (1988) found that protease and haemolysin activity promoted the development of lesions in fish, though there was another yet uncharacterized factor of the ECPs that was lethal for the fish. One ECP that has been proven to be lethal for fish is the toxin glycerophospholipid:cholesterol acyltransferase (GCAT) (Lee and Ellis, 1990). The GCAT protein sequence was present in all 101 sequenced *A. salmonicida* and displayed similar sequence

homogeneity as the A-layer (Manuscript III). This toxin can also aggregate with LPSs, creating a GCAT/LPS complex which is even more toxic than GCAT by itself (Lee and Ellis, 1990). Surprisingly, when the encoding genes of either GCAT or another reportedly important toxin (serine protease AspA) were mutated, virulence of the *A. salmonicida* mutant was not altered (Vipond et al., 1998).

Iron-regulated outer membrane proteins (IROMPs) are responsible for uptake of siderophore-iron complexes and heme (Hirst et al., 1994; Najimi et al., 2008, 2009). They are thus believed to play a significant role in virulence of *A. salmonicida*, since iron is an essential nutrient and acquisition of iron is necessary for survival within the host and maybe also the aquatic environments (Hirst et al., 1994; Najimi et al., 2008, 2009; Reith et al., 2008; Ebanks et al., 2013). Iron acquisition by *A. salmonicida* can be siderophore dependent, by which the siderophores remove iron from proteins of the host and enable entrance of the iron into the bacterial cell through IROMPs (Najimi et al., 2008; 2009). Though, iron can also be obtained by a siderophore independent system that functions as a way to remove iron from host hemoglobin (Ebanks et al., 2004; Najimi et al., 2008; 2009). With regard to *A. salmonicida*, IROMPs have also shown to have an inhibitory effect on host transferrin thereby improving the bacterial resistance against the host's phagocytes (Magnadottir, 2010). Investigation of genes involved in siderophore biosynthesis and IROMPs indicated that both systems seem to be conserved among the homogenous *A. salmonicida* (Fernandez et al., 1998; Najimi et al., 2009), as supported by the findings of Manuscript III, while other iron mechanism proteins sequences are not present in all *A. salmonicida* (Najimi et al., 2009).

Secretion systems have also been known for their significance for virulence and three have been described in *A. salmonicida*, which includes type II (T2SS), III (T3SS), and VI (T6SS) (Reith et al., 2008). T2SS is in charge of enzyme degradation and toxin secretion, while T6SS enables injection of effector proteins into the cytoplasm of the host cells (Reith et al., 2008). Much like T6SS, the main function of T3SS is also injection of effector proteins and toxins into the cytosol of host cells, but has also other functions including prevention of phagocytosis by leukocytes and establishing a systemic infection in the host (Burr et al., 2003; Stuber et al., 2003; Burr et al., 2005; Mota and Cornelis, 2005; Dacanay et al., 2006; Ebanks et al., 2006; Rasch et al., 2007; Fast et al., 2009; Dallaire-Dufresne et al., 2014). Interestingly, both T3SS and T6SS associated genes are moreover situated in the chromosome as well as plasmids (Reith et al., 2008; Fehr et al., 2006). Though, notably T3SS is the only virulence factor proven to be essential for virulence and toxicity of *A. salmonicida*, since inactivation of the T3SS structural proteins in *A. salmonicida* has always rendered the *A. salmonicida* mutants non-virulent in both *in vitro* and *in vivo* studies (Burr et al., 2002; Burr et al., 2003; Stuber et al., 2003; Burr et al., 2005; Dacanay et al., 2006; Froquet et al., 2007; Fast et al., 2009). Thus far there have been described five T3SS effector proteins in *A. salmonicida*: AexT (an ADP-ribosylating toxin encoded on the chromosome), AopH (a tyrosine phosphatase encoded on the plasmid pAsa5 along with its chaperone SycH), AopO (a serine/threonine kinase encoded on the plasmid pAsa5 along with its chaperone SycO), Ati2 (an inositol polyphosphate 5-phosphatase encoded on the plasmid pAsa5 along with its chaperone Ati1) and AopP (this toxin is involved in inhibition of I $\kappa$ B protein kinase activation and is encoded on plasmid pAsa11) (Dacanay et al., 2006; Fehr et al., 2006; Reith et al., 2008; Dallaire-Dufresne et al., 2013). Both the effector proteins and structural proteins of T3SS are thought to play a role in *A. salmonicida* survival within the host cell, though when

compared only the structural proteins proved to be vital for virulence (Dacanay et al., 2006; Fast et al., 2009). An additional factor supporting the important implication of T3SS in virulence is that the secretion system is widely spread among pathogenic Gram-negative bacteria, such as other *Aeromonas* spp., *Yersinia* spp., *Salmonella* spp., and *Pseudomonas* spp., where it has also been confirmed to be vital for virulence of the pathogens (Chacón et al., 2004; Vilches et al., 2004; Mota and Cornelis, 2005; Vilches et al., 2009). Interestingly, both certain T3SS effector protein and structural protein sequences e.g. the highly studied AscV, were absent in 24% of the 101 sequenced *A. salmonicida*; all of which were isolated from furunculosis outbreaks and were thus assumed to be virulent (Manuscript III).

## Chapter 2: Furunculosis

### 2.1 Historical background

Furunculosis is now spread worldwide, though the first time furunculosis was observed and documented among fish was in 1894 by Emmerich and Weibel. They observed swellings resembling boils as well as ulcerative lesions in brown trout (*Salmo trutta*) at a German freshwater hatchery. Another observation of Emmerich and Weibel was that the brown trout were held in low-quality water before being transported to a farm where other fish also became infected. Through experiments Emmerich and Weibel also discovered that by either intramuscularly injecting fish or adding culture to the water tank, the fish could become infected. Cohabitation experiments gave the same results, however, when examining healthy fish from these experiments, the bacterium was not found.


After the initial description by Emmerich and Weibel (1894), furunculosis was believed to be a hatchery associated infection until the studies of Plehn (1911) showed that furunculosis was also present among wild trout in Germany and others also observed the infection in several countries all over the world, including Great Britain who suffered great losses (Fuhrman, 1909; Pittet, 1910; Surbeck, 1911; Arkwright, 1912; Mettam, 1915; Christensen, 1980). In the United States, furunculosis was first described by Marsh (1902) at hatcheries in Michigan. Shortly thereafter, the infection was found in numerous salmon and trout hatcheries throughout the United States (Fish, 1937; Smith, 1942). The origin of furunculosis in the United States is uncertain, though the general theory is that either it was brought along with brown trout from Germany or it spread from rainbow trout farmed in the Western part of the United States (Fish, 1937). Signs of furunculosis were also seen in several fish species in Canada by Duff and Stewart (1933) and various trout farms in Japan (Furunculosis committee, 1933).

In Denmark furunculosis was first described in the 1950s at freshwater rainbow trout farms by Rasmussen (1964). In parallel with this discovery, a massive expansion in rainbow trout production started that continued its growth even further as production became established in seawater in the 1970s (Christensen, 1980). Moreover, a Bayesian temporal tree based on SNP analysis of 101 sequenced *A. salmonicida* showed that there have been four main introductions of *A. salmonicida* in Denmark, two of which occurred approximately the same time as the first expansion in rainbow trout production (~ 1950) and the other two during the second expansion in seawater (~ 1970) (Manuscript III). At present, it is in the seawater production during elevated temperatures that furunculosis is of great concern and causes huge financial losses (Larsen and Møllergaard, 1981; Dalsgaard and Madsen, 2000; Pedersen et al., 2008).

### 2.2 Clinical signs of disease

Fish infected with *A. salmonicida* do not necessarily show any clinical signs of disease; however, when fish become stressed or are compromised in some way, such that their immune system is lowered and a favorable condition within the fish is created for the pathogen, the infection can spread throughout the body and clinical signs can become visible (Cipriano et al. 1997; Hiney et al., 1997;

Hiney and Olivier, 1999; Austin and Austin, 2007; Noga, 2010). Typical clinical signs of the infection can include lethargy, lack of appetite, skin hyperpigmentation, boils and/or ulcers on the skin, lesions, internal hemorrhaging, enlargement of the spleen, septicemia and anemia (Fig. 2) (McCarthy, 1977; Ferguson and McCarthy, 1978; McCarthy and Roberts, 1980; Hiney et al., 1997; Hiney and Olivier, 1999; Austin and Austin, 2007; Noga, 2010).



**Figure 2.** Rainbow trout with signs of furunculosis. At the top: Boil and ulcer on the skin (photo by Morten Sichlau Bruun). To the left: Ulcer on the skin (Christensen, 1980). To the right: Enlargement of the spleen and hemorrhaging from internal organs (photo by Morten Sichlau Bruun).

### 2.3 Antibiotic treatment

In Denmark the antibiotics used in aquaculture have been sulfadiazine, trimethoprim, oxytetracycline and furazolidone (Dalsgaard et al., 1994) and since 1986, the only antibiotics licensed for use in aquaculture have been sulfadiazine/trimethoprim and oxolinic acid. The prevalence of antibiotic resistance genes (ARGs) among *A. salmonicida* in Denmark has been low, 5% in the study by Dalsgaard et al. (1994) and 9% in Manuscript III. Nevertheless, repeated treatment with antibiotics has proven to have many drawbacks, including induction of drug resistance in microorganisms, suppression of the immune system in fish, accumulation of residues in the fish, sediment and surrounding environment of the fish farms (Rijkers et al., 1981; Jacobsen and Berglund, 1988; Björklund et al., 1990; Aoki, 1997; Sørsum, 1998; Sørsum, 1999; Muziasari *et al.*, 2014). One major threat posed regarding antibiotic treatment is the ability of various genetic elements such as ARG

carrying plasmids (R plasmids) and integrons to disseminate multiple transferable ARGs (Aoki, 1997; L'Abée-Lund and Sørum, 2001; Berglund, 2015).

In the study by L'Abée-Lund and Sørum (2001), *A. salmonicida* and other bacteria originating from different locations around the world were investigated for the presence of a class 1 integron. Along with the integron, several ARGs were found: *aadA2*, *dfr16*, *aadA1*, *dfrIIc*, *qacG*, *orfD*, *tetA* and *tetE*, indicating that not only do class 1 integrons facilitate antibiotic resistance in marine environments, but also that ARGs can be transmitted between bacteria in various environments, since the found ARGs cassettes have also been associated with humans (L'Abée-Lund and Sørum, 2001). In agreement, Muziasari et al. (2014) found class 1 integrons and ARGs *sul1*, *sul2* and *dfrA1* in the sediment from farms located in the northern Baltic Sea and these same three ARGs, along with *aadA2* and *aadA1*, were also found in Danish *A. salmonicida* isolated from furunculosis outbreaks (Manuscript III). Kadlec et al. (2011) moreover found both class 1 integrons and ARGs against sulfonamide, trimethoprim and other antibiotics among *Aeromonas* species from Germany, where the only antibiotic therapy of fish is a combination of the two mentioned antibiotics.

The greatest concern with broad host range conjugative plasmids is that they can transfer ARGs across different bacterial genera and similar R plasmids have been isolated from separate ecological niches and across different environments (Sørum, 1998; L'Abée-Lund and Sørum, 2000; Sørum et al., 2003; Smillie et al., 2010). Sørum (1998) reported that after only 24 hours of mating between a fish pathogenic atypical *Aeromonas* carrying an R plasmid and *Escherichia coli*, the plasmid was directly transferred to every second *E. coli* cell. Direct transfer of the R plasmid from the atypical *Aeromonas* to human pathogens like *Salmonella enteritidis* and *Salmonella typhimurium* was also possible (Sørum, 1998). The atypical *Aeromonas* was also believed to be the origin of an R plasmid in *A. salmonicida* from a furunculosis outbreak (Sørum, 1998). Direct transfer of ARGs from pathogenic *A. salmonicida* to *E. coli* cells was also reported in the study by (Aoki et al., 1971). In the whole genome sequencing (WGS) study, none of the 101 *A. salmonicida* that were sequenced harbored any of the five investigated R plasmids (Manuscript III). However, eight *A. salmonicida* that also harbored multiple ARGs did show coverage (< 60%) of at least one of the R plasmids, indicating they could have acquired ARGs from the plasmids in the past through horizontal gene transfer and then subsequently lost the plasmid.

## 2.4 Vaccination

Unlike treatment with antibiotics, one does not have to worry about the bacterial pathogens developing resistance against vaccinations (Vinitnantharat et al., 1999), who provide a better alternative for future control of furunculosis. Immunization of fish against furunculosis by vaccine administration was already introduced experimentally in 1937, however, not until the early 1990's successful implementation of oil-adjuvanted vaccines in salmon aquaculture has there been made great advances in this field of research (Midtlyng, 1997). Fish can be immunized orally or by immersion or injection, though oral and immersion vaccines are less stressful for the fish than injection and would be preferred if their protection level would equal the one produced by injection (Vinitnantharat et al., 1999). Unfortunately, although many attempts have been made to produce an oral vaccine against furunculosis, thus far all have exhibited inconsistent antibody response and

protection (e.g. Krantz et al., 1964; Spence et al., 1965; Midtlyng et al., 1996), which in part could be caused by the fact that the vaccine components are poorly retained when administered orally (Press et al., 1996). Despite initially showing promising results (Cipriano et al., 1983; Johnson and Amend, 1984; Rodgers, 1990), immersion vaccines have also proved to be inadequate for use in aquaculture due to lack of long-term protection (Midtlyng et al., 1996; Midtlyng, 1997).

Over time, handling techniques improved for injection vaccines and automatic equipment, manuals and instruction videos for training became available, making it possible for injection vaccines to be administered on a large scale (Eithum, 1993; Midtlyng, 1997). Intraperitoneal injection of vaccines with oil adjuvant such as mineral oil, moreover induced much greater and longer protection compared to oral and immersion vaccines, making this vaccine administration superior to the others (Midtlyng, 1996; Midtlyng et al., 1996; Midtlyng, 1997). Even though numerous side effects for oil adjuvants have been observed, including lesions, pigmentation, granulomatous inflammation in the liver, autoimmune reactions and intra-abdominal adhesions, it is still recommended to use this administrative method to minimize loss of fish due to disease (Midtlyng, 1996; Midtlyng et al., 1996; Midtlyng, 1997; Håstein et al., 2005; Koppang et al., 2008; Satoh et al., 2011). The results of Mutoloki et al. (2006) using Atlantic salmon moreover showed that the combination of antigen and oil adjuvant is crucial and that it is their combined effect that is responsible for induction of a strong inflammatory reaction in the fish, thus highlighting the importance of choosing the correct antigen in order to develop an effective vaccine. Several antigen candidates have been suggested for stimulating early inflammatory reactions against *A. salmonicida*, which among others includes the A-layer, LPSs, IROMPs and ECPs (Midtlyng, 1997; Mutoloki et al., 2006).

The subspecies *A. salmonicida* is known to be very homogenous and this includes its virulence related A-layer proteins and LPSs located on the surface of the bacterium (Bjørnsdottir et al., 1992; Arnesen et al., 2010), which was also observed in Manuscript III. The combination of high similarity and location makes them very good antigen candidates and a positive correlation between vaccinating with A-layer proteins and protection by the immune system has been observed (Lund et al., 2003a; Arnesen et al., 2010). Research involving *A. salmonicida* IROMPs has shown that they also could have a potential as antigens included in vaccines, due to their *in vitro* bactericidal effect on both A-layer negative and positive *A. salmonicida* strains and *in vivo* protection of Atlantic salmon (Bricknell et al., 1999; O'Dowd et al., 1999). ECPs of *A. salmonicida* are already part of oil adjuvant vaccines against furunculosis, however, their contribution to the vaccine protection remains uncertain as studies using ECPs or their extracts as antigen showed varying results (Cipriano, 1982; Cipriano and Pyle, 1985; Prost, 2001). In fact, ECPs might not be important for inducing a protective immune responses (Lund et al., 2003a). There are even studies indicating that inclusion of ECPs in vaccines could have an adverse effect (Hirst and Ellis, 1994; Midtlyng et al., 1996; Lund et al., 2003a).

## **2.5 Transmission**

Furunculosis and its causative agent *A. salmonicida* have been investigated ever since the first discovery of furunculosis in 1894 by Emmerich and Weibel. Nevertheless the topic of transmission of *A. salmonicida* remains to be resolved. The primary focus and problem regarding transmission has

been carrier fish, whose existence was already indicated by several studies shortly after the discovery of furunculosis itself, by illustrating that the presence of *A. salmonicida* in fish does not necessarily lead to the development of furunculosis (e.g. Plehn, 1911; Mettam, 1915; Horne, 1928). Fish are thought to be capable of being infected with *A. salmonicida* for up to several months without showing any clinical signs of disease, in which time the infection is often said to be in a *latent* or *covert* phase and the fish are said to be “carriers” of the bacterium (Hiney et al., 1997).

Carrier fish have been recognized to play a significant role in the transmission of *A. salmonicida*, due to the ability of fish being able to shed bacteria in their surroundings and *A. salmonicida* being able to survive in water without a host (McCarthy, 1980; Rose et al., 1989b; Rose et al., 1990; Hastein and Lindstad, 1991; Nomura et al., 1992; Smith, 1992; Morgan et al., 1993; Nomura et al., 1993; Ogut and Reno, 2005). In a cohabitation study where Chinook salmon (*Oncorhynchus tshawytscha*) were injected with *A. salmonicida* and placed together with uninfected fish for ten days, prevalence of the bacterium among the initially healthy recipient fish was as high as 75% (Ogut and Reno, 2005). Mortality related to disease of the recipient fish moreover surpassed 50% and both bacterial concentrations in the water number of infected fish increased with time (Ogut and Reno, 2005). Rose et al. (1989b) studied Atlantic salmon experimentally infected with *A. salmonicida* in seawater and found bacteria were shed from dead and moribund fish at a high rate of  $10^5$  -  $10^8$  colony-forming units (CFU) per fish per hour.

Even though *A. salmonicida* initially has been thought of as an obligate pathogen not being able to survive in water without presence of fish (Popoff, 1984), others such as McCarthy (1980) found that the bacterium could survive up to 8 days. This was supported by the study of Nomura et al. (1992), where *A. salmonicida* was monitored in various type of water. In sterilized fresh water the bacterium could survive for sixty days while in non-sterile water, only around four days (Nomura et al., 1992). Survival in sea water was moreover shorter than in fresh water, though Nomura et al. (1992) concluded that *A. salmonicida* survival time in water was enough to infect other fish via this route. In a 21-day study of *A. salmonicida* in untreated lake water, Morgan et al. (1993) detected the bacterium in water samples by cultivation on TSA plates and with polymerase chain reaction (PCR) DNA amplification. Another interesting finding regarding survival of *A. salmonicida* in water is related to the increased hydrophobicity caused by the possession of the A-layer (Enger, 1997), whereby the A-layer positive *A. salmonicida* would become concentrated at the water surface and thus especially be a concern for the farms that use spray aeration. Nevertheless, even though no water samples were investigated, no gills were found positive from 20 fish showing no signs of disease that were sampled from three freshwater farms and one seawater farm in Denmark (Manuscript II).

Usually carrier fish start showing clinical manifestations of furunculosis when they become stressed, such as during rise in water temperature (Plehn, 1911; Bullock and Stuckey, 1975; McCarthy, 1977; Bernoth et al., 1997; Noga, 2010). This scenario is believed to be the key for transmission of *A. salmonicida* and cause of the furunculosis outbreaks among Danish fish farms. Initially, furunculosis was first observed by Rasmussen (1964) in freshwater, however, today outbreaks occur in seawater during periods of high temperature and especially causes great losses in sea reared rainbow trout production (Dalsgaard and Madsen, 2000; Pedersen et al., 2008). The theory is that fish at Danish



freshwater farms could be carriers of *A. salmonicida* and transfer the bacterium with them to the seawater farms, where the fish become stressed and develop the disease (Larsen and Mellergaard, 1981; Dalsgaard and Madsen, 2000; Pedersen et al., 2008). This theory is in agreement with results from Manuscript II, where most of the investigated fish were from a batch that was followed from freshwater to seawater. Some of these fish that showed no signs of disease at the freshwater farms were positive for *A. salmonicida* with quantitative real-time PCR (real-time PCR) and when transferred to a seawater farm, two furunculosis outbreaks occurred at the farm later on during high temperatures.

A different suggestion of *A. salmonicida* transmission has been that the bacterium is transmitted vertically (Wichardt et al., 1989). However, this was disputed through an extended study carried out by Bullock and Stuckey (1975), who followed carrier and artificially infected brood fish without finding any signs of vertical transmission. This notion was supported by McCarthy (1977), instead it was proposed that transmission is related to both contaminated equipment and infected fish. McCarthy (1977) and Rose et al. (1989b) also investigated transmission by feed through the gastrointestinal track, though without yielding any convincing results. Further proposed routes of transmission includes e.g. ciliated protozoans, who seemingly enhanced survival of *A. salmonicida* when the bacterium was co-cultured with it (King and Shotts, 1988).

## **2.6 Route of entry and colonization site(s)**

Although several suggestions have been made, there is currently still a great deal of uncertainty regarding both the route of entry and primary colonization site(s) of *A. salmonicida* and especially in carrier fish. The fins, gills, mucus layer as well as openings such as wounds have been proposed as possible entry sites for *A. salmonicida* (Hiney et al., 1997). In agreement, in an experimental study, artificially wounded Atlantic salmon infected with *A. salmonicida* showed higher mortality than salmon without the wounds (Svendsen and Børgwald, 1997). Svendsen and Børgwald (1997) further concluded that *A. salmonicida* can also adhere and most likely penetrate the mucous and skin of Atlantic salmon. The fins were suggested as a possible entry site for *A. salmonicida* by Hiney et al. (1994), who detected *A. salmonicida* in fins of pre-smolt Atlantic salmon. In another study with Atlantic salmon, the gills were indicated as a possible portal of entry into the fish (Svendsen et al., 1999). These findings are in congruence with Tatner et al. (1984) who investigated *A. salmonicida* infection in rainbow trout as well with results of Manuscript I, where the suggested main attachment sites for *A. salmonicida* after an experimental immersion were the dorsal and pectoral fins and gills.

In regards to possible colonization sites, there are several possible organs that have been investigated and proposed as the most probable site for *A. salmonicida* colonization. The first organ to be called the primary colonization site was the kidney (McCarthy, 1977; Popoff, 1984; Hiney et al., 1997). Though, *A. salmonicida* was also found in the gut as early as 1911 by Plehn. The gut was even the sole organ where *A. salmonicida* was found in naturally infected fish without showing clinical signs of disease by Willumsen (1990). By use of enzyme-linked immunosorbent assay (ELISA) *A. salmonicida* was also found in the gut by Rose et al. (1989b) and by use of PCR in the study of Gustafson et al. (1992). The bacterium was also found in the gut of experimentally infected rainbow trout in Manuscript I. Other organs such as the liver, heart, spleen or blood have also been used in

many studies as sampling sites for *A. salmonicida* (e.g. Daly and Stevenson, 1985; Cipriano, 1997; Hiney et al., 1997; Svendsen et al. 1999; Beaz-Hidalgo and Figueras, 2012). The spleen, along with the kidney, were also both positive for *A. salmonicida* by bacterial culturing in all experimentally infected fish in Manuscript I.

## 2.7 Susceptibility of fish species

Initially furunculosis was believed to be an exclusive disease of salmonids. Since then it has become known that *A. salmonicida* can also infect other fish species and other aquatic animals in freshwater and seawater e.g. catfish, carp, turbot, American eel, goby and wrasse (Bernoth et al., 1997). It has also become apparent that susceptibility to furunculosis varies among the host species (e.g. Plehn, 1911; Fish, 1937; McCarthy, 1977; Ellis and Stapleton, 1988; Perez et al., 1996). The first to document that species have different susceptibilities to *A. salmonicida* was Plehn (1911), who conducted experiments with infected rainbow trout and brown trout. The key result was that infected rainbow trout could remain unaffected and when placed in the same tank with brown trout and temperature was raised, brown trout developed disease and died, while the rainbow trout still remained unaffected (Plehn, 1911).


In general, fish belonging to the family Salmonidae are thought to be the most susceptible to furunculosis (McCarthy, 1977). Especially brown trout, brook trout (*Salvelinus fontinalis*) and Atlantic salmon have shown to be highly susceptible, while rainbow trout seemed to be more resistant as they needed to be wounded in a bath experiment before showing any signs of disease (McCarthy, 1977). The high degree of resistance which rainbow trout seem to possess against furunculosis compared to other farmed fish species was also illustrated by Cipriano et al. (1994a). Though it has to be mentioned that in their study, McCarthy and Roberts (1980) have argued that the presumed high susceptibility of salmonids to furunculosis might simply be related to the high degree of research that has been done on this family of fish due to their value as farmed fish.

Difference in susceptibility to furunculosis has been related to their immune system activity and especially their varying mucosal activity that is one of the main physical barriers and contains bioactive molecules such as lysosomes and other bacteriolytic enzymes (e.g. Cipriano and Heartwell, 1986; Cipriano et al., 1992; Cipriano et al., 1994a; Svendsen and Bøggwald, 1997). Teleost (bony) fish in general do exhibit a variation in their immune system wherein mucosal activity against pathogens is included (Dickerson, 2009). In agreement, a study by Cipriano and Heartwell (1986) showed that the fish species' mucus antibacterial activity directly correlated with their resistance towards furunculosis. This is further supported by results from Manuscript I, where the skin was not among the suggested primary attachment sites of *A. salmonicida* of rainbow trout that are known for their high resistance against *A. salmonicida*. Svendsen and Bøggwald (1997) also showed that mortality was higher for Atlantic salmon with an impaired skin mucous layer versus salmon with an intact mucous whereby it is indicated that skin mucous likely plays a role in the defense against *A. salmonicida*. Moreover, Cipriano et al. (1992) argued that it appears that several fish species actually lack an effective mucous protection layer against furunculosis.

## 2.8 *In vivo* imaging

One approach that could help elucidating the uncertainties about transmission, route of entry and colonization of *A. salmonicida*, as well as shed more light on the variation of species susceptibility to *A. salmonicida*, is tracking *A. salmonicida in vivo* in living fish. Tracking of *A. salmonicida* has been previously done by labeling the bacterium with radioactive isotopes (Svendsen et al., 1999). This resulted in findings such as a higher mortality in fish with artificial wounds and reduced epidermal mucus and also highlighted the importance of sampling time and the location of the bacterium. As an example, *A. salmonicida* was found in the blood after two hours, but not after 24 hours (Svendsen et al., 1999) .

In recent years much progress has been made regarding *in vivo* imaging and the two types that are most commonly used are bioluminescent and fluorescent reporters (Troy et al., 2004). Fluorescent proteins include the green fluorescent protein (GFP) (Chishima et al., 1997; Bouvet et al., 2002; Winnard et al., 2006) and DsRed (Baird et al., 2000; Dietrich and Maiss, 2002; Troy et al., 2004), however, there are numerous colors and near-infra red fluorescent dyes to choose from (Weissleder et al., 1999; Olenych et al., 2007; Day and Davidson, 2009; Filonov et al., 2011). Genes used in Bioluminescence imaging (BLI) originate from various luciferase proteins in bacteria, firefly, click beetles, and *Renilla* and the following components are involved in the light emission reaction: luciferase, luciferin, oxygen, and ATP (Troy et al., 2004). Although firefly and red click beetle luciferases are preferred in some studies due to their longer wavelength emission, bacterial luciferase is the only luciferase that does not require an injection of luciferin, which is a compound consisting of a long-chain aldehyde and a reduced riboflavin phosphate (FMNH<sub>2</sub>), in order to initiate the light-producing reaction (Troy et al., 2004). This is because the bacterial *lux* operon (*luxCDABE*) consists of five genes that encode both the luciferase enzyme and the aldehyde substrate (Fig. 3), while FMNH<sub>2</sub> is provided from the electron transport chain present in all bacteria (Troy et al., 2004; Lin and Meighen, 2009).



**Figure 3.** To the left: An illustration of genes of the bacterial *lux* operon. Genes *luxC*, *luxD*, and *luxE* code for a fatty acid reductase, a multicomplex enzyme that continuously supplies and regenerates the aldehyde substrate, while the genes *luxA* and *luxB* code for luciferase (Lin and Meighen, 2009). To the right: Bioluminescent *Photorhabdus luminescens* bacteria inside a nematode worm (Byrne, 2011).

When comparing the advantages and disadvantages of BLI versus fluorescence imaging, one has to take into consideration the type of imaging experiment one wants to execute. Luciferase proteins are also short lived, while fluorescence proteins are able to reside in the cell for hours (Burns et al. 2001; Troy et al. 2004), making fluorescence proteins more advantageous reporters for long term studies. Though, fluorescence proteins can take hours to become functional, while luciferase proteins mature rapidly following expression, making luciferase proteins more suited for shorter studies (Burns et al. 2001; Troy et al. 2004). Fluorescence imaging also has high “background noise”, meaning the animal itself emits fluorescence that interferes with the actual signal one is trying to visualize, while this kind of background noise is scarce in BLI (Burns et al. 2001; Troy et al. 2004). The background noise emitted by rainbow trout tissue in the immersion experiment conducted in Manuscript I was also scarce and could not be visualized. Luciferase proteins also have higher sensitivity and lower toxicity than fluorescence proteins (Burns et al. 2001; Troy et al. 2004). Bacterial luciferase is also an excellent choice for non-invasive studies, due to its ability of emitting light continuously without the need of adding any other substrate by e.g. an injection (Troy et al., 2004). However, although BLI three-dimensional techniques are available commercially (Virostko et al., 2008), BLI is usually used as a two-dimensional imaging technique and the spatial resolution is low compared to fluorescence imaging, making it difficult to separate photons produced by infected cells in two adjacent sites (Hutchens and Luker, 2007).

BLI has also been applied for monitoring *A. salmonicida* in Atlantic salmon by Ferguson et al. (1998). However, in these experiments an exogenous addition of the aldehyde substrate was needed in order to catalyze the light reaction and thereby visualize the bacterium. The incorporation of the luciferase genes into *A. salmonicida* moreover significantly lowered the virulence of the bacterium (Ferguson et al., 1998). Nevertheless, the experiment showed that *A. salmonicida* was shed from moribund and dead fish in the water column and were able to infect cohabitant fish, where the bacteria were mostly found in the gills and skin mucous. In recent studies of the fish pathogen *Edwardsiella ictaluri* by Karsi et al. (2006) and Menanteau-Ledouble et al. (2011), a bacterial luciferase *lux* operon from *Photobacterium luminescens* (Fig. 3) that emits light continuously by itself was used, enabling them to follow fish over several time points without the need for euthanization until the end of the experiments. Méndez and Guijarro (2013) also used this *lux* operon to successfully trace dissemination of *Yersinia ruckeri* in rainbow trout.

## 2.9 Manuscript I

Due to the fact that our focus for this study lay on the initial stages of the infection by *A. salmonicida*, meaning the route of entry and initial dissemination of *A. salmonicida* in fish, we choose to employ the same bacterial luciferase *lux* operon as the above mentioned authors in order to track *A. salmonicida* by *in vivo* BLI. This was done by transforming a highly virulent Danish *A. salmonicida* with a plasmid vector containing both GFP genes and the bacterial luciferase *lux* operon, which was subsequently used in immersion experiments where fish were followed over a 24-hour period. Although only luciferase was visualized in the immersion experiments, GFP enabled a practical way of visualizing the bacterial colonies through use of a fluorescence microscope. Results of these experiments showed that probable colonization sites of *A. salmonicida* are the gills and the dorsal and pectoral fins. The bacteria then progressed through the internal organs and seem to exit

via the anal opening. Modifications of this method are needed in order to attain more comprehensive knowledge regarding the route of entry and dissemination of *A. salmonicida*, nevertheless, this method does provide a possible tool for visualizing colonization of *A. salmonicida* and other bacterial pathogens in fish, as well as study host-pathogen interactions.

## Chapter 3: Detection

### 3.1 Detection of carrier fish

The main problem regarding identification of carrier fish has been the difficulty of detecting *A. salmonicida* within them (Bullock and Stuckey, 1975; Dalsgaard & Madsen 2000). Isolating *A. salmonicida* from fish showing signs of diseases has usually been done by bacterial culturing and the bacterium can also be detected in the fish by methods like histopathology, ELISA and PCR (Rose et al., 1989a; Gustafson et al., 1992; Dalsgaard and Madsen, 2000; Austin and Austin, 2007; Beaz-Hidalgo et al., 2013). There are several explanations for the struggle with detection of *A. salmonicida* and especially with bacterial culturing, which has been the most frequently used method for *A. salmonicida* detection throughout the years (Hiney et al., 1997; Austin and Austin, 2007). One of the two main notions is that the amount of *A. salmonicida* within the carriers might be too low for colonies being able to grow (Hiney et al., 1994). Another possibility is that the bacterium within carrier fish can be present in a non-culturable state, i.e. they are viable but non-culturable cells (VBNC) (Morgan et al., 1993; Ferguson et al., 1995; Nășcuțiu, 2010). The presence of VBNC *A. salmonicida* has, however, been a controversial subject due to the scepticism towards being able to revive these cells after their non-culturable state. Presence of VBNC *A. salmonicida* in fish would, nevertheless, correlate with the scenario in Denmark, where it has not been possible to detect the bacterium by culturing in rainbow trout from freshwater farms, but furunculosis outbreaks still occur during elevated temperatures after these fish are transferred out to seawater farms (Dalsgaard & Madsen 2000; Pedersen *et al.* 2008).

In order to improve detection of *A. salmonicida* in carrier fish, various methods have been applied including heat stress of fish, injection of corticosteroid into fish and pre-enrichment steps for bacterial culturing. The first to employ a stress test in order to discover carriers was Plehn (1911). In an experimental infection with *A. salmonicida*, fish infected with the bacterium were stressed by temperature increase, whereby some fish started showing signs of disease and soon thereafter died (Plehn, 1911). Though, Bullock and Stuckey (1975) reported that although heat stress by increase of temperature to 18°C did cause clinical signs of the disease and subsequently high mortality among carrier fish, detection from the fish still remained very low and the best way to increase detection in carriers was a combination of corticosteroid injection and heat stress, termed stress-inducible furunculosis (SIF) tests (Smith, 1991; Cipriano et al., 1997). The drawback of this method is that injected fish have to be held in heated tanks for about two weeks. Even though the method was modified in 1977 by McCarthy, the SIF tests are still time consuming and also require sacrifice of many fish for obtaining statistical significance (Hiney et al., 1994). Nevertheless, the high reliability of this method has consequently made it the primary examination method of salmonids in European aquaculture (Smith, 1991). In the study by Cipriano et al. (1997), performance of a SIF test was compared to another method suggested for usage of enabling detection of carriers, namely employing pre-enrichment steps before culturing (Daly and Stevenson, 1985; Cipriano et al., 1997). Though, statistical analysis revealed that the SIF test was more reliable than both pre-enrichment and direct culturing (Cipriano et al., 1997). One factor, however, needs to be considered when applying SIF tests to salmon, which is timing the test close to smoltification (Scallan and Smith, 1993).

### 3.2 PCR

In order to avoid the SIF tests, yet maintain the possibility of detecting carriers, molecular methods such as PCR assays have been developed for detecting *Aeromonas salmonicida* subspecies (Gustafson et al., 1992; Hiney et al., 1992; O'Brien et al., 1994; Mooney et al., 1995; Byers et al., 2002a, b; Altinok et al., 2008; Beaz-Hidalgo et al., 2008b). PCR has proven to be a powerful tool for amplification of nucleic acids, whereby a DNA sequence selected for amplification is exponentially increased in repeated cycles of synthesis by a thermostable DNA polymerase and two oligonucleotide primers that each hybridize to one strand of the double-stranded target DNA (Saiki et al., 1985; Saiki et al., 1988). This is usually followed by visualization of the amplified DNA by gel electrophoresis and a known DNA probe is used for determining size of the amplicon.

One of the first *Aeromonas salmonicida* subspecies specific PCRs was developed by Hiney et al. (1992), based on a DNA fragment (GenBank accession number X64214) of a 6.4 kb *A. salmonicida* cryptic plasmid. Sensitivity of detection for pure culture was two cells (Hiney et al., 1992) and no false positives were amplified when tested with numerous typed and clinical isolates of related aeromonads and other bacterial genera (Hiney et al., 1992). This target DNA was later used by multiple authors (e.g. Morgan et al., 1993; O'Brien et al., 1994; Byers et al., 2002a, b; Altinok et al., 2008) for detecting *Aeromonas salmonicida* subspecies from various fish tissues, feces and water. In all the studies no false positives were obtained by testing non-target bacterial DNA, however, not all *Aeromonas salmonicida* subspecies were identified. This is because the cryptic target plasmid that now has been identified as *A. salmonicida* plasmid pAsal1 sequenced by Fehr et al. (2006) (GenBank accession number AJ508382), is not universally present in all *A. salmonicida* (Nielsen et al., 1993; Attéré et al., 2015) and can be lost by culturing at above 22 - 25°C (Daher et al., 2011; Tanaka et al., 2012; Attéré et al., 2015). Though, in the 99 sequenced *A. salmonicida* that were isolated from furunculosis outbreaks in Denmark in Manuscript III, only 52% seemed to harbor pAsal1 and all were grown at 20°C.

Another target gene that has been used by more than one author is the *vapA* gene, which encodes the A-protein of the A-layer and was initially used as a PCR target by Gustafson et al. (1992). Sensitivity for detecting *Aeromonas salmonicida* subspecies in fish tissue was 10 CFU mg<sup>-1</sup> and only 1 CFU ml<sup>-1</sup> in pure culture. Nevertheless, it was concluded that enrichment steps were necessary for detecting potential carrier fish in order to avoid false negatives. As with the pAsal1 target, not all bacterial isolates were identified with use of the *vapA* target due to mutations in the *vapA* gene (Gustafson et al., 1992; Byers et al., 2002b). The pAsal1 target and *vapA* target were combined in the study by Byers et al. (2002b) and they correctly identified 93% and 94% of the *Aeromonas salmonicida* subspecies respectively. When used together, the two PCR assays identified 99% of the isolates (Byers et al., 2002b), though in another study conducted by the same authors, it was concluded that bacterial culturing was more reliable for detecting carrier fish than the PCR method and if PCR were to be used, pre-enrichment steps would be necessary (Byers et al., 2002a).

The chromosome encoded *fstA* gene has also been used as a PCR target in the study of Beaz-Hidalgo et al. (2008b). What separated this PCR from the others, according to Beaz-Hidalgo et al. (2008b), was that blood and mucus were used as sampling sites, making the PCR assay a non-destructive

diagnostic tool. Detection limit for experimentally infected mucus and blood samples were  $2.5 \times 10^2$  and  $1.5 \times 10^5$  CFU ml<sup>-1</sup> respectively. It is thought that the low sensitivity in blood samples was caused by heparin or other blood component interference that are able to compete with bacterial DNA in the assay. Nevertheless, when tested on potential carriers, 31 wild salmon with no signs of disease, six salmon were detected by the PCR while the culture method only detected one fish (Beaz-Hidalgo et al., 2008b).

### 3.3 Real-time PCR

The conventional method of PCR revolutionized molecular techniques in science, nevertheless, the fact remains that this method is laborious, difficult to automate, needs enrichment steps for enabling detection of carriers and is usually semi-quantitative at best (Kubista et al., 2006; Bustin et al., 2012). This is because conventional PCR is restrained by reagents and reaches a plateau where the product amount cannot be increased any longer (Bustin et al., 2012). This plateau varies from assay to assay and since the PCR is an endpoint assay analyzed after the reaction has reached its linear phase, the gel electrophoresis analysis shows roughly the same amount of DNA that was produced by the end of the PCR reaction regardless of what the initial amount of DNA was (Bustin et al., 2012). Although the same amplification principles apply to the real-time PCR method, as the name reveals, this method monitors the amplification process in “real time” and can be used for precise quantitative analysis by fluorescent reporter molecules (Higuchi et al., 1992; Kubista et al., 2006; Bustin et al., 2012). For real-time PCR, there are also different detection chemistries to choose from and the two major ones are either an intercalating dye such as SYBR green (Morrison et al., 1998) or a hydrolysis probe such as Taqman (Gibson et al., 1996). Both chemistries have their advantages and disadvantages though they share the same design, which is to generate fluorescence during the PCR reaction that is monitored in “real time”. Apart from enabling quantitative application, real-time PCR is also convenient, robust, simple, fast, sensitive and adapted to high throughput analysis; as a consequence it has become the most widely used molecular technique (Kubista et al., 2006; Bustin et al., 2012)

Real-time PCR has been used in several studies to detect *Aeromonas salmonicida* subspecies directly from fish tissue (e.g. Balcazar et al., 2007; Goodwin and Merry, 2009; Keeling et al., 2012; Gulla et al., 2015). In the study by Balcazar et al. (2007) a real-time PCR assay design was developed, which combined low costs with high sensitivity (Balcazar et al., 2007; Nazarenko et al., 2002a, 2002b). The assay was based on Light Upon eXtension (LUX) primer probes originally described by Nazarenko et al. (2002a, 2002b). The primers targeted the same *A. salmonicida* DNA sequence (gene *aopP*) located on the pAsal1 plasmid as originally developed by Hiney et al. (1992) for conventional PCR. In agreement with all previous PCR studies that had implemented this target, Balcazar et al. (2007) did not find any false positives. Though, on the contrary to the bulk of the previous studies, Balcazar et al. (2007) did obtain 100% correct identification of 16 isolates of both typical and atypical *Aeromonas salmonicida*. Balcazar et al. (2007) stressed that the reason behind this result is that the isolates used in his study were all from various disease outbreaks indicating that the presence of pAsal1 in the bacterium could be related to virulence of the bacterium. This hypothesis is supported by research of Goodwin and Merry (2009) who also used the same target for a real-time PCR assay to detect atypical *Aeromonas salmonicida* and also obtained 100% amplification of this species. In



our Manuscript II, all of the 20 tested *A. salmonicida* isolates from furunculosis outbreaks were also correctly identified by the real-time PCR originally developed by Balcazar et al. (2007). Nonetheless, the theory was later disputed by our research where 99 Danish *A. salmonicida* isolated from furunculosis outbreaks, were subjected to WGS and it was found that the AopP protein sequence encoded on pAsal1 was missing in 50% of the *A. salmonicida* isolates (Manuscript III).

In the two real-time PCR assays developed by Keeling et al. (2012) and Gulla et al. (2015) the same target sequence (gene *vapA*) was used as originally applied for conventional PCR by Gustafson et al. (1992). The assay by Keeling et al. (2012) was based on a molecular beacon and the assay was found to have 100% analytical specificity and an analytical sensitivity of  $5 \pm 0$  fg in pure culture and  $2.2 \times 10^4 \pm 1 \times 10^4$  CFU g<sup>-1</sup> for kidney tissue. Though, with enrichment steps as used in earlier research, the sensitivity for tissue increased to  $40 \pm 10$  CFU g<sup>-1</sup>. This assay was later modified by Gulla et al. (2015) for detecting atypical *Aeromonas salmonicida* from head kidney of cleaner fish in Norway, due to existence of multiple types of the A-layer caused by a single base variation resulting in a mismatch with the forward primer developed by Keeling et al. (2012). This variation in the A-layer was most likely responsible for why 6% of the *Aeromonas salmonicida* subspecies investigated in the study by Byers et al., (2002a) were not amplified by the *vapA* primers. After this modification, Gulla et al. (2015) obtained 100% analytical specificity of all presently recognized A-layer types of this species and a sensitivity of 7 - 8 bacterial genomes in pure culture and  $1.1 \times 10^3$  CFU mL<sup>-1</sup> in tissue.

### 3.4 Manuscript II

In order to develop a molecular method that would enable specific and highly sensitive detection directly from multiple fish tissue (even potential carrier fish), the previously designed real-time PCR by Balcazar et al. (2007) was tested on spleen, kidney, intestine, gill and brain tissues from 40 rainbow trout (n=200), sampled from three Danish freshwater farms and one seawater farm where no signs of disease were observed and one seawater farm after two furunculosis outbreaks. Prevalence of *A. salmonicida* obtained by the real-time PCR was compared to the one obtained by bacterial culturing. The real-time PCR was chosen primarily due to the bulk of available previous research that have used the target and the fact that the plasmid pAsal1 and its derivatives are ColE2-type replicons, meaning they are high copy number plasmids (Lilly and Camps, 2015) making the assay highly sensitive. Moreover, all 20 Danish *A. salmonicida* that were tested were amplified by the assay. The real-time PCR showed a better result than culturing (65% vs. 30% positive fish by real-time PCR and culturing, respectively). The assay also indicated possible presence of carrier fish harbouring VBNC *A. salmonicida* in Danish fresh- and seawater rainbow trout farms and that the spleen, brain and intestine, where the bacterium was detected by real-time PCR and not culturing in fish showing no signs of disease, could play an important role in *A. salmonicida* infection and persistence of VBNC. This means that in order to detect possible carriers, one should strive for testing several tissues from fish instead of just one (usually the kidney). Moreover, sensitivity of the assay was 1 - 2 genomic units per reaction and the real-time PCR had a high reproducibility and an excellent efficiency, thus providing a sensitive tool for detection of *A. salmonicida*. However, the major drawback of the assay, which was exemplified by our later findings in Manuscript III by WGS of 99 *A. salmonicida*, is that the target plasmid is absent in some *A. salmonicida* and seemingly half

of the representative Danish *A. salmonicida* population and the absence does not necessarily have to be associated with lack of virulence nor culturing above 22 - 25°C as indicated in earlier studies. In order to be certain of avoiding false negatives, another sensitive detection method with a different target would therefore need to be employed.

## Chapter 4: Epidemiology

### 4.1 Epidemiology and genetic variation

One major factor for establishing an effective prevention strategy, besides enabling effective treatment, is to determine the epidemiology and genetic variation of the Danish *A. salmonicida*. Numerous molecular techniques have been applied for deciphering the epidemiology and genetic variation of *A. salmonicida* and generally the results support the notion of *A. salmonicida* being a highly homogenous subspecies as previously indicated by its phenotypical characteristics (Toranzo et al., 1991; Dalsgaard et al., 1994; Nielsen et al., 1994a).

One of the first studies using a molecular method for investigating genetic similarity of *A. salmonicida* was Belland and Trust (1988). Eleven *A. salmonicida* isolated from different species of fish and geographical places around the world were subjected to a DNA:DNA reassociation technique using a radiolabeled probe and the resulting mean DNA sequence similarity was  $97 \pm 6.1\%$  (Belland and Trust; 1988). Other molecular methods have included restriction endonuclease fingerprinting analysis (McCormick et al., 1990), randomly amplified DNA polymorphism fingerprinting analysis (RAPD) (Miyata et al., 1995; Inglis et al., 1996; O'hici et al., 2000), amplified fragment length polymorphism (AFLP) (Lund et al., 2003b), plasmid profiling (Nielsen et al., 1993; Sørum et al., 1993), and ribotyping (Nielsen et al., 1994b). Although Nielsen et al. (1994b) and O'hici et al. (2000) did find some clustering using ribotyping and RAPD respectively; it was not enough to enable use of these methods for epidemiological studies except for maybe certain local geographical areas.

Pulsed-field gel electrophoresis (PFGE) has for a long time been the 'gold standard' for typing bacterial isolates, however, even this method did not have enough resolution for studying epidemiology of *A. salmonicida* (Garcia et al., 2000; O'hici et al., 2000; Cunningham and Colquhoun, 2002; Beaz-Hidalgo *et al.*, 2008). Garcia et al. (2000) examined 132 isolates from Denmark, Norway, Scotland, Ireland, the Faroe Islands, France, Canada and the USA and although PFGE was reportedly more useful to differentiate *A. salmonicida* than ribotyping, only slight differences were found and overall results suggested that a single clone of *A. salmonicida* was responsible for most of the worldwide furunculosis outbreaks. This theory is supported by findings of Nash et al. (2006). In order to identify virulence genes and possible vaccine candidates, Nash et al. (2006) developed a microarray-based comparative genomic hybridization (M-CGH) technique based on virulence associated genes from the genome sequence of *A. salmonicida* strain A449 (Reith et al., 2008). Nash et al. (2006) compared *A. salmonicida* and other *Aeromonas* species isolated from various fish species and geographic locations and found no correlation between host or geographic origin and the M-CGH patterns and a relatively low number of divergent genes in the *A. salmonicida* strains.

## 4.2 MLST-v

Currently, one of the most widely used molecular methods for typing microbial isolates is multilocus sequence typing (MLST) due to its great sensitivity, specificity and ease of use and data exchange (Enright and Spratt, 1998; Maiden et al., 1998; Platonov et al., 2000; Maiden, 2006; Martino et al., 2011; Martinez-Murcia et al., 2011). MLST is based on identifying polymorphic sites within DNA sequence fragments of multiple housekeeping genes. For each gene fragment, every sequence that differs is given its own unique allele no matter if they differ at one or several nucleotide sites. Each isolate is then characterized based on the combination of the alleles from all the gene fragments, whereby they can each be assigned a specific allelic profile or sequence type (ST) number (Maiden et al., 1998).

In the present thesis a modified MLST based on eight housekeeping genes (*dnaJ*, *rpoD*, *groL*, *gyrB*, *metG*, *ppsA*, *gltA* and *recA*) in combination with four virulence associated genes (*aexT*, *eno*, *vapA* and *fstB*) (Table 1) was applied on a preliminary representative collection of 23 Danish *A. salmonicida* isolates, one Scottish strain and the *A. salmonicida* type strain NCIMB 1102 in order to study their genetic variation using the publicly available WGS *A. salmonicida* strain A449 as reference (Reith et al., 2008) (Table 2). This kind of modified MLST, usually called MLST-v, has been used in previous studies where the method has proven to have a higher discriminatory power than MLST and has revealed important genetic information regarding virulence associated genes (Zhang et al., 2004; Chen et al., 2007; Doijad et al., 2014; Tankouo-Sandjong et al., 2007). A detailed materials and methods section for the MLST-v used in this thesis is described in Chapter 5: Methodological considerations.


The MLST-v scheme with allelic profiles for all *A. salmonicida* isolates was created, however, only five unique STs were identified (Table 2). The pattern of evolutionary descent of *A. salmonicida* based on the STs was also analyzed using eBURST (<http://eburst.mlst.net/3.asp>). Analysis showed that all isolates belong to the same clonal complex (CC), since no isolate differed by more than a single allelic mismatch. Nineteen of the Danish isolates and the type strain NCIMB 1102 belonged to the primary founder ST (ST 1) (Fig. 4A), which is the ST with the most single-locus variants (SLVs) (<http://eburst.mlst.net/3.asp>). This was supported with a bootstrap confidence level of 88% (percentage based on 1000 replicates). A Bayesian phylogeny tree based on the concatenated sequences from *A. salmonicida* MLST-v was also constructed, displaying the high homogeneity of the *A. salmonicida* (Fig. 4B). In conclusion, all analysis showed that the MLST-v was not an adequate tool for studying the epidemiology and genetic variation of Danish *A. salmonicida*. The results support the high genetic homogeneity of *A. salmonicida* found in previous studies using other typing methods (e.g. Belland and Trust; 1988; Beaz-Hidalgo et al., 2008; Nash et al., 2006).

**Table 1. Information about primers used for PCR amplification and sequencing of housekeeping and virulence associated genes for development of a MLST-v scheme.** The following is depicted in table form left to right: Primer names for each gene, nucleotide sequence for each primer, brief description of the protein product that the respective genes encode, length of the PCR amplicon of each gene fragment in base pairs (bp), length of the target sequence located within the PCR product of each gene fragment that is used for the MLST-v scheme, annealing temperature for the respective primer pairs used during PCR amplification, and reference article for each primer including this thesis as reference if primers were designed during the present thesis.

Primer	Sequence (5'-3')	Gene product	Length of PCR amplicon (bp)	Length of target sequence (bp)	Annealing temperature	Reference
<i>gyrB</i> -F	CATGTCTACGAGCA GACCTA	DNA gyrase (type II topoisomerase), subunit B	926	682	54 °C	(Martinez-Murcia et al., 2011)
<i>gyrB</i> -R	CTCCACGTTTCAGGA TCTTGCC					
<i>rpoD</i> -F	GAAGGCCGAAATCG ACATCGC	RNA polymerase, sigma 70 (sigma D) factor	700	649	55 °C	(Martinez-Murcia et al., 2011)
<i>rpoD</i> -R	ATGCTCATGCGRCG GTTGAT					
<i>groL</i> -F	CAAGGAAGTTGCTT CCAAGG	Chaperonin GroEL	782	604	57 °C	(Martino et al., 2011)
<i>groL</i> -R	CATCGATGATGGTG GTGTTC					
<i>dnaJ</i> -F	CGAGATCAAGAAG GCGTACAAG	Chaperone Hsp 40, co-chaperone with DnaK	934	814	54 °C	(Martinez-Murcia et al., 2011)
<i>dnaJ</i> -R	CACCACCTTGCACA TCAGATC					
<i>recA</i> -F	GCTGGGTCTAGATTG AAAAGC	Recombinase A	640	635	57 °C	(Martinez-Murcia et al., 2011)
<i>recA</i> -R	CTCGCCGTTATAGC TGTACC					
<i>gltA</i> -F	TTCGGTCTGCTCTC CAAGAT	Citrate synthase I	626	373	57 °C	(Martino et al., 2011)
<i>gltA</i> -R	TTCATGATGATGCC GGAGTA					
<i>metG</i> -F	TGGCAACTGATCCT CGTACA	Methionyl-tRNA synthetase	657	539	57 °C	(Martino et al., 2011)
<i>metG</i> -R	TCTTGTTGGCCATC TCTTCC					
<i>ppsA</i> -F	AGTCCAACGAGTA CGCCAA	Phosphoenolpyruvate synthase	619	502	61 °C	(Martino et al., 2011)
<i>ppsA</i> -R	CTCGGCCAGATAG AGCCAGGT					
<i>aexT</i> -F	TGCAGATTC AAGC AAACACC	ADP-ribosylating toxin	689	608	61 °C	This thesis
<i>aexT</i> -R	GCCAGCAACTTCTG CCTTTA					
<i>eno</i> -F	CGCCGACAACAAC GTCGACATC	Enolase	598	518	56 °C	(Martino et al., 2011)
<i>eno</i> -R	CTTGATGGCAGCCA GAGTTTCG					
<i>vapA</i> -F	CAACGGTTTCATTG TGTTGG	A-layer	630	556	56.5 °C	This thesis
<i>vapA</i> -R	TTGAAGGCAGAAA CATCACC					
<i>fstB</i> -F	GTTTTCCCGCTTTTC CTTGA	Ferric siderophore receptor B	594	512	58 °C	This thesis
<i>fstB</i> -R	GAAGATGCTGCGTT TGCTC					

**Table 2. Overview of *A. salmonicida* isolates used in the MLST-v and their respective sequence type (ST) and allelic profile.** For *A. salmonicida* isolates that were also used for whole genome sequencing (WGS), the same name is used as in manuscript III: The Danish isolates have a black color (freshwater farms) or a blue color (seawater farms) and are labeled by region of origin followed by year of isolation, with abbreviations Nj = Northern Jutland, Mj = Central Jutland, Sd = Southern Denmark, Sj = Zealand. The two Danish isolates that were not included in WGS are named Denmark followed by their year of isolation. There are five STs. Alleles *gltA*, *metG* and *vapA* are the only three alleles that vary in their sequence composition in at least one base pair (bp) among the isolates.

<i>A. salmonicida</i> isolate	ST	Alleles											
		<i>dnaJ</i>	<i>gltA</i>	<i>groL</i>	<i>gyrB</i>	<i>metG</i>	<i>ppsA</i>	<i>recA</i>	<i>rpoD</i>	<i>aexT</i>	<i>eno</i>	<i>vapA</i>	<i>fstB</i>
Denmark 1985	1	1	1	1	1	1	1	1	1	1	1	1	1
Mj13 1987	1	1	1	1	1	1	1	1	1	1	1	1	1
Mj16 2008	1	1	1	1	1	1	1	1	1	1	1	1	1
Sd3 1982	1	1	1	1	1	1	1	1	1	1	1	1	1
Sd6 (a) 2013	1	1	1	1	1	1	1	1	1	1	1	1	1
Mj21 1993	1	1	1	1	1	1	1	1	1	1	1	1	1
Mj5 1986	1	1	1	1	1	1	1	1	1	1	1	1	1
Sd1 2004	1	1	1	1	1	1	1	1	1	1	1	1	1
Sd2 1982	1	1	1	1	1	1	1	1	1	1	1	1	1
Sd2 1995	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj2 (a) 1993	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj2 (b) 1993	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj3 (b) 2009	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj5 1994	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj5 1995	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj6 (a) 1993	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj6 (a) 1996	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj6 (b) 1993	1	1	1	1	1	1	1	1	1	1	1	1	1
Sj6 (b) 1996	1	1	1	1	1	1	1	1	1	1	1	1	1
NCIMB 1102 (Type strain 1962)	1	1	1	1	1	1	1	1	1	1	1	1	1
Denmark 2009	2	1	1	1	1	1	1	1	1	1	1	2	1
Mj18 1986	2	1	1	1	1	1	1	1	1	1	1	2	1
Sj4 1982	2	1	1	1	1	1	1	1	1	1	1	2	1
Sd6 (b) 2013	3	1	1	1	1	1	1	1	1	1	1	3	1
Scotland	4	1	2	1	1	1	1	1	1	1	1	1	1
Reference A449 (France 1975)	5	1	1	1	1	2	1	1	1	1	1	1	1



**Figure 4.** (A) Pattern of evolutionary descent of *A. salmonicida* STs from the MLST-v, as shown by eBURST. Nineteen of the Danish isolates along with the reference type NCIMB 1102 belong to ST 1. The three Danish isolates Denmark 2009, Mj18 1986 and Sj4 1982 belonged to ST 2, the Danish isolate Sd6 (b) 2013 belongs to ST 3, the Scottish isolate belongs to ST 4 and A449 belongs to ST 5. Same isolate labels are used as in Table 2. (B) Bayesian phylogeny tree based on the concatenated sequences from *A. salmonicida* MLST-v. The *A. salmonicida* isolates have a clonal population structure. The three Danish isolates belonging to ST 2 form a monophyletic group with a bootstrap value of 57%. Same isolate labels are used as in Table 2.

### 4.3 Whole genome sequencing

Previous molecular methods such as RAPD, AFLP, PFGE, MLST and MLST-v focus on a small part of the genome, while WGS can avoid such bias and thus provides the best overview for studying a population (Foxman et al., 2005; Parkhill and Wren, 2011). Since the first time bacterial genomes were completely sequenced in 1995 (Fleischmann et al., 1995; Fraser et al., 1995), WGS has gone through immense progress. The most notable factor for this progression was development and implementation of next-generation sequencing (NGS) techniques, initially introduced by Roche 454 Life Sciences in 2005 (Henson et al., 2012), which reduced the price and increased the speed of sequencing by reducing the average read length compared to sanger-sequencing (Rothberg and Leamon, 2008; Metzker, 2010; Land et al., 2015; Loman and Pallen, 2015). As a result, the number of sequenced genomes has skyrocketed in recent years (Land et al., 2015; Loman and Pallen, 2015). Although this has been accompanied by an increase in the recommended coverage needed for a genome assembly and number of contigs needing closure before a genome could be completed (Land et al., 2015), Illumina sequencing platform seems to be a cost-effective, since assemblies can be generated that are almost complete genomes (Mavromatis et al., 2012). Due to the relatively low costs along with its high accuracy, yield of error-free reads and percentage base calls above Q30

(Henson et al., 2012; Ross et al., 2013), the Illumina sequencing platform is currently the most widely used WGS technique (Metzker, 2010; Land et al., 2015).

NGS technology has also proved to be a powerful tool in microbiology for studying epidemiology of bacterial outbreaks (Hiller et al., 2007; Shendure and Ji, 2008; Pandya et al., 2009; Lewis et al., 2010; Leekitcharoenphon et al., 2012; Bertelli and Greub, 2013; Salipante et al., 2015; Stucki et al., 2015). The advantage with WGS lies with its high discriminatory power that enables deciphering the evolution of bacterial isolates belonging to the same clonal lineage, whereby highly robust phylogenies can be generated and the origin and routes of transmission of infections can be revealed (Parkhill and Wren, 2011). WGS even revealed the patient-to-patient pattern of transfer of a bacterial pathogen within a hospital (Harris et al., 2010). In the study by Salipante et al. (2015) PFGE was compared to WGS and although the result for relatedness of the bacteria by both methods correlated with each other, the resolution of WGS was highly superior. The same was concluded by Leekitcharoenphon et al. (2012) when investigating the epidemiology of *Salmonella enterica* using both PFGE and WGS. Leekitcharoenphon et al. (2012) moreover compared several WGS bioinformatics approaches for the same data and determined that nucleotide difference and single nucleotide polymorphism (SNP) tools were superior to the other bioinformatics methods. Indeed, the SNP approach has been frequently used for WGS based epidemiological studies (e.g. Pandya et al., 2009; Lewis et al., 2010; Stucki et al., 2015). In agreement, WGS in combination with the SNP analysis was able to distinguish among 101 sequenced *A. salmonicida*, using the published genome of the French strain A449 (Reith et al., 2008) as reference (Manuscript III).

Recently genome sequence information of numerous bacterial fish pathogens has also become available from genome sequencing projects, which is believed to become a vital part in finding new intervention strategies against bacterial infections in fish (Sudheesh et al., 2012). The genome of *A. salmonicida* isolate (A449) from a brown trout in the Eure river in France was sequenced by Reith et al. (2008). This has provided insight into the genomics of *A. salmonicida* that could prove to be vital for discerning the pathogen's evolution and infection progress (Reith et al., 2008). The study provided basic genetic information regarding the strains chromosome size of 4,702,402 base pairs (bp) and its plasmid profile consisting of two large plasmids pAsa4 and pAsa5 and three small plasmids pAsa1, pAsa2 and pAsa3. All of these plasmids, except for pAsa4, were also present among the majority of the *A. salmonicida* isolates sequenced in Manuscript III, where plasmids pAsa1 and pAsa6 were also additionally present in many isolates. Though, one of the key findings by Reith et al. (2008) included identification of functional genes encoded on plasmid pAsa5 and the chromosome for T3SS that has been shown to be vital for virulence in *A. salmonicida* as described in detail in chapter 1. Most likely due to rearrangements in pAsa5, both T3SS effector proteins and structural proteins were, however, absent in 24% of the isolates in Manuscript III. T3SS genes are also absent in the genome sequenced *A. hydrophila* ATCC 7966T and are only present on the chromosome in other *A. hydrophila* (Sha et al., 2005). The genome of A449 moreover contains numerous of virulence associated genes including the A-layer, toxins, secreted enzymes, iron acquisition genes, quorum sensing genes and ARGs (Reith et al., 2008) that were also found among the sequenced *A. salmonicida* in Manuscript III. Comparisons with the genome of *A. hydrophila* ATCC 7966T (Seshadri et al., 2006) also highlighted their respective evolution associated with possible adaptation to their fish hosts (Reith et al., 2008). Nevertheless, two Danish isolates from



brown trout, the French strain A449 also isolated from brown trout, one Scottish strain and the type strain NCIMB 1102 isolated from Atlantic salmon all showed high SNP similarity with the 97 Danish *A. salmonicida* isolated from rainbow trout in the WGS study of Manuscript III.

#### **4.4 Manuscript III**

Due to the lacking ability of MLST-v to distinguish among the homogeneous *A. salmonicida* isolates and the above mentioned advantages of WGS, we sequenced 99 Danish *A. salmonicida* isolated from different geographical regions and years (1980 - 2014), the Scottish strain MT004 and the type strain NCIMB 1102 from England. WGS was successfully able to distinguish among the *A. salmonicida* isolates, though it revealed that *A. salmonicida* is very homogenous, since only 667 SNPs were found among the isolates within the 4,702,402 bp long sequence of the chromosome. Bayesian temporal phylogenetic reconstruction showed that four major introductions of *A. salmonicida* into Denmark have occurred. The introductions correlate with the initial expansion of Danish rainbow trout production and the beginning of production in seawater. There is also a possibility that the bacterium might have initially been transmitted from seawater to freshwater. We moreover found some variation in plasmids and virulence factors, especially those encoded on plasmids and nine *A. salmonicida* harbored worldwide known ARGs against several antibiotics. Overall, our WGS analysis provided valuable information regarding epidemiology as well as genetic and virulence variations among the Danish disease causing *A. salmonicida* population.

## Chapter 5: Methodological considerations

### 5.1 Materials and methods for MLST-v

#### Bacterial isolates

The following *A. salmonicida* isolates were selected for development a MLST-v scheme: 23 Danish *A. salmonicida* isolated between year 1982 and 2013 from various fresh- and seawater rainbow trout farms, one Scottish *A. salmonicida* isolated from Atlantic salmon and the type strain NCIMB 1102 isolated from Atlantic salmon (Table 2). The publicly available whole genome sequence of the French isolate A449 isolated from brown trout was used as a reference.

#### Primers

Primers for amplification of *A. salmonicida* gene fragments from eight housekeeping genes (*dnaJ*, *rpoD*, *groL*, *gyrB*, *metG*, *ppsA*, *gltA* and *recA*) and four virulence associated genes (*aexT*, *eno*, *vapA* and *fstB*) were either developed during the present thesis or obtained from previous publications (Martinez-Murcia et al., 2011; Martino et al., 2011). The list of housekeeping and virulence associated genes and all primers used for PCR amplifications and sequencing is listed in Table 1.

#### DNA extraction

All *A. salmonicida* were grown in VIB (Difco) for 48 h at 20°C. DNA was extracted using Qiagen QIAamp DNA mini kit (Qiagen) according to the manufacturer's protocol and stored at -20°C until further PCR amplification.

#### PCR reaction

PCR reaction was carried out in T3000 Thermocycler (Biometra) using a final volume of 50 µL containing: 5 µL GeneAmp® 10X Gold Buffer (150 mM Tris-HCl, pH 8.0, 500 mM KCl), 5 µL 25mM MgCl<sub>2</sub>, 1 µL 10µM deoxyribonucleotide (dNTP) mix, 1 µL of each forward and reverse primer (10 µM), 0.3 µL of AmpliTaq Gold® DNA Polymerase (5U µL<sup>-1</sup>) (Applied Biosystems), 34.7 µL of sterile water and 2 µL of 5ng DNA genomic DNA as the template. The reaction mixture was subjected to denaturation at 94°C for 3 min, followed by 35 cycles of amplification as follows: denaturation at 94°C for 1 min, annealing temperature depending on the primer (table 2 and 3) for 1 min and extension at 72°C for 1.30 min. At the end a final extension step at 72°C was achieved for 3 min.

#### Sequencing

PCR Amplified products were analyzed by electrophoresis on 2% agarose E-gels (Invitrogen) and visualized on a UV transilluminator. The products were then purified using the High Pure PCR Product Purification kit (Roche Applied Science, Germany), following the manufacturer's instructions. Nucleotide sequences for the MLST-v analysis were determined by bidirectional

sequencing using the Big Dye Terminator V3.1 Ready Reaction Cycle Sequencing Kit in ABI 3130 Genetic Analyzer (Applied Biosystems, USA), according to the manufacturer's instructions. Each gene fragment was sequenced using the same forward and reverse primer pair as used for PCR amplification.

### **Data processing and analysis**

Consensus sequences for twelve gene fragments from the *A. salmonicida* isolates used in the MLST were assembled via CLC Workbench software 6.5 (CLC Bio-Qiagen, Aarhus, Denmark). Alignment of all consensus sequences for each gene was done using the Clustal X program (Larkin et al., 2007). Trimming of alignments was done in Bioedit (Hall, 1999). All nucleotide variable positions and insertions/deletions found in the alignments were checked manually in raw sequence chromatograms by use of BioEdit (Hall, 1999) and CLC Workbench software 6.5. By use of nucleotide blast (Altschul *et al.*, 1990), all gene fragments (alleles) were found for the published isolate *A. salmonicida* A449 (Reith et al., 2008). Trimmed and edited sequences, including the *A. salmonicida* A449 sequences, were used to create a new MLST-v scheme for *A. salmonicida* in MLST plugin application in CLC workbench software 6.5 according to the program guidelines. Allele sequences that differed from each other by one or more polymorphisms were attributed to a unique allele number. Each unique allelic profile, as defined by the allele numbers of the twelve loci, was assigned a ST number. The same ST was assigned to isolates that shared the same allelic profile.

### **Analysis by eBURST**

Pattern of evolutionary descent of 25 *A. salmonicida* isolates and the reference strain A449 was analyzed using eBURST (<http://eburst.mlst.net/default.asp>). The program uses allelic profiles to identify potential CCs and founders by linking SLVs or double-locus variants (Feil et al., 2004). Parameters for the analysis were set to stringent default setting, meaning all STs had to be SLVs of some other ST in the population, which is illustrated as a diagram with a single CC i.e. a group where all STs are linked (<http://eburst.mlst.net/3.asp>). Here the parameter was eleven, since twelve alleles (gene fragments) were used in the MLST-v.

### **Bayesian phylogenetic tree**

Concatenated MLST-v sequences for 26 *A. salmonicida* isolates, including the reference strain A449, were subjected to Bayesian phylogenetic inference using Markov chain Monte Carlo (MCMC) methods in the software package MrBayes (Ronquist et al., 2012) available at: <http://mrbayes.csit.fsu.edu/download.php>. MCMC chains were simulated for one million generations with subsampling every 100 generations and an 'invgamma' molecular evolution model that allows a gamma distribution across positions of the alignment with invariable sites.

## **5.2 *In vivo* imaging (Manuscript I)**

Before the experimental infection of rainbow trout of this study was carried out, the following was investigated: 1) the lowest possible immersion time of fish in the diluted *A. salmonicida* *gfplux*

culture that would enable attachment and thus visualization of the bacterium on the fish for the first time point and 2) the difference between luminescence emittance of infected and non-infected fish i.e. the fish autoluminescence. Eight rainbow trout were experimentally infected in three liters of  $5 \times 10^7$  CFU ml<sup>-1</sup> *A. salmonicida gfplux* culture in well-aerated  $10 \pm 1^\circ\text{C}$  aquariums at our institute. Two fish were euthanized the following time points using an overdose of 3-aminobenzoic acid ethyl ester: 30 min, 1 hour, 2 hours and 3 hours post infection. All euthanized fish were washed twice in distilled water, dipped on a paper towel, and finally laid on a sterile Tissue Culture Dish (Greiner Bio-One, Germany). Fish were then transported to the Danish Cancer Institute in their respective culture dishes, where they were placed in the dark collection chamber of the IVIS spectrum imaging workstation (PerkinElmer) for bioluminescence image capture (30 s). One euthanized non-infected control fish was also scanned to compare autoluminescence emitted by non-infected fish with luminescence emitted from *A. salmonicida gfplux* infected fish. Luminescence emission from *A. salmonicida gfplux* was observed on all eight infected fish at each of the time points 30 min - 3 hours post challenge at one or several of the following sites: dorsal fin, pectoral fin, gills, oral and nasal cavity, and the eyes. While the attachment site patterns varied among individual fish, the most intensive luminescence emission was observed from fish infected for 2- and 3 hours respectively. Based upon this result, 2 hours of immersion time in *A. salmonicida gfplux* diluted culture was determined to be sufficient for the bacterium to attach to the fish and enable visualization of the bacterium in the IVIS. Measured autoluminescence from the non-infected control fish was scarce and could not be visualized, compared to the visualized intense luminescence emitted by the *A. salmonicida gfplux* from infected fish.

It would be recommended for future studies that an experimental infection set-up, similar as to the one described in Manuscript I, would be carried out. The differences would include fish being immersed for only 30 min and in lower concentrations of *A. salmonicida gfplux* in order to better mimic natural infections and the fish would be followed for a shorter and longer duration than 24 hours respectively, in order to allow visualization of the internal organs at various time points. Though, the details and optimal settings for the additional infection experiments would need to be investigated first. Only if the above mentioned experimental infections would be successfully employed, then one could also consider testing a co-habitation challenge using this method, which would be the most optimal way to mimic natural conditions of infection. The instability of the plasmid vector carrying the GFP and luciferase genes would also have been investigated more closely. One possibility would have been to do the transformation procedure of *A. salmonicida* over again in order to see if a more stable *A. salmonicida* transformant could be made. Another option could have been to try treating the fish with ampicillin before start of the experimental infection and thereby create a favorable environment inside the fish for *A. salmonicida gfplux* i.e. *A. salmonicida* carrying the plasmid vector with GFP, luciferase and ampicillin resistance.

### **5.3 Real-time PCR (Manuscript II)**

Initially, several extraction methods for extracting *A. salmonicida* DNA from seeded tissue were tested, as well as different amounts of tissue in order to avoid inhibition of the real-time PCR assay. The three extraction methods that were tested extensively were Maxwell® 16 LEV Blood DNA Purification Kit (Promega), QIAamp DNA Mini Kit (Qiagen) and InstaGene Matrix (Bio-Rad). The

Maxwell and QIAamp kits are both standard kits at our institute and have been used for extraction from various tissues and fluids of fish and other animals with slightly modified versions of the manufacturers' procedures. However, even after exhaustive testing with various modifications e.g. overnight lysis, different buffer and proteinase K combinations and lowering the amount of all tissues to 10 mg, there were still problems with the extractions and especially the spleen. The main issue with the QIAamp DNA Mini Kit was that the filters seemed to become clogged by the tissue even after extended lysis procedures. Difficulties extracting with the Maxwell® 16 LEV Blood DNA Purification Kit included carryover of particles during the extraction process, lower than expected DNA yield and inhibition of real-time PCR. Briefly the Maxwell® 16 Tissue DNA Purification Kit was also tested, though the same problems occurred as with the previous Maxwell kit. The InstaGene Matrix kit, which has been used for extracting DNA from fish tissues in a number of previous PCR and real-time PCR studies e.g. (Balcazar et al., 2007; Keeling et al., 2012; Beaz-Hidalgo et al., 2013), gave the most optimal results and was thus used henceforth.

Two other previously developed real-time PCR assay by Goodwin and Merry (2009) and Keeling et al. (2012) respectively, were also tested before continuing with the assay developed by (Balcazar et al., 2007). The assay by Goodwin and Merry (2009) included three different PCR primers used for conventional PCR assays in the study by Byers et al. (2002b). One of the primers had the same target (plasmid pAsal1) as the real-time PCR primers of Balcazar et al. (2007), though the main difference was that Goodwin and Merry (2009) did not change the PCR primers even though they amplified long (421 - 512 bp) DNA sequences, which is a standard size for conventional PCR amplicons, though the recommended optimal amplicon size for real-time PCR is usually 50 - 200 bp. This might have contributed to the fact why we were not able to reproduce the results by Goodwin and Merry (2009), as the primers did not amplify pure cultures of various *A. salmonicida* strains. When testing the assay by Keeling et al. (2012) with primers targeting the *vapA* gene, similar results were obtained as with the assay by Goodwin and Merry (2009). The only assay that showed promising results was the one originally developed by Balcazar et al. (2007) and was thus pursued further.

Given the current knowledge obtained by WGS of the 99 Danish *A. salmonicida* isolates in Manuscript III, a different target gene than *aopP* encoded on pAsal1 would have been used for the real-time PCR assay. Due to limited amount of time, developing and implementing new primers was not possible. Nevertheless, if new primers were to be developed in the future, it would be essential to obtain the same high sensitivity as the one obtained in this real-time PCR assay.

#### **5.4 Whole genome sequencing (Manuscript III)**

In retrospect, due to the inability of the developed MLST-v to distinguish among the Danish *A. salmonicida*, the best solution would have been to avoid this method and instead have moved straight to WGS. Plasmid profiling of the 101 sequenced *A. salmonicida* isolates would also be recommended in the future for enabling a better comparison with previous studies of *A. salmonicida* plasmids.

## Chapter 6: Discussion and future perspectives

### 6.1 Discussion

From a broader perspective, research conducted within the framework of this PhD project aimed at contributing to ongoing efforts towards resolving the current concerns of furunculosis in Danish rainbow trout production. As discussed in the following paragraphs, investigations of three key matters regarding this grave fish disease, done throughout this thesis, have contributed novel insights into each of these subjects.

The route of entry and dissemination of *A. salmonicida* was investigated by tracking the bacterium *in vivo* using BLI in Manuscript I. Although uncertainties regarding this topic remain, the *in vivo* study shed light on the initial host-pathogen relationship between rainbow trout and *A. salmonicida*. The key findings were that *A. salmonicida* mostly seemed to initially colonize the gills and the dorsal and pectoral fin and move on quite rapidly (within 24 hours) to internal organs such as the intestine, spleen and kidney. These findings, although more elaborate studies are needed to establish any firm conclusions, are in agreement with several previous studies regarding *A. salmonicida* (e.g. Willumsen, 1990; Hiney et al., 1997; Svendsen et al., 1999) and moreover highlight some key issues that need to be resolved in order to comprehend the initial stages of the *A. salmonicida* infections.

One of the important issues, stressed by the fact that the skin was not one of the suggested primary colonization sites of *A. salmonicida* in this study, was that fish species seem to have different levels of resistance to *A. salmonicida* that are correlated with the immune system and especially the mucosal activity of the species (McCarthy, 1977; Cipriano et al., 1994a; Dickerson, 2009). More attention should be paid to this important topic, especially since some studies have argued that the mucus of skin could be the best sampling site for carrier fish (Svendsen and Bøgwald, 1997). This might be true for more susceptible fish species like Atlantic salmon (McCarthy, 1977), but does not necessarily apply to rainbow trout due to their high resistance and the indicated lack of skin colonization observed in Manuscript I. Granted, the results from the *in vivo* study cannot be compared to carrier infections due to the high amount of bacteria used ( $5 \times 10^7$  CFU ml<sup>-1</sup> *A. salmonicida* *gflux* culture) for the immersion and non-natural infection conditions, nevertheless, it emphasizes that progression of *A. salmonicida* in fish is not uniform in all species, meaning one cannot rely on certain detection and prevention strategies for *A. salmonicida* being applicable for all farmed fish if only certain species have been investigated.

Another notable result in Manuscript I was the fast dissemination of *A. salmonicida* into the internal organs of the fish. Again it is important to emphasize that the fish were immersed in a high concentration of the bacterium, which could have contributed to this rapid dissemination. Nonetheless, it is an intriguing observation supported by other studies concerning digestion of various feed (e.g. Windell *et al.* 1969) that one should bear in mind in future research regarding this bacterium. Returning to the subject of carrier fish, this rapid dissemination pattern might also provide insight into possible colonization sites of the bacterium in carrier fish. If the pathogen is only attached to the outer surface of the fish for a short period of time, at least concerning rainbow trout,

one could thus speculate whether the focus should be shifted towards sampling of internal organs. This suggestion is supported by the real-time PCR findings of Manuscripts II, where *A. salmonicida* in 20 rainbow trout without signs of disease was found only in internal organs (the brain, spleen and intestine) and not in the gills. These findings also highlight the importance of sampling from more than one organ in order to detect *A. salmonicida*, whereby we move on to the second key subject of this thesis, namely development of a highly sensitive method for detection of the bacterium. On one hand this was accomplished, since the real-time PCR enabled detection of 1 - 2 genomic units per reaction and showed more sensitive detection of *A. salmonicida* than the culturing method. The difference in sensitivity was even more profound in fish without signs of disease (possible carriers), where only one out of 20 fish was found positive by culturing, while six fish were positive with real-time PCR. The major drawback of the real-time PCR, limiting the future applicability of this method for future field investigations in at least Denmark, is the fact that 50% of the representative *A. salmonicida* population in Denmark was missing the target sequence of the assay (Manuscript III). Previous studies have shown that the absence of the target (plasmid pAsal1) can be caused by culturing conditions above 22 - 25°C (Daher et al., 2011) and it has been argued that the absence could be related to lack of virulence of the bacterium (Balcazar et al., 2007). However, it was revealed by WGS in Manuscript III that these two conditions are not necessarily associated with the absence of pAsal1, since all the *A. salmonicida* isolates were cultured at 20°C and were isolated from furunculosis outbreaks.

One should nevertheless, keep in mind the two key factors that were highlighted through the findings of Manuscript II, which were the importance of sampling from multiple organs when trying to detect *A. salmonicida* and the indication of possible carriers being present in Danish freshwater farms. The most noteworthy aspects regarding the topic of sampling sites for *A. salmonicida*, is that the intestine might be a colonization site for *A. salmonicida* as indicated by findings in Manuscript I and that the brain could also be a possible colonization site. Both organs, along with the spleen, were the only positive organs in potential carrier fish, while the intestine is rarely used for sampling and to the best of our knowledge the brain has never been investigated in any previous studies of *A. salmonicida*. The indication of possible presence of carrier fish in Danish freshwater farms supports the widespread theory of freshwater to seawater (via carrier fish) *A. salmonicida* transmission in Denmark. In congruence, this transmission pattern of *A. salmonicida* was seen in Manuscript III for isolates harboring ARGs. Nevertheless, the four major introductions of *A. salmonicida* into Denmark that correlate with the initial expansion of Danish rainbow trout production and the beginning of production in seawater, indicated the possibility of the bacterium initially being transmitted from seawater to freshwater (Manuscript III). This brings us to the last key investigation of this thesis, namely obtaining knowledge on the epidemiology and genetic and virulence variation of the Danish disease causing *A. salmonicida*.

Although the Bayesian temporal phylogenetic reconstruction based on the SNPs obtained from WGS gave a general insight into the epidemiology of the Danish *A. salmonicida* population, it was difficult to find specific geographical correlations between the local fish farms, which might have been caused by a trade of fish for anglers and between individual farms. While WGS also proved to be a valuable tool for distinguishing among the homogenous Danish *A. salmonicida* isolates, overall it still supported the notion of high homogeneity, since only 667 SNPs were found among all 101

isolates (two from other countries). This finding also explained why almost all *A. salmonicida* used in the initially developed MLST-v had identical sequences.

Variation of *A. salmonicida* seemed mainly to be related to the plasmid profiles and virulence factors encoded on these plasmids, of which T3SS related virulence proteins seemed to be the most variable ones. This was associated with the high instability of pAsa11 and pAsa5 on which most of the T3SS virulence proteins are encoded. Though, *A. salmonicida* can harbor plasmids that share the same virulence genes e.g. plasmid pAsa6 that was included in the study (Manuscript III) and that harbor insertion sequences that can cause rearrangements and enable possible transfer of genes between plasmids and the chromosome (Daher et al., 2011; Dallaire-Dufresne et al., 2014). This intertwined system of virulence-associated genes makes it difficult to state anything concrete regarding this topic, without a more comprehensive analysis of the T3SS proteins. What can be suggested from the findings of Manuscript III regarding T3SS and the previously shown significance of its structural genes for *A. salmonicida* virulence (Burr et al., 2002; Burr et al., 2003; Stuber et al., 2003; Burr et al., 2005; Dacanay et al., 2006; Froquet et al., 2007), is that this theory needs to be investigated more thoroughly because the sequences encoding some of the structural proteins (e.g. *ascV*) were absent in 24% of the 101 sequenced *A. salmonicida*. The isolates that were missing these structural proteins have not been subjected to any *in vivo* studies that could provide information regarding their virulence or possible lack thereof, however, these *A. salmonicida* were isolated from furunculosis outbreaks. One can speculate whether this group of isolates could have lost these protein-coding sequences during culturing, since stressful culture conditions can cause rearrangements in plasmid pAsa5 whereby T3SS related genes can be lost (Stuber et al., 2003; Tanaka et al., 2012; Dallaire-Dufresne et al., 2014). On the contrary to this theory, all the Danish *A. salmonicida* were cultured at 20°C.

Another significant finding was that nine *A. salmonicida* harbor worldwide known ARGs against several antibiotics, among these were ARGs against trimethoprim and sulphonamide that are licensed antibiotics for treatment in Danish aquaculture. The prevalence of these ARGs was low (9%), nevertheless, there is a clear transmission pattern of some of the isolates harboring the ARGs from a freshwater farm to several seawater farms. The ARGs found in the Danish isolates have also been found in *Aeromonas* species isolated from different environments around the world (L'Abée-Lund and Sørum, 2001; Sørum et al., 2003; Kadlec et al., 2011; Muziasari et al., 2014), supporting the evidence of the widespread dissemination of ARGs. This also highlights the need for developing an effective vaccine. Based on the findings of Manuscript III, the A-layer protein and GCAT protein sequences seem to be highly homogeneous among the Danish *A. salmonicida*, which warrants further investigation into their potential use for a vaccine against *A. salmonicida*. On the other hand, further studies are needed to identify any specific isolate(s) and/or virulence factor(s) that could be recommended in implementation of a vaccine.

## 6.2 Conclusion and future perspectives

The bacterium *A. salmonicida* was discovered over 100 years ago, however, there are still many questions regarding this pathogen and the disease furunculosis that remain unanswered. The findings



of this project have provided some valuable information for research on *A. salmonicida*, however, there is much that could be improved and further elaborated.

The best way to gain more knowledge on the host-pathogen relationship of *A. salmonicida* is through *in vivo* imaging, a valuable approach that is rapidly advancing. Although fluorescent and bioluminescent reporters are still the most frequently used reporters for imaging, their limitations of resolution and range of depth prevent the possibility of obtaining a detailed picture of the host-pathogen relationship. Notably, one emerging method that promises to revolutionize imaging and surpasses both of the above methods in resolution and range of depth is photoacoustic imaging, which uses ultrasound waves for imaging (Xu and Wang, 2006). Thus far this technology has only been used for human biomedical research; however, in the near future this imaging method could become available in the veterinary field.

Another detection method for *A. salmonicida* that would have the same high sensitivity as the real-time PCR assay in Manuscript II, but would enable 100% detection of all *A. salmonicida*, could be developed by changing the target of the present assay. One possible target could be the high-copy number plasmid pAsa1 that thus far seems to be universally present and stable in all *A. salmonicida* (Att  r   et al., 2015).

Much data has been obtained by WGS of the 101 *A. salmonicida* isolates and only a fraction of this data has been utilized for analysis thus far. Indeed, much more valuable and in-depth knowledge could be found by applying some of the available bioinformatics tools, a promising goal for which the dataset created by our WGS analyses can provide a solid foundation.

Taking all the obtained results throughout this thesis into consideration, this PhD project has contributed novel and insightful information further promoting the current research on Danish disease causing *A. salmonicida* as well as the bacterium in general. There still seems to be a challenging ‘road ahead’ when trying to prevent furunculosis in the Danish rainbow trout production. On the positive side, one can hope that the findings of this project will inspire future research on *A. salmonicida* and that eventually prevention of furunculosis will be successful before even greater losses are seen due to the constant increase in intensity of farming combined with the major disease-promoting environmental stress factor – the overall raising global temperature.

## References

- Altinok, I., Capkin, E., and Kayis, S. (2008). Development of multiplex PCR assay for simultaneous detection of five bacterial fish pathogens. *Vet. Microbiol.* 131, 332–338. doi:10.1016/j.vetmic.2008.04.014.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. *J. Mol. Biol.* 215, 403–10. doi:10.1016/S0022-2836(05)80360-2.
- Anderson, D. P. (1972). Virulence and persistence of rough and smooth forms of *Aeromonas salmonicida* inoculated into coho salmon (*Oncorhynchus kisutch*). *J. Fish. Res. Board Canada* 29, 204–206. doi:10.1139/f72-034.
- Aoki, T. (1997). “Resistance plasmids and the risk of transfer,” in *Furunculosis: Multidisciplinary fish disease research*, eds. E. M. Bernoth, A. E. Ellis, P. J. Midtlyng, G. Olivier, and S. P (London, UK: Academic Press), 433 – 440.
- Aoki, T., Egusa, S., Kimura, T., and Watanabe, T. (1971). Detection of R factors in naturally occurring *Aeromonas salmonicida* strains. *Appl. Microbiol.* 22, 716–7. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=376391&tool=pmcentrez&rendertype=abstract>.
- Arkwright, J. (1912). An epidemic disease affecting salmon and trout in England during the summer of 1911. *J. Hyg. (Lond).* 12, 391–413.
- Arnesen, K. R., Mikkelsen, H., Schröder, M. B., and Lund, V. (2010). Impact of reattaching various *Aeromonas salmonicida* A-layer proteins on vaccine efficacy in Atlantic cod (*Gadus morhua*). *Vaccine* 28, 4703–4708. doi:10.1016/j.vaccine.2010.04.100.
- Attéré, S. A., Vincent, A. T., Trudel, M., Chanut, R., and Charette, S. J. (2015). Diversity and homogeneity among small plasmids of *Aeromonas salmonicida* subsp. *salmonicida* linked with geographical origin. *Front. Microbiol.* 6, 1274.
- Austin, B. (1993). Recovery of “atypical” isolates of *Aeromonas salmonicida*, which grow at 37 degrees C, from ulcerated non-salmonids in England. *J. Fish Dis.* 16, 165–168. doi:10.1111/j.1365-2761.1993.tb00862.x.
- Austin, B., and Austin, D. A. (2007). *Bacterial Fish Pathogens: Diseases of Farmed and wild fish*. 4th ed. Chichester, UK: Springer.
- Austin, B., and Austin, D. A. (2012). “Aeromonadaceae Representative (*Aeromonas salmonicida*),” in *Bacterial Fish Pathogens: Diseases of Farmed and Wild Fish* (Springer), 147–228.
- Austin, D. A., McIntosh, D., and Austin, B. (1989). Taxonomy of Fish Associated *Aeromonas* spp., with the Description of *Aeromonas salmonicida* subsp. *smithia* subsp. nov. *Syst. Appl. Microbiol.* 11, 277–290. doi:http://dx.doi.org/10.1016/S0723-2020(89)80026-8.
- Baird, G. S., Zacharias, D. A., and Tsien, R. Y. (2000). Biochemistry, mutagenesis, and oligomerization of DsRed, a red fluorescent protein from coral. *Proc. Natl. Acad. Sci. U. S. A.* 97, 11984–9. doi:10.1073/pnas.97.22.11984.
- Balcazar, J. L., Vendrell, D., de Blas, I., Ruiz-Zarzuola, I., Girones, O., and Luis Muzquiz, J. (2007). Quantitative detection of *Aeromonas salmonicida* in fish tissue by real-time PCR using self-quenched, fluorogenic primers. *J. Med. Microbiol.* 56, 323–328. doi:10.1099/jmm.0.46647-0.
- Beaz-Hidalgo, R., and Figueras, M. J. (2012). “Molecular Detection and Characterization of Furunculosis and Other *Aeromonas* Fish Infections,” in *Health and Environment in Aquaculture*, eds. E. D. Carvalho, G. S. David, and R. J. Silva (InTech), 97–132. doi:10.5772/2462.
- Beaz-Hidalgo, R., and Figueras, M. J. (2013). *Aeromonas* spp. whole genomes and virulence factors implicated in fish disease. *J. Fish Dis.* 36, 371–388. doi:10.1111/jfd.12025.
- Beaz-Hidalgo, R., Latif-Eugenín, F., and Figueras, M. J. (2013). The improved PCR of the *fstA* (ferric siderophore receptor) gene differentiates the fish pathogen *Aeromonas salmonicida* from other *Aeromonas* species. *Vet. Microbiol.* 166, 659–663.

doi:<http://dx.doi.org.globalproxy.cvt.dk/10.1016/j.vetmic.2013.06.028>.

- Beaz-Hidalgo, R., López-Romalde, S., Toranzo, A. E., and Romalde, J. L. (2008a). Polymerase chain reaction amplification of repetitive intergenic consensus and repetitive extragenic palindromic sequences for molecular typing of *Pseudomonas anguilliseptica* and *Aeromonas salmonicida*. *J. Aquat. Anim. Health* 20, 75–85. doi:10.1577/H07-007.1.
- Beaz-Hidalgo, R., Magi, G. E., Balboa, S., Barja, J. L., and Romalde, J. L. (2008b). Development of a PCR protocol for the detection of *Aeromonas salmonicida* in fish by amplification of the *fstA* (ferric siderophore receptor) gene. *Vet. Microbiol.* 128, 386–394. doi:10.1016/j.vetmic.2007.10.004.
- Belland, R. J., and Trust, T. J. (1988). DNA:DNA reassociation analysis of *Aeromonas salmonicida*. *J. Gen. Microbiol.* 134, 307–15. doi:10.1099/00221287-134-2-307.
- Berglund, B. (2015). Environmental dissemination of antibiotic resistance genes and correlation to anthropogenic contamination with antibiotics. *Infect. Ecol. Epidemiol.* 5, 28564. doi:10.3402/iee.v5.28564.
- Bernoth, E., Ellis, A. E., Midtlyng, P. J., Olivier, G., and Smith, P. eds. (1997). *Furunculosis: Multidisciplinary fish disease research*. London UK: Academic Press.
- Bertelli, C., and Greub, G. (2013). Rapid bacterial genome sequencing: methods and applications in clinical microbiology. *Clin. Microbiol. Infect.* 19, 803–13. doi:10.1111/1469-0691.12217.
- Björklund, H., Bondestam, J., and Bylund, G. (1990). Residues of oxytetracycline in wild fish and sediments from fish farms. *Aquaculture* 86, 359–367. doi:10.1016/0044-8486(90)90324-G.
- Björnsdóttir, R., Eggset, G., Nilsen, R., and Jørgensen, T. O. (1992). The A-layer protein of *Aeromonas salmonicida* further characterization and a new isolation procedure. *J. Fish Dis.* 15, 105–118. doi:DOI 10.1111/j.1365-2761.1992.tb00645.x.
- Boone, D., Castenholz, R., and Garrity, G. (2001). *Bergey's Manual of Systematic Bacteriology*. 2nd ed. , ed. G. Garrity New York: Springer.
- Bouvet, M., Wang, J., Nardin, S. R., Nassirpour, R., Yang, M., Baranov, E., et al. (2002). Real-time optical imaging of primary tumor growth and multiple metastatic events in a pancreatic cancer orthotopic model. *Cancer Res.* 62, 1534–1540.
- Brenner, D., Krieg, N., Garrity, G., Staley, J., and (Firm), K. (2005). *Bergey's Manual of Systematic Bacteriology. Volume Two, The Proteobacteria*. New York: Springer.
- Bricknell, I., King, J., Bowden, T., and Ellis, A. (1999). Duration of protective antibodies, and the correlation with protection in Atlantic salmon (*Salmo salar* L.), following vaccination with an *Aeromonas salmonicida* vaccine containing iron-regulated outer membrane proteins and secretory polysacch. *Fish Shellfish Immunol.* 9, 139–151. doi:10.1006/fsim.1998.0183.
- Bullock, G. L., and Stuckey, H. M. (1975). *Aeromonas salmonicida*: detection of asymptotically infected trout. *Progress. Fish Cult.* 37, 237–239.
- Burns, S. M., Joh, D., Francis, K. P., Shortliffe, L. D., Gruber, C. A., Contag, P. R., et al. (2001). Revealing the spatiotemporal patterns of bacterial infectious diseases using bioluminescent pathogens and whole body imaging. *Contrib. Microbiol.* 9, 71–88.
- Burr, S. E., Pugovkin, D., Wahli, T., Segner, H., and Frey, J. (2005). Attenuated virulence of an *Aeromonas salmonicida* subsp. *salmonicida* type III secretion mutant in a rainbow trout model. *Microbiology* 151, 2111–2118. doi:10.1099/mic.0.27926-0.
- Burr, S. E., Stuber, K., Wahli, T., and Frey, J. (2002). Evidence for a Type III Secretion System in *Aeromonas salmonicida* subsp. *salmonicida*. *J. Bacteriol.* 184, 5966–5970. doi:10.1128/JB.184.21.5966-5970.2002.
- Burr, S. E., Wahli, T., Segner, H., Pugovkin, D., and Frey, J. (2003). Association of Type III secretion genes with virulence of *Aeromonas salmonicida* subsp. *salmonicida*. *Dis. Aquat. Organ.* 57, 167–171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14735937>\nfile:///C:/Users/Joanna/Documents/my PhD/Literature/Burr et al 2003 aeromonas salm.pdf.

- Bustin, S. A., Zaccara, S., and Nolan, T. (2012). "An Introduction to the Real-time Polymerase Chain Reaction," in *Quantitative Real-time PCR in Applied Microbiology*, ed. M. Filion (Norfolk, UK: Academic Press), 3 – 25.
- Byers, H. K., Cipriano, R. C., Gudkovs, N., and Crane, M. S. (2002a). PCR-based assays for the fish pathogen *Aeromonas salmonicida*. II. Further evaluation and validation of three PCR primer sets with infected fish. *Dis. Aquat. Org.* 49, 139–144.
- Byers, H. K., Gudkovs, N., and Crane, M. S. (2002b). PCR-based assays for the fish pathogen *Aeromonas salmonicida*. I. Evaluation of three PCR primer sets for detection and identification. *Dis. Aquat. Organ.* 49, 129–138. doi:10.3354/dao049129.
- Byrne, J. (2011). *Photobacterium luminescens*: The Angel's Glow. *Naked Sci. (University Cambridge)*. Available at: <http://www.thenakedscientists.com/HTML/articles/article/angel-glow/> [Accessed March 7, 2016].
- Böhm, K. H., Fuhrmann, H., Schlotfeldt, H. J., and Körting, W. (1986). *Aeromonas salmonicida* from salmonids and cyprinids - serological and cultural identification. *J. Vet. Med. Ser. B* 33, 777–783. doi:10.1111/j.1439-0450.1986.tb00099.x.
- Chacón, M. R., Soler, L., Groisman, E. A., Guarro, J., and Figueras, M. J. (2004). Type III secretion system genes in clinical *Aeromonas* isolates. *J. Clin. Microbiol.* 42, 1285–1287. doi:10.1128/JCM.42.3.1285-1287.2004.
- Chapman, P. F., Cipriano, R. C., and Teska, J. D. (1991). Isolation and phenotypic characterization of an oxidase-negative *Aeromonas salmonicida* causing furunculosis in coho salmon (*Oncorhynchus kisutch*). *J. Wildl. Dis.* 27, 61–7. doi:10.7589/0090-3558-27.1.61.
- Chart, H., Shaw, D. H., Ishiguro, E. E., and Trust, T. J. (1984). Structural and immunochemical homogeneity of *Aeromonas salmonicida* lipopolysaccharide. *J. Bacteriol.* 158, 16–22.
- Chen, Y., Zhang, W., and Knabel, S. J. (2007). Multi-virulence-locus sequence typing identifies single nucleotide polymorphisms which differentiate epidemic clones and outbreak strains of *Listeria monocytogenes*. *J. Clin. Microbiol.* 45, 835–846. doi:10.1128/JCM.01575-06.
- Chishima, T., Miyagi, Y., Wang, X., Baranov, E., Tan, Y., Shimada, H., et al. (1997). Metastatic patterns of lung cancer visualized live and in process by green fluorescence protein expression. *Clin. Exp. Metastasis* 15, 547–552. doi:10.1023/A:1018431128179.
- Christensen, N. O. (1980). *Fiskesygdomme*. Copenhagen: Den Kgl. Veterinær- og Landbohøjskolen.
- Cipriano, R. (1982). Immunogenic potential of growth products extracted from cultures of *Aeromonas salmonicida* for brook trout (*Salvelinus fontinalis*). *Can. J. Fish. Aquat. Sci.* 39, 1512 – 1518.
- Cipriano, R. C. (1997). Differences in detection of *Aeromonas salmonicida* in covertly infected salmonid fishes by the stress-inducible furunculosis test and culture-based assays. *J. Aquat. Anim. Health* 9, 108 – 113.
- Cipriano, R. C., and Austin, B. (2011). "Furunculosis and other aeromonad diseases," in *Fish Diseases and Disorders, Volume 3*, eds. P. Woo and D. Bruno (Oxon: CABI Publishing), 424438.
- Cipriano, R. C., and Bertolini, J. (1988). Selection for Virulence in the Fish Pathogen *Aeromonas salmonicida* using Coomassie Brilliant Blue Agar. *J. Wildl. Dis.* 24, 672–678. doi:10.7589/0090-3558-24.4.672.
- Cipriano, R. C., Ford, L. A., and Jones, T. E. (1994). Relationship between resistance of salmonids to furunculosis and recovery of *Aeromonas salmonicida* from external mucus. *J. Wildl. Dis.* 30, 577–580. doi:10.7589/0090-3558-30.4.577.
- Cipriano, R. C., Ford, L. A., Smith, D. R., Schachte, J. H., and Petrie, C. J. (1997). Differences in detection of *Aeromonas salmonicida* in covertly infected salmonid fishes by the stress-inducible furunculosis test and culture-based assays. *J. Aquat. Anim. Health* 9, 108 – 113.
- Cipriano, R. C., Ford, L. A., Teska, J. D., and Hale, L. H. (1992). Detection of *Aeromonas salmonicida* in the mucus of salmonid fishes. *J. Aquat. Anim. Health* 4, 114–118.

- Cipriano, R. C., and Heartwell, C. M. (1986). Susceptibility of Salmonids to Furunculosis: Differences between Serum and Mucus Responses against *Aeromonas salmonicida*. *Trans. Am. Fish. Soc.* 115, 83–88. doi:10.1577/1548-8659(1986)115<83:sostf>2.0.co;2.
- Cipriano, R. C., Morrison, J. K., and Starliper, C. E. (1983). Immunization of Salmonids against the Fish Pathogen, *Aeromonas salmonicida*. *J. World Maric. Soc.* 14, 201–211. doi:10.1111/j.1749-7345.1983.tb00076.x.
- Cipriano, R. C., and Pyle, S. W. (1985). Adjuvant-dependent immunity and the agglutinin response of fishes against *Aeromonas salmonicida*, cause of furunculosis. *Can. J. Fish. Aquat. Sci.* 42, 1290–1295. doi:10.1139/f85-160.
- Colwell, R. R., Macdonell, M. T., and De Ley, J. (1986). Proposal to Recognize the Family *Aeromonadaceae* fam. nov. *Int. J. Syst. Bacteriol.* 36, 473–477. doi:10.1099/00207713-36-3-473.
- Cunningham, C. O., and Colquhoun, D. J. (2002). “Molecular diagnosis of *Aeromonas salmonicida* infections,” in *Molecular Diagnosis of Salmonid Diseases*, ed. C. D. Cunningham (Kluwer Academic Publishers), 81–99. doi:10.1007/978-94-017-2315-2.
- Dacanay, A., Knickle, L., Solanky, K. S., Boyd, J. M., Walter, J. a., Brown, L. L., et al. (2006). Contribution of the type III secretion system (TTSS) to virulence of *Aeromonas salmonicida* subsp. *salmonicida*. *Microbiology* 152, 1847–1856. doi:10.1099/mic.0.28768-0.
- Daher, R. K., Fillion, G., Tan, S. G. E., Dallaire-Dufresne, S., Paquet, V. E., and Charette, S. J. (2011). Alteration of virulence factors and rearrangement of pAsa5 plasmid caused by the growth of *Aeromonas salmonicida* in stressful conditions. *Vet. Microbiol.* 152, 353–60. doi:10.1016/j.vetmic.2011.04.034.
- Dallaire-Dufresne, S., Barbeau, X., Sarty, D., Tanaka, K. H., Denoncourt, A. M., Lague, P., et al. (2013). *Aeromonas salmonicida* Ati2 is an effector protein of the type three secretion system. *Microbiology* 159, 1937–1945. doi:mic.0.067959-0 [pii] 10.1099/mic.0.067959-0 [doi].
- Dallaire-Dufresne, S., Tanaka, K. H., Trudel, M. V., Lafaille, A., and Charette, S. J. (2014). Virulence, genomic features, and plasticity of *Aeromonas salmonicida* subsp. *salmonicida*, the causative agent of fish furunculosis. *Vet. Microbiol.* 169, 1–7. doi:10.1016/j.vetmic.2013.06.025.
- Dalsgaard, I., and Madsen, L. (2000). Bacterial pathogens in rainbow trout, *Oncorhynchus mykiss* (Walbaum), reared at Danish freshwater farms. *J. Fish Dis.* 23, 199 – 209.
- Dalsgaard, I., Nielsen, B., and Larsen, J. L. (1994). Characterization of *Aeromonas salmonicida* subsp. *salmonicida*: a comparative study of strains of different geographic origin. *J. Appl. Bacteriol.* 77, 21–30.
- Daly, J. G., and Stevenson, R. M. (1985). Importance of culturing several organs to detect *Aeromonas salmonicida* in salmonid fish. *Trans. Am. Fish. Soc.* 114, 909–910.
- DANMAP (2012). DANMAP 2012 - Use of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from food animals, food and humans in Denmark. Available at: [http://www.danmap.org/Downloads/~/media/Projekt sites/Danmap/DANMAP reports/DANMAP 2012/Danmap\\_2012](http://www.danmap.org/Downloads/~/media/Projekt sites/Danmap/DANMAP reports/DANMAP 2012/Danmap_2012).
- Day, R. N., and Davidson, M. W. (2009). The fluorescent protein palette: tools for cellular imaging. *Chem. Soc. Rev.* 38, 2887–921. doi:10.1039/b901966a.
- Dickerson, H. W. (2009). “The biology of teleost mucosal immunity,” in *Fish Defenses, Volume 2: Pathogens, Parasites and Predators*, eds. Z. Giacomo, C. Perrière, A. Mathis, and B. G. Kapoor (Enfield, NH, USA: Science Publishers), 2–42.
- Dietrich, C., and Maiss, E. (2002). Red fluorescent protein DsRed from *Discosoma* sp. as a reporter protein in higher plants. *Biotechniques* 32, 286–293.
- Doijad, S., Lomonaco, S., Poharkar, K., Garg, S., Knabel, S., Barbuddhe, S., et al. (2014). Multi-virulence-locus sequence typing of 4b *Listeria monocytogenes* isolates obtained from different sources in India over a 10-year period. *Foodborne Pathog. Dis.* 0, 1–6.

doi:10.1089/fpd.2013.1716.

- Duff, D. C. B., and Stewart, B. J. (1933). Studies on furunculosis of fish in British Columbia. *Contrib. to Can. Biol. Fish. new Ser. A, Gen.* 8, 103–122. Available at: <http://search.proquest.com/docview/20381547?accountid=14656>.
- Ebanks, R. O., Dacanay, A., Goguen, M., Pinto, D. M., and Ross, N. W. (2004). Differential proteomic analysis of *Aeromonas salmonicida* outer membrane proteins in response to low iron and in vivo growth conditions. *Proteomics* 4, 1074–1085. doi:10.1002/pmic.200300664.
- Ebanks, R. O., Goguen, M., Knickle, L., Dacanay, A., Leslie, A., Ross, N. W., et al. (2013). Analysis of a ferric uptake regulator (Fur) knockout mutant in *Aeromonas salmonicida* subsp. *salmonicida*. *Vet. Microbiol.* 162, 831–841. doi:10.1016/j.vetmic.2012.10.038.
- Ebanks, R. O., Knickle, L. C., Goguen, M., Boyd, J. M., Pinto, D. M., Reith, M., et al. (2006). Expression of and secretion through the *Aeromonas salmonicida* type III secretion system. *Microbiology* 152, 1275–1286. doi:10.1099/mic.0.28485-0.
- Eithum, I. (1993). Injection vaccination of fish. USA: Biomed Inc.
- Ellis, A. (1997). “The extracellular toxins of *Aeromonas salmonicida* subsp. *salmonicida*,” in *Furunculosis: Multidisciplinary fish disease research*, eds. E. Bernoth, A. Ellis, P. Midtlyng, G. Olivier, and P. Smith (London UK: Academic Press), 248–268.
- Ellis, A. E., Burrows, A. S., Hastings, T. S., and Stapleton, K. J. (1988). Identification of *Aeromonas salmonicida* extracellular protease as a protective antigen against furunculosis by passive immunization. *Aquaculture* 70, 207–218. doi:10.1016/0044-8486(88)90097-X.
- Ellis, A. E., Hastings, T. S., and Munro, A. L. S. (1981). The role of *Aeromonas salmonicida* extracellular products in the pathology of furunculosis. *J. Fish Dis.* 4, 41–51. doi:10.1111/j.1365-2761.1981.tb01108.x.
- Ellis, A. E., and Stapleton, K. J. (1988). Differential susceptibility of salmonid fishes to furunculosis correlates with differential serum enhancement of *Aeromonas salmonicida* extracellular protease activity. *Microb. Pathog.* 4, 299–304. doi:10.1016/0882-4010(88)90090-3.
- Emmerich, R., and Weibel, E. (1894). Über eine durch bakterien erzeugte seuche unter den forellen. *Arch. für Hyg.* 21, 1–21.
- Enger, Ø. (1997). “Survival and inactivation of *Aeromonas salmonicida* outside the host - a most superficial way of life,” in *Furunculosis: Multidisciplinary fish disease research*, eds. E. M. Bernoth, s A. E. Elli, P. J. Midtlyng, G. Olivier, and P. Smith (London, UK: Academic Press), 159–177.
- Enger, Ø., and Thorsen, B. K. (1992). Possible ecological implications of the high cell surface hydrophobicity of the fish pathogen *Aeromonas salmonicida*. *Can. J. Microbiol.* 38, 1048–1052. doi:10.1139/m92-172.
- Enright, M. C., and Spratt, B. G. (1998). A multilocus sequence typing scheme for *Streptococcus pneumoniae*: Identification of clones associated with serious invasive disease. *Microbiology* 144, 3049–3060. doi:10.1099/00221287-144-11-3049.
- Fast, M. D., Tse, B., Boyd, J. M., and Johnson, S. C. (2009). Mutations in the *Aeromonas salmonicida* subsp. *salmonicida* type III secretion system affect Atlantic salmon leucocyte activation and downstream immune responses. *Fish Shellfish Immunol.* 27, 721–728. doi:10.1016/j.fsi.2009.09.009.
- Fehr, D., Casanova, C., Liverman, A., Blazkova, H., Orth, K., Dobbelaere, D., et al. (2006). AopP, a type III effector protein of *Aeromonas salmonicida*, inhibits the NF-kappa B signalling pathway. *Microbiol.* 152, 2809 – 2818.
- Feil, E. J., Li, B. C., Aanensen, D. M., Hanage, W. P., and Spratt, B. G. (2004). eBURST: inferring patterns of evolutionary descent among clusters of related bacterial genotypes from multilocus sequence typing data. *J. Bacteriol.* 186, 1518–1530. doi:10.1128/JB.186.5.1518-1530.2004.
- Ferguson, H. W., and McCarthy, D. H. (1978). Histopathology of furunculosis in brown trout *Salmo trutta* L. *J. Fish Dis.* 1, 165–174. doi:10.1111/j.1365-2761.1978.tb00017.x.

- Ferguson, Y., Bricknell, I. R., Glover, L. A., MacGregor, D. M., and Prosser, J. I. (1998). Colonisation and transmission of lux-marked and wild-type *Aeromonas salmonicida* strains in Atlantic salmon (*Salmo salar* L.). *Fems Microbiol. Ecol.* 27, 251–260. doi:10.1016/s0168-6496(98)00071-3.
- Ferguson, Y., Glover, L. A., McGillivray, D. M., and Prosser, J. I. (1995). Survival and activity of lux-marked *Aeromonas salmonicida* in seawater. *Appl. Environ. Microbiol.* 61, 3494–3498.
- Fernandez, A. I. G., Fernández, A. F., Pérez, M. J., Nieto, T. P., and Ellis, A. E. (1998). Siderophore production by *Aeromonas salmonicida* subsp. *salmonicida*. Lack of strain specificity. *Dis. Aquat. Organ.* 33, 87–92. doi:10.3354/dao033087.
- Filonov, G. S., Piatkevich, K. D., Ting, L.-M., Zhang, J., Kim, K., and Verkhusha, V. V (2011). Bright and stable near-infrared fluorescent protein for in vivo imaging. *Nat. Biotechnol.* 29, 757–761. doi:10.1038/nbt.1918.
- Fish, F. F. (1937). Furunculosis in wild trout. *Copeia*, 37–40.
- Fleischmann, R. D., Adams, M. D., White, O., Clayton, R. A., Kirkness, E. F., Kerlavage, A. R., et al. (1995). Whole genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science (80- )*. 269, 496–512. doi:10.1126/science.7542800.
- Food and Agriculture Organization of the United Nations (2016). Aquaculture. Available at: <http://www.fao.org/aquaculture/en/> [Accessed January 12, 2015].
- Foxman, B., Zhang, L., Koopman, J. S., Manning, S. D., and Marrs, C. F. (2005). Choosing an appropriate bacterial typing technique for epidemiologic studies. *Epidemiol. Perspect. Innov.* 2, 10. doi:10.1186/1742-5573-2-10.
- Fraser, C. M., Gocayne, J. D., White, O., Adams, M. D., Clayton, R. A., Fleischmann, R. D., et al. (1995). The Minimal Gene Complement of *Mycoplasma genitalium*. *Science (80- )*. 270, 397–403. doi:10.1126/science.270.5235.397.
- Froquet, R., Cherix, N., Burr, S. E., Frey, J., Vilches, S., Tomas, J. M., et al. (2007). Alternative host model to evaluate *Aeromonas* virulence. *Appl. Environ. Microbiol.* 73, 5657–5659. doi:10.1128/AEM.00908-07.
- Fuhrman, O. (1909). La Furunculose. *Bell Suisse Pech. Piscic.* 12, 193–195.
- Furunculosis-committee (1933). The real furunculosis carrier. Recommendations. Editorial notes. *Salmon Trout Mag.* 75, 306–308.
- Ganusov, V. V., and Antia, R. (2003). Trade-offs and the evolution of virulence of microparasites: do details matter? *Theor. Popul. Biol.* 64, 211–220. doi:10.1016/S0040-5809(03)00063-7.
- Garcia, J. A., Larsen, J. L., Dalsgaard, I., and Pedersen, K. (2000). Pulsed-field gel electrophoresis analysis of *Aeromonas salmonicida* ssp *salmonicida*. *FEMS Microbiol. Lett.* 190, 163–166. doi:10.1016/S0378-1097(00)00322-0.
- Gibson, U. E., Heid, C. A., and Williams, P. M. (1996). A novel method for real time quantitative RT-PCR. *Genome Res.* 6, 995–1001. doi:10.1101/gr.6.10.995.
- Goodwin, A. E., and Merry, G. E. (2009). Are all koi ulcer cases associated with infection by atypical *Aeromonas salmonicida*? Polymerase chain reaction assays of koi carp skin swabs submitted by hobbyists. *J. Aquat. Anim. Health* 21, 98–103. doi:10.1577/H08-042.1.
- Griffin, P., Snieszko, S., and Friddle, S. (1953). A more comprehensive description of *Bacterium salmonicida*. *Trans. Am. Fish. Soc.*, 129–138.
- Gulla, S., Duodu, S., Nilsen, A., Fossen, I., and Colquhoun, D. J. (2015). *Aeromonas salmonicida* infection levels in pre- and post-stocked cleaner fish assessed by culture and an amended qPCR assay. *J. Fish Dis.*, [Epub ahead of print]. doi:10.1111/jfd.12420.
- Gustafson, C. E., Thomas, C. J., and Trust, T. J. (1992). Detection of *Aeromonas salmonicida* from fish by using polymerase chain reaction amplification of the virulence surface array protein gene. *Appl. Environ. Microbiol.* 58, 3816–3825.
- Hall, T. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 41, 95–98. doi:citeulike-article-id:691774.

- Harris, S. R., Feil, E. J., Holden, M. T. G., Quail, M. A., Nickerson, E. K., Chantratita, N., et al. (2010). Evolution of MRSA during hospital transmission and intercontinental spread. *Science* 327, 469–74. doi:10.1126/science.1182395.
- Harvell, C. D., Kim, K., Burkholder, J. M., Colwell, R. R., Epstein, P. R., Grimes, D. J., et al. (1999). Emerging marine diseases-climate links and anthropogenic factors. *Science* (80-. ). 285, 1505–1510. doi:10.1126/science.285.5433.1505.
- Haugarvoll, E., Bjerkås, I., Szabo, N. J., Satoh, M., and Koppang, E. O. (2010). Manifestations of systemic autoimmunity in vaccinated salmon. *Vaccine* 28, 4961–4969. doi:10.1016/j.vaccine.2010.05.032.
- Henson, J., Tischler, G., and Ning, Z. (2012). Next-generation sequencing and large genome assemblies. *Pharmacogenomics* 13, 901–915. doi:10.2217/pgs.12.72.
- Higuchi, R., Dollinger, G., Walsh, P. S., and Griffith, R. (1992). Simultaneous amplification and detection of specific DNA sequences. *Biotechnology. (N. Y.)* 10, 413–417. doi:10.1038/nbt0492-413.
- Hiller, N. L., Janto, B., Hogg, J. S., Boissy, R., Yu, S., Powell, E., et al. (2007). Comparative genomic analyses of seventeen *Streptococcus pneumoniae* strains: Insights into the pneumococcal supragenome. in *Journal of Bacteriology*, 8186–8195. doi:10.1128/JB.00690-07.
- Hiney, M., Dawson, M. T., Heery, D. M., Smith, P. R., Gannon, F., and Powell, R. (1992). DNA probe for *Aeromonas salmonicida*. *Appl. Environ. Microbiol.* 58, 1039 – 1042.
- Hiney, M., and Olivier, G. (1999). “Furunculosis (*Aeromonas salmonicida*),” in *Fish diseases and disorders, Vol 3. Viral, bacterial and fungal infections*, eds. P. Woo and D. Bruno (Oxon: CABI Publishing), 341–425.
- Hiney, M. P., Kilmartin, J. J., and Smith, P. R. (1994). Detection of *Aeromonas salmonicida* in atlantic salmon with asymptomatic furunculosis infections. *Dis. Aquat. Organ.* 19, 161 – 167.
- Hiney, M., Smith, P., and Bernoth, E. (1997). “Covert *Aeromonas salmonicida* infections,” in *Furunculosis: Multidisciplinary fish disease research*, eds. E. M. Bernoth, A. E. Ellis, P. J. Midtlyng, G. Olivier, and S. P (London UK: Academic Press), 54–97.
- Hirst, I. D., and Ellis, A. E. (1994). Iron-regulated outer membrane proteins of *Aeromonas salmonicida* are important protective antigens in Atlantic salmon against furunculosis. *Fish Shellfish Immunol.* 4, 29–45. doi:10.1006/fsim.1994.1004.
- Hirst, I. D., Hastings, T. S., and Ellis, A. E. (1994). Utilization of haem compounds by *Aeromonas salmonicida*. *J. Fish Dis.* 17, 365–373. doi:10.1111/j.1365-2761.1994.tb00232.x.
- Horne, J. H. (1928). Furunculosis in trout and the importance of carriers in the spread of the disease. *J. Hyg.* 28, 67–78.
- Hutchens, M., and Luker, G. D. (2007). Applications of bioluminescence imaging to the study of infectious diseases. *Cell Microbiol* 9, 2315–2322. doi:CMI995 [pii]r10.1111/j.1462-5822.2007.00995.x.
- Håstein, T., Gudding, R., and Evensen, O. (2005). Bacterial vaccines for fish - an update of the current situation worldwide. *Dev. Biol. (Basel)*. 121, 55–74.
- Håstein, T., and Lindstad, T. (1991). Diseases in wild and cultured salmon: possible interaction. *Aquaculture* 98, 277–288. doi:10.1016/0044-8486(91)90392-K.
- Inglis, V., Colquhoun, D., Pearson, M. D., Miyata, M., and Aoki, T. (1996). Analysis of DNA relationships among *Aeromonas* species by RAPD (randomly amplified polymorphic DNA) typing. *Aquac. Int.* 4, 43–53. doi:10.1007/BF00175220.
- Ishiguro, E., Ainsworth, T., and Trust, T. J. (1985). Congo red agar a differential medium for *Aeromonas salmonicida* detects the presence of the cell surface protein array involved in virulence. *J. Bacteriol.* 164, 1233 – 1237.
- Ishiguro, E. E., Kay, W. W., Ainsworth, T., Chamberlain, J. B., Austen, R. A., Buckley, J. T., et al. (1981). Loss of virulence during culture of *Aeromonas salmonicida* at high temperature. *J. Bacteriol.* 148, 333–340.



- Jacobsen, P., and Berglind, L. (1988). Persistence of oxytetracycline in sediments from fish farms. *Aquaculture* 70, 365–370. doi:10.1016/0044-8486(88)90120-2.
- Johnson, C. M., Tatner, M. F., and Horne, M. T. (1985). Autoaggregation and extracellular A-layer protein in *Aeromonas salmonicida*. *Aquaculture* 46, 163–166. doi:10.1016/0044-8486(85)90200-5.
- Johnson, K. A., and Amend, D. F. (1984). Potential for immersion vaccination against *Aeromonas salmonicida*. *J. Fish Dis.* 7, 101–105. doi:10.1111/j.1365-2761.1984.tb00912.x.
- Kadlec, K., von Czapiewski, E., Kaspar, H., Wallmann, J., Michael, G. B., Steinacker, U., et al. (2011). Molecular basis of sulfonamide and trimethoprim resistance in fish-pathogenic *Aeromonas* isolates. *Appl. Environ. Microbiol.* 77, 7147–7150. doi:10.1128/AEM.00560-11.
- Karsi, A., Menanteau-Ledouble, S., and Lawrence, M. L. (2006). Development of bioluminescent *Edwardsiella ictaluri* for noninvasive disease monitoring. *FEMS Microbiol. Lett.* 260, 216–223. doi:10.1111/j.1574-6968.2006.00310.x.
- Kay, W., and Trust, T. (1997). “The surface of *Aeromonas salmonicida*: what does it look like and what does it do?,” in *Furunculosis: Multidisciplinary fish disease research*, eds. E. Bernoth, A. E. Ellis, P. J. Midtlyng, G. Olivier, and P. Smith (London UK: Academic Press), 235–247.
- Keeling, S. E., Brosnahan, C. L., Johnston, C., Wallis, R., Gudkovs, N., and McDonald, W. L. (2012). Development and validation of a real-time PCR assay for the detection of *Aeromonas salmonicida*. *J. Fish Dis.* 36, 495–503. doi:10.1111/jfd.12014.
- King, C. H., and Shotts, E. B. (1988). Enhancement of *Edwardsiella tarda* and *Aeromonas salmonicida* through ingestion by the ciliated protozoan *Tetrahymena pyriformis*. *FEMS Microbiol. Lett* 51, 95–100.
- Koppang, E. O., Bjerkås, I., Haugarvoll, E., Chan, E. K. L., Szabo, N. J., Ono, N., et al. (2008). Vaccination-induced systemic autoimmunity in farmed Atlantic salmon. *J. Immunol.* 181, 4807–4814. doi:181/7/4807 [pii].
- Koppang, E. O., Fjølstad, M., Melgård, B., Vigerust, M., and Ørum, H. S. (2000). Non-pigment-producing isolates of *Aeromonas salmonicida* subspecies *salmonicida*: isolation, identification, transmission and pathogenicity in Atlantic salmon, *Salmo salar* L. *J. Fish Dis.* 23, 39–48. doi:10.1046/j.1365-2761.2000.00205.x.
- Kozinska, A., Figueras, M. J., Chacon, M. R., and Soler, L. (2002). Phenotypic characteristics and pathogenicity of [i]Aeromonas genomospecies isolated from common carp (*Cyprinus carpio* L.). *J. Appl. Microbiol.* 93, 1034–1041. doi:10.1046/j.1365-2672.2002.01784.x.
- Krantz, G. E., Reddecliff, J. M., and Heist, C. E. (1964). Immune Response of Trout to *Aeromonas Salmonicida* Part II. Evaluation of Feeding Techniques. *Progress. Fish-Culturist* 26, 65–69. doi:10.1577/1548-8659(1964)26[65:IROTTA]2.0.CO;2.
- Kubista, M., Andrade, J. M., Bengtsson, M., Forootan, A., Jonák, J., Lind, K., et al. (2006). The real-time polymerase chain reaction. *Mol. Aspects Med.* 27, 95–125. doi:10.1016/j.mam.2005.12.007.
- L’Abée-Lund, T. M., and Sørum, H. (2000). Functional Tn5393-like transposon in the R plasmid pRAS2 from the fish pathogen *Aeromonas salmonicida* subspecies *salmonicida* isolated in Norway. *Appl. Environ. Microbiol.*, 5533–5535.
- L’Abée-Lund, T. M., and Sørum, H. (2001). Class 1 integrons mediate antibiotic resistance in the fish pathogen *Aeromonas salmonicida* worldwide. *Microb. Drug Resist.* 7, 263–72. doi:10.1089/10766290152652819.
- Lafferty, K. D. (2009). The ecology of climate change and infectious diseases. *Ecology* 90, 888–900. doi:10.1890/08-0079.1.
- Land, M., Hauser, L., Jun, S.-R., Nookaew, I., Leuze, M. R., Ahn, T.-H., et al. (2015). Insights from 20 years of bacterial genome sequencing. *Funct. Integr. Genomics* 15, 141–61. doi:10.1007/s10142-015-0433-4.
- Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., Mcgettigan, P. A., McWilliam, H., et al.

- (2007). Clustal W and Clustal X version 2.0. *Bioinformatics* 23, 2947–2948. doi:10.1093/bioinformatics/btm404.
- Larsen, J. L., and Møllgaard, S. (1981). Microbiological and hygienic problems in marine aquaculture: furunculosis and vibriosis in rainbow trout (*Salmo gairdneri*). *Bull. Eur. Ass. Fish Pathol.* 1, 29–32.
- Lee, K. K., and Ellis, A. E. (1990). Glycerophospholipid:cholesterol acyltransferase complexed with lipopolysaccharide (LPS) is a major lethal exotoxin and cytotoxin of *Aeromonas salmonicida*: LPS stabilizes and enhances toxicity of the enzyme. *J. Bacteriol.* 172, 5382–5393.
- Leekitcharoenphon, P., Lukjancenko, O., Friis, C., Aarestrup, F. M., and Ussery, D. W. (2012). Genomic variation in *Salmonella enterica* core genes for epidemiological typing. *BMC Genomics* 13, 88. doi:10.1186/1471-2164-13-88.
- Lewis, T., Loman, N. J., Bingle, L., Jumaa, P., Weinstock, G. M., Mortiboy, D., et al. (2010). High-throughput whole-genome sequencing to dissect the epidemiology of *Acinetobacter baumannii* isolates from a hospital outbreak. *J. Hosp. Infect.* 75, 37–41. doi:10.1016/j.jhin.2010.01.012.
- Lilly, J., and Camps, M. (2015). Mechanisms of theta plasmid replication. *Microbiol. Spectr.* 3, PLAS-0029–2014. doi:10.1128/microbiolspec.PLAS-0029-2014.
- Lin, L., and Meighen, E. (2009). Bacterial bioluminescence biochemistry and molecular biology. Available at: <http://photobiology.info/Lin.html> [Accessed January 4, 2016].
- Loman, N. J., and Pallen, M. J. (2015). Twenty years of bacterial genome sequencing. *Nat. Rev. Microbiol.*, 1–9. doi:10.1038/nrmicro3565.
- Lund, V., Arne Arnesen, J., Coucheron, D., Modalsli, K., and Syvertsen, C. (2003a). The *Aeromonas salmonicida* A-layer protein is an important protective antigen in oil-adjuvanted vaccines. *Fish Shellfish Immunol.* 15, 367–372. doi:10.1016/S1050-4648(02)00180-8.
- Lund, V., Espelid, S., and Mikkelsen, H. (2003b). Vaccine efficacy in spotted wolffish *Anarhichas minor*: Relationship to molecular variation in a-layer protein of atypical *Aeromonas salmonicida*. *Dis. Aquat. Organ.* 56, 31–42. doi:10.3354/dao056031.
- Magnadottir, B. (2010). Immunological control of fish diseases. *Mar. Biotechnol.* 12, 361–379. doi:10.1007/s10126-010-9279-x.
- Maiden, M. C., Bygraves, J. A., Feil, E., Morelli, G., Russell, J. E., Urwin, R., et al. (1998). Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc. Natl. Acad. Sci. U. S. A.* 95, 3140–5. doi:10.1073/pnas.95.6.3140.
- Maiden, M. C. J. (2006). Multilocus sequence typing of bacteria. *Annu. Rev. Microbiol.* 60, 561–588. doi:10.1146/annurev.micro.59.030804.121325.
- Marcogliese, D. J. (2008). The impact of climate change on the parasites and infectious diseases of aquatic animals. *Sci. Technol.* 27, 467–484.
- Marcos-López, M., Gale, P., Oidtmann, B. C., and Peeler, E. J. (2010). Assessing the Impact of Climate Change on Disease Emergence in Freshwater Fish in the United Kingdom. *Transbound. Emerg. Dis.* doi:10.1111/j.1865-1682.2010.01150.x.
- Markwardt, N., Gocha, Y., and Klontz, G. (1989). A new application for Coomassie Brilliant Blue agar detection of *Aeromonas salmonicida* in clinical samples. *Dis. Aquat. Organ.* 6, 231–233. doi:10.3354/dao006231.
- Marsh, M. C. (1902). *Bacterium truttae* a new species of bacterium pathogenic to trout. *Sci. N.S.* 16, 706–707.
- Martinez-Murcia, A. J., Monera, A., Saavedra, M. J., Oncina, R., Lopez-Alvarez, M., Lara, E., et al. (2011). Multilocus phylogenetic analysis of the genus *Aeromonas*. *Syst. Appl. Microbiol.* 34, 189–199. doi:10.1016/j.syapm.2010.11.014.
- Martino, M. E., Fasolato, L., Montemurro, F., Rosteghin, M., Manfrin, A., Patarnello, T., et al. (2011). Determination of microbial diversity of *Aeromonas* strains on the basis of multilocus sequence typing, phenotype, and presence of putative virulence genes. *Appl. Environ.*

- Microbiol.* 77, 4986–5000. doi:10.1128/AEM.00708-11.
- Mavromatis, K., Land, M. L., Brettin, T. S., Quest, D. J., Copeland, A., Clum, A., et al. (2012). The Fast Changing Landscape of Sequencing Technologies and Their Impact on Microbial Genome Assemblies and Annotation. *PLoS One* 7. doi:10.1371/journal.pone.0048837.
- McCarthy, D. H. (1977). “Some ecological aspects of the bacterial fish pathogen - *Aeromonas salmonicida*,” in *Aquatic Microbiology. Soc. Appl. Bacteriol. Symp. Ser. 6*, eds. F. A. Skinner and J. M. Shewan (London, UK: Academic Press), 299–324.
- McCarthy, D. H. (1980). “Some ecological aspects of the bacterial fish pathogen- *Aeromonas salmonicida*,” in *Aquatic microbiology: symposium of the Society of Applied Bacteriology*, 299–324.
- McCarthy, D. H., and Roberts, R. J. (1980). Furunculosis of fish - the present state of our knowledge. *Adv. Aquat. Microbiol.*, 293–342.
- McCormick, W. A., Stevenson, R. M., and MacInnes, J. I. (1990). Restriction endonuclease fingerprinting analysis of Canadian isolates of *Aeromonas salmonicida*. *Can. J. Microbiol.* 36, 24–32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2159364>.
- Menanteau-Ledouble, S., Karsi, A., and Lawrence, M. L. (2011). Importance of skin abrasion as a primary site of adhesion for *Edwardsiella ictaluri* and impact on invasion and systematic infection in channel catfish *Ictalurus punctatus*. *Vet. Microbiol.* 148, 425–430. doi:10.1016/j.vetmic.2010.08.022.
- Méndez, J., and Guijarro, J. A. (2013). In vivo monitoring of *Yersinia ruckeri* in fish tissues: progression and virulence gene expression. *Environ. Microbiol. Rep.* 5, 179–185. doi:10.1111/1758-2229.12030.
- Mettam, A. E. (1915). Report on the outbreak of furunculosis in the River Liffey in 1913. *Sci. Invest. Dept. Agr. Tech. Instr.*, 19.
- Metzker, M. L. (2010). Sequencing technologies - the next generation. *Nat. Rev. Genet.* 11, 31–46. doi:10.1038/nrg2626.
- Midtlyng, P. J. (1996). A field study on intraperitoneal vaccination of Atlantic salmon (*Salmo salar* L.) against furunculosis. *Fish Shellfish Immunol.* 6, 553–565. doi:10.1006/fsim.1996.0011.
- Midtlyng, P. J. (1997). “Vaccination Against Furunculosis,” in *Furunculosis: Multidisciplinary fish disease research*, ed. E. Bernoth (London UK: Academic Press), 382–405.
- Midtlyng, P. J., Reitan, L. J., Lillehaug, A., and Ramstad, A. (1996). Protection, immune responses and side effects in Atlantic salmon (*Salmo salar* L.) vaccinated against furunculosis by different procedures. *Fish Shellfish Immunol.* 6, 599–613. doi:10.1006/fsim.1996.0055.
- Miljø- og Fødevareministeriet (2014). Strategi for bæredygtig udvikling af akvakultursektoren i Danmark 2014-2020. Available at: <http://naturerhverv.dk/fiskeri/akvakultur/strategi-for-baeredygtig-udvikling-af-akvakultursektoren-i-danmark-2014-2020/#c29059> [Accessed February 15, 2016].
- Miljø- og Fødevareministeriet (2015). Danskerne spiser sundere. Available at: <http://altomkost.dk/nyheder/nyhed/nyhed/danskerne-spiser-sundere/> [Accessed January 9, 2016].
- Miyata, M., Aoki, T., Inglis, V., Yoshida, T., and Endo, M. (1995). RAPD analysis of *Aeromonas salmonicida* and *Aeromonas hydrophila*. *J. Appl. Bacteriol.* 79, 181–185. doi:10.1111/j.1365-2672.1995.tb00933.x.
- Mooney, J., Powell, E., Clabby, C., and Powell, R. (1995). Detection of *Aeromonas salmonicida* in wild Atlantic salmon using a specific DNA probe test. *Dis. Aquat. Organ.* 21, 131–135. doi:10.3354/dao021131.
- Morgan, J., Rhodes, G., and Pickup, R. W. (1993). Survival of nonculturable *Aeromonas salmonicida* in lake water. *Appl. Environ. Microbiol.* 59, 874 – 880.
- Morrison, T. B., Weis, J. J., and Wittwer, C. T. (1998). Quantification of lowcopy transcripts by continuous SYBR Green I monitoring during amplification. *Biotechniques* 24, 954–962.

- Mota, L. J., and Cornelis, G. R. (2005). The bacterial injection kit: type III secretion systems. *Ann. Med.* 37, 234–249. doi:10.1080/07853890510037329.
- Munro, A., and Hastings, T. S. (1993). “Furunculosis,” in *Bacterial Diseases of Fish*, eds. V. Inglis, R. J. Roberts, and N. Bromage (Oxford: Blackwell Scientific Publishers), 122–142.
- Mutoloki, S., Brudeseth, B., Reite, O. B., and Evensen, Ø. (2006). The contribution of *Aeromonas salmonicida* extracellular products to the induction of inflammation in Atlantic salmon (*Salmo salar* L.) following vaccination with oil-based vaccines. *Fish Shellfish Immunol.* 20, 1–11. doi:10.1016/j.fsi.2005.01.005.
- Mutoloki, S., Cooper, G. a, Marjara, I. S., Koop, B. F., and Evensen, Ø. (2010). High gene expression of inflammatory markers and IL-17A correlates with severity of injection site reactions of Atlantic salmon vaccinated with oil-adjuvanted vaccines. *BMC Genomics* 11, 336. doi:10.1186/1471-2164-11-336.
- Muziasari, W. I., Managaki, S., Pärnänen, K., Karkman, A., Lyra, C., Tamminen, M., et al. (2014). Sulphonamide and trimethoprim resistance genes persist in sediments at Baltic Sea aquaculture farms but are not detected in the surrounding environment. *PLoS One* 9. doi:10.1371/journal.pone.0092702.
- Najimi, M., Lemos, M. L., and Osorio, C. R. (2008). Identification of heme uptake genes in the fish pathogen *Aeromonas salmonicida* subsp. *salmonicida*. *Arch. Microbiol.* 190, 439–449. doi:10.1007/s00203-008-0391-5.
- Najimi, M., Lemos, M. L., and Osorio, C. R. (2009). Identification of iron regulated genes in the fish pathogen *Aeromonas salmonicida* subsp. *salmonicida*: Genetic diversity and evidence of conserved iron uptake systems. *Vet. Microbiol.* 133, 377–382. doi:10.1016/j.vetmic.2008.07.008.
- Născuțiu, A. M. (2010). Bacterii viabile necultivabile. *Bacteriol. Virusol. Parazitol. Epidemiol.* 55, 11 – 18.
- Nash, J. H. E., Findlay, W. A., Luebbert, C. C., Mykytczuk, O. L., Foote, S. J., Taboada, E. N., et al. (2006). Comparative genomics profiling of clinical isolates of *Aeromonas salmonicida* using DNA microarrays. *BMC Genomics* 7, 43. doi:10.1186/1471-2164-7-43.
- Nazarenko, I., Lowe, B., Darfler, M., Ikononi, P., Schuster, D., and Rashtchian, A. (2002a). Multiplex quantitative PCR using self-quenched primers labeled with a single fluorophore. *Nucleic Acids Res.* 30.
- Nazarenko, I., Pires, R., Lowe, B., Obaidy, M., and Rashtchian, A. (2002b). Effect of primary and secondary structure of oligodeoxyribonucleotides on the fluorescent properties of conjugated dyes. *Nucleic Acids Res.* 30, 2089 – 2095.
- Nielsen, B., Dalsgaard, I., Brown, D. J., and Larsen, J. L. (1994a). *Aeromonas salmonicida* subsp. *salmonicida* : correlation of protein-patterns, antibiotic resistance, exoprotease activity, hemolysis and pathological lesions produced *in vivo*. *J. Fish Dis.* 17, 387–397. doi:DOI 10.1111/j.1365-2761.1994.tb00234.x.
- Nielsen, B., Olsen, J. E., and Larsen, J. L. (1993). Plasmid profiling as an epidemiological marker within *Aeromonas salmonicida*. *Dis. Aquat. Organ.* 15, 129–135. doi:10.3354/dao015129.
- Nielsen, B., Olsen, J., and JL, L. (1994b). Ribotyping of *Aeromonas salmonicida* subsp. *salmonicida*. *Dis. Aquat. Organ.* 18, 155 – 158.
- Noga, E. J. (2010). *Fish Diseases*. 2nd ed. Singapore: Wiley-Blackwell.
- Nomura, T., Yoshimizu, M., and Kimura, T. (1992). The Epidemiological Study of Furunculosis in Salmon Propagation. Available at: <http://hdl.handle.net/2115/39004>.
- Nomura, T., Yoshimizu, M., and Kimura, T. (1993). An epidemiological study of furunculosis in salmon propagation in Japanese rivers. *Fish. Res.* 17, 137–146. doi:10.1016/0165-7836(93)90013-W.
- O’Brien, D., Mooney, J., Ryan, D., Powell, E., Hiney, M., Smith, P. R., et al. (1994). Detection of *Aeromonas-salmonicida*, causal agent of furunculosis in salmonid fish, from the tank effluent of

- hatchery-reared atlantic salmon smolts. *Appl. Environ. Microbiol.* 60, 3874 – 3877.
- O'Dowd, A. M., Bricknell, I. R., Secombes, C. J., and Ellis, A. E. (1999). The primary and secondary antibody responses to IROMP antigens in Atlantic salmon (*Salmo salar* L) immunised with A+ and A- *Aeromonas salmonicida* bacterins. *Fish Shellfish Immunol.* 9, 125–138.
- O'hici, B., Olivier, G., and Powell, R. (2000). Genetic diversity of the fish pathogen *Aeromonas salmonicida* demonstrated by random amplified polymorphic DNA and pulsed-field gel electrophoresis analyses. *Dis. Aquat. Organ.* 39, 109–119. doi:10.3354/dao039109.
- Ogut, H., and Reno, P. W. (2005). Evaluation of an experimental *Aeromonas salmonicida* epidemic in chinook salmon, *Oncorhynchus tshawytscha* (Walbaum). *J. Fish Dis.* 28, 263–269. doi:10.1111/j.1365-2761.2005.00626.x.
- Olenych, S. G., Claxton, N. S., Ottenberg, G. K., and Davidson, M. W. (2007). The fluorescent protein color palette. *Curr. Protoc. Cell Biol.* Chapter 21, Unit 21.5. doi:10.1002/0471143030.cb2105s36.
- Olivier, G. (1990). Virulence of *Aeromonas salmonicida*: Lack of Relationship with Phenotypic Characteristics. *J. Aquat. Anim. Health* 2, 119–127. doi:10.1577/1548-8667(1990)002<0119:VOSLOR>2.3.CO;2.
- Pandya, G. a, Holmes, M. H., Petersen, J. M., Pradhan, S., Karamycheva, S. a, Wolcott, M. J., et al. (2009). Whole genome single nucleotide polymorphism based phylogeny of *Francisella tularensis* and its application to the development of a strain typing assay. *BMC Microbiol.* 9, 213. doi:10.1186/1471-2180-9-213.
- Parkhill, J., and Wren, B. W. (2011). Bacterial epidemiology and biology-lessons from genome sequencing. *Genome Biol.* 12, 230. doi:10.1186/gb-2011-12-10-230.
- Pedersen, K., Skall, H. F., Lassen-Nielsen, A. M., Nielsen, T. F., Henriksen, N. H., and Olesen, N. J. (2008). Surveillance of health status on eight marine rainbow trout, *Oncorhynchus mykiss* (Walbaum), farms in Denmark in 2006. *J. Fish Dis.* 31, 659–67. doi:10.1111/j.1365-2761.2008.00941.x.
- Perez, M. J., Fernandez, A. I. G., Rodriguez, L. A., and Nieto, T. P. (1996). Differential susceptibility to furunculosis of turbot and rainbow trout and release of the furunculosis agent from furunculosis-affected fish. *Dis. Aquat. Org.* 26, 133–137. doi:10.3354/dao026133.
- Pharmaq (2016). Injectable vaccines. Available at: <http://www.pharmaq.no/products/injectable/> [Accessed February 8, 2016].
- Phipps, B. M., Trust, T. J., Ishiguro, E. E., and Kay, W. W. (1983). Purification and characterization of the cell surface virulent A protein from *Aeromonas salmonicida*. *Biochemistry* 22, 2934–2939. doi:10.1021/bi00281a023.
- Pittet, L. (1910). La furunculose observe sur la Bordeliere, le Nase, et le Barbeau. *Bull. Suisse Pech. Pisciculture* 2, 171–173.
- Platonov, A. E., Shipulin, G. A., and Platonova, O. V (2000). Multilocus sequence typing: A new method and the first results in the genotyping of bacteria. *Russ. J. Genet.* 36, 481–487.
- Plehn, M. (1911). Die Furunculose der Salmoniden. *Zentr. Bakt. Parasitenk., I Org.* 60, 609–624.
- Popoff, M. (1984). “Family II. *Vibrionaceae*,” in *Bergey's Manual of Systematic Bacteriology, Vol. 1*, eds. N. Krieg and J. Holt (Baltimore: Willams and Wilkins), 545–550.
- Press, C. M., Evensen, Ø., Reitan, L. J., and Landsverk, T. (1996). Retention of furunculosis vaccine components in Atlantic salmon, *Salmo salar* L., following different routes of vaccine administration. *J. Fish Dis.* 19, 215–224. doi:10.1111/j.1365-2761.1996.tb00128.x.
- Prost, M. (2001). Extracellular proteases ECP of *Aeromonas salmonicida* and their significance in the pathogenesis and immuno-prophylaxis of fish furunculosis. *Med. Wet.* 57, 79–82.
- Pulkkinen, K., Suomalainen, L.-R., Read, A. F., Ebert, D., Rintamäki, P., and Valtonen, E. T. (2010). Intensive fish farming and the evolution of pathogen virulence: the case of columnaris disease in Finland. *Proc. Biol. Sci.* 277, 593–600. doi:10.1098/rspb.2009.1659.

- Rasch, M., Kastbjerg, V. G., Bruhn, J. B., Dalsgaard, I., Givskov, M., and Gram, L. (2007). Quorum sensing signals are produced by *Aeromonas salmonicida* and quorum sensing inhibitors can reduce production of a potential virulence factor. *Dis. Aquat. Organ.* 78, 105–113. doi:10.3354/dao01865.
- Rasmussen, C. J. (1964). Furunkulose hos damørreder (Furunculosis in farmed trout) (in Danish). *Scand. J. Vet. Sci.* 16, 463–472.
- Reith, M. E., Singh, R. K., Curtis, B., Boyd, J. M., Bouevitch, A., Kimball, J., et al. (2008). The genome of *Aeromonas salmonicida* subsp. *salmonicida* A449: insights into the evolution of a fish pathogen. *BMC Genomics* 9.
- Rijkers, G. T., Van Oosterom, R., and Van Muiswinkel, W. B. (1981). The immune system of cyprinid fish. Oxytetracycline and the regulation of humoral immunity in carp (*Cyprinus carpio*). *Vet. Immunol. Immunopathol.* 2, 281–290. doi:10.1016/0165-2427(81)90029-5.
- Rodgers, C. J. (1990). Immersion vaccination for control of fish furunculosis. *Dis. Aquat. Organ.* 8, 69–72. doi:10.1017/S0008/0069/\$03.00.
- Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., et al. (2012). MrBayes 3.2: Efficient bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–542. doi:10.1093/sysbio/sys029.
- Rose, A. S., Ellis, A. E., and Adams, A. (1989a). An assessment of routine *Aeromonas salmonicida* carrier detection by ELISA. *Bull. Eur. Ass. Fish Pathol.* 9, 65–67.
- Rose, A. S., Ellis, A. E., and Munro, A. L. S. (1989b). The infectivity by different routes of exposure and shedding rates of *Aeromonas salmonicida* subsp. *salmonicida* in Atlantic salmon, *Salmo salar* L., held in sea water. *J. Fish Dis.* 12, 573–578. doi:10.1111/j.1365-2761.1989.tb00566.x.
- Rose, A. S., Ellis, A. E., and Munro, A. L. S. (1990). The survival of *Aeromonas salmonicida* subsp. *salmonicida* in sea water. *J. Fish Dis.* 13, 205–214. doi:10.1111/j.1365-2761.1990.tb00775.x.
- Ross, M. G., Russ, C., Costello, M., Hollinger, A., Lennon, N. J., Hegarty, R., et al. (2013). Characterizing and measuring bias in sequence data. *Genome Biol.* 14, R51. doi:10.1186/gb-2013-14-5-r51.
- Rothberg, J. M., and Leamon, J. H. (2008). The development and impact of 454 sequencing. *Nat. Biotechnol.* 26, 1117–1124. doi:10.1038/nbt1485.
- Saiki, R. K., Gelfand, D. H., Stoffel, S., Scharf, S. J., Higuchi, R., Horn, G. T., et al. (1988). Primer directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239, 487–491. doi:10.1126/science.2448875.
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A., et al. (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science (80-. )*. 230, 1350–1354.
- Salipante, S. J., SenGupta, D. J., Cummings, L. A., Land, T. A., Hoogstraat, D. R., and Cookson, B. T. (2015). Application of whole-genome sequencing for bacterial strain typing in molecular epidemiology. *J. Clin. Microbiol.* 53, 1072–1079. doi:10.1128/JCM.03385-14.
- Satoh, M., Bjerckås, I., Haugarvoll, E., Chan, E. K. L., Szabo, N. J., Jirillo, E., et al. (2011). Polyclonal hypergammaglobulinemia and autoantibody production induced by vaccination in farmed Atlantic salmon. *Fish Shellfish Immunol.* 30, 1080–1086. doi:10.1016/j.fsi.2011.02.006.
- Scallan, A., and Smith, P. R. (1993). Importance of sampling time in detecting stress inducible furunculosis in Atlantic salmon smolts. *Bull. Eur. Assoc. Fish Pathol.* 13, 77–78.
- Seshadri, R., Joseph, S. W., Chopra, A. K., Sha, J., Shaw, J., Graf, J., et al. (2006). Genome sequence of *Aeromonas hydrophila* ATCC 7966T: Jack of all trades. *J. Bacteriol.* 188, 8272–8282. doi:10.1128/JB.00621-06.
- Sha, J., Pillai, L., Fadl, A. A., Galindo, C. L., Erova, T. E., and Chopra, A. K. (2005). The type III secretion system and cytotoxic enterotoxin alter the virulence of *Aeromonas hydrophila*. *Infect. Immun.* 73, 6446–6457. doi:10.1128/IAI.73.10.6446-6457.2005.
- Shendure, J., and Ji, H. (2008). Next-generation DNA sequencing. *Nat Biotechnol* 26, 1135–1145.

doi:10.1038/nbt1486.

- De Silva, S., and Soto, D. (2009). "Climate change and aquaculture: potential impacts, adaptation and mitigation," in *Climate change implications for fisheries and aquaculture: Overview of current scientific knowledge*, eds. K. Cochrane, C. De Young, D. Soto, and T. Bahri (Rome: FAO Fisheries and Aquaculture Technical Paper), 151–212.
- Smillie, C., Garcillan-Barcia, M. P., Francia, M. V., Rocha, E. P. C., and de la Cruz, F. (2010). Mobility of Plasmids. *Microbiol. Mol. Biol. Rev.* 74, 434–452. doi:10.1128/MMBR.00020-10.
- Smith, P. R. (1991). Stress test picks up furunculosis carriers. *Fish Farmer* 14, 18.
- Smith, P. R. (1992). "Antibiotics and the alternatives," in *Aquaculture and the environment*, eds. N. De Pauw and J. Joyce (European Aquaculture Society Spec. Publ. No. 16), 223–234.
- Smith, W. W. (1942). Action of alkaline acriflavine solution on *Bacterium salmonicida* and trout eggs. *Froc. Soc. Exp. Biol. Med* 51, 324–326.
- Snieszko, S. (1957). "Genus IV. *Aeromonas* (Kluyver and van Niel, 1936)," in *Bergey's Manual of Determinative Bacteriology*, eds. R. Breed, E. Murray, and N. Smith (Baltimore: Williams and Wilkins), 189–193.
- Spence, K. D., Fryer, J. L., and Pilcher, K. S. (1965). Active and passive immunization of certain salmonid fishes against *Aeromonas salmonicida*. *Can. J. Microbiol.* 11, 397–405. doi:10.1139/m65-053.
- Stuber, K., Burr, S. E., Braun, M., Wahli, T., and Frey, J. (2003). Type III Secretion Genes in *Aeromonas salmonicida* subsp. *salmonicida* Are Located on a Large Thermolabile Virulence Plasmid. *J. Clin. Microbiol.* 41, 3854–3856. doi:10.1128/JCM.41.8.3854-3856.2003.
- Stucki, D., Ballif, M., Bodmer, T., Coscolla, M., Maurer, A. M., Droz, S., et al. (2015). Tracking a tuberculosis outbreak over 21 years: Strain-specific single-nucleotide polymorphism typing combined with targeted whole-genome sequencing. in *Journal of Infectious Diseases*, 1306–1316. doi:10.1093/infdis/jiu601.
- Studer, N., Frey, J., and Vanden Bergh, P. (2013). Clustering subspecies of *Aeromonas salmonicida* using IS630 typing. *BMC Microbiol.* 13, 13–36. doi:10.1186/1471-2180-13-36.
- Sudheesh, P. S., Al-Ghabshi, A., Al-Mazrooei, N., and Al-Habsi, S. (2012). Comparative pathogenomics of bacteria causing infectious diseases in fish. *Int. J. Evol. Biol.* 2012, 457264. doi:10.1155/2012/457264.
- Surbeck, G. (1911). Die Furunkulosis der Fische in freien Gewässern, Schweiz. *Fischerei Zeitg.* 19, 125–132.
- Svendsen, Y. S., and Bøggwald, J. (1997). Influence of artificial wound and non-intact mucus layer on mortality of Atlantic salmon (*Salmo salar* L.) following a bath challenge with *Vibrio anguillarum* and *Aeromonas salmonicida*. *Fish Shellfish Immunol.* 4, 317–325. doi:10.1006/fsim.1997.0087.
- Svendsen, Y. S., Dalmo, R. A., and Bøggwald, J. (1999). Tissue localization of *Aeromonas salmonicida* in Atlantic salmon, *Salmo salar* L., following experimental challenge. *J. Fish Dis.* 22, 125–131. doi:10.1046/j.1365-2761.1999.00153.x.
- Sørum, H. (1998). Mobile drug resistance genes among fish bacteria. *Apmis, Suppl.* 106, 74 – 76.
- Sørum, H. (1999). Antibiotic resistance in aquaculture. *Acta Vet. Scand. Suppl.* 92, 29 – 36.
- Sørum, H., Kvello, J. H., and Hastein, T. (1993). Occurrence and stability of plasmids in *Aeromonas salmonicida* ss *salmonicida* isolated from salmonids with furunculosis. *Dis. Aquat. Organ.* 16, 199 – 206.
- Sørum, H., L'Abée-Lund, T. M., Solberg, A., and Wold, A. (2003). Integron-containing IncU R plasmids pRAS1 and pAr-32 from the fish pathogen *Aeromonas salmonicida*. *Antimicrob. Agents Chemother.* 47, 1285–1290. doi:10.1128/AAC.47.4.1285-1290.2003.
- Tan, Y., Srinivasa, P., Zhang, Y., Hew, C., and Leung, K. (2002). "Identification of virulence genes in bacterial fish pathogens: a genomic and proteomic approach," in *Aquatic Genomics: Steps Toward a Great Future*, eds. N. Shimizu, T. Aoki, I. Hirono, and F. Takashima (Tokyo

- Heidelberg New York: Springer), 340–343. doi:10.1007/978-4-431-65938-9.
- Tanaka, K. H., Dallaire-Dufresne, S., Daher, R. K., Frenette, M., and Charette, S. J. (2012). An insertion sequence-dependent plasmid rearrangement in *Aeromonas salmonicida* causes the loss of the type three secretion system. *PLoS One* 7, e33725. doi:10.1371/journal.pone.0033725.
- Tankouo-Sandjong, B., Sessitsch, A., Liebana, E., Kornschöber, C., Allerberger, F., Haechler, H., et al. (2007). MLST-v, multilocus sequence typing based on virulence genes, for molecular typing of *Salmonella enterica* subsp. *enterica* serovars. *J. Microbiol. Methods* 69. doi:10.1016/j.mimet.2006.11.013.
- Tatner, M. F., Johnson, C. M., and Horne, M. T. (1984). The tissue localization of *Aeromonas salmonicida* in rainbow trout, *Salmo gairdneri* Richardson, following three methods of administration. *J. Fish Biol.* 25, 95–108. doi:10.1111/j.1095-8649.1984.tb04854.x.
- Teska, J. D., and Cipriano, R. C. (1993). Nonselective nature of Coomassie Brilliant Blue agar for the presumptive identification of *Aeromonas salmonicida* in clinical specimens. *Dis. Aquat. Organ.* 16, 239–242. doi:10.3354/dao016239.
- Toranzo, A. E., Santos, Y., Nunez, S., and Barja, J. L. (1991). Biochemical and serological characteristics, drug resistance and plasmid profiles of Spanish isolates of *Aeromonas salmonicida*. *Fish Pathol.* 26, 55–60. doi:10.3147/jsfp.26.55.
- Troy, T., Jekic-McMullen, D., Sambucetti, L., and Rice, B. (2004). Quantitative comparison of the sensitivity of detection of fluorescent and bioluminescent reporters in animal models. *Mol. Imaging* 3, 9–33.
- Trust, T. J., Kay, W. W., and Ishiguro, E. E. (1983). Cell surface hydrophobicity and macrophage association of *Aeromonas salmonicida*. *Curr. Microbiol.* 9, 315–318. doi:10.1007/BF01588826.
- Udey, L., and Fryer, J. (1978). Immunization of fish with bacterins of *Aeromonas salmonicida*. *Mar. Fish. Rev.* 40, 12 – 17.
- Vargas-Chacoffa, L., Martíneza, D., Oyarzúna, R., Nualartb, D., Olavarriab, V., Yáñezb, A., et al. (2014). Combined effects of high stocking density and *Piscirickettsia salmonis* treatment on the immune system, metabolism and osmoregulatory responses of the Sub-Antarctic Notothenioid fish *Eleginops maclovinus*. *Fish Shellfish Immunol.* 40, 424–434. doi:10.1016/j.fsi.2014.07.024.
- Vilches, S., Jimenez, N., Tomás, J. M., and Merino, S. (2009). *Aeromonas hydrophila* AH-3 type III secretion system expression and regulatory network. *Appl. Environ. Microbiol.* 75, 6382–6392. doi:10.1128/AEM.00222-09.
- Vilches, S., Urgell, C., Merino, S., Chacón, M. R., Soler, L., Castro-Escarpulli, G., et al. (2004). Complete type III secretion system of a mesophilic *Aeromonas hydrophila* strain. *Appl. Environ. Microbiol.* 70, 6914–6919. doi:10.1128/AEM.70.11.6914-6919.2004.
- Vinitnantharat, S., Gravningen, K., and Greger, E. (1999). Fish vaccines. *Adv. Vet. Sci. Comp. Med.* 41, 539 – 550.
- Vipond, R., Bricknell, I. R., Durant, E., Bowden, T. J., Ellis, A. E., Smith, M., et al. (1998). Defined deletion mutants demonstrate that the major secreted toxins are not essential for the virulence of *Aeromonas salmonicida*. *Infect. Immun.* 66, 1990–1998.
- Virostko, J. M., Powers, A. C., and Jansen, E. D. (2008). Validation of luminescent source reconstruction using spectrally resolved bioluminescence images. *Proc. Spie - Int. Soc. Opt. Eng.* 6849. doi:10.1117/12.771416.
- Weissleder, R., Tung, C. H., Mahmood, U., and Bogdanov, A. (1999). In vivo imaging of tumors with protease-activated near-infrared fluorescent probes. *Nat. Biotechnol.* 17, 375–378. doi:10.1038/7933.
- Wichardt, U. P., Johansson, N., and Ljungber, O. (1989). Occurrence and distribution of *Aeromonas salmonicida* infection on Swedish fish farms, 1951-1987. *J. Aquat. Anim. Health* 1, 187–196. doi:10.1577/1548-8667(1989)001.
- Wiklund, T., and Dalsgaard, I. (1998). Occurrence and significance of atypical *Aeromonas salmonicida* in non-salmonid and salmonid fish species: a review. *Dis. Aquat. Organ.* 32, 49–



69. doi:10.3354/dao032049.

Williamson, I. J. F. (1928). Furunculosis of the Salmonidae. *Fish. Board Scotland, salmon Fish.* 5, 17.

Willumsen, B. (1990). *Aeromonas salmonicida* subsp. *salmonicida* isolated from Atlantic cod and coalfish. *Bull. Eur. Assoc. Fish Pathol.* 10, 62–63.

Wilson, A., and Horne, M. T. (1986). Detection of A-protein in *Aeromonas salmonicida* and some effects of temperature on A-layer assembly. *Aquaculture* 56, 23–27. doi:10.1016/0044-8486(86)90287-5.

Windell, J. T., Norris, D. O., Kitchell, J. F., and Norris, J. S. (1969). Digestive response of rainbow trout, *Salmo gairdneri*, to pellet diets. *J. Fish. Res. Board Canada* 26, 1801–1812. doi:10.1139/f69-164.

Winnard, P. T., Kluth, J. B., and Raman, V. (2006). Noninvasive optical tracking of red fluorescent protein-expressing cancer cells in a model of metastatic breast cancer. *Neoplasia* 8, 796–806. doi:10.1593/neo.06304.

Xu, M., and Wang, L. V. (2006). Photoacoustic imaging in biomedicine. *Rev. Sci. Instrum.* 77. doi:10.1063/1.2195024.

Zhang, W., Jayarao, B. M., and Knabel, S. J. (2004). Multi-Virulence-Locus Sequence Typing of *Listeria monocytogenes*. *Appl. Environ. Microbiol.* 70, 913–920. doi:10.1128/AEM.70.2.913-920.2004.





**Accompanying manuscripts I - III**





## **Manuscript I**

**Infection routes of *Aeromonas salmonicida* in rainbow trout monitored *in vivo* by real-time bioluminescence imaging**



**Infection routes of *Aeromonas salmonicida* in rainbow trout monitored  
*in vivo* by real-time bioluminescence imaging**

1 **Simona Bartkova<sup>\*</sup>, Branko Kokotovic, Inger Dalsgaard**

2

3 National Veterinary Institute, Section for Bacteriology and Pathology, Technical University of  
4 Denmark, Frederiksberg C, Denmark

5

6 **\* Correspondence:** Simona Bartkova, National Veterinary Institute, Section for Bacteriology and  
7 Pathology, Technical University of Denmark, Bülowsvej 27, 1870 Frederiksberg C, Denmark

8

9 [sibar@vet.dtu.dk](mailto:sibar@vet.dtu.dk)

10

11 **Running title:** *Aeromonas salmonicida* bioluminescence imaging

12

13 **Keywords:** *Aeromonas salmonicida*, furunculosis, rainbow trout, bioluminescence imaging,  
14 *luxCDABE* genes



## 15 **Abstract**

16

17 Recent development of imaging tools has facilitated studies of pathogen infections *in vivo* in real  
18 time. This trend can be exemplified by advances in bioluminescence imaging (BLI), an approach  
19 that helps to visualize dissemination of pathogens within the same animal over several time points.  
20 Here we employ bacterial BLI for examining routes of entry and spread of *Aeromonas salmonicida*  
21 *susbp. salmonicida* in rainbow trout. A virulent Danish *A. salmonicida* strain was tagged with  
22 pAK*gfplux1*, a dual-labeled plasmid vector containing the mutated *gfpmut3a* gene from *Aequorea*  
23 *victoria* and the *luxCDABE* genes from the bacterium *Photorhabdus luminescens*. The resulting *A.*  
24 *salmonicida* transformant exhibited growth properties and virulence identical to the wild type *A.*  
25 *salmonicida*, which made it suitable for an experimental infection, mimicking natural conditions.  
26 Fish were infected with pAK*gfplux1* tagged *A. salmonicida* via immersion bath. Colonization and  
27 subsequent tissue dissemination was followed over a 24-hour period using the IVIS Spectrum  
28 imaging workstation. Results suggest the pathogen's colonization sites are the dorsal and pectoral  
29 fin and the gills, followed by a progression through the internal organs and an ensuing exit via the  
30 anal opening. The present work provides a tool for visualizing colonization of *A. salmonicida* and  
31 other bacterial pathogens in fish.

32

## 33 **Introduction**

34

35 *Aeromonas salmonicida susbp. salmonicida*, the causative agent of the disease furunculosis, is one  
36 of the major bacterial pathogens in aquaculture throughout the world. Furunculosis is a septicemic  
37 infection that was first described in Denmark in freshwater rainbow trout (*Oncorhynchus mykiss*)  
38 farms during the 1950s (Rasmussen 1964). Currently, furunculosis causes the greatest problems in

39 Danish rainbow trout production in sea water, where outbreaks occur during stress-associated  
40 periods with elevated temperatures around 20°C in July and August (Larsen & Møllgaard 1981;  
41 Dalsgaard & Madsen 2000; Pedersen, Skall, Lassen-Nielsen, Nielsen, Henriksen & Olesen 2008).  
42 This indicates a possibility of *A. salmonicida* being spread from freshwater into the sea via rainbow  
43 trout carriers (Dalsgaard & Madsen, 2000). However, attempts to isolate bacteria from these  
44 presumed carrier fish have not been successful so far (Dalsgaard & Madsen 2000). Consequently,  
45 discerning the sites of entry and dissemination pattern of *A. salmonicida* within fish could be an  
46 important factor for optimizing future sampling procedures and detection of the bacterium.

47  
48 Given the advances in the field of *in vivo* imaging in recent years, real-time monitoring of  
49 pathogens with various fluorescence and luciferase protein reporters has emerged as a promising  
50 strategy for following the pathogens' dissemination within their hosts (Contag, Contag, Mullins,  
51 Spilman, Stevenson & Benaron 1995; Rocchetta, Boylan, Foley, Iversen, Letourneau, McMillian,  
52 Contag, Jenkins & Parr 2001; Karsi, Menanteau-Ledouble & Lawrence 2006; Karsi & Lawrence  
53 2007; Zinn, Chaudhuri, Szafran, O'Quinn, Weaver, Dugger, Lamar, Kesterson, Wang & Frank  
54 2008; Menanteau-Ledouble, Karsi & Lawrence 2011). In comparison to fluorescence reporters,  
55 luciferase proteins that exhibit bioluminescence (BLI) show higher sensitivity, lower toxicity and  
56 faster response to changing environments (Burns, Joh, Francis, Shortliffe, Gruber, Contag & Contag  
57 2001; Troy, Jekic-McMullen, Sambucetti & Rice 2004). Moreover, there is scarcely any  
58 autoluminescence (BLI background noise) emitted by animal tissues, as opposed to other light  
59 sources where the background disturbance emitted by tissues usually affects the light-emitting  
60 reporters' sensitivity and overall applicability (Troy *et al.* 2004).

61

62 BLI has been applied for monitoring *A. salmonicida* in dead Atlantic salmon (*Salmo salar*)  
63 (Ferguson, Bricknell, Glover, MacGregor & Prosser 1998). However, requirement for an exogenous  
64 addition of aldehyde for BLI visualization and incorporation of luciferase genes into *A. salmonicida*  
65 significantly lowered the virulence of the bacterium (Ferguson *et al.* 1998). First application of BLI  
66 for real-time monitoring of bacteria in live fish was reported by Karsi *et al.* (2006) who used the  
67 method to investigate dissemination of *Edwardsiella ictaluri* inside channel catfish (*Ictalurus*  
68 *punctatus*). Recently, Méndez & Guijarro (2013) used BLI to trace dissemination of *Yersinia*  
69 *ruckeri* in rainbow trout. Both studies used a bacterial luciferase operon from *Photobacterium*  
70 *luminescens* consisting of five genes (*luxCDABE*) encoding the luciferase and fatty acid reductase  
71 enzyme complex, enabling emission of luminescence without the addition of any cofactors or  
72 exogenous substrates (Meighen 1993; Burns *et al.* 2001; Troy *et al.* 2004).

73

74 The aim of this study was to explore the routes of entry and subsequent tissue dissemination of *A.*  
75 *salmonicida* by using *in vivo* imaging and luciferase coding operon (Karsi *et al.* 2006; Méndez &  
76 Guijarro 2013). In short, a virulent *A. salmonicida* was tagged with a dual-labeled reporter plasmid  
77 containing a mutated green fluorescence protein (GFP) gene and the BLI coding genes (*luxCDABE*)  
78 under the same *lacZ* promoter that requires only oxygen for constitutive expression of both protein  
79 reporters (Karsi & Lawrence 2007). Expression of GFP provided a marker for verifying the  
80 presence of plasmid in the tagged bacterium and proper function of the *lacZ* promoter, while  
81 expression of BLI provided a marker for *in vivo* monitoring of a bacterium in the host after  
82 experimental infection.

83

## 84 **Materials and Methods**

85

86 **Bacterial strains and plasmid transfer**

87

88 *Aeromonas salmonicida* subsp. *salmonicida* 090710-1/23 (further referred to as *A. salmonicida*  
89 WT) is a virulent strain from a disease outbreak in Denmark. This strain has been used several times  
90 for various infection experiments in our and other Danish laboratories (e.g. Chettri, Skov, Jaafar,  
91 Krossøy, Kania, Dalsgaard & Buchmann 2015). *Escherichia coli* B/K 12 (Addgene, USA) is a  
92 donor strain that has an incorporated plasmid vector pAKgfplux1 consisting of the plasmid  
93 pBBR1MCS4 with an inserted: 1) *gfpmut3a* mutant gene from *Aequorea Victoria*, 2) *luxCDABE*  
94 operon from *Photobacterium luminescens*, and 3) an ampicillin resistance gene. The pAKgfplux1  
95 plasmid was first isolated from the *E. coli* B/K 12 donor strain by QIAprep Spin Miniprep Kit  
96 (Qiagen, Valencia, CA), according to the manufactures instructions. The plasmid was then  
97 transformed into *A. salmonicida* WT by conjugal mating using a natural kanamycin resistant *E. coli*  
98 strain SM10  $\lambda$ pir (Biomedal, Spain) as described by Karsi *et al.* (2007).

99

100 In details, natural kanamycin resistant SM10  $\lambda$ pir was transformed with pAKgfplux1 plasmid  
101 carrying ampicillin resistance by electroporation using a Gene Pulser instrument (Bio Rad) at 25  
102  $\mu$ FD, 200  $\Omega$ , 1.8 kV and with a time constant (tau value) of 5 msec. SM10  $\lambda$ pir ampicillin and  
103 kanamycin resistant colonies were grown overnight in 2 ml of Luria Bertani (LB) medium (Difco)  
104 at 37°C with shaking at 225 rpm. The recipient was grown separately for 48 h in Veal Infusion  
105 Broth (VIB) (Difco) at 20°C. Subsequently, 750  $\mu$ l of the donor *E. coli* SM10  $\lambda$ pir and 1.5 ml of  
106 recipient culture were centrifuged separately at 8700 g for 2 minutes and the supernatant was  
107 removed. Collected bacteria were then washed by resuspending the pellets in 750  $\mu$ l of Brain Heart  
108 Infusion (BHI) (Difco). Washing procedure was repeated three times in order to remove antibiotics  
109 used during broth culture. In 1.5 ml centrifuge tubes, 50  $\mu$ l of donor and 100  $\mu$ l of recipient washed

110 cells were mixed and bacteria were centrifuged as previously. Supernatant was poured off and  
111 donor and host mixture was suspended in 5-10  $\mu\text{l}$  of BHI. Punched and sterilized 0.45  $\mu\text{M}$  filters  
112 (Life GE Healthcare Life Sciences) were placed on blood agar plates (Columbia agar base (Oxoid)  
113 with 5% calf blood) without antibiotics and 5  $\mu\text{l}$  of bacterial mixture was transferred on the filter as  
114 a spot. Plates were returned to incubator set to suitable temperature for the recipient (20°C) and  
115 conjugation continued for 48 h. At the end of the conjugation period, filters were dropped into  
116 sterile 1.5 ml Eppendorf tubes and bacteria were washed away by adding 750  $\mu\text{l}$  of VIB with 100 $\mu\text{g}$   
117  $\text{ml}^{-1}$  ampicillin (Sigma-Aldrich). Ten microliters of bacterial suspension were mixed with 990  $\mu\text{l}$  of  
118 ddH<sub>2</sub>O and 25  $\mu\text{l}$  of the mixture was spread onto selective blood agar plates containing 100  $\mu\text{g ml}^{-1}$   
119 ampicillin and incubated for 48 h at 20°C. Incubation temperature 20°C ensured optimal growth of  
120 *A. salmonicida* and ampicillin selected for *A. salmonicida* with incorporated pAK*gfplux*1 (further  
121 referred to as *A. salmonicida gfplux*). *A. salmonicida gfplux* colonies showing the strongest  
122 fluorescence under an Axio imager M1 (Zeiss, Germany), were transferred into 5 ml VIB and  
123 incubated for 48 h at 20°C. The bacteria were subcultivated two times under these conditions and  
124 then two times using blood agar plates without ampicillin, from which a single *A. salmonicida*  
125 *gfplux* colony was selected for storage in glycerol stocks (600  $\mu\text{L}$  of 50% glycerol with 900  $\mu\text{L}$  of  
126 the 48 h grown bacterial culture in VIB) at -80°C until further use.

127

## 128 **Experimental fish**

129

130 Fertilized eggs of rainbow trout from Fousing Trout Farm (Jutland, Denmark) were brought to our  
131 institute, where disinfection, hatching and rearing were carried out under pathogen-free conditions.  
132 Fish were held at  $10 \pm 1^\circ\text{C}$  in 180-liter tanks containing a flow-through system with non-chlorinated  
133 tap water and air supply. Fish were fed dry commercial feed (Inicio Plus; BioMar A/S, Denmark) at

134 1% of biomass per day. Average weight and length of rainbow trout used for all challenges was 8.8  
135  $\pm 2.7$  g and  $9.3 \pm 1.7$  cm. Experimental infections were carried out in accordance with the accepted  
136 guidelines for the care and use of laboratory animals in research and with regulations set forward by  
137 the Danish Ministry of Justice and Animal Protection committees by Danish Animal Experiments  
138 Inspectorate permit number 2013-15-2934-00976.

139

#### 140 **Sensitivity of IVIS for detecting *A. salmonicida gfplux***

141

142 To determine the threshold detection limit for visualization of *A. salmonicida gfplux* using an IVIS  
143 spectrum imaging workstation (PerkinElmer) with an exposure time of 30 s, four separate two-fold  
144 serial dilutions in 0.9% saline solution ranging from  $8 \times 10^4$  -  $2 \times 10^2$  colony-forming-units (CFU)  
145  $\text{ml}^{-1}$  of the bacterium grown in VIB for 48 h at  $20^\circ\text{C}$ , were made in a black 96-well microtiter plate  
146 (Thermo Scientific) using a volume of 0.1 ml per well. To measure background noise  
147 (autoluminescence), aliquots of 0.1 ml 0.9% saline solution were also added to four wells as  
148 controls. The plate was scanned for 30 s. Relative intensity of luminescence emission for each well  
149 was estimated by IVIS software and represented with a pseudo-colour scale of counts  $\text{s}^{-1}$ . Each  
150 dilution series was then cultivated on blood agar plates to confirm  $\text{CFU ml}^{-1}$  and after 48 h growth  
151 at  $20^\circ\text{C}$ , all plates were visualized using an Axio imager M1 in order to observe fluorescence  
152 emitted from the bacterial colonies. Mean autoluminescence was subtracted from luminescence  
153 values of all *A. salmonicida gfplux* wells and correlation between  $\text{CFU ml}^{-1}$  and relative intensity of  
154 luminescence emission was determined.

155

#### 156 **Plasmid stability and effect of transformation on bacterial growth *in vitro***

157

158 Plasmid stability was previously investigated by Karsi *et al.* (2006) and Karsi and Lawrence (2007).  
159 Results from these studies suggested that the broad host range vector employed in this study can be  
160 transferred and stably maintained in Gram-negative bacteria.

161

162 In order to test whether the introduction of plasmid affected growth properties of the wild-type  
163 strain, bacterial growth of *A. salmonicida* WT and *A. salmonicida gflux* were compared: 1)  
164 indirectly using procedure by Karsi *et al.* (2006) for measuring optical density of bacterial culture  
165 and 2) directly by plating serial dilutions onto blood agar to obtain CFU ml<sup>-1</sup>. Readings of OD<sub>625</sub>  
166 were analyzed by a Student's paired t test using the Microsoft Excel statistical package. After the 48  
167 h period, a serial dilution was made from each bacterial culture and plated on blood agar to  
168 determine CFU ml<sup>-1</sup> and fluorescence emitted from the bacterial colonies was observed using an  
169 Axio imager M1. Determined CFU ml<sup>-1</sup> was compared by a Student's t test analysis using Microsoft  
170 Excel. The experiment was repeated three times.

171

#### 172 **Effect of transformation on *in vivo* virulence**

173

174 *In vivo* virulence of *A. salmonicida* WT and *A. salmonicida gflux* was compared by determination  
175 of CFU required to infect 50% of the fish population. Bacteria were grown in VIB for 48 h at 20°C.  
176 Subsequently, ten-fold serial dilutions of bacteria, containing 10<sup>8</sup> - 10<sup>4</sup> CFU ml<sup>-1</sup> were made in 0.9%  
177 saline solution. Each dilution was then used for inoculation of six fish. Each fish was inoculated  
178 intraperitoneally with 0.1 ml of the corresponding dilution, so the infection doses administered  
179 ranged from 10<sup>7</sup> - 10<sup>3</sup> CFU. One control group with six fish was injected with 0.1ml sterile veal  
180 infusion broth, and one control group with six fish was left uninjected. Fish were held at 10 ± 1°C  
181 in 10-liter tanks containing a flow-through system with non-chlorinated tap water and air supply.

182 Fish were observed several times per day and moribund fish were euthanized by immersion in 250  
183 mg l<sup>-1</sup> of 3-aminobenzoic acid ethyl ester (MS-222) (Sigma-Aldrich) until it was certain that  
184 swimming and gill movement had ceased. After two weeks the experiment was terminated. In order  
185 to confirm bacterial infection and to estimate plasmid stability during infection of the host, spleen,  
186 kidney and brain specimens were cultivated on blood agar and LB agar containing 100 µg ml<sup>-1</sup>  
187 ampicillin. Fluorescence emission in the isolated bacteria was observed by using an Axio imager  
188 M1. Calculation of ID50 was done according to Reed & Muench (1938) and Student's t test in  
189 Microsoft Excel was used to assess significance.

190

#### 191 **Experimental setup for real-time monitoring *A. salmonicida gfplux* in vivo**

192

193 Two fish were experimentally infected with *A. salmonicida gfplux* by separate immersion in two  
194 well-aerated 19 ± 1°C 10-liter tanks containing three liters of 5 × 10<sup>7</sup> CFU ml<sup>-1</sup> *A. salmonicida*  
195 *gfplux* for two hours. For visualization each fish was anesthetized with MS-222, immersed twice in  
196 distilled water and dipped on a paper towel before finally being laid on a Tissue Culture Dish  
197 (Greiner Bio-One, Germany) and scanned in the IVIS for 30 s to estimate relative intensity of  
198 luminescence emission (counts s<sup>-1</sup>). After scanning, each fish was placed in a new well-aerated 10-  
199 liter tank containing three liters of distilled water for recovery. Visualization was performed at the  
200 following time points: 2, 4, 6 and 24 h after immersion in the infection bath. Fish were euthanized  
201 before the last visualization time-point (24 h). Scanning in the IVIS was done on the whole fish and  
202 for the last scanning also on fish that were cut open to expose internal organs. Finally, for  
203 bacteriology examination, kidney and spleen samples were taken from each fish and streaked on  
204 blood agar that was incubated for 48 h at 20°C. Subsequently all plates were examined in an Axio  
205 imager M1 microscope for fluorescence emission. One non-infected (control) fish was also scanned



206 in the IVIS as whole and cut open for monitoring of autoluminescence (counts s<sup>-1</sup>) emitted from  
207 different external as well as internal areas. Obtained autoluminescence was subtracted from all  
208 luminescence readings of infected fish in order to acquire the correct luminescence (counts s<sup>-1</sup>) for  
209 each scanning, which could then be correlated to a CFU count and be represented with a pseudo-  
210 colour scale. The experiment was repeated three times using one fish per tank and two times using  
211 two fish per tank. A total of fourteen infected fish and five non-infected control fish were examined.

212

## 213 **Results**

214

### 215 **Detection limit of *A. salmonicida gfplux***

216

217 Visualization and measurement of the *A. salmonicida gfplux* two-fold dilutions showed that for the  
218 exposure time of 30 s, the threshold detection limit for visualization of *A. salmonicida gfplux*  
219 appears to be 4 x 10<sup>4</sup> CFU ml<sup>-1</sup>. Correlation between CFU ml<sup>-1</sup> and measured relative intensity of  
220 luminescence emissions was linear (R<sup>2</sup> = 0.977) over the range of 8 x 10<sup>4</sup> - 2 x 10<sup>2</sup> CFU ml<sup>-1</sup> (Fig.  
221 1), indicating luminescence emission should present the accurate CFU values for a given sample.  
222 Average percent of *A. salmonicida gfplux* colonies emitting fluorescence on a blood agar plate was  
223 95% (data not shown).

224

### 225 **[Figure 1]**

226

### 227 **Effect of transformation on bacterial growth *in vitro***

228

229 No significant difference was observed between *A. salmonicida* WT and *A. salmonicida gfplux*  
230 growth as measured by the indirect method of OD<sub>625</sub> when analysed with a Student's paired t test (p  
231 > 0.05). Bacterial CFU ml<sup>-1</sup> after 48 h of growth at 20°C for each experiment was analysed by a  
232 Student's t test and showed no significant difference (p > 0.05) in all experimental repeats (Table  
233 1). Average percent of *A. salmonicida gfplux* colonies emitting fluorescence on a blood agar plate in  
234 all experimental repeats was 96%, while no fluorescence emission was observed in any *A.*  
235 *salmonicida* WT colonies (data not shown).

236

237 **[Table 1]**

238

239 **Effect of transformation on *in vivo* virulence**

240

241 The ID50 value for *A. salmonicida* WT was 5 x 10<sup>5</sup> CFU and for *A. salmonicida gfplux* 6 x 10<sup>5</sup>  
242 CFU. Insertion of pAK*gfplux*1 into *A. salmonicida* WT did not seem to affect *in vivo* virulence of  
243 the bacterium. *A. salmonicida gfplux* colonies were re-isolated from all three sampled organs i.e.  
244 kidney, spleen and brain in euthanized fish. The number of colonies emitting fluorescence on blood  
245 agar plates gradually decreased over the course of the experiment and after 10 days fluorescence  
246 emission could not be detected any longer (Table 2). Colonies grown on LB agar with ampicillin  
247 showed consistently strong fluorescence but their number rapidly decreased. From day 5 of the  
248 experiment only few colonies grew on LB agar and after the day 8 no growth was recorded (Table  
249 2).

250

251 **[Table 2]**

252

253 **Experimental infection for real-time monitoring *A. salmonicida* *gfplux* in vivo**

254

255 For the experimental infection, luminescence signal from *A. salmonicida gfplux* was observed in  
256 overall twelve of the fourteen examined fish following a two-hour immersion time (Table 3). At the  
257 2-hour time point, a luminescence signal was detected on eight of the twelve positive fish. Bacteria  
258 were visualized on the following sites: the dorsal, pectoral, caudal and anal fin, anal opening, gills,  
259 oral and nasal cavity and eyes (Table 3; Fig. 2).

260

261 **[Figure 2]**

262

263 For the 4- and 6-hour time points, luminescence emission could be seen in three fish (Table 3). In  
264 two fish, the gills where observed luminescence was found at the 2-hour time point were still  
265 emitting luminescence at the 4-hour time point. Luminescence was also observed in the body organ  
266 area in one fish at the 6-hour time point, which was presumed to be emitted from the inside of the  
267 fish.

268

269 **[Table 3]**

270

271 After 24 hours, luminescence was found in four of the total of twelve positive fish (Table 3).  
272 Among the ten fish where the luminescent bacteria were found, in three fish the luminescence  
273 signal was strong enough to be seen through the skin in whole fish (Table 3; Fig. 3). In one of the  
274 three fish, the luminescence signal was located around the anal opening and after being cut open;  
275 the signal was still located around that area and the lower intestine (Fig. 3). The seven remaining  
276 fish needed to be cut open in order to be able to detect a luminescence signal. In all ten positive fish

277 at the 24-hour time point, the signal was located in the intestine (Table 3). In one fish the signal was  
278 also located in the stomach (Table 3 and Fig. 3). In two fish the signal seemed to be located both in  
279 the intestine and spleen. Bacteriological examination of kidney and spleen samples was positive for  
280 *A. salmonicida gfplux* in all fourteen fish. However, very faint or no fluorescence emission was  
281 observed from these colonies (data not shown). In all five non-infected control fish no bacteria were  
282 isolated and measured autoluminescence emission was scarce compared to measured luminescence  
283 emission from infected fish (negative data not shown).

284

285 **[Figure 3]**

286

## 287 **Discussion**

288

289 Experimental infections with *A. salmonicida gfplux* provided an indication of potentially important  
290 colonization sites of *A. salmonicida*. However, colonization and dissemination of *A. salmonicida*  
291 *gfplux* in fish could only be visualized in twelve of the fourteen experimentally infected fish and  
292 after 24 hours bacteria were only visualized in the digestive system, while bacteria were re-isolated  
293 from the spleen and kidney in all fourteen fish after 24 hours. The lack of visualization in other  
294 organs could be due to the bacterial amount being below the threshold detection limit and/or a lack  
295 of plasmid stability. The later scenario is supported by the fact that the re-isolated bacteria on blood  
296 agar plates either emitted very little or no luminescence at all. Previous studies using plasmids with  
297 the same luciferase coding operon as in this study did not report on any difficulties regarding  
298 plasmid stability within fish (Karsi *et al.* 2006; Menanteau-Ledouble *et al.* 2011; Méndez &  
299 Guijarro 2013). It is not possible to explain the reason(s) for the observed instability of the plasmid  
300 within fish in this study.

301

302 When comparing the threshold detection limit of BLI to previous studies, the limit of  $4 \times 10^4$  CFU  
303  $\text{ml}^{-1}$  in this study is higher than the  $10^3$  CFU  $\text{ml}^{-1}$  as reported by both Karsi *et al.* (2006) and  
304 Méndez & Guijarro (2013). The luminescence signal is proportional to exposure time i.e. the  
305 duration a sample is scanned for (Caliper Life Sciences). In this study fish were scanned for 30 s as  
306 oppose to 1 min in the previous studies (Karsi *et al.* 2006; Méndez & Guijarro 2013). We have  
307 chosen a shorter exposure time taking into the consideration: 1) uncertainty of the anaesthetic  
308 effectiveness over longer durations than 30 s, 2) minimizing chance of obtaining false positives and  
309 3) the overall well-being of the fish.

310

311 In this study, fins were suggested as one of the key colonization sites of *A. salmonicida*. This results  
312 is in agreement with the findings by Hiney, Kilmartin & Smith (1994), who used ELISA to detect  
313 *A. salmonicida* in pre-smolt Atlantic salmon with stress-inducible furunculosis infections. Fins  
314 were also found as major colonization sites in other fish pathogens (Martinez, Casado & Enriquez  
315 2004; Harmache, LeBerre, Droineau, Giovannini & Brémont 2006; Menanteau-Ledouble *et al.*  
316 2011). One reason why fins and especially the dorsal fin seem to be an important attachment site  
317 could be related to bite wounds (Jobling, Jørgensen & Christiansen 1993). Consistently Svendsen &  
318 Bøgwald (1997) found *A. salmonicida* infected salmon with artificial wounds exhibiting higher  
319 mortality than infected salmon with no wounds. In contrast, fish used in our study did not have any  
320 injuries around the fins during the experiment.

321

322 No luminescence emission from the skin was observed in this study and the role of skin as possible  
323 colonization site of *A. salmonicida* found in previous studies by Svendsen & Bøgwald (1997),  
324 Ferguson *et al.* (1998), Cipriano, Ford, Teska & Hale (1992) and Cipriano, Ford, Schachte & Petrie

325 (1994) could not be confirmed. Still, given the relatively high threshold limit of luminescence  
326 detection, an attachment of a low number of bacteria to the skin cannot be ruled out. An alternative  
327 explanation is that teleost (bony) fish feature variation in their immune system (Svendsen, Dalmo &  
328 Bøggwald 1999), including varying mucosal activity against pathogens (Dickerson 2009). Rainbow  
329 trout could thus have a better mucosal protection against pathogens compared to many other farmed  
330 fish species. This hypothesis is supported by other studies where rainbow trout showed the highest  
331 degree of resistance against furunculosis compared to other farmed fish species (Cipriano &  
332 Heartwell 1986).

333

334 A strong BLI signal was also seen around the gills, indicating that this might be an important  
335 colonization site as well. This finding is consistent with both Tatner, Johnson & Horne (1984) who  
336 investigated *A. salmonicida* infection in rainbow trout and with Svendsen *et al.* (1999) who studied  
337 *A. salmonicida* infection in Atlantic salmon. The gills also seem to be an important colonization site  
338 for other bacterial fish pathogens like *Yersinia ruckeri*, demonstrated by Ohtani, Villumsen, Strøm  
339 & Raida (2014) who observed infection of the gill epithelial cells as early as one minute post  
340 infection.

341

342 Our study also indicates the oral and nasal cavity and the eyes might be colonization sites for *A.*  
343 *salmonicida*. These three sites are all 'open', i.e. lacking the primary barrier of the skin as a defense  
344 against pathogens (Roberts & Ellis 2012). The mouth has also been proposed as a possible entry  
345 route for *A. salmonicida* by Svendsen & Bøggwald (1997). Moreover all three sites were seen to be  
346 probable colonization sites for *Novirhabdovirus* in a bioluminescence experiment conducted on  
347 juvenile trout (Harmache *et al.* 2006). Finally, Karsi *et al.* (2006) reported that bioluminescent *E.*  
348 *ictaluri* became visible around the eye and mouth area during early disease progression.

349

350 At the 4- and 6-hour time points, luminescence was detected in only three of the infected fish,  
351 which could, at least in part, be due to the relatively low sensitivity of the method. In two fish, the  
352 gills that were found positive at the 2-hour time point were still positive. A luminescence emission  
353 signal was observed at the 6-hour time point at the location of the digestive system in one fish,  
354 which had not been detected at the 2- and 4-hour time point. Due to the limitation of only being able  
355 to acquire two-dimensional images by this method, it was not possible to ascertain whether the  
356 luminescence signal came from the inside or the outside of the fish. Though it is likely that the  
357 signal came from inside the fish, since the fish had been transferred to clean water after the 2-hour  
358 infection immersion time and it would be expected that the bacteria can progress into the fish after a  
359 few hours as seen in the study with *Y. ruckeri* (Méndez & Guijarro 2013). In support of this, the  
360 luminescence signal intensified at the same location from the 6-hour time point to the 24-hour time  
361 point and when the abdominal area was opened revealing the internal organs, a luminescence signal  
362 was observed in the intestine (Table 3).

363

364 At the final 24-hour time point, bacteria were seen by imaging in the digestive system and in two  
365 fish, also in the spleen. Bacteria were re-isolated from spleen and kidney in all fourteen fish. The  
366 reason why bacteria in the kidney and spleen were not detected by imaging could be that the  
367 bacteria had lost the pAK*gfplux1* plasmid, as supported by the lack of fluorescence emission by the  
368 re-isolated colonies on blood agar plates, or the bacterial amount was below the threshold detection  
369 limit. Dissemination of *A. salmonicida* in the two organs is in agreement with Svendsen *et al.*  
370 (1999) who found *A. salmonicida* in blood already after 2 hours post infection and thereafter in  
371 kidney and a strong correlation between bacterial amounts in blood and gill tissue samples  
372 (Svendsen *et al.* 1999). It is known that bacteria spread from gills to the blood (Dickerson 2009),

373 which is then filtered by the kidney and spleen (Tatner *et al.* 1984; Hadidi, Glenney, Welch,  
374 Silverstein & Wiens 2008).

375

376 The imaging results at the 24-hour time point revealed consistently signal from organs after  
377 dissection, indicating dissemination of *A. salmonicida* in fish through the digestive system and that  
378 this can happen after 24 hours. This scenario is similar to the gut dissemination pattern seen in the  
379 study with *Y. ruckeri* (Méndez & Guijarro 2013). Radiolabelled *A. salmonicida* have also  
380 previously been found in tissue of the gut (Svendsen *et al.* 1999). In four fish we did not find any  
381 luminescent bacteria by 24 hours. Though, at the 2-hour time-point bacteria were seen in three of  
382 these fish, indicating that the initial bacteria that entered the digestive system could have passed  
383 through the fish in less than 24 hours. This is in agreement with research on digestive responses to  
384 feed pellets in rainbow trout, where gastric evacuation was seen after only 4 - 6 hours (Windell,  
385 Norris, Kitchell & Norris 1969). Further investigations are needed to shed more light on the role of  
386 digestive system in dissemination of *A. salmonicida*.

387

388 In summary, our results indicate that the dorsal and pectoral fin and gills are important colonization  
389 sites for *A. salmonicida* in rainbow trout. Novel information regarding *A. salmonicida* tissue  
390 dissemination pattern was also revealed, including the possible significant role of the digestive  
391 system. The bioluminescence-based *A. salmonicida* *gfplux* model used in the present study provides  
392 a valuable tool for *in vivo* real-time imaging of *A. salmonicida* and studying host-pathogen  
393 interaction.

394

395 **Acknowledgements**

396



397 This work was supported by The Danish Council for Strategic Research under ProFish project  
398 (Grant no. DSF: 11-116252) and the National Veterinary Institute (DTU). The authors would like to  
399 thank all ProFish partners. Mark Lee Lawrence and Simon Menanteau-Ledouble are thanked for  
400 their advice on plasmid incorporation and bioluminescence experiments with fish. The authors are  
401 also grateful to Martin Weiss Nielsen, Mette Boye, Lisbeth Schade Hansen and Anastasia Isbrand at  
402 the National Veterinary Institute (DTU) for their technical support, as well as Kamilla Ellerman  
403 Jensen and Petra Hamerlik at the Danish Cancer Society for their collaboration and introduction to  
404 the IVIS Spectrum Imaging Workstation.

405

## 406 **References**

407

- 408 Burns S.M., Joh D., Francis K.P., Shortliffe L.D., Gruber C.A., Contag P.R. & Contag C.H. (2001)  
409       Revealing the spatiotemporal patterns of bacterial infectious diseases using bioluminescent  
410       pathogens and whole body imaging. *Contributions To Microbiology* **9**, 71–88.
- 411 Caliper Life Sciences *Living Image® Software User's Manual version 4.2*.
- 412 Chettri J.K., Skov J., Jaafar R.M., Krossøy B., Kania P.W., Dalsgaard I. & Buchmann K. (2015)  
413       Comparative evaluation of infection methods and environmental factors on challenge success:  
414       *Aeromonas salmonicida* infection in vaccinated rainbow trout. *Fish & Shellfish Immunology*  
415       **44**, 485–95.
- 416 Cipriano R.C., Ford L.A., Schachte J.H. & Petrie C. (1994) Evaluation of mucus as a valid site to  
417       isolate *Aeromonas salmonicida* among asymptomatic populations of lake trout (*Salvelinus*  
418       *namaycush*). *Biomedical Letters* **49**, 229–233.
- 419 Cipriano R.C., Ford L.A., Teska J.D. & Hale L.H. (1992) Detection of *Aeromonas salmonicida* in  
420       the mucus of salmonid fishes. *Journal of Aquatic Animal Health* **4**, 114–118.

- 421 Cipriano R.C. & Heartwell C.M. (1986) Susceptibility of Salmonids to Furunculosis: Differences  
422 between Serum and Mucus Responses against *Aeromonas salmonicida*. *Transactions of the*  
423 *American Fisheries Society* **115**, 83–88.
- 424 Contag C.H., Contag P.R., Mullins J.I., Spilman S.D., Stevenson D.K. & Benaron D.A. (1995)  
425 Photonic detection of bacterial pathogens in living hosts. *Molecular Microbiology* **18**, 593–  
426 603.
- 427 Dalsgaard I. & Madsen L. (2000) Bacterial pathogens in rainbow trout, *Oncorhynchus mykiss*  
428 (Walbaum), reared at Danish freshwater farms. *Journal of Fish Diseases* **23**, 199 – 209.
- 429 Dickerson H.W. (2009) The biology of teleost mucosal immunity. In: *Fish Defenses, Volume 2:*  
430 *Pathogens, Parasites and Predators*. (ed. by Z. Giacomo, C. Perrière, A. Mathis & B.G.  
431 Kapoor), pp. 2–42. CRC Press.
- 432 Ferguson Y., Bricknell I.R., Glover L.A., MacGregor D.M. & Prosser J.I. (1998) Colonisation and  
433 transmission of lux-marked and wild-type *Aeromonas salmonicida* strains in Atlantic salmon  
434 (*Salmo salar* L.). *Fems Microbiology Ecology* **27**, 251–260.
- 435 Hadidi S., Glenney G.W., Welch T.J., Silverstein J.T. & Wiens G.D. (2008) Spleen size predicts  
436 resistance of rainbow trout to *Flavobacterium psychrophilum* challenge. *Journal of*  
437 *Immunology* **180**, 4156–4165.
- 438 Harmache A., LeBerre M., Droineau S., Giovannini M. & Brémont M. (2006) Bioluminescence  
439 imaging of live infected salmonids reveals that the fin bases are the major portal of entry for  
440 *Novirhabdovirus*. *Journal of Virology* **80**.
- 441 Hiney M.P., Kilmartin J.J. & Smith P.R. (1994) Detection of *Aeromonas salmonicida* in atlantic  
442 salmon with asymptomatic furunculosis infections. *Diseases of Aquatic Organisms* **19**, 161 –  
443 167.
- 444 Jobling M., Jørgensen E.H. & Christiansen J.S. (1993) Growth performance of salmonids exposed

445 to different flow regimes. In: *Fish Farming Technology. Proceedings of the First International*  
446 *Conference of Fish Farming*. (ed. by H. Reinertsen, L.A. Dahle, L. Jørgensen & K.  
447 Tvinnereim), pp. 301–302. A.A. Balkema, Rotterdam, Netherlands.

448 Karsi A. & Lawrence M.L. (2007) Broad host range fluorescence and bioluminescence expression  
449 vectors for Gram-negative bacteria. *Plasmid* **57**, 286–295.

450 Karsi A., Menanteau-Ledouble S. & Lawrence M.L. (2006) Development of bioluminescent  
451 *Edwardsiella ictaluri* for noninvasive disease monitoring. *FEMS Microbiology Letters* **260**,  
452 216–223.

453 Larsen J.L. & Møllergaard S. (1981) Microbiological and hygienic problems in marine aquaculture:  
454 furunculosis and vibriosis in rainbow trout (*Salmo gairdneri*). *Bull. Eur. Ass. Fish Pathol.*, 29–  
455 32.

456 Martínez J.L., Casado A. & Enriquez R. (2004) Experimental infection of *Flavobacterium*  
457 *psychrophilum* in fins of Atlantic salmon *Salmo salar* revealed by scanning electron  
458 microscopy. *Diseases of Aquatic Organisms* **59**, 79–84.

459 Meighen E.A. (1993) Bacterial bioluminescence - organization, regulation, and application of the lux  
460 genes. *Faseb Journal* **7**, 1016–1022.

461 Menanteau-Ledouble S., Karsi A. & Lawrence M.L. (2011) Importance of skin abrasion as a  
462 primary site of adhesion for *Edwardsiella ictaluri* and impact on invasion and systematic  
463 infection in channel catfish *Ictalurus punctatus*. *Veterinary Microbiology* **148**, 425–430.

464 Méndez J. & Guijarro J.A. (2013) In vivo monitoring of *Yersinia ruckeri* in fish tissues: progression  
465 and virulence gene expression. *Environmental Microbiology Reports* **5**, 179–185.

466 Ohtani M., Villumsen K.R., Strøm H.K. & Raida M.K. (2014) 3D Visualization of the Initial  
467 *Yersinia ruckeri* Infection Route in Rainbow Trout (*Oncorhynchus mykiss*) by Optical  
468 Projection Tomography. *PLoS ONE* **9**.

469 Pedersen K., Skall H.F., Lassen-Nielsen A.M., Nielsen T.F., Henriksen N.H. & Olesen N.J. (2008)  
470 Surveillance of health status on eight marine rainbow trout, *Oncorhynchus mykiss* (Walbaum),  
471 farms in Denmark in 2006. *Journal of Fish Diseases* **31**, 659–67.

472 Rasmussen C.J. (1964) Furunkulose hos damørreder (Furunculosis in farmed trout) (in Danish).  
473 *Scandinavian Journal of Veterinary Science* **16**, 463–472.

474 Reed L.J. & Muench H.H. (1938) A simple method of estimating fifty percent endpoints. *American*  
475 *Journal of Hygiene* **27**, 493–497.

476 Roberts R.J. & Ellis A.E. (2012) The Anatomy and Physiology of Teleosts. In: *Fish Pathology*. (ed.  
477 by R.J. Roberts), pp. 17–62. Wiley - Blackwell, UK.

478 Rocchetta H.L., Boylan C.J., Foley J.W., Iversen P.W., Letourneau D.L., McMillian C.L., Contag  
479 P.R., Jenkins D.E. & Parr T.R. (2001) Validation of a noninvasive, real-time imaging  
480 technology using bioluminescent *Escherichia coli* in the neutropenic mouse thigh model of  
481 infection. *Antimicrobial Agents and Chemotherapy* **45**, 129–137.

482 Svendsen Y.S. & Bøggwald J. (1997) Influence of artificial wound and non-intact mucus layer on  
483 mortality of Atlantic salmon (*Salmo salar* L.) following a bath challenge with *Vibrio*  
484 *anguillarum* and *Aeromonas salmonicida*. *Fish and Shellfish Immunology*, **4**, 317–325.

485 Svendsen Y.S., Dalmo R.A. & Bøggwald J. (1999) Tissue localization of *Aeromonas salmonicida* in  
486 Atlantic salmon, *Salmo salar* L., following experimental challenge. *Journal of Fish Diseases*  
487 **22**, 125–131.

488 Tatner M.F., Johnson C.M. & Horne M.T. (1984) The tissue localization of *Aeromonas salmonicida*  
489 in rainbow trout, *Salmo gairdneri* Richardson, following three methods of administration.  
490 *Journal of Fish Biology* **25**, 95–108.

491 Troy T., Jekic-McMullen D., Sambucetti L. & Rice B. (2004) Quantitative comparison of the  
492 sensitivity of detection of fluorescent and bioluminescent reporters in animal models.

493 *Molecular Imaging* **3**, 9–33.

494 Windell J.T., Norris D.O., Kitchell J.F. & Norris J.S. (1969) Digestive response of rainbow trout,  
495 *Salmo gairdneri*, to pellet diets. *Journal of the Fisheries Research Board of Canada* **26**, 1801–  
496 1812.

497 Zinn K.R., Chaudhuri T.R., Szafran A.A., O’Quinn D., Weaver C., Dugger K., Lamar D., Kesterson  
498 R.A., Wang X. & Frank S.J. (2008) Noninvasive Bioluminescence Imaging in Small Animals.  
499 *Institute for Laboratory Animal Research Journal* **49**, 103–115.

500 **Tables**

501

502 **Table 1.** Bacterial CFU ml<sup>-1</sup> comparison of wild type *A. salmonicida* WT and *A. salmonicida gfplux*  
 503 after 48 h growth at 20°C, including p-values of a Student's t test, for three experimental repeats.

Experiment	Strain	log <sub>10</sub> CFU ml <sup>-1</sup> ± SD	p-value
1	<i>A. salmonicida</i> WT	8.85 ± 0.08	0.972
	<i>A. salmonicida gfplux</i>	8.85 ± 0.10	
2	<i>A. salmonicida</i> WT	8.51 ± 0.12	0.393
	<i>A. salmonicida gfplux</i>	8.45 ± 0.06	
3	<i>A. salmonicida</i> WT	8.69 ± 0.06	0.105
	<i>A. salmonicida gfplux</i>	8.80 ± 0.10	

504

505

506

507

508

509

510

511 **Table 2.** Percent *A. salmonicida gfplux* colonies emitting fluorescence on blood agar plates and  
 512 Luria Bertani (LB) plates with 100µg ml<sup>-1</sup> ampicillin. Colonies were re-isolated on agar plates from  
 513 the kidney, spleen, and brain in fish during the ID50 experiment where fish were injected with  
 514 different dilutions of *A. salmonicida gfplux* culture.

Blood agar plates		LB plates with 100µg ml <sup>-1</sup> ampicillin	
Colonies emitting fluorescence <sup>α</sup>	Period (day)	Colonies emitting fluorescence <sup>α</sup>	Period (day)
30-60 %	3-4	100 %	3-4
10-30 %	5-8	100 % *	5-8
10 > %	9-10	NA #	9-10
0 %	11-14	NA #	11-14

<sup>α</sup> Average percent of fluorescence emitting colonies on a plate, isolated from each organ i.e. kidney, spleen and the brain

\* There were very few colonies on the LB plates compared to the number of colonies on the respective blood agar plates

# There was no growth on the plates

515 **Table 3.** Overview of results from the experimental infection. A total of 14 fish infected with *A.*  
516 *salmonicida gfplux* were scanned for 30 s in an IVIS spectrum imaging workstation for detection of  
517 luminescence emission at four time points: 2 h, 4 h, 6 h, and 24 h post infection in five independent  
518 experiments. Fish from the same experiment are grouped together and given an identification  
519 number. All areas of the fish where luminescence was observed at least once are displayed on the  
520 left. Observed luminescence signal is presented with an ‘X’. No luminescence signal was observed  
521 from fish 1.1 and 2.3.

Time point	Area	Fish #													
		1.1	2.1	1.2	2.2	1.3	2.3	1.4a	1.4b	2.4a	2.4b	1.5a	1.5b	2.5a	2.5b
2 h	Caudal fin		X												
	Anal fin											X			
	Dorsal fin		X		X			X				X	X		X
	Pectoral fin		X	X	X								X		
	Anal opening							X							
	Gills		X		X			X	X						
	Nasal cavity		X										X		
	Oral cavity		X										X		
	Eyes		X												
4 h	Gills				X			X							
6 h	Inside of fish					X									
24 h uncut	Anal opening							X							
	Inside of fish					X							X	X	
24 h organs	Stomach					X									
	Upper Intestine				X	X				X			X	X	
	Middle Intestine			X	X	X				X	X	X	X	X	
	Lower Intestine			X		X		X		X			X	X	
	Spleen			X*	X*										

\* In a few fish, organ structure at the 24 h time point was deteriorated, making it difficult to distinguish which organ the luminescence signal was coming from, however, it was believed that in two fish the luminescence signal could be emitted from the spleen.

522

523

524

525

526

527

528 **Figure legends**

529

530 **Figure 1.** Correlation between measured relative intensity of luminescence emission (counts s<sup>-1</sup>)  
531 and CFU ml<sup>-1</sup> for *A. salmonicida gfplux* serial dilutions ranging from 8 x 10<sup>4</sup> - 2 x 10<sup>2</sup> CFU ml<sup>-1</sup>  
532 after being scanned for 30 s in an IVIS spectrum imaging workstation. Correlation between CFU  
533 ml<sup>-1</sup> and relative intensity of luminescence emission for *A. salmonicida gfplux* was determined to be  
534 linear ( $R^2 = 0.977$ ) over the range of 8 x 10<sup>4</sup> - 2 x 10<sup>2</sup> CFU ml<sup>-1</sup>. The correlation coefficient, slope  
535 and intercept of the linear regression curve are shown.


536

537 **Figure 2.** Three bioluminescence imaging illustrations from the 2-hour time point of the  
538 experimental infection of rainbow trout with *A. salmonicida gfplux*. Illustrations show *A.*  
539 *salmonicida gfplux* colonization of the dorsal and pectoral fin and gills of the following three fish  
540 from Table 3: 1.4b, 2.2 and 2.5a.


541

542 **Figure 3.** Six bioluminescence imaging illustrations from the 24-hour time point of the  
543 experimental infection of rainbow trout with *A. salmonicida gfplux*. After 24 hours fish were  
544 euthanized and visualized in the IVIS as whole fish and were then cut open. In uncut fish 1.3,  
545 luminescence signal was observed in the body organ area and when cut open, the signal was  
546 observed in the stomach and upper, middle and lower intestine. In uncut fish 1.5b, luminescence  
547 signal was observed in the body organ area and when cut open, the signal was observed in the  
548 upper, middle and lower intestine. In uncut fish 1.4a, luminescence signal was observed around the  
549 anal opening and when cut open, the signal was observed around the anal opening and in the lower  
550 intestine.






**Figure 1.** Correlation between measured relative intensity of luminescence emission (counts s<sup>-1</sup>) and CFU ml<sup>-1</sup> for *A. salmonicida gfplux* serial dilutions ranging from 8 x 10<sup>4</sup> - 2 x 10<sup>2</sup> CFU ml<sup>-1</sup> after being scanned for 30 s in an IVIS spectrum imaging workstation. Correlation between CFU ml<sup>-1</sup> and relative intensity of luminescence emission for *A. salmonicida gfplux* was determined to be linear (R<sup>2</sup> = 0.977) over the range of 8 x 10<sup>4</sup> - 2 x 10<sup>2</sup> CFU ml<sup>-1</sup>. The correlation coefficient, slope and intercept of the linear regression curve are shown.



**Figure 2.** Three bioluminescence imaging illustrations from the 2-hour time point of the experimental infection of rainbow trout with *A. salmonicida gfplx*. Illustrations show *A. salmonicida gfplx* colonization of the dorsal and pectoral fin and gills of the following three fish from Table 3: 1.4b, 2.2 and 2.5a.



**Figure 3.** Six bioluminescence imaging illustrations from the 24-hour time point of the experimental infection of rainbow trout with *A. salmonicida gfplux*. After 24 hours fish were euthanized and visualized in the IVIS as whole fish and were then cut open. In uncut fish 1.3, luminescence signal was observed in the body organ area and when cut open, the signal was observed in the stomach and upper, middle and lower intestine. In uncut fish 1.5b, luminescence signal was observed in the body organ area and when cut open, the signal was observed in the upper, middle and lower intestine. In uncut fish 1.4a, luminescence signal was observed around the anal opening and when cut open, the signal was observed around the anal opening and in the lower intestine.





## **Manuscript II**

**Detection and quantification of *Aeromonas salmonicida* in fish tissue  
by real-time PCR**





## Detection and quantification of *Aeromonas salmonicida* in fish tissue by real-time PCR

S Bartkova<sup>1</sup>, B Kokotovic<sup>1</sup>, H F Skall<sup>2</sup>, N Lorenzen<sup>2</sup> and I Dalsgaard<sup>1</sup>

1 National Veterinary Institute, Section for Bacteriology and Pathology, Technical University of Denmark, Frederiksberg C, Denmark

2 Department of Animal Science, Aarhus University, Aarhus N, Denmark

### Abstract

Furunculosis, a septicaemic infection caused by the bacterium *Aeromonas salmonicida* subsp. *salmonicida*, currently causes problems in Danish seawater rainbow trout production. Detection has mainly been achieved by bacterial culture, but more rapid and sensitive methods are needed. A previously developed real-time PCR assay targeting the plasmid encoded *aopP* gene of *A. salmonicida* was, in parallel with culturing, used for the examination of five organs of 40 fish from Danish freshwater and seawater farms. Real-time PCR showed overall a higher frequency of positives than culturing (65% of positive fish by real-time PCR compared to 30% by a culture approach). Also, no real-time PCR-negative samples were found positive by culturing. *A. salmonicida* was detected by real-time PCR, though not by culturing, in freshwater fish showing no signs of furunculosis, indicating possible presence of carrier fish. In seawater fish examined after an outbreak and antibiotics treatment, real-time PCR showed the presence of the bacterium in all examined organs (1–482 genomic units mg<sup>-1</sup>). With a limit of detection of 40 target copies (1–2 genomic units) per reaction, a high reproducibility and an excellent efficiency, the present real-time PCR assay provides a sensitive tool for the detection of *A. salmonicida*.

**Keywords:** *Aeromonas salmonicida*, *aopP*, furunculosis, pAsal1, rainbow trout, real-time PCR.

**Correspondence** S Bartkova, National Veterinary Institute, Section for Bacteriology and Pathology, Technical University of Denmark, Bülowsvej 27, 1870 Frederiksberg C, Denmark (e-mail: sibar@vet.dtu.dk)

### Introduction

*Aeromonas salmonicida* subsp. *salmonicida* is the causative agent of furunculosis, a septicaemic infection that over the years has caused worldwide losses in aquaculture (O'Brien, Mooney, Ryan, Powell, Hiney, Kilmartin & Smith 1994; Beaz-Hidalgo & Figueras 2012). In Denmark, the infection was first described in freshwater during the 1950s by Rasmussen (1964). Today, problems with furunculosis in Denmark occur mainly in sea-reared rainbow trout (*Oncorhynchus mykiss*) production under stressful conditions and high temperatures (Dalsgaard & Madsen 2000; Pedersen *et al.* 2008). The presence of *A. salmonicida* in fish does not necessarily lead to the development of furunculosis, although some fish may be carriers that transfer *A. salmonicida* from freshwater to the sea (Jarp *et al.* 1993; Dalsgaard & Madsen 2000).

Detection of *A. salmonicida* has usually been performed by the use of bacterial cultivation (Dalsgaard & Madsen 2000; Austin & Austin 2007). However, the detection of the bacterium from supposed carrier fish with use of this method has not been successful so far (Dalsgaard & Madsen 2000). Although employing pre-enrichment steps or subjecting the fish to stress improves the detection of *A. salmonicida* by culturing (Cipriano *et al.* 1997), more sensitive methods are needed for the evaluation of the presence of *A. salmonicida* in carriers.

Real-time PCR has been used in several studies to detect *A. salmonicida* in fish tissue (Balcazar *et al.* 2007; Goodwin & Merry 2009; Keeling



*et al.* 2012; Gulla *et al.* 2015). In the study by Balcazar *et al.* (2007), an assay was developed with 100% specificity and a sensitivity of 16 CFU per reaction. The primers target an *A. salmonicida* DNA probe sequence from a 6.4-kb *A. salmonicida* plasmid named pAsal1 by Fehr *et al.* (2006), which has been the most frequently used target for species-specific *A. salmonicida* PCR and real-time PCR assays to date (Hiney *et al.* 1992; Morgan, Rhodes & Pickup 1993; O'Brien *et al.* 1994; Mooney *et al.* 1995; Byers, Gudkovs & Crane 2002; Balcazar *et al.* 2007; Goodwin & Merry 2009).

The objective was to investigate whether the real-time PCR assay developed by Balcazar *et al.* (2007), employed on multiple organs, would provide a more sensitive tool than bacterial culturing for determining *A. salmonicida* prevalence in rainbow trout from freshwater and seawater farms showing no signs of disease and from one seawater farm after a furunculosis outbreak. Samples included spleen, kidney, intestine, gill and brain tissues from 40 fish, sampled at five freshwater and seawater farms over the course of 2 years.

## Materials and methods

### Bacterial strains

To examine distribution of the pAsal1 plasmid and the target gene *aopP* in natural populations of *A. salmonicida* in Denmark, 20 Danish strains, isolated from outbreaks in fresh and seawater farms at different geographical locations in the period 1984–2014, were included in the study (data not shown). The strains were grown in Veal Infusion Broth (VIB) (Difco) for 48 h at 20°C. DNA was extracted with Qiagen QIAamp DNA mini kit (Qiagen) according to the manufacturer's protocol and immediately stored at –20°C until further use.

### Development of standard

An *A. salmonicida* DNA standard used for production of the standard curve was made from cloned PCR fragment of the target gene (*aopP*). PCR primers 5' TAGCTGGTTCCATAAGAA GC 3' and 5' TCCAAGAGGCAACTAAAGAAG 3' flanking the target sequence of the real-time PCR LUX primers developed by Balcazar *et al.* (2007), ensuring that both the real-time PCR

LUX target and primer sequences would be included in the amplified fragment, were generated from *A. salmonicida* pAsal1 plasmid sequence (GenBank accession no. AJ508382) and used to amplify a 340-bp fragment of the *aopP* gene from extracted DNA of *A. salmonicida* type strain ATCC 33658. The PCR product was then purified using the High Pure PCR Product Purification kit (Roche Applied Science) and cloned using a pGEM<sup>®</sup>-T and pGEM<sup>®</sup>-T Easy Vector Systems cloning kit (Promega) according to the manufacturer's instructions. Subsequently, the vector plasmid was purified with Qiagen Plasmid Midi Kit (Qiagen) and DNA concentration was determined using a Qubit 2.0 fluorometer and Quant-iT dsDNA BR kit (Invitrogen). Insertion of the 340-bp *aopP* gene fragment into the plasmid vector was verified by amplification with pUC/M13 universal primers (Promega) and sequencing of the amplicon using an ABI 3130 Genetic Analyser and Big Dye Terminator V 3.1 Cycle Sequencing Kit (Applied Biosystems) according to manufacturer's instruction.

### Real-time PCR

Real-time PCR was carried out in a total of 25 µL volume, containing 12.5 µL of 2X JumpStart Taq ReadyMix for quantitative PCR (Sigma-Aldrich), 1.5 µL of 25 mM MgCl<sub>2</sub>, 0.5 µL of 10 µM solution of each LUX PCR primer (Balcazar *et al.* 2007), 8 µL of nuclease-free water and 2 µL of DNA template. The thermal cycling conditions included an initial step at 95°C for 2 min, followed by 40 cycles consisting of 95°C for 15 s, 55°C for 30 s and 72°C for 30 s. Amplification and data analysis were performed using a Rotor-Gene Q system (Qiagen) and software version 2.0.2. All real-time PCR runs included template-free negative controls and positive *A. salmonicida* DNA standards consisting of selected dilutions. Samples were considered negative if no amplification signal was produced. Samples were considered positive if an amplification signal was produced in at least one of the triplicates and verified with melting point analysis.

### Sensitivity, reproducibility and dynamic range of real-time PCR

In order to create a standard curve, a solution of plasmid vector containing 14.3 ng µL<sup>-1</sup> of DNA

was used to prepare nine 10-fold serial dilutions in nuclease-free water down to  $1.43 \text{ fg } \mu\text{L}^{-1}$ , corresponding to a dilution range of approximately  $4 \times 10^9$  to four plasmid copies per  $\mu\text{L}$ . Two microlitres of each dilution was used as template in three independent real-time PCRs with three replicates of each dilution.

To determine limit of detection (LOD) and quantification limit (QL), 2  $\mu\text{L}$  of plasmid vector containing 80, 40, 20 and 10 copies of plasmid, respectively, was amplified. The obtained data were then used to graph the linear regression for the standard curve plot with the LOD point using RStudio (2015). The LOD was defined as the lowest DNA concentration (target gene copies per reaction) at which 95% of the positive samples were detected. The QL was defined as the lowest DNA concentration that remained within the linear region of the standard curve concentrations. Given an average of 20–50 pAsal1 plasmid copies per cell (Fehr *et al.* 2006; Attéré *et al.* 2015), quantitative real-time PCR data were transformed to express the results as genomic copies (genomic units, GU).

Reproducibility of the real-time PCR was assessed by calculating the interassay variance coefficient (CV%) in MS Excel (Microsoft). Calculation was based on the mean log of DNA copies per reaction, generated in three independent real-time PCR runs on the eight 10-fold serial dilution of plasmid vector DNA.

### Analysis of tissue inhibition in real-time PCR

An effect of possible co-purified inhibitors from the host tissues on sensitivity of the PCR assay was examined by amplification of gill, intestine, brain, kidney and spleen tissues seeded with *A. salmonicida* ATCC 33658.

Rainbow trout fry originating from eggs from Fousing Trout Farm that were disinfected and hatched at AquaBaltic and brought to our institute for rearing were used for collection of tissue. Six fish were killed in  $5 \text{ g L}^{-1}$  of Tricaine Methanesulfonate MS-222 (Sigma-Aldrich) in accordance with regulations set forward by the Danish Ministry of Justice and Animal Protection committees and under the Danish Animal Experiments Inspectorate permit number 2012-15-2924-00629. Spleen, gills, intestine, brain and kidney were collected under aseptic conditions and immediately stored at  $-20^\circ\text{C}$  until further use.

Seeding of tissues was performed as follows: *A. salmonicida* ATCC 33658 was grown in VIB at  $20^\circ\text{C}$  for 48 h. Serial 10-fold dilutions of bacterial cells containing  $5 \times 10^8$  to  $5 \times 10^2 \text{ CFU mL}^{-1}$  were prepared in 0.9% sterile saline solutions and counted on blood agar plates. In addition, two extra dilutions containing  $2.5 \times 10^2$  and  $1.25 \times 10^2 \text{ CFU mL}^{-1}$  were prepared, making up a total of nine different concentrations of bacterial cells. Subsequently, 1 mL of each bacterial cell concentration was mixed with 40 mg of gill, intestine, brain or kidney tissue or 15 mg of spleen tissue. All samples were then homogenized at 30 Hz for 20 s in a TissueLyser II (Qiagen), and DNA was prepared for amplification using the InstaGene Matrix (Bio-Rad) according to the manufacturer's instruction. Dilutions of bacterial cells without addition of any tissue and homogenized tissue with 0.9% saline solution were also included as positive and negative controls, respectively. The whole seeding experiment was repeated three times.

The real-time PCR data were analysed using Rotor-Gene Q Series Software (Qiagen) and used to graph the linear regression for organ tissue dilution series using RStudio (2015). The LOD and QL for each organ tissue were defined as for pure bacterial culture. For each seeding experiment, obtained log copies of seeded specimens and bacterial culture not mixed with any tissue were compared by ANOVA using RStudio (2015).

### Detection of *A. salmonicida* in fish

Rainbow trout ( $n = 40$ ) with and without signs of disease were collected for testing with bacterial culturing and real-time PCR from November 2013 through November 2015 at three freshwater and two seawater farms in Denmark. Twenty of the 40 fish were collected from one of the seawater farms (no. 5) 5 days after an antibiotic treatment of furunculosis and again at slaughtering 3 months after the first sampling. Between the two samplings, antibiotic treatment of fish was repeated. Sampling from farm no. 1, 2, 3 and 5 consisted of the same batch of fish that had been followed throughout the 2 years.

The spleen, gills, intestine, brain and kidney were removed from all fish and placed in individual Eppendorf tubes. Each sample was inoculated onto a blood agar plate (Columbia agar base [Oxoid] with 5% calf blood) by dipping an

inoculation loop into the tissue and then plating the content onto the blood agar, which was incubated at 20°C for 48–72 h (Dalsgaard & Madsen 2000). The remaining organ tissues were immediately stored at –20°C until DNA extraction for real-time PCR.

For DNA extraction, 20–40 mg specimens of the gill, intestine, brain and kidney and 10–15 mg from the spleen were homogenized with 1 mL of nuclease-free water at 30 Hz for 20 s in a TissueLyser II (Qiagen). Extraction was performed using the InstaGene Matrix (Bio-Rad) kit according to the manufacturer's protocol. Extracted DNA was immediately stored at –20°C until further use. All samples and positive and negative control samples were run in triplicates with real-time PCR. Rotor-Gene Q Series Software, version 2.0.2 (Qiagen), was used for analysis of the data.

## Results


### Specificity, sensitivity and reproducibility of real-time PCR

All examined Danish *A. salmonicida* strains ( $n = 20$ ) isolated from the outbreaks of furunculosis over a 30-year period, tested positive with real-time PCR (results not shown).


Amplification of standard DNA templates showed a broad linear dynamic range, spanning from  $8 \times 10^9$  to  $8 \times 10^1$  target copies, with a slope of  $-3.28$  ( $R^2 = 0.994$ ) and an efficiency of 102% (Fig. 1). The LOD was found to be 40 target copies per reaction with a quantification cycle ( $C_q$ ) of  $38.32 \pm 0.73$ , which is equivalent to 1–2 *A. salmonicida* GU per reaction. The QL was found to be 80 target copies per reaction ( $C_q$   $37.66 \pm 0.56$ ) equivalent to 2–4 *A. salmonicida* GU per reaction. The interassay CV for three independent real-time PCR runs was 0.94%.

### Analysis of tissue inhibition in real-time PCR

No evidence of inhibition was found for 40 mg of organs (15 mg for spleen) over the broad range of *A. salmonicida* ATCC 33658 concentrations. Dilution series for all five organs showed a linear trend from  $5 \times 10^8$  to  $2.5 \times 10^2$  CFU mL<sup>-1</sup> (Fig. 2), and assay results corresponded well to mean CFU mL<sup>-1</sup> values (results not shown). No significant difference was observed when comparing log copies per reaction obtained by real-time



**Figure 1** Standard curve created by plotting the mean quantification cycle ( $C_q$ ) values as a function of log target copies per reaction of the plasmid vector dilutions. Ten-fold serial dilutions ranging from  $14.3 \text{ ng } \mu\text{L}^{-1}$  to  $1.43 \text{ fg } \mu\text{L}^{-1}$  of the plasmid vector were analysed in three independent runs performed on the Rotor-Gene Q with three replicates per dilution concentration. The correlation coefficient, slope and intercept of the regression curve are shown. Error bars represent standard deviations (SD) of *Aeromonas salmonicida* standard log copies per reaction. The limit of detection (LOD) with a mean  $C_q$  of 38.32 and  $1.61 \pm 0.22$  SD log target copies per reaction is plotted, but lies outside of the linear region of the standard curve and is therefore not connected to the linear regression line.



**Figure 2** Regression curves for the *Aeromonas salmonicida* real-time PCR assay for five seeded organ tissues, created by plotting the mean quantification cycle ( $C_q$ ) values as a function of log target copies per reaction of the *A. salmonicida* ATCC concentrations added to the respective tissues. Serial dilutions ranging from  $5 \times 10^8$  to  $2.5 \times 10^2$  CFU mL<sup>-1</sup> of *A. salmonicida* ATCC 33658 were added to five different tissues: intestine, kidney, brain, spleen and the gills. DNA was extracted and analysed in three independent runs performed on the Rotor-Gene Q with three replicates per dilution concentration. The combined mean of the correlation coefficients, slopes and intercepts of the five regression curves is shown. Error bars represent standard deviations (SD) of *A. salmonicida* ATCC 33658 log target copies per reaction.

PCR from dilutions of *A. salmonicida* ATCC 33658 broth culture vs. dilutions of ATCC 33658 culture added to tissue using ANOVA ( $P > 0.05$ ). All negative control tissue samples were found negative by the real-time PCR.

### Detection of *A. salmonicida* in fish

In fish from the three freshwater farms (no. 1–3), no *A. salmonicida* were detected by the culture method (Table 1). However, *A. salmonicida* was detected in four of the 18 examined fish by real-time PCR (Table 1). In two cases, *A. salmonicida* was detected in both the brain and spleen and in the two other cases the bacterium was detected in either the brain or spleen. Plasmid copy number was in the range of 103–402  $\text{mg}^{-1}$  which equals from 2 up to 20 GU  $\text{mg}^{-1}$ .

At seawater farm no. 4, one of the two examined fish was positive for *A. salmonicida* in one organ (the brain) by culturing (Table 1). The real-time PCR detected *A. salmonicida* in both fish and several organs: the spleen and intestine of one fish and the spleen, intestine and brain of the other fish with a plasmid copy number in the range of 137–736  $\text{mg}^{-1}$ , which equals from 2 up to 37 GU  $\text{mg}^{-1}$ .

At fish farm no. 5 (Table 2) sampled 5 days after antibiotics treatment, *A. salmonicida* was detected in nine of 16 fish by culturing, but was detected in all 16 fish and in 73 of 80 organ samples by real-time PCR. Plasmid copy numbers were in the range of 21–9638  $\text{mg}^{-1}$ , which equals from 1 up to 482 GU  $\text{mg}^{-1}$ . At the second sampling, 3 months after the first sampling, *A. salmonicida* was detected in two of the four examined fish by culturing. All fish and organ samples were, however, found positive by real-time PCR with a plasmid copy number in the range of 77–2364  $\text{mg}^{-1}$ , which equals from 1 up to 118 GU  $\text{mg}^{-1}$ .

Overall, culturing detected *A. salmonicida* in 12 of the 40 examined fish, while real-time PCR detected *A. salmonicida* in 26 fish.

### Discussion

Specificity for the assay target gene *aopP* located on the plasmid pAsal1 was found to be 100% in the studies by Balcazar *et al.* (2007), who examined 16 *A. salmonicida* and 26 non-*A. salmonicida* bacterial strains, and by Goodwin & Merry (2009) that tested six *A. salmonicida* strains. In

the present study, all 20 Danish *A. salmonicida* strains that were examined with the assay were amplified, thereby suggesting a 100% prevalence of the *aopP* gene. Earlier studies have shown that plasmid pAsal1 is absent from some *A. salmonicida* strains (Morgan *et al.* 1993; Sørum, Kvello & Hastein 1993; O'Brien *et al.* 1994; Mooney *et al.* 1995; Byers *et al.* 2002), including 25% of 57 examined Danish strains (Nielsen, Olsen & Larsen 1993). Some authors explain the missing plasmid by stressful culturing conditions above 22–25°C, which was shown to activate an pAsal1 insertion sequence element ISAS11 that leads to the loss of the plasmid (Daher *et al.* 2011; Tanaka *et al.* 2012; Attéré *et al.* 2015). Others, such as Balcazar *et al.* (2007), have argued that the presence of pAsal1 could be associated with virulence of the bacterium and *A. salmonicida* lacking this plasmid might thus be less virulent. However, neither stressful culturing nor lack of virulence is necessarily associated with the absence of pAsal1 (Boyd *et al.* 2003; Fehr *et al.* 2006; Attéré *et al.* 2015), and one must keep this in mind when using *aopP* as the target gene.

Plasmid pAsal1 is present in 20–50 copies per bacterial cell (Fehr *et al.* 2006; Attéré *et al.* 2015), and the high abundance of the target facilitates detection by real-time PCR. We were able to reliably detect 1–2 *A. salmonicida* GU per reaction. Four positive samples had lower values than 1 GU per reaction (Table 2), but were still considered positive based on correct melting point of the obtained PCR product, despite being slightly below the 95% reliability. Although the plasmid copy variation per cell presents an obstacle for precise quantification of GU, it has little importance for practical application of the assay. The drawback of pAsal1 is that the plasmid is not universally present in all isolates of *A. salmonicida*. However, our results indicate that the pAsal1 plasmid is present in isolates causing new outbreaks in Danish fish farms and the frequent application of this target for *A. salmonicida* detection around the world enables comparison of results with the bulk of the published literature (Hiney *et al.* 1992; Morgan *et al.* 1993; O'Brien *et al.* 1994; Mooney *et al.* 1995; Byers *et al.* 2002; Balcazar *et al.* 2007; Goodwin & Merry 2009).

Compared to real-time PCR by Balcazar *et al.* (2007), sensitivity obtained in the present study was higher. Balcazar *et al.* (2007) achieved a LOD/QL of 16 *A. salmonicida* CFU per reaction

**Table 1** Detection and quantification of *Aeromonas salmonicida* from 12 fish sampled from three freshwater (no. 1–3) and one sea-water fish farm (no. 4) in Denmark during 2013–2015

Fish farm	Sampling date (yyyy.mm.dd)	Fish ID	Tissue	Bacteriology	Real-time PCR			
					Target copies mg <sup>-1</sup>	Log target copies mg <sup>-1</sup> ± SD	C <sub>q</sub> ± SD	
No. 1	2013.11.04	3	Gills	–	–	–	–	
			Spleen	–	–	–	–	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
			Brain	–	–	–	–	
		4	Gills	–	–	–	–	
			Spleen	–	–	–	–	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
			Brain	–	–	–	–	
		11	Gills	–	–	–	–	
			Spleen	–	–	–	–	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
			Brain	–	–	–	–	
		15	Gills	–	–	–	–	
			Spleen	–	–	–	–	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
			Brain	–	–	–	–	
		26	Gills	–	–	–	–	
			Spleen	–	–	–	–	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
Brain	–		–	–	–			
35	Gills	–	–	–	–			
	Spleen	–	–	–	–			
	Intestine	–	–	–	–			
	Kidney	–	–	–	–			
	Brain	–	–	–	–			
No. 2	2014.02.18	48	Gills	–	–	–	–	
			Spleen	–	–	–	–	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
			Brain	–	–	–	–	
		76	Gills	–	–	–	–	
			Spleen	–	–	–	–	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
			Brain	–	–	–	–	
		2014.05.14	83	Gills	–	–	–	–
				Spleen	–	–	–	–
	Intestine			–	–	–	–	
	Kidney			–	–	–	–	
	Brain			–	–	–	–	
	95		Gills	–	–	–	–	
			Spleen	–	402	2.60 ± 0.08	36.87 ± 0.26	
			Intestine	–	–	–	–	
			Kidney	–	–	–	–	
			Brain	–	107	2.03 ± 0.22 <sup>b</sup>	38.06 ± 0.72 <sup>b</sup>	
	105		Gills	–	–	–	–	
			Spleen	–	198	2.30 ± 0.13 <sup>ab</sup>	38.06 ± 0.41 <sup>ab</sup>	
		Intestine	–	–	–	–		
		Kidney	–	–	–	–		
Brain		–	159	2.20 ± 0.17	37.44 ± 0.57 <sup>b</sup>			
113	Gills	–	–	–	–			
	Spleen	–	–	–	–			
	Intestine	–	–	–	–			
	Kidney	–	–	–	–			

Table 1 Continued

Fish farm	Sampling date (yyyy.mm.dd)	Fish ID	Tissue	Bacteriology	Real-time PCR				
					Target copies mg <sup>-1</sup>	Log target copies mg <sup>-1</sup> ± SD	C <sub>q</sub> ± SD		
No. 3	2014.11.03	122	Brain	–	–	–	–		
			Gills	–	–	–	–		
			Spleen	–	–	–	–		
			Intestine	–	–	–	–		
			Kidney	–	–	–	–		
	148	2015.03.24	170	Brain	–	103	2.02 ± 0.27 <sup>b</sup>	37.95 ± 0.89 <sup>b</sup>	
				Gills	–	–	–	–	
				Spleen	–	–	–	–	
				Intestine	–	–	–	–	
				Kidney	–	–	–	–	
	184	2015.03.24	170	Brain	–	–	–	–	
				Gills	–	–	–	–	
				Spleen	–	–	–	–	
				Intestine	–	–	–	–	
				Kidney	–	–	–	–	
	185	2015.03.24	184	Brain	–	–	–	–	
				Gills	–	–	–	–	
				Spleen	–	–	–	–	
				Intestine	–	–	–	–	
				Kidney	–	–	–	–	
198	2015.03.24	185	Brain	–	–	–	–		
			Gills	–	–	–	–		
			Spleen	–	–	–	–		
			Intestine	–	–	–	–		
			Kidney	–	–	–	–		
No. 4	2015.11.04	1	Brain	–	–	–	–		
			Gills	–	–	–	–		
			Spleen	–	443	2.65 ± 0.26	37.11 ± 0.86		
			Intestine	–	137	2.14 ± 0.21	37.54 ± 0.69		
			Kidney	–	–	–	–		
		3	2015.11.04	198	Brain	–	–	–	–
					Gills	–	–	–	–
					Spleen	–	763	2.88 ± 0.07	36.34 ± 0.22
					Intestine	–	258	2.41 ± 0.22	37.01 ± 0.71
					Kidney	–	–	–	–
Brain	+	231	2.33 ± 0.03	36.61 ± 0.10					

–, Examined with negative result.

<sup>a</sup>Only two of three sample replicates produced an amplification signal.

<sup>b</sup>Quantification cycle (C<sub>q</sub>) value below quantitation limit (QL). Reported quantification value of copies mg<sup>-1</sup> could be imprecise.

in pure culture as well as inoculated tissue. In comparison, the QL and LOD of the present assay were 80 and 40 target genes copies per reaction, respectively, equivalent to, respectively, 2–4 and 1–2 *A. salmonicida* GU per reaction. An explanation for the difference in sensitivity between the present study and Balcazar *et al.* (2007), even though the same assay is applied in both studies, could be that another real-time PCR instrument and kit was used in this study and the standard curve was generated from a cloned PCR product, while Balcazar *et al.* (2007) used

extracted DNA from pure cultures of *A. salmonicida* NCIMB 1102.

The most frequently examined organ for *A. salmonicida* in previous real-time PCR studies has been the kidney (Balcazar *et al.* 2007; Goodwin & Merry 2009; Keeling *et al.* 2012; Gulla *et al.* 2015). In the present study, four additional organs were included: spleen, intestine, gills and the brain. Interestingly, in all four real-time PCR-positive freshwater farm fish, the brain and/or the spleen were found positive but not the kidney (Table 1). In comparison, no *A. salmonicida* was

**Table 2** Detection and quantification of *Aeromonas salmonicida* from 20 fish sampled 11th of August and 11th of November 2015 at seawater farm no. 5. These samples were taken 5 days and 3 months after antibiotic treatment of furunculosis

Fish farm	Sampling date (yyyy.mm.dd)	Fish ID	Tissue	Bacteriology	Real-time PCR		
					Target copies mg <sup>-1</sup>	Log target copies mg <sup>-1</sup> ± SD	C <sub>q</sub> ± SD
No. 5	2015.08.11	9	Gills	–	171	2.23 ± 0.55	36.32 ± 1.80
			Spleen	–	220	2.34 ± 0.13	37.44 ± 0.43
			Intestine	–	510	2.71 ± 0.20	34.61 ± 0.65
			Kidney	–	473	2.67 ± 0.18	35.17 ± 0.58
		13	Brain	–	417	2.62 ± 0.25	34.89 ± 0.82
			Gills	+	3732	3.57 ± 0.06	37.73 ± 0.21
			Spleen	–	2276	3.36 ± 0.02	33.57 ± 0.06
			Intestine	–	–	–	–
		15	Kidney	–	1577	3.20 ± 0.27	32.95 ± 0.87
			Brain	–	669	2.83 ± 0.48	33.54 ± 1.58
			Gills	–	92	1.96 ± 0.11	37.15 ± 1.00
			Spleen	–	2526	3.40 ± 0.01	33.77 ± 0.77
		17	Intestine	–	562	2.75 ± 0.16	34.34 ± 0.79
			Kidney	–	283	2.45 ± 0.05	35.26 ± 0.16
			Brain	–	54	1.73 <sup>b</sup>	37.28 <sup>b</sup>
			Gills	+	968	2.68 ± 0.20	35.70 ± 0.66
		22	Spleen	–	2050	3.31 ± 0.12	33.29 ± 0.38
			Intestine	–	14	1.15 <sup>bd</sup>	39.23 <sup>bd</sup>
			Kidney	–	825	2.92 ± 0.09	33.74 ± 0.30
			Brain	–	702	2.85 <sup>b</sup>	33.68 <sup>b</sup>
		23	Gills	–	38	1.51 ± 0.35 <sup>ac</sup>	37.99 ± 1.15 <sup>ac</sup>
			Spleen	–	1071	3.02 ± 0.13 <sup>a</sup>	34.58 ± 0.42 <sup>a</sup>
			Intestine	–	198	2.3 <sup>b</sup>	35.34 <sup>b</sup>
			Kidney	–	–	–	–
		25	Brain	–	277	2.44 ± 0.19	34.99 ± 0.62
			Gills	+	146	2.17 ± 0.44	36.44 ± 1.45
			Spleen	–	766	2.88 ± 0.21	35.42 ± 0.70
			Intestine	–	473	2.67 ± 0.23	34.97 ± 0.76
		26	Kidney	–	7286	3.86 ± 0.20	32.11 ± 0.64
			Brain	–	432	2.64 ± 0.32	34.83 ± 1.06
			Gills	–	–	–	–
			Spleen	–	722	2.86 ± 0.12	36.03 ± 0.39
		28	Intestine	–	34	1.53 ± 0.21 <sup>ad</sup>	39.03 ± 0.67 <sup>ad</sup>
			Kidney	–	227	2.36 ± 0.34	37.02 ± 1.11
			Brain	–	49	1.69 ± 0.05 <sup>ad</sup>	38.63 ± 0.17 <sup>ad</sup>
			Gills	–	44	1.63 <sup>bd</sup>	38.69 <sup>bd</sup>
		29	Spleen	–	203	2.31 ± 0.17 <sup>c</sup>	38 ± 0.57 <sup>c</sup>
			Intestine	–	21	1.32 <sup>bd</sup>	39.99 <sup>bd</sup>
			Kidney	–	122	2.09 ± 0.26 <sup>c</sup>	37.89 ± 0.85 <sup>c</sup>
			Brain	–	–	–	–
		30	Gills	–	–	–	–
			Spleen	–	933	2.97 ± 0.18	35.51 ± 0.58
Intestine	–		–	–	–		
Kidney	–		1189	3.08 ± 0.13	34.46 ± 0.41		
31	Brain	–	1344	3.13 ± 0.12	33.63 ± 0.40		
	Gills	+	86	1.93 ± 0.26	36.68 ± 0.84		
	Spleen	+	882	2.95 ± 0.05	35.06 ± 0.15		
	Intestine	–	24	1.37 ± 0.23 <sup>ad</sup>	39.07 ± 0.76 <sup>ad</sup>		
32	Kidney	+	943	2.97 ± 0.46	34.12 ± 1.51		
	Brain	–	591	2.77 ± 0.21	34.35 ± 0.68		
	Gills	–	493	2.69 ± 0.26	34.80 ± 0.84		
	Spleen	–	5106	3.71 ± 0.35	32.54 ± 1.13		
33	Intestine	–	134	2.13 ± 0.43	36.58 ± 1.42		
	Kidney	–	1436	3.00 ± 0.56 <sup>a</sup>	34.21 ± 1.82 <sup>a</sup>		
	Brain	+	1907	3.28 ± 0.15	32.71 ± 0.48		
	Gills	+	173	2.24 ± 0.06	36.59 ± 0.21		
34	Spleen	–	9638	3.98 ± 0.07	31.87 ± 0.23		
	Intestine	–	345	2.54 ± 0.20	35.33 ± 0.66		
	Kidney	–	957	2.98 ± 0.23	34.73 ± 0.76		

Table 2 Continued

Fish farm	Sampling date (yyyy.mm.dd)	Fish ID	Tissue	Bacteriology	Real-time PCR		
					Target copies mg <sup>-1</sup>	Log target copies mg <sup>-1</sup> ± SD	C <sub>q</sub> ± SD
		32	Brain	–	755	2.88 ± 0.17	34.25 ± 0.57
			Gills	+	241	2.38 ± 0.15	36.14 ± 0.49
			Spleen	+	3611	3.56 ± 0.18	33.59 ± 0.60
			Intestine	–	1185	3.07 ± 0.15	33.66 ± 0.50
			Kidney	+	821	2.91 ± 0.17	34.62 ± 0.56
			Brain	+	3463	3.54 ± 0.11	32.23 ± 0.36
		37	Gills	–	–	–	–
			Spleen	–	2565	3.41 ± 0.21	34.30 ± 0.70
			Intestine	–	288	2.46 ± 0.20	36.29 ± 0.67
			Kidney	–	110	2.04 ± 0.16 <sup>c</sup>	37.58 ± 0.53 <sup>c</sup>
			Brain	–	300	2.48 ± 0.32	35.89 ± 1.04
		39	Gills	+	65	1.81 ± 0.62 <sup>c</sup>	37.86 ± 2.04 <sup>c</sup>
			Spleen	–	7930	3.90 ± 0.07	32.63 ± 0.24
			Intestine	–	631	2.80 ± 0.28	34.76 ± 0.92
			Kidney	–	448	2.65 ± 0.13	35.45 ± 0.41
			Brain	–	299	2.48 ± 0.30	35.69 ± 0.98
		42	Gills	+	288	2.46 ± 0.21	35.81 ± 0.70
			Spleen	+	1336	3.13 ± 0.33	34.82 ± 1.09
			Intestine	–	168	2.23 ± 0.14	36.52 ± 0.47
			Kidney	+	1937	3.29 ± 0.11	33.74 ± 0.35
			Brain	–	338	2.53 ± 0.07	35.51 ± 0.23
	2015.11.11	3	Gills	–	185	2.27 ± 0.24	37.16 ± 0.85
			Spleen	–	1019	3.01 ± 0.16	36.24 ± 0.56
			Intestine	–	299	2.48 ± 0.11	36.39 ± 0.37
			Kidney	–	143	2.16 ± 0.46	37.60 ± 1.61
			Brain	–	216	2.33 ± 0.20	37.13 ± 0.69
		8	Gills	+	222	2.35 ± 0.1	36.94 ± 0.34
			Spleen	–	441	2.64 <sup>b</sup>	36.69 <sup>b</sup>
			Intestine	–	456	2.66 ± 0.04	35.71 ± 0.13
			Kidney	+	1322	3.12 ± 0.15	34.87 ± 0.52
			Brain	–	341	2.53 ± 0.40	36.06 ± 1.41
		23	Gills	–	183	2.26 ± 0.53	37.49 ± 1.85
			Spleen	–	2111	3.32 ± 0.20	34.48 ± 0.69
			Intestine	–	77	1.89 ± 0.57 <sup>ad</sup>	38.45 ± 1.98 <sup>ad</sup>
			Kidney	+	162	2.21 ± 0.16 <sup>c</sup>	38.06 ± 0.56 <sup>c</sup>
			Brain	–	190	2.28 ± 0.34	36.95 ± 1.18
		27	Gills	–	126	2.10 ± 0.21 <sup>ac</sup>	37.62 ± 0.75 <sup>ac</sup>
			Spleen	–	2364	3.37 ± 0.08	34.22 ± 0.28
			Intestine	–	670	2.83 ± 0.11	35.90 ± 0.38
			Kidney	–	481	2.68 ± 0.15	36.14 ± 0.51
			Brain	–	785	2.90 ± 0.19	34.80 ± 0.66

–, Examined with negative result.

<sup>a</sup>Only two of three sample replicates produced an amplification signal.

<sup>b</sup>Only one of three sample replicates produced an amplification signal.

<sup>c</sup>Quantification cycle (C<sub>q</sub>) value below quantitation limit (QL). Reported quantification value of copies mg<sup>-1</sup> could be imprecise.

<sup>d</sup>Quantification cycle (C<sub>q</sub>) value below limit of detection (LOD). Reported quantification value of copies mg<sup>-1</sup> has to be taken tentatively.

isolated by bacterial culturing in any of the 18 fish from freshwater farms as well as in none of the additional 182 fish sampled from the same three freshwater farms (data not shown). None of the freshwater farm fish showed any signs of furunculosis, indicating that the four fish found positive by real-time PCR could be carriers of viable but non-culturable cells (VBNC) (Morgan *et al.* 1993; Ferguson *et al.* 1995; Nascuti 2010). Due to the high sensitivity of the present real-time PCR, one

could speculate whether the four fish could be false positives by contamination of the surrounding water. However, in that case one would presume the gills would be the primarily infected organs, while in the present study; the gills were negative in all fish showing no signs of disease. In addition, none of the previous studies that have employed the pAsal1 target have detected other bacterial DNA than that of *A. salmonicida* (Hiney *et al.* 1992; Morgan *et al.* 1993; O'Brien *et al.*



1994; Mooney *et al.* 1995; Byers *et al.* 2002; Balcazar *et al.* 2007; Goodwin & Merry 2009).

It was reported by Gustafson, Thomas & Trust (1992) that in farmed brown trout (*Salmo trutta*) up to 80% of the fish are thought to be carriers of *A. salmonicida* and VBNC *A. salmonicida* have also been revived (Austin, Austin & Colwell 1984). The presence of VBNC *A. salmonicida* would also explain why the bacterium has not been found by culturing in the Danish rainbow trout freshwater farms, although furunculosis outbreaks occur during elevated temperatures after fish are transferred to the sea (Dalsgaard & Madsen 2000; Pedersen *et al.* 2008). Our results support this hypothesis, as the fish sampled at three freshwater farms represented the same batch of fish. Initially, fish were hatched at freshwater farm no. 1, then transferred to freshwater farm no. 2 where *A. salmonicida* became detectable by real-time PCR, then further transferred to freshwater farm no. 3 still being *A. salmonicida* positive by real-time PCR, and finally transferred to seawater farm no. 5 at 2 years age, where furunculosis outbreak occurred later that year. Further studies are underway to shed a light on possible transmission of the disease.

In fish reared in sea water, *A. salmonicida* was detected by culturing in 12 of 22 fish and 20 of 110 organs, while the bacterium was detected in all 22 fish and 98 organs by real-time PCR (Table 1 and 2). This substantial difference in detection between the culturing and real-time PCR approach might be explained, at least in part, by the fact that 20 fish accounting for 93 of the positive organs were obtained from farm no. 5. At this farm, there had been a furunculosis outbreak, but unfortunately, it was not possible to do the sampling until 5 days after the fish had been treated with antibiotics. The treatment most likely either reduced or killed *A. salmonicida* cells in the fish, thereby reducing or eliminating the probability of detecting the bacterium by culturing while maintaining the chance for real-time PCR to detect DNA from either dead or VBNC bacteria still present within the fish. The possible presence of both viable, dead and VBNC bacteria in variable frequencies in these fish would also explain the lack of direct correlation between culturing and GU counts as quantified by real-time PCR (Table 2).

The same seawater farm (no. 5) was sampled again approximately 3 months after antibiotic treatment, where *A. salmonicida* was detected in two of

the four sampled fish by culturing, while real-time PCR detected *A. salmonicida* in all four fish and all 20 organs (Table 2). The question remains here whether the real-time PCR mostly detected DNA from dead bacteria or whether there were low amounts of viable *A. salmonicida* present in all fish and most of the organs, which were only detected in two fish (three organs) by culturing.

In the other seawater farm (no. 4), one of the two examined fish was found positive by culturing, while both fish were found positive by real-time PCR (Table 1). The organs where *A. salmonicida* was detected by real-time PCR were the spleen and intestine in both fish, along with the brain in one fish. Detection of *A. salmonicida* in the spleen and brain correlates with the findings of the bacterium by real-time PCR in fish from freshwater farms. However, it is interesting that the intestine was also found positive in both fish; in accordance, Hiney *et al.* (1994) stated that the intestine may be the primary location of *A. salmonicida* in salmon. Results from this seawater farm and the freshwater farms also bring attention to the importance of sampling from more than one or two organs in order to be sure of avoiding false negatives regarding the detection of *A. salmonicida* in fish.

In summary, our present findings indicate that there may be carrier fish harbouring VBNC *A. salmonicida* in Danish freshwater and seawater rainbow trout farms and that the spleen, brain and intestine could play an important role in *A. salmonicida* infection and persistence of VBNC. Further studies are needed for obtaining more knowledge about VBNC and carrier fish. The real-time PCR assay showed higher sensitivity for the detection of *A. salmonicida* than the culture method and exhibited a high reproducibility and efficiency. The real-time PCR assay presents a proficient tool for the detection of *A. salmonicida* in fish. One must keep in mind though that not all *A. salmonicida* seem to possess the target plasmid pAsal1. In order to be certain of avoiding false negatives, another sensitive detection method with a different target would need to be employed.

## Acknowledgements

This work was supported by The Danish Council for Strategic Research under PROFISH project (Grant no. DSF: 11-116252) and the National Veterinary Institute (DTU). The authors would

like to thank all project partners and Lisbeth Schade Hansen and Niccolò Vendramin at the National Veterinary Institute (DTU), Lene Nørskov at Aarhus University, and Niels Henrik Henriksen at Danish Aquaculture for their assistance in field sampling and technical support.

## References

- Attéré S.A., Vincent A.T., Trudel M., Chanut R. & Charette S.J. (2015) Diversity and homogeneity among small plasmids of *Aeromonas salmonicida* subsp. *salmonicida* linked with geographical origin. *Frontiers in Microbiology* **6**, 1274.
- Austin B. & Austin D.A. (2007) *Bacterial Fish Pathogens: Diseases of Farmed and Wild Fish*. 4th edn. Springer, Chichester, UK.
- Austin D.A., Austin B. & Colwell R.R. (1984) Survival of *Aeromonas salmonicida* in river water. *FEMS Microbiology Letters* **21**, 143–146.
- Balcazar J.L., Vendrell D., de Blas I., Ruiz-Zarzuola I., Girones O. & Luis Muzquiz J. (2007) Quantitative detection of *Aeromonas salmonicida* in fish tissue by real-time PCR using self-quenched, fluorogenic primers. *Journal of Medical Microbiology* **56**, 323–328.
- Beaz-Hidalgo R. & Figueras M.J. (2012) Molecular detection and characterization of furunculosis and other *Aeromonas* fish infections. In: *Health and Environment in Aquaculture*. (ed. by E.D. Carvalho, G.S. David & R.J. Silva), pp. 97–132. InTech, Rijeka, Croatia.
- Boyd J., Williams J., Curtis B., Kozera C., Singh R. & Reith M. (2003) Three small, cryptic plasmids from *Aeromonas salmonicida* subsp. *salmonicida* A449. *Plasmid* **50**, 131–144.
- Byers H.K., Gudkovs N. & Crane M.S. (2002) PCR-based assays for the fish pathogen *Aeromonas salmonicida*. I. Evaluation of three PCR primer sets for detection and identification. *Diseases of Aquatic Organisms* **49**, 129–138.
- Cipriano R.C., Ford L.A., Smith D.R., Schachte J.H. & Petrie C.J. (1997) Differences in detection of *Aeromonas salmonicida* in covertly infected salmonid fishes by the stress-inducible furunculosis test and culture-based assays. *Journal of Aquatic Animal Health* **9**, 108–113.
- Daher R.K., Filion G., Tan S.G.E., Dallaire-Dufresne S., Paquet V.E. & Charette S.J. (2011) Alteration of virulence factors and rearrangement of pAsa5 plasmid caused by the growth of *Aeromonas salmonicida* in stressful conditions. *Veterinary Microbiology* **152**, 353–360.
- Dalsgaard I. & Madsen L. (2000) Bacterial pathogens in rainbow trout, *Oncorhynchus mykiss* (Walbaum), reared at Danish freshwater farms. *Journal of Fish Diseases* **23**, 199–209.
- Fehr D., Casanova C., Liverman A., Blazkova H., Orth K., Dobbelaere D., Frey J. & Burr S.E. (2006) AopP, a type III effector protein of *Aeromonas salmonicida*, inhibits the NF- $\kappa$ B signalling pathway. *Microbiology (Reading)* **152**, 2809–2818.
- Ferguson Y., Glover L.A., McGillivray D.M. & Prosser J.I. (1995) Survival and activity of lux-marked *Aeromonas salmonicida* in seawater. *Applied and Environmental Microbiology* **61**, 3494–3498.
- Goodwin A.E. & Merry G.E. (2009) Are all koi ulcer cases associated with infection by atypical *Aeromonas salmonicida*? Polymerase chain reaction assays of koi carp skin swabs submitted by hobbyists. *Journal of Aquatic Animal Health* **21**, 98–103.
- Gulla S., Duodu S., Nilsen A., Fossen I. & Colquhoun D.J. (2015) *Aeromonas salmonicida* infection levels in pre- and post-stocked cleaner fish assessed by culture and an amended qPCR assay. *Journal of Fish Diseases* doi: 10.1111/jfd.12420.
- Gustafson C.E., Thomas C.J. & Trust T.J. (1992) Detection of *Aeromonas salmonicida* from fish by using polymerase chain reaction amplification of the virulence surface array protein gene. *Applied and Environmental Microbiology* **58**, 3816–3825.
- Hiney M., Dawson M.T., Heery D.M., Smith P.R., Gannon F. & Powell R. (1992) DNA probe for *Aeromonas salmonicida*. *Applied and Environmental Microbiology* **58**, 1039–1042.
- Hiney M.P., Kilmartin J.J. & Smith P.R. (1994) Detection of *Aeromonas salmonicida* in atlantic salmon with asymptomatic furunculosis infections. *Diseases of Aquatic Organisms* **19**, 161–167.
- Jarp J., Tangen K., Willumsen F.V., Djupvik H.O. & Tveit A.M. (1993) Risk factors for infection with *Aeromonas salmonicida* subsp. *salmonicida* in Norwegian freshwater hatcheries. *Diseases of Aquatic Organisms* **17**, 81–86.
- Keeling S.E., Brosnahan C.L., Johnston C., Wallis R., Gudkovs N. & McDonald W.L. (2012) Development and validation of a real-time PCR assay for the detection of *Aeromonas salmonicida*. *Journal of Fish Diseases* **36**, 495–503.
- Mooney J., Powell E., Clabby C. & Powell R. (1995) Detection of *Aeromonas salmonicida* in wild Atlantic salmon using a specific DNA probe test. *Diseases of Aquatic Organisms* **21**, 131–135.
- Morgan J., Rhodes G. & Pickup R.W. (1993) Survival of nonculturable *Aeromonas salmonicida* in lake water. *Applied and Environmental Microbiology* **59**, 874–880.
- Nascutiu A.M. (2010) Bacterii viabile necultivabile. *Bacteriologia, Virusologia, Parazitologia, Epidemiologia (bucharest, Romania: 1990)* **55**, 11–18.
- Nielsen B., Olsen J.E. & Larsen J.L. (1993) Plasmid profiling as an epidemiological marker within *Aeromonas salmonicida*. *Diseases of Aquatic Organisms* **15**, 129–135.
- O'Brien D., Mooney J., Ryan D., Powell E., Hiney M., Smith P.R. & Powell R. (1994) Detection of *Aeromonas salmonicida*, causal agent of furunculosis in salmonid fish, from the tank effluent of hatchery-reared atlantic salmon smolts. *Applied and Environmental Microbiology* **60**, 3874–3877.
- Pedersen K., Skall H.F., Lassen-Nielsen A.M., Nielsen T.F., Henriksen N.H. & Olesen N.J. (2008) Surveillance of health status on eight marine rainbow trout, *Oncorhynchus mykiss* (Walbaum), farms in Denmark in 2006. *Journal of Fish Diseases* **31**, 659–667.
- Rasmussen C.J. (1964) Furunkulose hos damørreder (Furunculosis in farmed trout) (in Danish). *Scandinavian Journal of Veterinary Science* **16**, 463–472.
- RStudio (2015) R: A language and environment for statistical computing.

Sørum H., Kvello J.H. & Hastein T. (1993) Occurrence and stability of plasmids in *Aeromonas salmonicida* ss *salmonicida* isolated from salmonids with furunculosis. *Diseases of Aquatic Organisms* **16**, 199–206.

Tanaka K.H., Dallaire-Dufresne S., Daher R.K., Frenette M. & Charette S.J. (2012) An insertion sequence-dependent

plasmid rearrangement in *Aeromonas salmonicida* causes the loss of the type three secretion system. *PLoS One* **7**, e33725.

*Received: 1 February 2016*

*Revision received: 21 April 2016*

*Accepted: 21 April 2016*





## **Manuscript III**

**Epidemiology and genetics of *Aeromonas salmonicida* using  
whole genome sequencing**



# **Epidemiology and genetics of *Aeromonas salmonicida* using whole genome sequencing**

1 **Simona Bartkova<sup>1\*</sup>, Pimlapas Leekitcharoenphon<sup>2</sup>, Frank M. Aarestrup<sup>2</sup>, Inger Dalsgaard<sup>1</sup>**

2

3 <sup>1</sup> National Veterinary Institute, Section for Bacteriology and Pathology, Technical University of  
4 Denmark, Frederiksberg C, Denmark

5

6 <sup>2</sup> National Food Institute, Division for Epidemiology and Microbial Genomics, Technical  
7 University of Denmark, Kgs. Lyngby, Denmark

8

9 **\* Correspondence:** Simona Bartkova, National Veterinary Institute, Section for Bacteriology and  
10 Pathology, Technical University of Denmark, Bülowsvej 27, 1870 Frederiksberg C, Denmark

11

12 sibar@vet.dtu.dk

13

14 **Running title: Epidemiology of *Aeromonas salmonicida***



15 **Abstract**

16

17 Sea-reared rainbow trout production in Denmark currently struggles with furunculosis, a septicemic  
18 infection caused by the bacterium *Aeromonas salmonicida* subsp. *salmonicida*. Developing an  
19 effective control strategy is vital for future production, but this requires having knowledge of the  
20 epidemiology, as well as the genetic and virulent variability of the Danish *A. salmonicida* isolates.  
21 In order to obtain this, the genomes of 101 *A. salmonicida*, including 99 Danish isolates, one  
22 Scottish strain and the type strain NCIMB 1102, were sequenced using the Illumina HiSeq  
23 platform. Isolates were *de novo* assembled, examined for presence of plasmids, virulence and iron  
24 acquisition proteins and antibiotic resistance genes. Single Nucleotide Polymorphisms were aligned  
25 and subjected to Bayesian temporal phylogenetic and maximum likelihood tree reconstruction using  
26 the published genome of *A. salmonicida* A449 as reference. Bayesian temporal phylogenetic  
27 reconstruction suggests that four major introductions of *A. salmonicida* into Denmark have  
28 occurred. The introductions correlate with the freshwater and subsequent seawater expansion of  
29 rainbow trout production. Initial transmission of the bacterium could have been from seawater to  
30 freshwater or vice versa, both scenarios are open and most minor clades include a mixture of strains  
31 from different fresh- and seawater farms. Genetic variation of *A. salmonicida* is mostly associated  
32 with their plasmids and plasmid encoded virulence factors. Nine *A. salmonicida* harbored  
33 worldwide known antibiotic resistance genes against several antibiotics. These findings provide  
34 novel information regarding the Danish *A. salmonicida* population and demonstrate that whole  
35 genome sequencing is a highly useful tool for studying homogenous bacteria such as *A.*  
36 *salmonicida*.

37

38 **Keywords:** *Aeromonas salmonicida*, furunculosis, rainbow trout, whole genome sequencing,  
39 **SNP analysis, BEAST, virulence factors**

## 40 1. Introduction

41

42 *Aeromonas salmonicida* subsp. *salmonicida*, the causative agent of a septicemic infection  
43 furunculosis, was first described by Emmerich and Weibel (1894) at a German freshwater brown  
44 trout hatchery. Although the first rainbow trout (*Oncorhynchus mykiss*) hatchery in Denmark was  
45 already established in 1858 (Christensen, 1980), signs of furunculosis among fish were first  
46 described in the 1950s at freshwater rainbow trout farms (Rasmussen 1964). At this point the  
47 Danish freshwater rainbow trout production had begun its massive expansion. In the late 1970s,  
48 production was extended to seawater and dry pellet feed was introduced instead of the common wet  
49 feed consisting of herring, whiting, sand-eels and other marine fish not used for human  
50 consumption (Christensen, 1980). Both actions increased the growth of the Danish rainbow trout  
51 production even further.

52

53 Currently, it is in the expanded Danish sea-reared rainbow trout production that *A. salmonicida* is  
54 responsible for great financial losses. Despite fish being vaccinated before transfer from freshwater  
55 to seawater farms, furunculosis has occurred repeatedly during situations with elevated water  
56 temperatures (Larsen and Møllgaard, 1981; Dalsgaard and Madsen, 2000; Pedersen *et al.*, 2008).  
57 This situation, along with previous research, has led to the belief that *A. salmonicida* could be  
58 spread from freshwater to the sea by carrier fish that harbor the bacterium without showing any  
59 signs of disease (Larsen and Møllgaard, 1981; Dalsgaard and Madsen, 2000). Verifying this  
60 would be critical for developing an effective prevention strategy against furunculosis.

61

62 Several methods exist for molecular typing of bacterial isolates. The ‘gold standard’ for typing has  
63 long been Pulsed-field gel electrophoresis (PFGE), however, this method is time-consuming and  
64 lacks resolution power to distinguish the highly homogenous *A. salmonicida* on the subspecies level  
65 (Cunningham and Colquhoun, 2002; Beaz-Hidalgo *et al.*, 2008). Although other methods such as  
66 restriction fragment length polymorphism (RFLP) and multilocus sequence typing (MLST) analysis  
67 can match the resolution of PFGE and can often provide results faster (Cunningham and  
68 Colquhoun, 2002; van Belkum *et al.*, 2007; Beaz-Hidalgo *et al.*, 2008) most of the methods are  
69 laborious and expensive (van Belkum *et al.* 2007). Recently a MLST-V based on eight fragments of  
70 housekeeping genes and four fragments of virulence associated genes was applied on 25 Danish  
71 isolates, though it did not have enough discriminatory power for distinguishing *A. salmonicida* at  
72 the subspecies level (authors’ unpublished results).

73

74 Whole-genome sequencing (WGS) using next generation sequencing technology has over the past  
75 few years drastically decreased in cost and increased in speed, enabling its usage for studying  
76 everything from specific genes and virulence factors to epidemiology and long term evolution of  
77 various bacteria on a regular basis (Parkhill and Wren, 2011). Moreover, WGS provides the best  
78 overview of a studied population, since it avoids bias present in other molecular methods such as  
79 MLST, which only investigate a small part of the genome (Foxman *et al.* 2005; Parkhill and Wren,  
80 2011).

81

82 In order to create an overview of the variation in genetics and virulence factors, as well as the  
83 epidemiology and evolution of Danish *A. salmonicida* isolates, a representative collection of 99  
84 Danish *A. salmonicida* isolates varying in isolation years 1980 - 2014 and geographical regions, a  
85 Scottish strain and the type strain NCIMB 1102 were sequenced using the Illumina HiSeq platform.  
86 Sequences of all isolates were *de novo* assembled and analyzed using the published genome of *A.*  
87 *salmonicida* A449 (Reith *et al.*, 2008) as reference.

88

## 89 **2. Methods**

90

### 91 **2.1 Bacterial isolates**

92

93 Ninety-nine Danish *A. salmonicida* isolated from furunculosis outbreaks between 1980 and 2014  
94 were selected. The collection consisted of 42 *A. salmonicida* isolated from various freshwater  
95 farms, of which 40 were from rainbow trout and two from brown trout (*Salmo trutta*). Fifty-seven  
96 of the *A. salmonicida* were isolated from rainbow trout at various seawater farms, of which 14  
97 isolates (isolated between 1981 - 2014) belonged to one large seawater farm and nine isolates  
98 (isolated between 1989 - 2010) to another large seawater farm named Sj4 and Sj3 respectively in  
99 this study. The Scottish *A. salmonicida* strain MT004 from Atlantic salmon (*Salmo salar* L.) is  
100 according to literature isolated around 1980. The *A. salmonicida* type strain NCIMB 1102 from  
101 England was isolated from an Atlantic salmon in year 1962. Extracted genomic DNA from all  
102 *A. salmonicida* was used for sequencing.

103

### 104 **2.2 Sample preparation**

105

106 All *A. salmonicida* were grown in Veal Infusion Broth (VIB) (Difco) at 20°C for 48 hours and then  
107 inoculated on blood agar plates (Columbia agar base (Oxoid) with 5% calf blood at 20°C for 48 - 72  
108 hours. Genomic DNA was extracted from bacterial colonies using a QIAGEN QIAamp DNA mini  
109 kit (QIAGEN, Valencia, CA, USA) according to the manufacturer's protocol. DNA quality was  
110 determined by NanoDrop ND-1000 (Thermo Scientific, Waltham, MA, USA) and DNA  
111 concentration by Qubit 2.0 fluorometer and Quant-iT dsDNA BR kit (Invitrogen, Carlsbad, CA,  
112 USA). All DNA extractions were immediately stored at -20°C until further use.

113

### 114 **2.3 Whole genome sequencing, de novo assembly, and antibiotic resistance genes**

115

116 Genomic DNA was prepared for Illumina pair-end sequencing using the Illumina (Illumina, Inc.,  
117 San Diego, CA) NexteraXT® Guide 150319425031942 following the protocol revision C  
118 ([http://support.illumina.com/downloads/nextera\\_xt\\_sample\\_preparation\\_guide\\_15031942.html](http://support.illumina.com/downloads/nextera_xt_sample_preparation_guide_15031942.html)). A  
119 sample of the pooled NexteraXT Libraries was loaded onto an Illumina HiSeq reagent cartridge  
120 using HiSeq Reagent Kit v2 and 500 cycles with a Standard Flow Cell. The libraries were  
121 sequenced using an Illumina platform and HiSeq Control Software 2.3.0.3. All isolates were pair-  
122 end sequenced. Raw sequence data have been submitted to the European Nucleotide Archive  
123 (<http://www.ebi.ac.uk/ena>) under study accession no.: xxxxxxxx. The raw reads were *de novo*

124 assembled using the assemble pipeline (version 1.0) available from the Center for Genomic  
125 Epidemiology (CGE) <https://cge.cbs.dtu.dk/services/Assembler/> which is based on the Velvet  
126 algorithms for *de novo* short reads assembly (Zerbino and Birney, 2008). Full genomic data can be  
127 retrieved from the supplementary information (SI) appendix, Table S1.

128

129 Identification of acquired antibiotic resistance genes (ARGs) was performed through assembled  
130 genomes using the pipeline ResFinder (version 2.1) (Zankari *et al.*, 2012) available from Center for  
131 Genomic Epidemiology (<http://cge.cbs.dtu.dk/services/>). Threshold for presence of an ARG in an  
132 isolate was set to 75% similarity expressed as percent sequence identity (ID) and 60% of alignment  
133 length (coverage) of resistance gene.

134

## 135 **2.4 Single Nucleotide Polymorphisms (SNPs)**

136

137 SNPs were determined using the pipeline; CSI phylogeny (Leekitcharoenphon *et al.*, 2012; Kaas *et al.*,  
138 2014) available on the CGE ([www.genomicepidemiology.org](http://www.genomicepidemiology.org)). In principle, the paired-end  
139 reads were mapped to the reference chromosome, the French *A. salmonicida* strain A449 isolated  
140 year 1975 from a brown trout (accession number CP000644, chromosome length 4,702,402 bp)  
141 using Burrows-Wheeler Aligner (BWA) version 0.7.2 (Li and Durbin, 2009). The ‘mpileup’ module  
142 in SAMTools version 0.1.18 (Li *et al.*, 2009) was used to identify SNPs. Qualified SNPs were  
143 determined when fulfilling the following criteria: 1) a minimum distance of 10 bps between each  
144 SNP, 2) a minimum of 10% of the relative depth at SNP positions, 3) the mapping quality was more  
145 above 25, 4) the SNP quality was more than 30 and 5) all indels were excluded. The SNPs from  
146 each genome were concatenated to a single alignment corresponding to position of the reference  
147 genome. The concatenated sequences were subjected to maximum likelihood tree using Fastree  
148 (Price *et al.*, 2009)

149

## 150 **2.5 Temporal Bayesian Phylogenetic tree**

151

152 SNPs were subjected to Bayesian temporal phylogenetic reconstruction using BEAST (Bayesian  
153 Evolutionary Analysis Sampling Trees) version 1.7 (Drummond and Rambaut, 2007; Drummond *et al.*,  
154 2012) to estimate mutation rate and divergence time. Combinations of population size change  
155 and molecular clock were evaluated to identify the best-fit model (exponential clock and coalescent  
156 Bayesian skyline). The Bayesian temporal tree was constructed using the best-fit model. The  
157 BEAST MCMC chains were simulated for 300 million steps and subsampled every 10,000 steps.  
158 The final single maximum clade credibility (MCC) was examined using TreeAnnotator (Drummond  
159 *et al.*, 2012) with 10% of the MCMC steps discarded as burn-in. Statistical confidence was  
160 represented by the 95% highest posterior density (HPD) interval.

161

## 162 **2.6 Virulence and iron acquisition proteins**

163

164 To compare presence of virulence and iron acquisition proteins among all *A. salmonicida* isolates, a  
165 blastp search (Altschul *et al.*, 1990) was performed with 78 known virulence associated and iron

166 acquisition protein sequences (SI appendix, Table S2) found in the NCBI protein database against  
167 the assembled *A. salmonicida* genomes. Threshold limit for presence of protein in an isolate was set  
168 to 75% ID.

## 169 **2.7 Plasmid profiles**

170

171 The plasmid content of each *A. salmonicida* was analyzed by using blastn (Altschul *et al.*, 1990)  
172 with 11 known *A. salmonicida* plasmid sequences found in the NCBI database (SI appendix, Table  
173 S3) against the assembled *A. salmonicida* genomes. Threshold limit for presence of plasmid in an  
174 isolate was set to 75% ID and 60% coverage of plasmid due to the long length of plasmid  
175 sequences. Acquired ARGs present in the 11 plasmid sequences were determined using the pipeline  
176 ResFinder (version 2.1) (Zankari *et al.*, 2012) with the above settings for threshold limit.

177

## 178 **3. Results**

179

### 180 **3.1 Phylogeny**

181

182 A total of 667 SNPs were identified in the chromosome from the *A. salmonicida* isolates. The  
183 French reference strain A449 displayed an average SNP difference of 147 to the rest of the isolates.  
184 The Scottish isolate and the type culture NCIMB 1102 had an average SNP difference of 115 and  
185 41 respectively, while two Danish *A. salmonicida* isolated from brown trout (Mj2 1990 and Sd8  
186 1992) had an average difference of 50 and 42 respectively. The average SNP difference among the  
187 Danish isolates was 47 and 46 SNPs between Danish isolates from freshwater versus isolates from  
188 seawater. The three Danish isolates with the highest average SNP difference were Sj7 1980 with 92  
189 SNPs, Mj12 2014 with 67 SNPs and Mj4 2008 with 61 SNPs. Based on the alignment of the 667  
190 SNPs, two trees were constructed: a Bayesian temporal tree (Fig. 1) with a Bayesian Skyline  
191 population size change and an exponential clock rate as the best fit combination model for the *A.*  
192 *salmonicida* population and a maximum likelihood tree (SI appendix, Fig. S1) for topology  
193 confirmation. The two trees showed similar topology and the Bayesian tree (Figure 1) was  
194 illustrated with obtained genetic information regarding acquired ARGs, virulence and iron  
195 acquisition proteins and plasmid profiles of each *A. salmonicida* isolate for further analysis.

196

197 The mutation rate of *A. salmonicida* isolates was estimated to be  $1.93 \times 10^{-7}$  substitutions/site/year,  
198 which corresponds to 0.91 SNPs/genome/year. The most recent common ancestor of the *A.*  
199 *salmonicida* isolates was estimated to have emerged in ~1915 (95% HPD interval 1764 - 1947).  
200 There are two major clades originating back to ~1926 (95% HPD interval 1881 - 1950) that each  
201 branched out further into two more clades in ~1936 (95% HPD interval 1922 - 1958) and ~1970  
202 (95% HPD interval 1934 - 1974) respectively, resulting in roughly four main introductions of *A.*  
203 *salmonicida* in Denmark: ~1973 (95% HPD interval 1958 - 1979), ~1973 (95% HPD interval 1964  
204 - 1981), ~1948 (95% HPD interval 1934 - 1964) and ~1946 (95% HPD interval 1939 - 1961). From  
205 approximately 1975 - 1995 the Danish *A. salmonicida* population experienced a massive clonal  
206 expansion. There was a correlation of local geographical transmission among the Danish freshwater

207 isolates grouped together in the upper clade of the tree. There was another transmission link  
208 between isolates from a freshwater farm Mj10 and isolates from two seawater farms that had  
209 received fish from this farm.

210

### 211 **3.2 Antibiotic resistance**

212

213 All sequenced *A. salmonicida* isolates harbored three ARGs against beta-lactam antibiotics encoded  
214 on the chromosome (Table 1). Nine Danish *A. salmonicida* isolates also harbored several other  
215 plasmid encoded resistance genes against trimethoprim, sulphonamide and aminoglycoside  
216 antibiotics (Table 1). All three isolates from freshwater farm Mj10 sampled during different years  
217 harbored ARGs against several of the above mentioned antibiotics. The same ARGs against  
218 multiple antibiotics were found in isolates sampled from three seawater farms (Mj8 1997, Mj11  
219 2014 and Mj3 2014) located in the same bay that all received fish from the freshwater farm Mj10.  
220 The French reference strain A449 also harbored ARGs against beta-lactam, sulphonamide,  
221 aminoglycoside, phenicol and tetracycline antibiotics and more resistance genes are described by  
222 (Reith *et al.*, 2008).

223

### 224 **3.3 Virulence and iron acquisition**

225

226 Out of 78 investigated protein sequences, 22 were considered as absent (<65% ID) in one or more  
227 isolates (Fig. 2). The Type Three Secretion System (T3SS) Effector protein AopP encoded on  
228 plasmid pAsa11 by the *aopP* gene was absent in 50% the *A. salmonicida* isolates, including the  
229 reference strain A449. A cluster of 15 T3SS related proteins were absent in 25 isolates. In nine of  
230 the 25 isolates, the T3SS putative tyrosine phosphatase AopH and its chaperone that are encoded on  
231 pAsa5 and have homologs encoded on pAsa6 were also absent. Three of the isolates were also  
232 missing the T3SS putative serine/threonine kinase AopO and its chaperone that are encoded on  
233 pAsa5, while isolate Mj12 2014 was missing the extracellular nuclease protein (48% ID) coded by  
234 the gene *nucH* on the chromosome. Isolate Sj4 1998, which is not included in above mentioned  
235 group of 25 isolates, did not possess the tetragonal surface virulence array protein VapA (A-layer)  
236 encoded on the chromosome. Lastly, the chromosome encoded ABC-type ferric siderophore  
237 transporter permease protein only showed 75% ID in all sequenced *A. salmonicida* as well as the  
238 reference strain A449.

239

### 240 **3.4 Plasmid profiles**

241

242 All examined *A. salmonicida* isolates displayed presence of multiple plasmids (SI appendix, Table  
243 S4). Seven plasmids were present in one or more isolates, while four plasmids: pAr-32, pRAS1,  
244 pRAS3.1 and pRAS3.2 were not present in any isolates. The only plasmid found in all isolates was  
245 pAsa1, although pAsa2 showed high stability with a presence of 99% among the isolates. Plasmids  
246 pAsa5 and pAsa6 were present in 90% and 85% respectively. The two plasmids pAsa3 and  
247 pAsa11 were present in 76% and 52% of the isolates respectively, while pAsa4 was only present in  
248 the reference strain A449. Twelve different plasmid profiles were detected among the isolates, with

249 one profile consisting of pAsa1, pAsa2, pAsa3, pAsa5, pAsa6 and pAsa11 representing 44% of the  
250 isolates (Table 2).

251

252 Five plasmids are known for harboring ARGs (R plasmids): pAsa4, pAr-32, pRAS1, pRAS3.1 and  
253 pRAS3.2 (SI appendix Table S3) and only plasmid pAsa4 was present in any of the investigated *A.*  
254 *salmonicida* (the reference strain A449).

255

256 All 51 *A. salmonicida* isolates in which the AopP protein encoded on pAsa11 was absent (Fig. 2),  
257 were missing plasmid pAsa11 (SI appendix Table S4). Of the 25 *A. salmonicida* isolates that were  
258 missing a cluster of 15 T3SS proteins encoded on pAsa5, ten were also missing the plasmid pAsa5.  
259 The remaining 15 isolates all displayed <80% coverage of the pAsa5. All nine *A. salmonicida*  
260 isolates that lacked the protein AopH and its chaperone that are encoded on pAsa5 and have  
261 homologs on pAsa6 showed <80% coverage for pAsa5 and were missing pAsa6.

262

## 263 **4. Discussion**

264

### 265 **4.1 Phylogeny**

266

267 *A. salmonicida* subspecies is known to be a highly homogenous group that is considered clonal  
268 (Wiklund and Dalsgaard, 1998; Garcia *et al.*, 2000; Cunningham and Colquhoun, 2002; Beaz-  
269 Hidalgo *et al.*, 2008). The fact that only a total of 667 SNPs were found in the entire 4,702,402 bp  
270 long chromosome among the investigated *A. salmonicida* varying in year of isolation (span of 34  
271 years), geographical region, and host fish species only confirms this further. The highest average  
272 SNP difference was found between the French reference strain A449 and the rest of the *A.*  
273 *salmonicida* isolates (average of 147 SNPs), which is not a large difference when considering that  
274 A449 was isolated in France and from a brown trout, while almost all (97 out of 99) of the Danish  
275 isolates were isolated from rainbow trout. When comparing the average SNP difference between the  
276 Scottish strain from Atlantic salmon and the rest of the isolates, the results were even lower  
277 (average of 115 SNPs). The two Danish *A. salmonicida* isolated from brown trout (Mj2 1990 and  
278 Sd8 1992) and the type strain NCIMB 1102 from Atlantic salmon also grouped together with  
279 Danish *A. salmonicida* from rainbow trout in one of the four major clades (Fig. 1) and only have an  
280 average of 41, 50 and 52 SNP difference, which challenges the theory of *A. salmonicida* genome  
281 adapting to the environment of their specific hosts species (Reith *et al.*, 2008). More *A. salmonicida*  
282 isolates from various fish species would, however, need to be sequenced in order to shed more light  
283 on this theory.

284

285 Noticeably there appears to be four major *A. salmonicida* introductions to Denmark, giving rise to  
286 four major clades (Fig. 1). The two introductions that occurred in ~1973 (95% HPD interval 1958 -  
287 1979) and ~1973 (95% HPD interval 1964 - 1981) and gave rise to the two upper clades (Fig. 1),  
288 seemingly took place right before the massive clonal expansion during 1975 - 1995, which all four  
289 clades underwent. The two introduction points in 1973 and the expansion period of all four clades  
290 correspond to the time period where rainbow trout farming in Denmark began expanding out to

291 seawater and intensifying their production. The two bottom clades were introduced further in the  
292 past ~1948 (95% HPD interval 1934 - 1964) and ~ 1946 (95% HPD interval 1939 - 1961)  
293 respectively and also contain *A. salmonicida* with older isolation years (average year of isolation  
294 1991), than the two upper clades that include *A. salmonicida* with an average isolation year of 2001  
295 (Fig. 1).

296  
297 The introduction of the two bottom clades correlate with the end of the Second World War and the  
298 beginning of an expansion of rainbow trout production in Danish freshwater. When examining the  
299 branches of each of the four major clades, there is a possibility that *A. salmonicida* might have been  
300 introduced into Denmark through seawater and was from thereon spread to freshwater. One  
301 explanation for this possible scenario is the fact that wet feed, comprised of marine fish including  
302 sand-eels, was used at all Danish fish farms until the late 1970s (Christensen, 1980). *A.*  
303 *salmonicida*, although the atypical type, has been isolated from sand-eels caught in the surrounding  
304 seawater of Denmark (Dalsgaard and Paulsen, 1986). It is also known that *A. salmonicida* can be  
305 harbored by various farmed and wild freshwater and seawater aquatic animals (Bernoth *et al.*,  
306 1997). It could be thus be hypothesized that the wet feed could be the cause behind a possible initial  
307 transmission of the bacterium from seawater to freshwater. However, the bacterium could also have  
308 been transmitted from freshwater to seawater, which is the widespread theory i.e. that *A.*  
309 *salmonicida* is present in freshwater fish showing no signs of diseases (carriers) and are then  
310 transferred out to seawater with the fish, where outbreaks occur during high temperatures (Larsen  
311 and Møllgaard, 1981; Dalsgaard and Madsen, 2000; Pedersen *et al.*, 2008).

312  
313 The local transmission pattern of *A. salmonicida* among the Danish farms also suggests that  
314 transmission of isolates from freshwater to seawater farms have occurred, as exemplified by a  
315 minor clade where ARGs were transmitted from a freshwater farm to seawater farms, though this  
316 will be discussed under the antibiotic resistance section. Isolates from different freshwater farms are  
317 moreover mixed with different isolates from seawater in most of the minor clades. Though, in  
318 general it is hard to find specific geographical correlations between the fish farms. One of the main  
319 causes for this could be the widespread trade of fingerlings for anglers in Denmark throughout the  
320 years as well as local trade among fish farmers. There is nonetheless a correlation among the group  
321 of freshwater farms isolates in the top clade. Mj13 is located upstream to Mj16 in a stream that runs  
322 out into a river named Guden Å. Two other farms (Mj12 and Mj2) are also located at streams that  
323 lead out to Guden Å and one of these (Mj12) produces brown trout.

324  
325 Finally, it was also found that isolates grouped in the two bottom clades were missing a higher  
326 amount of virulence associated proteins (average of 1.0 protein per isolate), compared to isolates  
327 grouped in the two upper clades where the average absence of virulence associated proteins was 0.6  
328 per isolate (Fig. 1). This indicates that the most recently common ancestor of the bottom two clades  
329 from ~1936 (95% HPD interval 1922 - 1958), presumably harbored itself and gave rise to two older  
330 lineages that harbored fewer virulence associated proteins than the more recently introduced two  
331 lineages (the two upper clades in the phylogeny tree). Considering these results, it could be



332 suggested that the upper two clades might consist of more virulent *A. salmonicida* that were  
333 introduced from a more recent and more virulent ancestor around 1970.

334

## 335 **4.2 Antibiotic resistance**

336

337 Interestingly, all investigated *A. salmonicida* isolates possessed three beta-lactam ARGs. Since the  
338 genes are encoded on the chromosome, it seems that either they have always been a part of the *A.*  
339 *salmonicida* genome repertoire, or they must have been acquired at least around 67 - 250 years ago.  
340 Nine Danish *A. salmonicida* also harbored resistance genes against trimethoprim, sulphonamide and  
341 aminoglycoside antibiotics, which are all plasmid encoded. Trimethoprim and sulphonamide are  
342 also two of the scarce number of antibiotics allowed to be used for treatment of bacterial diseases in  
343 Danish fish farms. All three *A. salmonicida* isolates from freshwater farm Mj10, isolated year 1982,  
344 2009 and 2010 harbored resistance genes against at least two of the above mentioned antibiotics  
345 (Table 1). Fish from farm Mj10 have always been transferred out to a bay, where several seawater  
346 farms are located. In *A. salmonicida* isolated during 2014 from two of these seawater farms, the  
347 same set of resistance genes were detected as those seen in *A. salmonicida* from the freshwater farm  
348 Mj10 during 2009 and 2010. Fittingly, the two freshwater isolates form a minor clade with the two  
349 seawater isolates in the Bayesian temporal tree, according to which the isolates spread from the  
350 freshwater to seawater (Fig. 1). In seawater farm Mj8, which is also located in the bay, an *A.*  
351 *salmonicida* isolate from 1997 did harbor ARGs against the mentioned antibiotics as well.  
352 However, these ARGs were slightly different than those seen in the Mj10 1982 isolate, where one  
353 otherwise would expect the resistance genes had originated (Table1). This could be associated with  
354 the fact that trimethoprim was not licensed for use in Denmark until 1983 and the isolate from 1982  
355 had therefore not acquired ARGs against this antibiotic.

356

357 None of the nine Danish isolates harbored any of the five investigated R plasmids, though eight did  
358 show coverage (< 60%) of at least one of the R plasmids (SI appendix, Table S4), indicating they  
359 could have acquired ARGs from the plasmids in the past through horizontal gene transfer and then  
360 lost the plasmid. Isolate Sj4 2014 that showed zero coverage of all R plasmids harbored ARGs *strA*  
361 and *strB*, which were present in R plasmid pRAS2 isolated from *A. salmonicida* in salmon from  
362 Norway (L'Abée-Lund and Sørnum, 2000). This plasmid was not included in the analysis, since only  
363 a couple of gene sequences from this plasmid are available in Genbank. However, pRAS2 could be  
364 or have been present among the Danish *A. salmonicida* population. Noteworthy, the only other  
365 isolate harboring ARGs *strA* and *strB* was isolated in 2011 from farm Nj1, though there has not  
366 been any transfer of fish from this farm to Sj4, whereby the two occurrences of the ARGs *strA* and  
367 *strB* are incidental. When looking at the broader picture, there are many highly similar broad host R  
368 plasmids that have been isolated from *Aeromonas* species in various environments all over the  
369 world that harbor the same ARGs found in the Danish *A. salmonicida* (L'Abée-Lund and Sørnum,  
370 2001; Sørnum *et al.*, 2003; Kadlec *et al.*, 2011; Muziasari *et al.*, 2014). Present findings only provide  
371 further evidence of this widespread dissemination of R plasmids and ARGs, although the  
372 prevalence of these ARGs seems to be similar (9%) to the low prevalence of 5% found by  
373 Dalsgaard *et al.*, (1994).

374

### 375 **4.3 Iron acquisition**

376

377 Iron acquisition has been proven to be an important factor for virulence in almost all bacterial  
378 pathogens, including *A. salmonicida* where it also seems to be linked to survival in aquatic  
379 environments (Reith *et al.*, 2008). Iron is also thought to be linked to the innate immune response in  
380 the host, which in turn attempts limiting iron availability in order to lower virulence and access of  
381 the pathogen into the host (Ganz, 2009; Lee *et al.*, 2014). Due to the important nature of iron  
382 acquisition and the fact that all the investigated proteins related to iron acquisition were encoded on  
383 the chromosome, it was expected that all 102 *A. salmonicida* isolates possessed all the investigated  
384 iron acquisition proteins (SI appendix, Table S2). Though, it has to be noted that the ABC-type  
385 ferric siderophore transporter permease protein only showed 75% ID in all *A. salmonicida* isolates.

386

### 387 **4.4 Virulence**

388

389 In addition to iron acquisition proteins, numerous of potential virulence factors have been identified  
390 in *A. salmonicida*, including extracellular proteases, lipases, adhesins, and functional secretion  
391 systems (Burr *et al.*, 2002; Rasch *et al.*, 2007; Reith *et al.*, 2008; Dallaire-Dufresne *et al.*, 2014).  
392 The majority of potential virulence proteins investigated in this study was present in all sequenced  
393 *A. salmonicida* and the reference strain. Though, two proteins encoded on the chromosome and 20  
394 encoded on plasmids were missing in at least one isolate (SI appendix, Table S2). In agreement with  
395 the present results, DNA micro array study of virulence genes in *Aeromonas* species including  
396 several subspecies *salmonicida* isolates by (Nash *et al.*, 2006) also showed high degree of  
397 variability among genes associated with plasmids, whereas genes encoded on the chromosome did  
398 not vary significantly.

399

400 All of the above mentioned 20 plasmid encoded proteins were related to the functional type three  
401 secretion system (T3SS). This secretion system is wide spread among Gram-negative bacteria and  
402 has several functions, including: disrupting host cells by translocating toxins (effector proteins) into  
403 their cytoplasm, preventing phagocytosis by leukocytes, and establishing systemic infection (Burr *et al.*,  
404 2003; Stuber *et al.*, 2003; Burr *et al.*, 2005; Dacanay *et al.*, 2006; Rasch *et al.*, 2007; Dallaire-  
405 Dufresne *et al.*, 2014). T3SS is also the only virulence factor proven to be essential for virulence of  
406 *A. salmonicida*, as all *in vitro* and *in vivo* studies involving inactivation of T3SS structural proteins  
407 in *A. salmonicida* strains have resulted in non-virulent *A. salmonicida* mutants (Burr *et al.*, 2002;  
408 Burr *et al.*, 2003; Stuber *et al.*, 2003; Burr *et al.*, 2005; Dacanay *et al.*, 2006; Froquet *et al.*, 2007).  
409 Nevertheless, among the 20 missing T3SS related proteins in this study were T3SS structural  
410 proteins, while all the *A. salmonicida* in this study are isolated from furunculosis outbreaks,  
411 whereby one would assume that all the *A. salmonicida* isolates are virulent.

412

413 There are 36 T3SS encoding genes located on the large plasmid pAsa5 (Reith *et al.*, 2008; Najimi *et al.*  
414 *et al.*, 2009; Tanaka *et al.*, 2012; Vincent *et al.*, 2016) and 19 of them that were investigated in this  
415 study were missing in at least three isolates (SI appendix, Table S2). Initially these 36 genes were

416 found to be located on a 140 kb plasmid named pASvirA (Stuber *et al.*, 2003) and while pASvirA  
417 and pAsa5 are almost the same size, it remains unclear whether they are variants of the same  
418 plasmid (Najimi *et al.*, 2009). Though, both plasmids become unstable under stressful conditions  
419 like being subjected to growth in temperature above 22 - 25°C (Stuber *et al.*, 2003; Tanaka *et al.*,  
420 2012; Dallaire-Dufresne *et al.*, 2014). While pASvirA is seemingly lost by *A. salmonicida* during  
421 the stressful conditions (Stuber *et al.*, 2003), pAsa5 is thought to undergo genetic rearrangement  
422 resulting in the loss of its T3SS region caused by activation of ISAS11 insertion sequence (IS)  
423 elements (Tanaka *et al.*, 2012). This could explain the fact that all the 25 *A. salmonicida* isolates  
424 missing the cluster of T3SS proteins, encoded on pAsa5 in our study still harbored the plasmid, but  
425 displayed <80% coverage. The only issue with this justification is that all *A. salmonicida* cultures in  
426 our laboratory are always grown at 20°C, meaning it is unlikely that growth at high temperature  
427 triggered the activation of ISAS11. It is unclear what other cause for the rearrangement of pAsa5  
428 could have been.

429  
430 Plasmid pAsa6 also shares most of its sequences, including sequences for AopH and its chaperone,  
431 with pAsa5 with a close to 100% nucleotide sequence similarity (Najimi *et al.*, 2009). In agreement,  
432 all nine *A. salmonicida* isolates in our study that lacked AopH and its chaperone showed <80%  
433 coverage of pAsa5 and were missing pAsa6 (SI appendix, Table S4). Presence of AopH and  
434 chaperone homologs on pAsa6 would also explain why the sequences for these two proteins were  
435 not missing in all of the above mentioned 25 isolates i.e. the two protein sequences in all but nine  
436 isolates could still be present on pAsa6. Though, it remains unclear why only three of the 25  
437 isolates were also missing the AopO protein and its chaperone, since both proteins are only encoded  
438 on pAsa5. One reason could be that they were encoded in a different region and were thus not  
439 rearranged along with the other 15 T3SS protein sequences.

440  
441 Unlike most of the T3SS proteins, the AopP protein is encoded on plasmid pAsal1 and the protein  
442 was missing in 50% of the *A. salmonicida* isolates investigated in this study (Fig. 2). Interestingly,  
443 pAsal1 was present in isolate Sj 1981 and the Scottish isolate, both of which did not harbor the  
444 AopP protein. The isolates did possess the nucleotide sequence for the *aopP* gene; however, both  
445 sequences had an identical frameshift mutation caused by point deletions (data not showed) that  
446 presumably lead to an incorrect translation of the AopP protein sequence. Apart from the two  
447 mentioned isolates, all *A. salmonicida* that were missing the AopP protein sequence were also  
448 missing the plasmid pAsal1. Previous studies have shown that pAsal1 is lost due to activation of the  
449 same insertion sequence as in pAsa5 (Daher *et al.* 2011; Tanaka *et al.* 2012; Attéré *et al.*, 2015;  
450 Vincent *et al.*, 2016), though there was no correlation between a lowered coverage of pAsa5 and  
451 absence of pAsal1. Though the plasticity of *A. salmonicida* pAsal1 is complex, as the precise  
452 mechanism responsible for loss of pAsal1 remains unknown (Attéré *et al.*, 2015) and at least three  
453 larger variants of the plasmid exist: pAsal1B, pAsal1C and pAsal1D (Trudel *et al.*, 2013; Attéré *et al.*,  
454 2015). All the variants harbor another insertion sequence element called ISAS5 that in pAsal1C  
455 and pAsal1D disrupts the ISAS11, which hence cannot be activated leading to the prevention of loss  
456 of these two variants of pAsal1 during stressful conditions (Attéré *et al.*, 2015). Although the ISAS5

457 does not disrupt ISAS11 in pAsa1B, according to Attéré *et al.*, (2015) it could still prevent  
458 activation of ISAS11.

459

460 The last two missing virulence associated proteins, the A-layer protein encoded by the *vapA* gene  
461 and the extracellular nuclease protein encoded by the gene *nucH*, were missing in isolate Sj4 1998  
462 and Mj12 2014 respectively. The absence of the extracellular protein in isolate Mj12 2014  
463 correlates with its unusually high average SNP difference of 67 compared to other Danish *A.*  
464 *salmonicida*. Isolate Sj4 1998 that is missing *vapA* does not have a higher SNP difference (47)  
465 compared to the other Danish *A. salmonicida*. Interestingly, isolate Sj7 1980 that has the highest  
466 average SNP difference of 92 SNPs among the Danish isolates, had an ID percentage of 87% for  
467 *vapA* and when the isolates' *vapA* gene sequence was manually investigated with the program  
468 BioEdit (Hall, 1999), it was discovered that the first half of the nucleotide gene sequence differs  
469 significantly from the *vapA* nucleotide sequences of the other *A. salmonicida* isolates (data not  
470 shown). As illustrated on Fig. 2, there was otherwise an overall high similarity among all isolates  
471 regarding all the chromosome encoded virulence associated protein sequences. The high prevalence  
472 and similarity of the A-layer (VapA) in *A. salmonicida* discovered in this study, along with previous  
473 findings of *A. salmonicida* surface structures in contact with host defenses having a high antigenic  
474 conservation (Chart *et al.*, 1984), could also provide valuable knowledge for future vaccine  
475 development.

476

#### 477 **4.5 Plasmid profiles**

478

479 Out of the twelve plasmid profiles found in this study, the most abundant profile consisting of  
480 pAsa1, pAsa2, pAsa3, pAsa5, pAsa6 and pAsa11 represented 44% of the *A. salmonicida* isolates  
481 (Table 2). In the study by Nielsen *et al.* (1993) of *A. salmonicida* from various geographical  
482 locations using DNA restriction fragment plasmid profiling, a plasmid profile with the following  
483 five plasmids: pAsa1, pAsa2, pAsa3, pAsa5 and pAsa11 was the most common profile among  
484 Danish *A. salmonicida* isolates, representing 32% of the 57 investigated Danish strains in that study.  
485 Nielsen *et al.* (1993) also investigated *A. salmonicida* NCIMB 1102, which belonged to the plasmid  
486 profile group mentioned above and was thus missing pAsa6. Seemingly pAsa6 (molecular weight  
487 of 18.5 kb) was not found in any of the 124 *A. salmonicida* strains investigated by Nielsen *et al.*  
488 (1993). On the contrary, present results showed that pAsa6 was present in 87 *A. salmonicida*  
489 including NCIMB 1102. Possible explanations for the disagreeing results regarding pAsa6 could be  
490 that pAsa6 was not observed on gel by Nielsen *et al.* (1993) due being present in a low copy  
491 number, or pAsa6 could integrate into the *A. salmonicida* chromosome due to the abundance of IS  
492 elements within the plasmid (Najimi *et al.*, 2009).

493

494 Present findings support previous results by Nielsen *et al.* (1993), Boyd *et al.* (2003) and Attéré *et*  
495 *al.* (2015) regarding high stability of plasmid pAsa1 and pAsa2 and instability of plasmid pAsa3  
496 and pAsa11. Attéré *et al.* (2015) suggested an explanation for the stability of pAsa1 and instability  
497 of pAsa11 could be that pAsa1 and pAsa3 encode genes for a type II toxin-antitoxin (TA) system  
498 that kills all daughter cells that do not receive the plasmids (Boyd *et al.*, 2003), while the TA system

499 has not been found in plasmids pAsa2 and pAsa1 (Shao *et al.*, 2011). Though there is still the issue  
500 regarding stability of pAsa2 that does not encode a TA system and the instability of pAsa3 that does  
501 have a TA system (Attéré *et al.*, 2015); an issue accentuated by the present findings. It might be  
502 possible that some clonal lineages do not acquire pAsa3 and pAsa1 (Attéré *et al.*, 2015), which  
503 cannot be ruled out according to our results, where out of 24 *A. salmonicida* that did not harbor  
504 pAsa3 and pAsa1 (plasmid profiles nine and ten), 17 clustered together in four minor clades (Fig.  
505 1, four minor clades with a red ring).

506

## 507 **5. Conclusion**

508

509 The present findings have provided novel insight into the epidemiology of the disease causing  
510 Danish *A. salmonicida*, revealing four main introductions in consistency with the historical  
511 expansion of the Danish aquaculture production that could have been transmitted either from  
512 freshwater to seawater or vice versa. There was also transmission of isolates harboring ARGs from  
513 a freshwater farm to seawater farms, supporting the theory of *A. salmonicida* being spread from  
514 freshwater to seawater via carrier fish. The mixture of freshwater and seawater isolates from  
515 different farms in almost every minor clade and the lack of geographical connections among farms  
516 also indicates that the widespread trade of fingerlings and other fish could have played a role in the  
517 local dissemination of *A. salmonicida* in Denmark. The genome based analysis moreover showed  
518 genetic and virulence variability among the highly homogenous *A. salmonicida* population in  
519 Denmark, which consisted of isolates with varying plasmid profiles and plasmid encoded virulence  
520 proteins, especially those related to T3SS. Overall, WGS proved to be a highly useful tool for  
521 investigating Danish *A. salmonicida* and presented important new information about this bacterium.

522

## 523 **5. Acknowledgements**

524

525 This work was supported by The Danish Council for Strategic Research under ProFish project  
526 (Grant no. DSF: 11-116252) and National Veterinary Institute (DTU). The authors would like to  
527 thank all ProFish project partners. The authors are also grateful to Lisbeth Schade Hansen at the  
528 National Veterinary Institute (DTU) and Inge Marianne Hansen at the National Food Institute  
529 (DTU) for their technical support.

530

## 531 **6. Contributions**

532

533 Conceived and designed the experiments: ID, FMA. Contributed reagents/materials/analysis tools:  
534 SB, PL, ID, FMA. Performed field sampling: ID. Performed the experiments: SB, ID. Analyzed the  
535 data: SB, PL, ID, FMA. Wrote the paper: SB, PL. Critically revised the paper: FMA, ID.

536

## 537 **7. References**

538

539 Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment  
540 search tool. *J. Mol. Biol.* 215, 403–10. doi:10.1016/S0022-2836(05)80360-2.

- 541 Attéré, S. A., Vincent, A. T., Trudel, M., Chanut, R., and Charette, S. J. (2015). Diversity and  
542 homogeneity among small plasmids of *Aeromonas salmonicida* subsp. *salmonicida* linked with  
543 geographical origin. *Front. Microbiol.* 6, 1274.
- 544 Beaz-Hidalgo, R., López-Romalde, S., Toranzo, A. E., and Romalde, J. L. (2008). Polymerase  
545 chain reaction amplification of repetitive intergenic consensus and repetitive extragenic  
546 palindromic sequences for molecular typing of *Pseudomonas anguilliseptica* and *Aeromonas*  
547 *salmonicida*. *J. Aquat. Anim. Health* 20, 75–85. doi:10.1577/H07-007.1.
- 548 van Belkum, A., Tassios, P. T., Dijkshoorn, L., Haeggman, S., Cookson, B., Fry, N. K., et al.  
549 (2007). Guidelines for the validation and application of typing methods for use in bacterial  
550 epidemiology. *Clin. Microbiol. Infect.* 13 Suppl 3, 1 – 46.
- 551 Bernoth, E., Ellis, A. E., Midtlyng, P. J., Olivier, G., and Smith, P. eds. (1997). *Furunculosis:*  
552 *Multidisciplinary fish disease research*. London UK: Academic Press.
- 553 Boyd, J., Williams, J., Curtis, B., Kozera, C., Singh, R., and Reith, M. (2003). Three small, cryptic  
554 plasmids from *Aeromonas salmonicida* subsp. *salmonicida* A449. *Plasmid* 50, 131–144.  
555 doi:10.1016/S0147-619X(03)00058-1.
- 556 Burr, S. E., Pugovkin, D., Wahli, T., Segner, H., and Frey, J. (2005). Attenuated virulence of an  
557 *Aeromonas salmonicida* subsp. *salmonicida* type III secretion mutant in a rainbow trout model.  
558 *Microbiology* 151, 2111–2118. doi:10.1099/mic.0.27926-0.
- 559 Burr, S. E., Stuber, K., Wahli, T., and Frey, J. (2002). Evidence for a Type III Secretion System in  
560 *Aeromonas salmonicida* subsp. *salmonicida*. *J. Bacteriol.* 184, 5966–5970.  
561 doi:10.1128/JB.184.21.5966-5970.2002.
- 562 Burr, S. E., Wahli, T., Segner, H., Pugovkin, D., and Frey, J. (2003). Association of Type III  
563 secretion genes with virulence of *Aeromonas salmonicida* subsp. *salmonicida*. *Dis. Aquat.*  
564 *Organ.* 57, 167–171. Available at:  
565 <http://www.ncbi.nlm.nih.gov/pubmed/14735937>\nfile:///C:/Users/Joanna/Documents/my  
566 PhD/Literature/Burr et al 2003 aeromonas salm.pdf.
- 567 Chart, H., Shaw, D. H., Ishiguro, E. E., and Trust, T. J. (1984). Structural and immunochemical  
568 homogeneity of *Aeromonas salmonicida* lipopolysaccharide. *J. Bacteriol.* 158, 16–22.
- 569 Christensen, N. O. (1980). *Fiskesygdome*. Copenhagen: Den Kgl. Veterinær- og  
570 Landbohøjskolen.
- 571 Cunningham, C. O., and Colquhoun, D. J. (2002). “Molecular diagnosis of *Aeromonas salmonicida*  
572 infections,” in *Molecular Diagnosis of Salmonid Diseases*, ed. C. D. Cunningham (Kluwer  
573 Academic Publishers), 81–99. doi:10.1007/978-94-017-2315-2.
- 574 Dacanay, A., Knickle, L., Solanky, K. S., Boyd, J. M., Walter, J. a., Brown, L. L., et al. (2006).  
575 Contribution of the type III secretion system (TTSS) to virulence of *Aeromonas salmonicida*  
576 subsp. *salmonicida*. *Microbiology* 152, 1847–1856. doi:10.1099/mic.0.28768-0.
- 577 Daher, R. K., Filion, G., Tan, S. G. E., Dallaire-Dufresne, S., Paquet, V. E., and Charette, S. J.  
578 (2011). Alteration of virulence factors and rearrangement of pAsa5 plasmid caused by the  
579 growth of *Aeromonas salmonicida* in stressful conditions. *Vet. Microbiol.* 152, 353–60.  
580 doi:10.1016/j.vetmic.2011.04.034.
- 581 Dallaire-Dufresne, S., Tanaka, K. H., Trudel, M. V., Lafaille, A., and Charette, S. J. (2014).  
582 Virulence, genomic features, and plasticity of *Aeromonas salmonicida* subsp. *salmonicida*, the  
583 causative agent of fish furunculosis. *Vet. Microbiol.* 169, 1–7.  
584 doi:10.1016/j.vetmic.2013.06.025.
- 585 Dalsgaard, I., and Madsen, L. (2000). Bacterial pathogens in rainbow trout, *Oncorhynchus mykiss*  
586 (Walbaum), reared at Danish freshwater farms. *J. Fish Dis.* 23, 199 – 209.
- 587 Dalsgaard, I., Nielsen, B., and Larsen, J. L. (1994). Characterization of *Aeromonas salmonicida*  
588 subsp. *salmonicida*: a comparative study of strains of different geographic origin. *J. Appl.*

589 *Bacteriol.* 77, 21–30.

590 Dalsgaard, I., and Paulsen, H. (1986). Atypical *Aeromonas salmonicida* isolated from diseased  
591 sand-eels, *Ammodytes lancea* (Cuvier) and *Hyperoplus lanceolatus* (Lesauvege). *J. Fish Dis.* 9,  
592 361–364.

593 Drummond, A. J., and Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling  
594 trees. *Bmc Evol. Biol.* 7.

595 Drummond, A. J., Suchard, M. A., Xie, D., and Rambaut, A. (2012). Bayesian phylogenetics with  
596 BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 29, 1969–73. doi:10.1093/molbev/mss075.

597 Emmerich, R., and Weibel, E. (1894). Über eine durch bakterien erzeugte seuche unter den forellen.  
598 *Arch. für Hyg.* 21, 1–21.

599 Foxman, B., Zhang, L., Koopman, J. S., Manning, S. D., and Marrs, C. F. (2005). Choosing an  
600 appropriate bacterial typing technique for epidemiologic studies. *Epidemiol. Perspect. Innov.*  
601 2, 10. doi:10.1186/1742-5573-2-10.

602 Froquet, R., Cherix, N., Burr, S. E., Frey, J., Vilches, S., Tomas, J. M., et al. (2007). Alternative  
603 host model to evaluate *Aeromonas* virulence. *Appl. Environ. Microbiol.* 73, 5657–5659.  
604 doi:10.1128/AEM.00908-07.

605 Ganz, T. (2009). Iron in innate immunity: starve the invaders. *Curr. Opin. Immunol.* 21, 63–7.  
606 doi:10.1016/j.coi.2009.01.011.

607 Garcia, J. A., Larsen, J. L., Dalsgaard, I., and Pedersen, K. (2000). Pulsed-field gel electrophoresis  
608 analysis of *Aeromonas salmonicida* ssp *salmonicida*. *FEMS Microbiol. Lett.* 190, 163–166.  
609 doi:10.1016/S0378-1097(00)00322-0.

610 Hall, T. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program  
611 for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 41, 95–98. doi:citeulike-article-id:691774.

612 Kadlec, K., von Czapiewski, E., Kaspar, H., Wallmann, J., Michael, G. B., Steinacker, U., et al.  
613 (2011). Molecular basis of sulfonamide and trimethoprim resistance in fish-pathogenic  
614 *Aeromonas* isolates. *Appl. Environ. Microbiol.* 77, 7147–7150. doi:10.1128/AEM.00560-11.

615 Kaas, R. S., Leekitcharoenphon, P., Aarestrup, F. M., and Lund, O. (2014). Solving the problem of  
616 comparing whole bacterial genomes across different sequencing platforms. *PLoS One* 9,  
617 e104984. doi:10.1371/journal.pone.0104984.

618 L'Abée-Lund, T. M., and Sørum, H. (2000). Functional Tn5393-like transposon in the R plasmid  
619 pRAS2 from the fish pathogen *Aeromonas salmonicida* subspecies *salmonicida* isolated in  
620 Norway. *Appl. Environ. Microbiol.*, 5533–5535.

621 L'Abée-Lund, T. M., and Sørum, H. (2001). Class 1 integrons mediate antibiotic resistance in the  
622 fish pathogen *Aeromonas salmonicida* worldwide. *Microb. Drug Resist.* 7, 263–72.  
623 doi:10.1089/10766290152652819.

624 Larsen, J. L., and Mellergaard, S. (1981). Microbiological and hygienic problems in marine  
625 aquaculture: furunculosis and vibriosis in rainbow trout (*Salmo gairdneri*). *Bull. Eur. Ass. Fish*  
626 *Pathol.* 1, 29–32.

627 Lee, J.-H., Mohd-Adnan, A., Pooley, N. J., and Martin, S. A. M. (2014). Cloning and  
628 Characterisation of Multiple Ferritin Isoforms in the Atlantic Salmon (*Salmo salar*). *PLoS One*  
629 9.

630 Leekitcharoenphon, P., Kaas, R. S., Thomsen, M., Friis, C., Rasmussen, S., and Aarestrup, F. M.  
631 (2012). snpTree - a web-server to identify and construct SNP trees from whole genome  
632 sequence data. *BMC Genomics* 13. doi:doi:10.1186/1471-2164-13-S7-S6.

633 Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler  
634 transform. *Bioinformatics* 25, 1754–60. doi:10.1093/bioinformatics/btp324.

635 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The Sequence  
636 Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–9.

637 doi:10.1093/bioinformatics/btp352.

638 Muziasari, W. I., Managaki, S., Pärnänen, K., Karkman, A., Lyra, C., Tamminen, M., et al. (2014).  
639 Sulphonamide and trimethoprim resistance genes persist in sediments at Baltic Sea aquaculture  
640 farms but are not detected in the surrounding environment. *PLoS One* 9.  
641 doi:10.1371/journal.pone.0092702.

642 Najimi, M., Balado, M., Lemos, M. L., and Osorio, C. R. (2009). Genetic characterization of pAsa6,  
643 a new plasmid from *Aeromonas salmonicida* subsp. *salmonicida* that encodes a type III  
644 effector protein AopH homolog. *Plasmid* 61, 176–81. doi:10.1016/j.plasmid.2009.01.001.

645 Nash, J. H. E., Findlay, W. A., Luebbert, C. C., Mykytczuk, O. L., Foote, S. J., Taboada, E. N., et  
646 al. (2006). Comparative genomics profiling of clinical isolates of *Aeromonas salmonicida*  
647 using DNA microarrays. *BMC Genomics* 7, 43. doi:10.1186/1471-2164-7-43.

648 Nielsen, B., Olsen, J. E., and Larsen, J. L. (1993). Plasmid profiling as an epidemiological marker  
649 within *Aeromonas salmonicida*. *Dis. Aquat. Organ.* 15, 129–135. doi:10.3354/dao015129.

650 Parkhill, J., and Wren, B. W. (2011). Bacterial epidemiology and biology-lessons from genome  
651 sequencing. *Genome Biol.* 12, 230. doi:10.1186/gb-2011-12-10-230.

652 Pedersen, K., Skall, H. F., Lassen-Nielsen, A. M., Nielsen, T. F., Henriksen, N. H., and Olesen, N.  
653 J. (2008). Surveillance of health status on eight marine rainbow trout, *Oncorhynchus mykiss*  
654 (Walbaum), farms in Denmark in 2006. *J. Fish Dis.* 31, 659–67. doi:10.1111/j.1365-  
655 2761.2008.00941.x.

656 Price, M. N., Dehal, P. S., and Arkin, A. P. (2009). FastTree: computing large minimum evolution  
657 trees with profiles instead of a distance matrix. *Mol. Biol. Evol.* 26, 1641–50.  
658 doi:10.1093/molbev/msp077.

659 Rasch, M., Kastbjerg, V. G., Bruhn, J. B., Dalsgaard, I., Givskov, M., and Gram, L. (2007).  
660 Quorum sensing signals are produced by *Aeromonas salmonicida* and quorum sensing  
661 inhibitors can reduce production of a potential virulence factor. *Dis. Aquat. Organ.* 78, 105–  
662 113. doi:10.3354/dao01865.

663 Rasmussen, C. J. (1964). Furunkulose hos damørreder (Furunculosis in farmed trout) (in Danish).  
664 *Scand. Jouranl Vet. Sci.* 16, 463–472.

665 Reith, M. E., Singh, R. K., Curtis, B., Boyd, J. M., Bouevitch, A., Kimball, J., et al. (2008). The  
666 genome of *Aeromonas salmonicida* subsp. *salmonicida* A449: insights into the evolution of a  
667 fish pathogen. *BMC Genomics* 9.

668 Shao, Y., Harrison, E. M., Bi, D., Tai, C., He, X., Ou, H. Y., et al. (2011). TADB: A web-based  
669 resource for Type 2 toxin-antitoxin loci in bacteria and archaea. *Nucleic Acids Res.* 39.  
670 doi:10.1093/nar/gkq908.

671 Stuber, K., Burr, S. E., Braun, M., Wahli, T., and Frey, J. (2003). Type III Secretion Genes in  
672 *Aeromonas salmonicida* subsp. *salmonicida* Are Located on a Large Thermolabile Virulence  
673 Plasmid. *J. Clin. Microbiol.* 41, 3854–3856. doi:10.1128/JCM.41.8.3854-3856.2003.

674 Sørum, H., L'Abée-Lund, T. M., Solberg, A., and Wold, A. (2003). Integron-containing IncU R  
675 plasmids pRAS1 and pAr-32 from the fish pathogen *Aeromonas salmonicida*. *Antimicrob.*  
676 *Agents Chemother.* 47, 1285–1290. doi:10.1128/AAC.47.4.1285-1290.2003.

677 Tanaka, K. H., Dallaire-Dufresne, S., Daher, R. K., Frenette, M., and Charette, S. J. (2012). An  
678 insertion sequence-dependent plasmid rearrangement in *Aeromonas salmonicida* causes the  
679 loss of the type three secretion system. *PLoS One* 7, e33725.  
680 doi:10.1371/journal.pone.0033725.

681 Trudel, M. V., Tanaka, K. H., and Geneviève, F. (2013). Insertion sequence AS5 (ISAS5) is  
682 involved in the genomic plasticity of *Aeromonas salmonicida*. *Mob. Genet. Elem.* 3.

683 Vincent, A. T., Trudel, M. V., Freschi, L., Nagar, V., Gagné-Thivierge, C., Levesque, R. C., et al.  
684 (2016). Increasing genomic diversity and evidence of constrained lifestyle evolution due to



685 insertion sequences in *Aeromonas salmonicida*. *BMC Genomics* 17, 44. doi:10.1186/s12864-  
686 016-2381-3.

687 Wiklund, T., and Dalsgaard, I. (1998). Occurrence and significance of atypical *Aeromonas*  
688 *salmonicida* in non-salmonid and salmonid fish species: A review. *Dis. Aquat. Organ.* 32, 49–  
689 69. doi:10.3354/dao032049.

690 Zankari, E., Hasman, H., Cosentino, S., Vestergaard, M., Rasmussen, S., Lund, O., et al. (2012).  
691 Identification of acquired antimicrobial resistance genes. *J. Antimicrob. Chemother.* 67, 2640–  
692 4. doi:10.1093/jac/dks261.

693 Zerbino, D. R., and Birney, E. (2008). Velvet: algorithms for de novo short read assembly using de  
694 Bruijn graphs. *Genome Res.* 18, 821–9. doi:10.1101/gr.074492.107.

695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732

## 733 8. Tables

734

735

736

737

738

739

740

**Table 1. Overview of acquired antibiotic resistance genes among the 101 *A. salmonicida* sequenced isolates and the reference A449.** Threshold for presence of a resistance gene in an isolates was set to 75% similarity expressed as percent sequence identity (ID) and 60% of alignment length (coverage) of the resistance gene. Isolates are labeled according to region and year of isolation as in Figure 1.

<i>A. salmonicida</i> isolate	Beta-lactam			Trimethoprim	Sulphonamide	Aminoglycoside			Phenicol	Tetracycline
<b>Reference A449 (France 1975)</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>aadA1</i>	<i>cat</i>	<i>tet(E)</i>
<b>Sd5 2005</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>aadA1</i>	-	-
<b>Mj10 2009</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>aadA1</i>	-	-
<b>Mj10 2010</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>aadA1</i>	-	-
<b>Nj1 2011</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>strB</i>	<i>strA</i>	-
<b>Mj11 2014</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>aadA1</i>	-	-
<b>Mj3 2014</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>aadA1</i>	-	-
<b>Sj4 2014</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	<i>dfra14</i>	-	<i>sul2</i>	<i>strB</i>	-	<i>strA</i>	-
<b>Mj10 1982</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	<i>aadA2</i>	-	-	-
<b>Mj8 1997</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	<i>dfra1</i>	<i>sul1</i>	-	<i>aadA1</i>	-	-
<b>Nj2 2001</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj22 2002</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Nj4 2002</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj11 2003</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj3 2003</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd1 2004</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd4 2004</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj3 2004</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj7 2008</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj16 2008</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj4 2008</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj3 2008</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 (a) 2009</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj3 (a) 2009</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj3 (b) 2009</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 (b) 2009</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj3 2010</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 (a) 2010</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 (b) 2010</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 2011</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj9 2012</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj12 2014</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd6 (a) 2013</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd6 (b) 2013</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj14 2014</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd4 2014</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd10 2014</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj7 1980</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj24 1980</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj18 1981</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj7 1981</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 1981</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj16 1981</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd3 1982</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 1982</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd2 1982</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 1983</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj6 1983</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj15 1983</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sd2 1983</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj7 1984</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj15 1984</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Nj3 1984</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj16 1985</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj5 1986</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj18 1986</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj16 1986</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj11 1987</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj16 1987</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Mj13 1987</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj5 1988</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-
<b>Sj4 1988</b>	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-

Sd9 1988	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj11 1988	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj1 1989	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj3 1989	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj2 1990	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj3 1990	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj20 1990	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj5 1990	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Scotland	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj3 1991	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Nj5 1991	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj5 1991	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj18 1991	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj4 1992	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj1 1992	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sd8 1992	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj6 (a) 1993	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj6 (b) 1993	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj2 (a) 1993	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj2 (b) 1993	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj21 1993	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj19 1994	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj5 1994	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Nj3 1995	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj12 1995	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj1 1995	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj4 1995	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj5 1995	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sd2 1995	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj23 (a) 1996	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj23 (b) 1996	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj6 (a) 1996	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj6 (b) 1996	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sd7 1997	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj20 1997	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj4 1998	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj4 (a) 1999	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Mj17 1999	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
Sj4 (b) 1999	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-
NCIMB 1102 (Type strain 1962)	<i>blaFOX-2</i>	<i>ampS</i>	<i>blaCEPH-A3</i>	-	-	-	-	-	-	-	-	-	-

741  
742  
743  
744  
745  
746  
747

**Table 2. Overview of plasmid profiles among the 101 *A. salmonicida* sequenced isolates and the reference A449.** Plasmid profile number is displayed to the right, as well as the number of *A. salmonicida* that have the respective profile. Presence and absence of a plasmid for the given profile is presented with a plus (present) and minus (absent) sign respectively, below each plasmid name.

Profile No.	No. of <i>A. salmonicida</i> isolates	Plasmids											
		pAsa1	pAsa2	pAsa3	pAsa4	pAsa5	pAsa6	pAsal1	pAr_32	pRAS1	pRAS3.1	pRAS3.2	
1	45	+	+	+	-	+	+	+	-	-	-	-	-
2	1	+	-	+	-	+	+	+	-	-	-	-	-
3	2	+	+	+	-	-	+	+	-	-	-	-	-
4	4	+	+	+	-	-	-	+	-	-	-	-	-
5	1	+	+	+	-	+	-	+	-	-	-	-	-
6	3	+	+	+	-	-	-	-	-	-	-	-	-
7	1	+	+	+	-	-	+	-	-	-	-	-	-
8	4	+	+	+	-	+	-	-	-	-	-	-	-
9	2	+	+	-	-	+	-	-	-	-	-	-	-
10	22	+	+	-	-	+	+	-	-	-	-	-	-
11	16	+	+	+	-	+	+	-	-	-	-	-	-
12	1	+	+	+	+	+	-	-	-	-	-	-	-

748 **9. Figure legends**

749

750 **Figure 1. Phylogeny of *A. salmonicida*.** Bayesian temporal phylogenetic tree based on the  
751 alignment of 667 SNPs found among the 101 *A. salmonicida* sequenced isolates and the reference  
752 A449. The tree shows the most recent common ancestor of the *A. salmonicida* isolates dates back to  
753 ~1915 (95% HPD interval 1764 - 1947) and that there have been four main introductions of *A.*  
754 *salmonicida* in Denmark: ~1973 (95% HPD interval 1958 -1979), ~1973 (95% HPD interval 1964 -  
755 1981), ~1948 (95% HPD interval 1934 - 1964) and ~ 1946 (95% HPD interval 1939 - 1961). The  
756 four main clades are each shaded with a color SNP differences between major clades are shown  
757 above the estimated year of emergence. The three non-Danish *A. salmonicida* each have their own  
758 color and have the following labels: Scotland, NCIMB 1102 (type strain 1962), Reference A449  
759 (France 1975). The Danish isolates either have a black color (freshwater farms) or a blue color  
760 (seawater farms) and they are labeled by region of origin followed by year of isolation. Following  
761 abbreviations are used for regions in Denmark: Nj = Northern Jutland, Mj = Central Jutland, Sd =  
762 Southern Denmark, Sj = Zealand. A heatmap illustration with information regarding acquired  
763 ARGs, virulence and iron acquisition proteins and plasmid profile numbers of each *A. salmonicida*  
764 isolate is shown to the right of the tree. Presence and absence of protein sequences are illustrated by  
765 presence and absence of a red square, respectively. Plasmid profile number is shown and isolates  
766 that harbor ARGs against multiple antibiotics are labeled with “res”. Four minor clades marked  
767 with a red ring consist solely of isolates without plasmid pAsa3 and pAsa11.

768

769 **Figure 2. Heatmap illustrating presence and absence of 78 virulence associated and iron**  
770 **acquisition protein sequences found in the NCBI protein database among the 101 *A.***  
771 ***salmonicida* sequenced isolates and the reference A449.** Isolates are displayed on the right and  
772 sequence protein names on the bottom. Threshold limit for presence of protein in an isolate was set  
773 to 75% similarity, expressed as percent sequence identity (ID). Red color represents > 95% ID, pink  
774 color > 85% ID, dark blue > 75% ID and light blue > 65% ID.

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796 **10. Supplementary information appendix**

797

798 **Table S1. Full genomic data of the 101 sequenced *A. salmonicida* isolates.** Isolates are labeled as  
799 stated in Figure 1.

800

801 **Table S2. Overview of 78 virulence associated and iron acquisition protein sequences found in**  
802 **the NCBI protein database.** The following is shown in the table: name of protein sequence, short  
803 description, location of the protein coding gene, Genbank accession number and the number of *A.*  
804 *salmonicida* isolates harboring the given protein sequence. Isolates are labeled as stated in Figure 1.

805

806 **Table S3. Overview of 11 *A. salmonicida* plasmids found in the NCBI nucleotide database.** The  
807 following is shown in the table: name of plasmid, short description, antibiotic resistance genes  
808 encoded on plasmid, length of plasmid (in base pairs), and Genbank accession number. Isolates are  
809 labeled as stated in Figure 1.


810

811 **Table S4. Plasmid content of the 101 *A. salmonicida* sequenced isolates and the reference**  
812 **A449.** Numbers under each plasmid name represent percent coverage of that plasmid (in base pairs)  
813 for a given isolate. If percent coverage is higher than 60%, the plasmid is said to be present in the  
814 isolate and the color of the cell is green. Threshold limit for presence of plasmid in an isolate was  
815 set to 75% similarity expressed as percent sequence identity (ID) and 60% of alignment length  
816 (coverage) of the plasmid. Following abbreviations are used for percent coverage: 100 = 100%  
817 coverage, <100 = from 80% up to 99% coverage, <80 = from 60% up to 79% coverage, <60 = from  
818 10 up to 59% coverage, <10 = from 1% up to 9% coverage, 0 = 0% coverage. Isolates are labeled as  
819 stated in Figure 1.

820


821 **Figure S1. Phylogeny of *A. salmonicida*.** Maximum likelihood tree based on the alignment of 667  
822 SNPs found among the 101 *A. salmonicida* sequenced isolates and the reference A449. Isolates are  
823 labeled as stated in Figure 1.





**Figure 1. Phylogeny of *A. salmonicida*.** Bayesian temporal phylogenetic tree based on the alignment of 667 SNPs found among the 101 *A. salmonicida* sequenced isolates and the reference A449. The tree shows the most recent common ancestor of the *A. salmonicida* isolates dates back to ~1915 (95% HPD interval 1764 - 1947) and that there have been four main introductions of *A. salmonicida* in Denmark: ~1973 (95% HPD interval 1958 -1979), ~1973 (95% HPD interval 1964 - 1981), ~1948 (95% HPD interval 1934 - 1964) and ~ 1946 (95% HPD interval 1939 - 1961). The four main clades are each shaded with a color SNP differences between major clades are shown above the estimated year of emergence. The three non-Danish *A. salmonicida* each have their own color and have the following labels: Scotland, NCIMB 1102 (type strain 1962), Reference A449 (France 1975). The Danish isolates either have a black color (freshwater farms) or a blue color (seawater farms) and they are labeled by region of origin followed by year of isolation. Following abbreviations are used for regions in Denmark: Nj = Northern Jutland, Mj = Central Jutland, Sd = Southern Denmark, Sj = Zealand. A heatmap illustration with information regarding acquired ARGs, virulence and iron acquisition proteins and plasmid profile numbers of each *A. salmonicida* isolate is shown to the right of the tree. Presence and absence of protein sequences are illustrated by presence and absence of a red square, respectively. Plasmid profile number is shown and isolates that harbor ARGs against multiple antibiotics are labeled with “res”. Four minor clades marked with a red ring consist solely of isolates without plasmid pAsa3 and pAsa11.





**Figure 2. Heatmap illustrating presence and absence of 78 virulence associated and iron acquisition protein sequences found in the NCBI protein database among the 101 *A. salmonicida* sequenced isolates and the reference A449.** Isolates are displayed on the right and sequence protein names on the bottom. Threshold limit for presence of protein in an isolate was set to 75% similarity, expressed as percent sequence identity (ID). Red color represents > 95% ID, pink color > 85% ID, dark blue > 75% ID and light blue > 65% ID.

## Supplementary information appendix

**Table S1. Full genomic data of the 101 sequenced *A. salmonicida* isolates.** Isolates are labeled as stated in Figure 1.

Isolate	Coverage	N50	Contigs	Longest contig	Average contig coverage	Total bps in contigs	Pct. of expected size	Pct. reads in as	Insert Size	Country	Accession no.
Nj2 2001	103	99385	184	381279	304	4696711	94	99	489	Denmark	XXXXXXXXXX
Mj22 2002	99	110666	201	381769	264	4750938	95	98	493	Denmark	XXXXXXXXXX
Nj4 2002	128	105305	193	381512	calc. error	4786335	96	98	431	Denmark	XXXXXXXXXX
Mj11 2003	107	110599	194	381412	486	4697103	94	98	492	Denmark	XXXXXXXXXX
Sj3 2003	98	115031	207	381370	312	4758893	95	98	500	Denmark	XXXXXXXXXX
Sd1 2004	91	105305	199	381627	255	4753998	95	99	424	Denmark	XXXXXXXXXX
Sd4 2004	142	110691	213	381366	calc. error	4759628	95	99	469	Denmark	XXXXXXXXXX
Sj3 2004	101	102467	202	381541	300	4750816	95	98	484	Denmark	XXXXXXXXXX
Sd5 2005	86	110641	213	381106	238	4793498	96	99	483	Denmark	XXXXXXXXXX
Mj7 2008	55	102530	188	381640	193	4690018	94	98	513	Denmark	XXXXXXXXXX
Mj16 2008	83	110691	159	381687	211	4740131	95	99	469	Denmark	XXXXXXXXXX
Mj4 2008	108	89390	223	381675	calc. error	4747695	95	98	482	Denmark	XXXXXXXXXX
Sj3 2008	94	106301	197	381477	249	4751762	95	98	492	Denmark	XXXXXXXXXX
Sj4 (a) 2009	105	99384	200	381109	290	4750997	95	98	479	Denmark	XXXXXXXXXX
Sj3 (a) 2009	120	103045	202	381434	calc. error	4758324	95	99	458	Denmark	XXXXXXXXXX
Sj3 (b) 2009	98	102584	214	381231	288	4757155	95	98	461	Denmark	XXXXXXXXXX
Sj4 (b) 2009	99	102333	209	381317	313	4756966	95	99	461	Denmark	XXXXXXXXXX
Mj10 2009	55	99385	216	381398	164	4771942	95	99	449	Denmark	XXXXXXXXXX
Mj10 2010	88	102393	204	381260	242	4768899	95	98	456	Denmark	XXXXXXXXXX
Sj3 2010	87	102561	206	381380	calc. error	4758588	95	99	438	Denmark	XXXXXXXXXX
Sj4 (a) 2010	93	102544	217	381709	309	4757978	95	99	431	Denmark	XXXXXXXXXX
Sj4 (b) 2010	101	107109	203	381641	calc. error	4753148	95	99	439	Denmark	XXXXXXXXXX
Nj1 2011	150	110550	228	381259	calc. error	4932368	99	98	463	Denmark	XXXXXXXXXX
Sj4 2011	98	110673	206	381376	310	4760058	95	99	467	Denmark	XXXXXXXXXX
Mj9 2012	144	102679	198	381659	402	4753140	95	99	439	Denmark	XXXXXXXXXX
Mj12 2014	85	110681	166	381442	262	4676044	94	99	463	Denmark	XXXXXXXXXX
Sd6 (a) 2013	126	110711	203	381684	371	4758827	95	99	451	Denmark	XXXXXXXXXX
Sd6 (b) 2013	135	102582	209	381428	376	4761737	95	99	459	Denmark	XXXXXXXXXX
Mj14 2014	78	99257	209	381297	calc. error	4844616	97	97	513	Denmark	XXXXXXXXXX
Sd4 2014	147	102543	200	381302	457	4755925	95	98	438	Denmark	XXXXXXXXXX
Mj11 2014	112	102566	209	381692	361	4772818	95	98	427	Denmark	XXXXXXXXXX
Sd10 2014	117	106303	199	381207	314	4750421	95	99	457	Denmark	XXXXXXXXXX
Mj3 2014	149	107109	204	381362	411	4769894	95	98	463	Denmark	XXXXXXXXXX
Sj4 2014	81	110654	212	381386	489	4757435	95	98	513	Denmark	XXXXXXXXXX
Sj7 1980	29	114971	144	381478	99	4665337	93	98	533	Denmark	XXXXXXXXXX

Mj24 1980	36	115031	144	381653	177	4641600	93	98	475	Denmark	XXXXXXXXXX
Mj18 1981	84	110681	166	381396	297	4680471	94	98	531	Denmark	XXXXXXXXXX
Sj7 1981	130	102554	191	381632	401	4700129	94	99	476	Denmark	XXXXXXXXXX
Sj4 1981	41	115042	163	381518	194	4661527	93	98	421	Denmark	XXXXXXXXXX
Mj16 1981	28	96255	170	381112	82	4698702	94	99	448	Denmark	XXXXXXXXXX
Sd3 1982	31	110578	187	381371	94	4703018	94	99	488	Denmark	XXXXXXXXXX
Sj4 1982	44	110479	158	381370	129	4669963	93	99	495	Denmark	XXXXXXXXXX
Sd2 1982	61	110574	181	381663	187	4655791	93	99	460	Denmark	XXXXXXXXXX
Mj10 1982	35	110592	179	381556	147	4717685	94	98	510	Denmark	XXXXXXXXXX
Sj4 1983	47	110643	189	381565	185	4745156	95	98	444	Denmark	XXXXXXXXXX
Mj6 1983	59	110685	171	381612	calc. error	4678164	94	99	422	Denmark	XXXXXXXXXX
Mj15 1983	52	110673	162	381503	158	4668980	93	99	486	Denmark	XXXXXXXXXX
Sd2 1983	39	102684	180	381349	126	4740974	95	99	470	Denmark	XXXXXXXXXX
Sj7 1984	37	102725	189	381428	133	4687108	94	98	433	Denmark	XXXXXXXXXX
Mj15 1984	43	110650	160	381326	132	4669787	93	99	459	Denmark	XXXXXXXXXX
Nj3 1984	59	102317	191	381545	208	4702292	94	98	465	Denmark	XXXXXXXXXX
Mj16 1985	92	102683	186	381167	352	4693921	94	99	449	Denmark	XXXXXXXXXX
Mj5 1986	51	110600	205	381324	154	4723248	94	99	483	Denmark	XXXXXXXXXX
Mj18 1986	54	106303	184	381326	201	4704747	94	98	451	Denmark	XXXXXXXXXX
Mj16 1986	32	99374	208	381361	93	4760365	95	98	478	Denmark	XXXXXXXXXX
Mj11 1987	58	102503	183	381392	167	4686542	94	99	426	Denmark	XXXXXXXXXX
Mj16 1987	56	110581	197	381546	170	4723289	94	99	453	Denmark	XXXXXXXXXX
Mj13 1987	45	110665	204	381556	121	4759024	95	99	488	Denmark	XXXXXXXXXX
Sj5 1988	85	115073	162	381442	248	4648311	93	99	512	Denmark	XXXXXXXXXX
Sj4 1988	84	115001	191	381621	336	4698058	94	98	506	Denmark	XXXXXXXXXX
Sd9 1988	77	102439	236	381519	calc. error	4778057	96	98	478	Denmark	XXXXXXXXXX
Mj11 1988	112	110588	180	381329	344	4712958	94	99	435	Denmark	XXXXXXXXXX
Sj1 1989	97	99385	193	381207	403	4696354	94	98	451	Denmark	XXXXXXXXXX
Sj3 1989	82	118192	147	381415	298	4709626	94	98	497	Denmark	XXXXXXXXXX
Mj2 1990	105	110638	138	381483	calc. error	4712403	94	99	451	Denmark	XXXXXXXXXX
Sj3 1990	123	115044	153	381353	392	4708363	94	99	450	Denmark	XXXXXXXXXX
Mj20 1990	97	99384	206	381475	396	4745926	95	98	488	Denmark	XXXXXXXXXX
Sj5 1990	147	110649	180	381458	487	4735891	95	99	376	Denmark	XXXXXXXXXX
Scotland	68	99386	161	381321	234	4672419	93	99	438	Scotland	XXXXXXXXXX
Sj3 1991	144	110534	187	381129	461	4686788	94	99	407	Denmark	XXXXXXXXXX
Nj5 1991	145	110468	179	381427	450	4744445	95	99	394	Denmark	XXXXXXXXXX
Sj5 1991	110	115054	150	381276	368	4708174	94	99	471	Denmark	XXXXXXXXXX
Mj18 1991	89	110645	170	381628	299	4700329	94	99	453	Denmark	XXXXXXXXXX
Sj4 1992	109	90171	192	381434	calc. error	4744289	95	99	472	Denmark	XXXXXXXXXX
Sj1 1992	78	111005	171	381498	248	4700594	94	99	491	Denmark	XXXXXXXXXX

<b>Sd8 1992</b>	84	102569	187	381652	250	4693600	94	99	523	Denmark	XXXXXXXXXX
<b>Sj6 (a) 1993</b>	104	110683	184	381352	503	4711783	94	98	481	Denmark	XXXXXXXXXX
<b>Sj6 (b) 1993</b>	81	102839	210	381188	264	4756685	95	99	420	Denmark	XXXXXXXXXX
<b>Sj2 (a) 1993</b>	173	102788	201	381562	564	4757469	95	99	367	Denmark	XXXXXXXXXX
<b>Sj2 (b) 1993</b>	119	102537	209	381325	348	4756068	95	99	443	Denmark	XXXXXXXXXX
<b>Mj21 1993</b>	107	105305	200	381727	323	4760383	95	99	478	Denmark	XXXXXXXXXX
<b>Mj19 1994</b>	77	105305	187	381317	206	4693576	94	99	482	Denmark	XXXXXXXXXX
<b>Sj5 1994</b>	49	99385	207	381506	146	4757079	95	98	521	Denmark	XXXXXXXXXX
<b>Nj3 1995</b>	90	102828	202	381519	263	4758748	95	98	465	Denmark	XXXXXXXXXX
<b>Mj12 1995</b>	133	110503	167	381263	431	4737917	95	99	408	Denmark	XXXXXXXXXX
<b>Mj1 1995</b>	126	110655	170	381310	calc. error	4716005	94	99	373	Denmark	XXXXXXXXXX
<b>Mj4 1995</b>	117	102745	197	381303	423	4746620	95	98	431	Denmark	XXXXXXXXXX
<b>Sj5 1995</b>	89	102519	200	381605	274	4758268	95	98	472	Denmark	XXXXXXXXXX
<b>Sd2 1995</b>	129	102730	205	381625	395	4754891	95	99	430	Denmark	XXXXXXXXXX
<b>Mj23 (a) 1996</b>	121	102588	175	381681	388	4737728	95	98	466	Denmark	XXXXXXXXXX
<b>Mj23 (b) 1996</b>	124	110588	166	381327	388	4688488	94	99	508	Denmark	XXXXXXXXXX
<b>Sj6 (a) 1996</b>	62	110688	202	381738	calc. error	4755441	95	98	560	Denmark	XXXXXXXXXX
<b>Sj6 (b) 1996</b>	101	105305	197	381146	calc. error	4697453	94	99	430	Denmark	XXXXXXXXXX
<b>Mj8 1997</b>	164	99757	208	381552	598	4779438	96	99	404	Denmark	XXXXXXXXXX
<b>Sd7 1997</b>	59	102690	182	381336	164	4741454	95	99	503	Denmark	XXXXXXXXXX
<b>Mj20 1997</b>	108	102547	205	381430	calc. error	4749870	95	99	374	Denmark	XXXXXXXXXX
<b>Sj4 1998</b>	77	110635	187	381495	387	4697323	94	98	480	Denmark	XXXXXXXXXX
<b>Sj4 (a) 1999</b>	72	110581	160	484118	193	4735436	95	99	485	Denmark	XXXXXXXXXX
<b>Mj17 1999</b>	93	110527	186	381251	calc. error	4697920	94	98	516	Denmark	XXXXXXXXXX
<b>Sj4 (b) 1999</b>	86	99385	197	381581	228	4707087	94	99	496	Denmark	XXXXXXXXXX
<b>NCIMB 1102 (Type strain 1962)</b>	91	103414	195	381570	552	4713735	94	97	482	England	XXXXXXXXXX

**Table S2. Overview of 78 virulence associated and iron acquisition protein sequences found in the NCBI protein database.** The following is shown in the table: name of protein sequence, short description, location of the protein coding gene, Genbank accession number and the number of *A. salmonicida* isolates harboring the given protein sequence. Isolates are labeled as stated in Figure 1.

Protein	Short description of protein function	Gene location	Genbank accession no.	Present in no. of isolates
AopP	T3SS effector protein	pAsa1	YP_009062872.1	51
AscV	T3SS inner membrane export apparatus	pAsa5	EHI50383.1	77
AscC	T3SS protein of the outer membrane ring	pAsa5	ABD48952.1	77
AscG	T3SS chaperone	pAsa5	ABO92527.1	77
AscP	T3SS needle length control; ruler protein; regulation of secretion; substrate specificity switch	pAsa5	EHI50374.1	77
AscH	T3SS regulator needle assembling	pAsa5	EHI50401.1	77
AscF	T3SS early substrate, needle subunit	pAsa5	EHI50399.1	77
AscX	T3SS translocator needle subunit chaperoned by AscY	pAsa5	ABO92546.1	77
AcrH	T3SS chaperone for AopB/AopD	pAsa5	EHI50387.1	77
Acr2	T3SS chaperone	pAsa5	ABO92547.1	77
AcrV	T3SS middle substrate; tip translocon; hydrophilic translocators; protective antigen; anti-host factor	pAsa5	ABO92541.1	77
AopB	T3SS translocon; hydrophobic translocators; pore in host cell	pAsa5	EHI50388.1	77
AopD	T3SS translocon; hydrophobic translocators; pore in host cell	pAsa5	EHI50389.1	77
AopN	T3SS secretion control of translocators and immune suppressor	pAsa5	EHI50378.1	77
Ati2	T3SS inositol polyphosphate 5-phosphatase	pAsa5	ABO92519.1	77
Ati1	T3SS translocator needle subunit (Ati2 chaperone)	pAsa5	ABO92520.1	77
AopH	T3SS putative tyrosine phosphatase	pAsa5	ABD48950.1	93
SycH	T3SS AopH chaperone	pAsa5	ABO92484.1	93
AopO	T3SS putative serine/threonine kinase	pAsa5	ABD48951.1	99
AopO chaperone	T3SS AopO chaperone	pAsa5	ABO92569.1	99
VapA (A-layer protein)	Tetragonal virulence array protein; coats surface of bacteria; confers virulence to bacteria	Chromosome	EHI51039.1	101
Extracellular nuclease	Secreted enzyme; cell communication	Chromosome	ABO90268.1	101
AexT	T3SS secreted toxin; ADP-ribosyltransferase	Chromosome	ABD48949.1	102
FlpC	Type IV pilus secretin	Chromosome	ABD57354.1	102
IcmF	T6SS protein	Chromosome	KFN20003.1	102
VgrG	T6SS effector protein	pAsa4 (homolog on chromosome)	KIX26565.1	102
S-protein	T2SS S-protein secretion component E	Chromosome	ABO89526.1	102
Aerolysin	Aerolysin toxin	Chromosome	ABO91859.1	102
Aerolysin B	Aerolysin B toxin	Chromosome	ABO90867.1	102
EF-G	Elongation factor G	Chromosome	A4SHV8.2	102

EF-Tu	Elongation factor Tu	Chromosome	A4SHU2.1	102
DnaK	Molecular chaperone	Chromosome	EHI53225.1	102
HtpG	Heat shock protein; ATPase activity	Chromosome	ABO89902.1	102
PNPase	mRNA degradation; 3'-5'-exoribonuclease activity; magnesium ion binding	Chromosome	A4SJR9.1	102
Aminopeptidase N	Aminopeptidase activity; metallopeptidase activity; zinc ion binding	Chromosome	EHI52287.1	102
Methionine gamma-lyase	Lyase activity; pyridoxal phosphate binding	Chromosome	ABO90396.1	102
Phosphate acetyltransferase	Phosphate acetyltransferase activity; acetyl-CoA biosynthetic process	Chromosome	ABO91377.1	102
Extracellular lipase	Triglyceride lipase activity	Chromosome	EHI50452.1	102
TagA	ToxR-regulated lipoprotein; metalloendopeptidase activity	Chromosome	EHI53531.1	102
Ahe2	Serine-type endopeptidase activity	Chromosome	EHI54143.1	102
N-acetylglucosamine	N-acetylglucosamine-binding protein A; viral capsid	Chromosome	KHF01495.1	102
Enolase	Magnesium ion binding; phosphopyruvate hydratase activity; glycolytic process	Chromosome	EHI51131.1	102
Outer membrane	Outer membrane protein	Chromosome	ABO89374.1	102
OmpK40	Outer membrane porin II	Chromosome	EHI50654.1	102
DegQ	Serine-type endopeptidase activity	Chromosome	EHI52649.1	102
Hemolysin	Hemolysis in other organism; toxin	Chromosome	EHI53183.1	102
Alpha-amylase extracellular	Alpha-amylase activity; cation binding; carbohydrate metabolic process	Chromosome	ABO89392.1	102
Alpha-amylase	Alpha-amylase activity; cation binding; carbohydrate metabolic process	Chromosome	ABO91424.1	102
Secreted metalloprotease	Metallopeptidase activity	Chromosome	EHI52510.1	102
collagenase	Serine-type endopeptidase activity; zinc ion binding	Chromosome	EHI52177.1	102
GCAT	Lipase activity; transferase activity, transferring acyl groups	Chromosome	ABO88676.1	102
Chitinase	Carbohydrate binding ; chitinase activity	Chromosome	ABO90207.1	102
Endochitinase	Carbohydrate binding ; chitinase activity	Chromosome	ABO91302.1	102
Pullulanase	Carbohydrate binding; pullulanase activity	Chromosome	EHI52980.1	102
RTX	Calcium ion binding	Chromosome	ABO88975.1	102
Cytolytic delta-endotoxin	Insecticidal protein; pathogenesis	Chromosome	EHI53783.1	102
Patatin family phospholipase	Lipid metabolic process	Chromosome	EHI53765.1	102
Phospholipase	phospholipase C precursor	Chromosome	EHI52973.1	102
ABC-type ferric transporter	ABC-type ferric siderophore transporter; periplasmic binding protein	Chromosome	ABO92284.1	102
Ferric siderophore receptor B	Iron ion binding; receptor activity; siderophore transport	Chromosome	ABO92283.1	102
Hydroxamate-type ferric transporter	ABC-type hydroxamate-type ferric siderophore transporter; periplasmic binding protein	Chromosome	ABO92281.1	102
Outer membrane ferric receptor	Iron ion binding; receptor activity; siderophore transport	Chromosome	ABO91837.1	102
ABC-type ferric permease	Transporter activity	Chromosome	ABO92286.1	102
Outer-membrane heme receptor	Receptor activity; transporter activity	Chromosome	ABO91308.1	102
Hemin receptor	Receptor activity; transporter activity	Chromosome	ABO90714.1	102

Colicin receptor	Receptor activity; transport	Chromosome	ABO89926.1	102
N-acyl homoserine lactone synthase	N-acyl homoserine lactone synthase activity, quorum sensing	Chromosome	EHI52135.1	102
LuxS	Iron ion binding; S-ribosylhomocysteine lyase activity; quorum sensing	Chromosome	A4SIY8.1	102
AsaR	Transcriptional activator; DNA binding	Chromosome	ABO91724.1	102
LuxU	Phosphorelay protein; signal transducer activity	Chromosome	ABO90799.1	102
LuxO	Phosphorelay signal transduction system; ATP binding; sequence-specific DNA binding	Chromosome	ABO91277.1	102
CpaB	Flp pilus assembly protein	Chromosome	EHI53195.1	102
FlgE	flagellar hook protein	Chromosome	EHI52692.1	102
Oligopeptidase A	Metal ion binding; metalloendopeptidase activity	Chromosome	ABO92225.1	102
M13 family peptidase	Metalloendopeptidase activity	Chromosome	EHI53386.1	102
TRAP	Transporter solute receptor; TAXI family	Chromosome	ABO91932.1	102
LasA	Metalloendopeptidase activity	Chromosome	ABO89393.1	102
Immune inhibitor A	Metalloendopeptidase activity	Chromosome	ABO88993.1	102



**Table S3. Overview of 11 *A. salmonicida* plasmids found in the NCBI nucleotide database.** The following is shown in the table: name of plasmid, short description, antibiotic resistance genes encoded on plasmid, length of plasmid in base pairs (bp), and Genbank accession number. Isolates are labeled as stated in Figure 1.

<b>Plasmid</b>	<b>Short description</b>	<b>Antibiotic resistance genes</b>	<b>Length (bp)</b>	<b>Genbank accession no.</b>
<b>pAsa1</b>	Small cryptic plasmid		5424	NC_004923.1
<b>pAsa2</b>	Small cryptic plasmid		5247	NC_004925.1
<b>pAsa3</b>	Small cryptic plasmid		5616	NC_004924.1
<b>pAsa4</b>	Large plasmid encoding three T6SS proteins	<i>aadA1, cat, sul1, tet(E)</i>	166749	NC_009349.1
<b>pAsa5</b>	Large plasmid encoding the majority of T3SS proteins		155098	NC_009350.1
<b>pAsa6</b>	Medium plasmid encoding homologs of the T3SS Effector protein AopH and AopH chaperone		18536	NC_009352.2
<b>pAsa11</b>	Small plasmid encoding the T3SS Effector protein AopP		6371	NC_004338.1
<b>pAr_32</b>	Sequence from the pAr-32 resistance-determining region	<i>aadA2, catA2, sul1</i>	9340	AJ517791.1
<b>pRAS1</b>	Sequence from the pRAS1 resistance-determining region	<i>sul1, tet(A), dfrA16</i>	11663	AJ517790.2
<b>pRAS3.1</b>	Non-conjugative Tet C plasmid	<i>tet(C)</i>	11851	NC_003123.1
<b>pRAS3.2</b>	Non-conjugative Tet C plasmid (variant of pRAS3.1)	<i>tet(C)</i>	11823	NC_003124.1


**Table S4. Plasmid content of the 101 *A. salmonicida* sequenced isolates and the reference A449.** Numbers under each plasmid name represent percent coverage of that plasmid (in base pairs) for a given isolate. If percent coverage is higher than 60%, the plasmid is said to be present in the isolate and the color of the cell is green. Threshold limit for presence of plasmid in an isolate was set to 75% similarity expressed as percent sequence identity (ID) and 60% of alignment length (coverage) of the plasmid. Following abbreviations are used for percent coverage: 100 = 100% coverage, <100 = from 80% up to 99% coverage, <80 = from 60% up to 79% coverage, <60 = from 10 up to 59% coverage, <10 = from 1% up to 9% coverage, 0 = 0% coverage. Isolates are labeled as stated in Figure 1.

<i>A. salmonicida</i> isolate	pAsa1	pAsa2	pAsa3	pAsa4	pAsa5	pAsa6	pAsaI1	pAr-32	pRAS1	pRAS3.1	pRAS3.2	Plasmid profile no.
Sd4 2004	100	< 80	100	< 10	< 100	< 80	100	0	0	0	0	1
Mj7 2008	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Nj1 2011	100	< 100	< 100	< 60	< 100	< 80	100	0	0	0	0	1
Sd6 (a) 2013	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sd6 (b) 2013	100	100	< 100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sd4 2014	100	< 80	< 100	< 10	< 100	< 80	100	0	0	0	0	1
Mj16 1981	100	< 100	< 100	< 10	< 80	< 80	100	0	< 10	0	0	1
Nj3 1984	100	< 100	100	< 10	< 80	< 80	100	0	< 10	0	0	1
Mj16 1987	100	< 100	< 100	< 10	< 80	< 80	100	0	0	0	0	1
Sd9 1988	100	< 100	100	< 10	100	< 80	100	0	0	0	0	1
Nj3 1995	100	< 80	< 100	< 10	< 100	< 80	100	0	< 10	0	0	1
Mj1 1995	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Mj4 1995	100	< 100	100	< 10	< 100	< 80	100	0	0	0	0	1
Mj23 (a) 1996	100	< 100	< 100	< 10	< 100	< 80	100	0	< 10	0	0	1
Mj23 (b) 1996	100	< 80	< 100	< 10	< 100	< 80	100	0	0	0	0	1
Mj17 1999	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Mj11 2003	100	< 100	100	0	< 100	< 80	100	0	< 10	0	0	1
Sj3 2003	100	< 100	100	< 10	< 100	< 80	100	0	0	0	0	1
Sj3 (a) 2009	100	< 100	< 100	< 10	< 100	< 80	100	0	0	0	0	1
Sj3 (b) 2009	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sj4 (b) 2009	100	100	< 100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sj3 2010	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sj4 (a) 2010	100	< 80	100	< 10	< 100	< 80	100	0	0	0	0	1
Sj4 2011	100	100	< 100	< 10	< 100	< 80	100	0	< 10	0	0	1
Mj14 2014	100	< 100	100	< 60	< 100	< 80	100	0	0	0	0	1

Sj4 2014	100	< 80	100	< 10	< 100	< 80	100	0	0	0	0	1
Sj4 1981	100	< 100	100	< 10	< 80	< 80	100	0	0	0	0	1
Sj4 1983	100	< 80	100	< 10	< 100	< 80	100	0	0	0	0	1
Sj7 1984	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Mj5 1986	100	< 100	100	< 10	< 80	< 80	100	0	0	0	0	1
Mj18 1986	100	< 100	100	< 10	< 80	< 80	100	0	0	0	0	1
Sj4 1988	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sj1 1989	100	< 100	100	< 10	< 100	< 80	100	0	0	0	0	1
Mj20 1990	100	< 100	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sj6 (a) 1993	100	< 100	100	< 10	< 80	< 80	100	0	0	0	0	1
Sj6 (b) 1993	100	< 80	< 100	< 10	< 100	< 80	100	0	0	0	0	1
Sj2 (a) 1993	100	< 100	< 100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sj2 (b) 1993	100	100	< 80	< 10	< 100	< 80	< 100	0	< 10	0	0	1
Mj21 1993	100	< 100	100	< 10	< 100	< 80	100	0	0	0	0	1
Sj5 1994	100	100	< 100	< 10	< 100	< 80	100	0	< 10	0	0	1
Sj5 1995	100	< 80	< 100	< 10	< 100	< 80	100	0	0	0	0	1
Sj6 (b) 1996	100	< 80	100	< 10	< 100	< 80	100	0	< 10	0	0	1
Mj8 1997	100	< 100	100	< 10	< 100	< 80	100	0	< 10	< 60	< 60	1
Sj4 1998	100	< 100	100	< 10	< 100	< 80	100	0	0	0	0	1
NCIMB 1102 (Type strain 1962)	100	< 100	100	< 10	< 80	< 80	100	0	0	0	0	1
Mj4 2008	100	< 10	100	< 10	< 100	< 80	< 100	0	0	0	0	2
Sj7 1981	100	< 100	< 100	< 10	< 60	< 80	100	0	0	0	0	3
Scotland	100	< 100	< 100	< 10	< 60	< 80	100	0	0	0	0	3
Mj24 1980	100	< 80	100	< 10	< 60	< 60	100	0	0	0	0	4
Mj10 1982	100	< 100	100	< 10	< 60	< 60	100	< 60	< 60	0	0	4
Mj6 1983	100	< 100	100	< 10	< 60	< 60	100	0	0	0	0	4
Mj18 1981	100	< 80	100	< 10	< 60	< 60	100	0	0	0	0	4
Mj16 1986	100	< 80	< 100	< 10	< 100	< 60	100	0	0	0	0	5
Mj15 1983	100	< 100	100	< 10	< 60	< 60	< 60	0	0	0	0	6
Mj15 1984	100	< 100	100	< 10	< 60	< 60	< 60	0	0	0	0	6
Sj4 1982	100	< 80	100	< 10	< 60	< 60	< 60	0	0	0	0	6
Mj12 2014	100	< 80	100	< 10	< 60	< 80	< 60	0	< 10	0	0	7
Sj4 (a) 2009	< 100	100	< 80	< 10	< 100	< 60	< 60	0	0	0	0	8
Sj7 1980	100	< 100	100	< 10	< 80	< 60	< 60	0	0	0	0	8

Sj4 1992	100	< 100	100	< 10	< 100	< 60	< 60	0	0	0	0	8
Sj1 1992	100	< 80	100	< 10	< 80	< 60	< 60	0	0	0	0	8
Sj4 (b) 2010	< 100	< 80	< 60	< 10	< 100	< 60	< 60	0	< 10	0	0	9
Sj4 (a) 1999	< 100	< 100	0	< 10	< 80	< 60	< 60	0	0	0	0	9
Mj22 2002	< 100	< 100	< 60	< 10	< 100	< 80	< 60	0	0	0	0	10
Nj4 2002	< 100	< 100	< 60	< 10	100	< 80	< 60	0	< 10	0	0	10
Sd5 2005	< 100	< 80	< 10	< 10	< 100	< 80	< 60	< 60	< 60	0	0	10
Mj16 2008	< 100	< 100	0	< 10	100	< 100	< 60	0	0	0	0	10
Mj10 2009	< 100	< 100	< 10	< 10	< 100	< 80	< 60	< 10	0	< 60	< 60	10
Mj10 2010	< 100	100	0	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Mj9 2012	< 80	< 100	0	< 10	< 80	< 80	< 60	0	0	0	0	10
Sd10 2014	< 100	100	< 10	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Mj13 1987	< 100	< 100	< 10	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Mj2 1990	< 100	< 80	0	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Mj12 1995	< 80	< 100	< 60	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Sd1 2004	< 80	< 100	0	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Sj3 2004	< 100	< 100	< 60	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Sj3 2008	< 80	100	< 60	< 10	< 100	< 80	< 60	0	0	0	0	10
Mj11 2014	< 80	< 100	0	< 10	< 100	< 80	< 60	0	0	6	6	10
Mj3 2014	< 100	< 100	< 10	< 10	< 100	< 80	< 60	< 60	< 60	0	0	10
Sj5 1988	< 100	< 80	< 60	< 10	< 80	< 80	< 60	0	0	0	0	10
Mj18 1991	< 100	< 80	< 60	< 10	< 80	< 80	< 60	0	0	0	0	10
Sd2 1995	< 100	< 80	0	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Sj6 (a) 1996	< 100	< 100	< 60	< 10	< 100	< 80	< 60	0	0	0	0	10
Mj20 1997	< 80	< 100	< 60	< 10	< 100	< 80	< 60	0	< 10	0	0	10
Sj4 (b) 1999	< 100	< 100	0	< 10	< 100	< 80	< 60	0	0	0	0	10
Nj2 2001	100	< 100	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Sd3 1982	100	< 80	100	< 10	< 80	< 80	< 60	0	< 10	0	0	11
Mj16 1985	100	< 100	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Nj5 1991	100	< 100	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Sd8 1992	100	< 100	100	< 10	< 100	< 80	< 60	0	0	0	0	11
Mj19 1994	100	< 100	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Sd7 1997	100	< 100	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Sd2 1982	100	< 80	100	< 10	< 80	< 80	< 60	0	0	0	0	11

Sd2 1983	100	< 100	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Mj11 1987	100	< 80	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Mj11 1988	100	< 100	100	< 10	< 80	< 80	< 60	0	0	0	0	11
Sj3 1989	100	< 80	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Sj3 1990	100	< 80	100	< 10	< 100	< 80	< 60	0	< 10	0	0	11
Sj5 1990	100	< 100	100	< 10	< 100	< 80	< 60	0	0	0	0	11
Sj3 1991	100	< 100	100	< 10	< 100	< 80	< 60	0	0	0	0	11
Sj5 1991	100	< 100	100	< 10	< 100	< 80	< 60	0	0	0	0	11
Reference A449 (France 1975)	< 100	< 100	100	100	< 100	0	0	0	0	0	0	12
TOTAL	102	101	78	1	92	87	53	0	0	0	0	12



**Figure S1. Phylogeny of *A. salmonicida*.** Maximum likelihood tree based on the alignment of 667 SNPs found among the 101 *A. salmonicida* sequenced isolates and the reference A449. Isolates are labeled as stated in Figure 1.