Technical University of Denmark



Modeling of transport phenomena in battery cathode materials

García Lastra, Juan Maria; Mekonnen, Yedilfana Setarge; Loftager, Simon; Melander, Marko; Mathiesen, Nikolai Rask ; Christensen, Rune; Vegge, Tejs

Publication date: 2016

Document Version Peer reviewed version

Link back to DTU Orbit

Citation (APA):

García Lastra, J. M., Mekonnen, Y. S., Loftager, S., Melander, M., Mathiesen, N. R., Christensen, R., & Vegge, T. (2016). Modeling of transport phenomena in battery cathode materials. Abstract from 67th Annual Meeting of the International Society of Electrochemistry, The Hague, Netherlands.

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Modeling of transport phenomena in battery cathode materials

Juan Maria García Lastra, Yedilfana Setarge Mekonnen, Simon Loftager, Marko Melander, Nikolai Rask

Mathiesen, Rune Christensen, and Tejs Vegge

Department of Energy Conversion and Storage, Technical University of Denmark.

Fysikvej 309, 2800 Kgs. Lyngby, Denmark

jmgla@dtu.dk

Transport phenomena in the cathode materials of batteries, both ionic and electronic, play a key role in the overall performance of the device. An intense modeling activity of different aspects of these transport phenomena has taken place during the last two decades, which covers different time and length scales. In this talk we will focus on modeling at the atomic scale of ionic and electronic transport in the cathode materials of lithium-ion batteries and aprotic metal-air batteries.

The ionic transport of intercalated lithium in different transition metal oxides has been extensively modeled in the literature by combining density functional theory (DFT) and the nudged elastic band (NEB) method to obtain the activation barriers for lithium diffusion within these materials.¹ Lately the method is starting to be applied also to the ionic transport between coating materials (e.g. graphite) and the cathode and also at the interfaces that are created between different phases of the same cathode material². During the talk we will present some examples of lithium transport through these interfaces.

The electronic transport in lithium-ion batteries typically occurs through a polaronic hopping mechanism, which is conventionally modeled using DFT and NEB as in the case of the ionic transport. This way of modeling the electronic transport is valid in the purely adiabatic limit, omitting all the non-adiabatic features of the charge transfer. Here we will briefly introduce our recent developments in the implementation of a DFT-based Marcus theory model which is able to capture the non-adiabatic effects.

In contrast with lithium-ion batteries a conversion process takes place in metal-air batteries. In this conversion process an oxide is formed at the cathode through the reaction between atmospheric oxygen and alkali or alkali-earth ions. We will talk about modelling the formation of these oxides and its relation with the overpotentials observed in these batteries³.

Finally we will address the electronic transport at metal-air battery cathodes, which can take place through polaronic hopping⁴ as in the case of lithium-ion batteries, but also can occur through electron tunneling⁵. We will discuss which of the two processes is dominant depending on the different conditions of the charge/discharge process.

(1) G. K. P. Dathar, D. Sheppard, K. J. Stevenson and G. Henkelman, , *Chem. Mater.* 23, 4032-4037 (2011).

(2) W.T. Weng, T. Ohno, J.Phys. Chem. C. 117, 276 (2012)

(3) J. S. Hummelshoj, J. Blomqvist, S. Datta, T. Vegge, J. Rossmeisl, K. S. Thygesen, A. C. Luntz, K. W. Jacobsen and J. K. Norskov, *J. Chem. Phys* **132**, 071101 (2010).

(4) J. M. García-Lastra, J.S.G. Myrdal, R. Christensen, K.S. Thygesen and T. Vegge, *J.Phys. Chem. C.*, **117**, 5568 (2013)

(5) V. Viswanathan, K. S. Thygesen, J. S. Hummelshoj, J. K. Norskov, G. Girishkumar, B. D. McCloskey and A. C. Luntz, *J. Chem. Phys.* **135**, 214704 (2011).