

Technical University of Denmark



## Genome Sequence of *Talaromyces atroseus*, Which Produces Red Colorants for the Food Industry

**Thrane, Ulf; Rasmussen, Kasper Bøwig; Petersen, Bent; Rasmussen, Simon; Sicheritz-Pontén, Thomas; Mortensen, Uffe Hasbro**

*Published in:*  
Genome Announcements

*Link to article, DOI:*  
[10.1128/genomeA.01736-16](https://doi.org/10.1128/genomeA.01736-16)

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Thrane, U., Rasmussen, K. B., Petersen, B., Rasmussen, S., Sicheritz-Pontén, T., & Mortensen, U. H. (2017). Genome Sequence of *Talaromyces atroseus*, Which Produces Red Colorants for the Food Industry. *Genome Announcements*, 5(9), [e01736-16]. DOI: 10.1128/genomeA.01736-16

## DTU Library

Technical Information Center of Denmark


---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Genome Sequence of *Talaromyces atrovireus*, Which Produces Red Colorants for the Food Industry

Ulf Thrane,<sup>a</sup> Kasper Bøwig Rasmussen,<sup>a\*</sup> Bent Petersen,<sup>b</sup> Simon Rasmussen,<sup>b</sup> Thomas Sicheritz-Pontén,<sup>b</sup> Uffe Hasbro Mortensen<sup>a</sup>

Department of Biotechnology and Biomedicine, Technical University of Denmark, Kongens Lyngby, Denmark<sup>a</sup>;  
Department of Bio and Health Informatics, Technical University of Denmark, Kongens Lyngby, Denmark<sup>b</sup>

**ABSTRACT** *Talaromyces atrovireus* is a known producer of *Monascus* colorants suitable for the food industry. Furthermore, genetic tools have been established that facilitate elucidation and engineering of its biosynthetic pathways. Here, we report the draft genome of a potential fungal cell factory, *T. atrovireus* IBT 11181 (CBS 123796).

The genus *Talaromyces* primarily contains saprophytic fungi and encompasses medically and industrially relevant species such as the opportunistic human pathogen *T. marneffei* (formerly *Penicillium marneffei*), species with high production of cellulolytic enzymes, i.e., *T. cellulolyticus* (1), as well as the interesting pigment-producing species *T. atrovireus* (2). Several strains of *T. atrovireus* and closely related species are recognized as potential cell factories for *Monascus* pigment production, as they may serve as mycotoxin-free alternatives to *Monascus* spp. (2–4).

*T. atrovireus* IBT 11181 was originally isolated from red sweet bell pepper bought in a Danish supermarket and is deposited in the CBS collection at CBS-KNAW, Utrecht, the Netherlands, as CBS 123796 and CBS 238.95. We intend to implement this isolate as a model for *T. atrovireus* by investigating its growth physiology (5), by establishing genetic tools (6), and by reporting here the full-genome sequence of *T. atrovireus* IBT 11181.

Genomic DNA was extracted from the mycelium with a slightly modified protocol of the cetyltrimethylammonium bromide method used by Fulton et al. (7). The *T. atrovireus* IBT 11181 genome was sequenced using an Illumina HiSeq 2000 platform on a 180-bp paired-end library and a 6-kb mate-paired library both with reads of  $2 \times 100$  bp by Beijing Genome Institute (BGI), Hong Kong. Sequencing depth was  $193\times$ , and assembly of the genome was performed with the ALLPATHS-LG algorithm (8). The final assembly resulted in 48 scaffolds with a G+C content of 44.35% and a total assembly size of 30.85 Mb corresponding to 93% of the estimated genome size from *k*-mer spectral analysis. The minimum number of sequences making up 50% of the genome assembly was seven, and the  $N_{50}$  length was 1,577,401 bp. The CEGMA pipeline (9) identified 242 of the 248 core eukaryotic genes, assessing the genome assembly completeness to be 97.58%. This indicated that the draft genome assembly was good with a high completeness and was valid to use for whole-genome analysis.

Gene-calling of the genome was performed using a pipeline of first masking the genome with RepeatMasker version 4.0.5 (Institute for Systems Biology, Seattle, WA, USA; <http://www.repeatmasker.org>), and then gene-calling with AUGUSTUS version 3.0.3 (10, 11), FGENESH version 3.1.2 (SoftBerry) (12), and GeneMark-ES (13). The individual *ab initio* gene predictions were merged into a consensus gene prediction using EVIDENCEModeler (14), resulting in a total of 9,519 protein-encoding genes serving as the final gene prediction. The genome sequence reported here represents a

Received 22 December 2016 Accepted 26 December 2016 Published 2 March 2017

**Citation** Thrane U, Rasmussen KB, Petersen B, Rasmussen S, Sicheritz-Pontén T, Mortensen UH. 2017. Genome sequence of *Talaromyces atrovireus*, which produces red colorants for the food industry. *Genome Announc* 5:e01736-16. <https://doi.org/10.1128/genomeA.01736-16>.

**Copyright** © 2017 Thrane et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ulf Thrane, [uthrane@gmail.com](mailto:uthrane@gmail.com).

\* Present address: Kasper Bøwig Rasmussen, Novo Nordisk A/S, Gentofte, Denmark.

useful resource for further research into the metabolism of *T. atroseus* and its potential as a cell factory for colorant production.

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [LFMY000000000](https://www.ncbi.nlm.nih.gov/nuclink/LFMY000000000). The version described in this paper is the first version, LFMY01000000.

## ACKNOWLEDGMENT

This work was supported by grant 09-064967 from the Danish Council for Independent Research, Technology, and Production Sciences.

## REFERENCES

- Inoue H, Decker SR, Taylor LE 2nd, Yano S, Sawayama S. 2014. Identification and characterization of core cellulolytic enzymes from *Talaromyces cellulolyticus* (formerly *Acremonium cellulolyticus*) critical for hydrolysis of lignocellulosic biomass. *Biotechnol Biofuels* 7:151. <https://doi.org/10.1186/s13068-014-0151-5>.
- Frisvad JC, Yilmaz N, Thrane U, Rasmussen KB, Houbraken J, Samson RA. 2013. *Talaromyces atroseus*, a new species efficiently producing industrially relevant red pigments. *PLoS One* 8:e84102. <https://doi.org/10.1371/journal.pone.0084102>.
- Mapari SAS, Nielsen KF, Larsen TO, Frisvad JC, Meyer AS, Thrane U. 2005. Exploring fungal biodiversity for the production of water-soluble pigments as potential natural food colorants. *Curr Opin Biotechnol* 16: 231–238. <https://doi.org/10.1016/j.copbio.2005.03.004>.
- Mapari SAS, Meyer AS, Thrane U, Frisvad JC. 2009. Identification of potentially safe promising fungal cell factories for the production of polyketide natural food colorants using chemotaxonomic rationale. *Microb Cell Fact* 8:24. <https://doi.org/10.1186/1475-2859-8-24>.
- Mapari SAS, Thrane U, Meyer AS. 2010. Fungal polyketide azaphilone pigments as future natural food colorants? *Trends Biotechnol* 28: 300–307. <https://doi.org/10.1016/j.tibtech.2010.03.004>.
- Nielsen ML, Isbrandt T, Rasmussen KB, Thrane U, Hoof JB, Larsen TO, Mortensen UH. 2017. Genes linked to production of secondary metabolites in *Talaromyces atroseus* revealed using CRISPR-Cas9. *PLoS One* 12:e0169712. <https://doi.org/10.1371/journal.pone.0169712>.
- Fulton TM, Chunwongse J, Tanksley SD. 1995. Microprep protocol for extraction of DNA from tomato and other herbaceous plants. *Plant Mol Biol Rep* 13:207–209. <https://doi.org/10.1007/BF02670897>.
- Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A* 108:1513–1518. <https://doi.org/10.1073/pnas.1017351108>.
- Parra G, Bradnam K, Korf I. 2007. CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics* 23: 1061–1067. <https://doi.org/10.1093/bioinformatics/btm071>.
- Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics* 7:62. <https://doi.org/10.1186/1471-2105-7-62>.
- Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically mapped cDNA alignments to improve *de novo* gene finding. *Bioinformatics* 24:637–644. <https://doi.org/10.1093/bioinformatics/btn013>.
- Salamov AA, Solovyev VV. 2000. Ab initio gene finding in *drosophila* genomic DNA. *Genome Res* 10:516–522. <https://doi.org/10.1101/gr.10.4.516>.
- Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M. 2008. Gene prediction in novel fungal genomes using an ab initio algorithm with unsupervised training. *Genome Res* 18:1979–1990. <https://doi.org/10.1101/gr.081612.108>.
- Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR. 2008. Automated eukaryotic gene structure annotation using EvidenceModeler and the program to assemble spliced alignments. *Genome Biol* 9:R7. <https://doi.org/10.1186/gb-2008-9-1-r7>.