



Title	Using <i>Nassarius stolatus</i> as a potential heavy metal biomonitor
Author(s)	Khidkhan, Kraisiri; Imsilp, Kanjana; Poapolathep, Amnart; Poapolathep, Saranya; Jermnak, Usuma; Tanhan, Phanwimol
Citation	Japanese Journal of Veterinary Research, 64(4), 277-280
Issue Date	2016-11
DOI	10.14943/jjvr.64.4.277
Doc URL	http://hdl.handle.net/2115/63758
Type	bulletin (article)
File Information	059.277-280 SC2 KHIDKHAN.pdf



[Instructions for use](#)

Using *Nassarius stolatus* as a potential heavy metal biomonitor

Kraisiri Khidkhan, Kanjana Imsilp, Amnart Poapolathep,
Saranya Poapolathep, Usuma Jermnak and Phanwimol Tanhan^{*)}

Department of Pharmacology, Faculty of Veterinary Medicine, Kasetsart University, Thailand

Received for publication, September 26, 2016; accepted, October 6, 2016

Abstract

Nassarius stolatus was evaluated as the potential heavy metal (Cd, Fe, Mn, Ni and Pb) accumulator in Don Hoi Lot sandbar, Samut Songkhram province, Thailand. This selected species belongs to the Gastropoda, which is widely distributed in the coastal areas from the upper Gulf through the southern part of Thailand. From our findings, the heavy metal accumulations in *N. stolatus* tissues were Fe > Pb > Mn > Ni > Cd. The retrieved bioaccumulation factor (BAF) indicated that *N. Stolatus* has high potential to be a biomonitor for the contaminations of Fe and Mn in water and Cd, Ni and Pb in sediment.

Key Words: biomonitor, heavy metal, *Nassarius stolatus*

Don Hoi Lot sandbar are located in Samut Songkhram province, Thailand. It is one of the most coastal biodiversity in the upper Gulf of Thailand¹⁷. This area is known as the important aquaculture sites, famous tourist attractions as well as many kinds of industries. These increasing anthropogenic activities can expose a large amount of heavy metal contaminations to the coastal environments, thus, effecting on the living organisms' health. The Pollution Control Department of Thailand reported the major heavy metals contamination found in this area including cadmium (Cd), iron (Fe), manganese (Mn), nickel (Ni), and lead (Pb)^{12,15}. Our previous report showed the sediment contamination

consisted of Cd ($0.85 \pm 0.28 \mu\text{g/g}$), Fe ($5806.53 \pm 333.51 \mu\text{g/g}$), Mn ($409.40 \pm 26.23 \mu\text{g/g}$), Ni ($8.05 \pm 0.28 \mu\text{g/g}$) and Pb ($25.84 \pm 0.68 \mu\text{g/g}$)⁹. To assess the degrees of heavy metal contaminations, many organisms that bioaccumulate pollutants were used as the biomonitors^{5,10}. The main purpose of using biomonitor is the reduction in time and cost used for determination of heavy metals in environmental samples (water and sediment). However, the biomonitors or bioaccumulators must have the basic requirements that they are widely distributed and available all year round; they accumulate and tolerate to the high concentrations of heavy metals present in their environments without being killed; they are

^{*)}Corresponding author: Phanwimol Tanhan, Ph.D., Department of Pharmacology, Faculty of Veterinary Medicine, Kasetsart University
Phone: +66 2 579 7537. Fax: +66 2 579 7537. E-mail: fvetpmt@ku.ac.th
doi: 10.14943/jjvr.64.4.277

Table 1. Average heavy metal accumulations (mean \pm SEM) in sediment, water and *N. stolatus* tissues

Heavy metal	Average concentration		
	Sediment ($\mu\text{g/g}$)	Water ($\mu\text{g/mL}$)	<i>N. stolatus</i> ($\mu\text{g/g}$)
Cd	0.86 \pm 0.05	0.02 \pm 0.00	8.14 \pm 0.82
Fe	5891.07 \pm 61.72	0.06 \pm 0.00	210.24 \pm 11.88
Mn	499.20 \pm 16.18	0.02 \pm 0.00	135.17 \pm 15.33
Ni	8.25 \pm 0.11	0.08 \pm 0.00	49.52 \pm 5.29
Pb	25.81 \pm 0.18	0.02 \pm 0.01	205.50 \pm 24.26

easily collected and preserved³). From all above requirements, *Nassarius stolatus* is a non-edible gastropod normally found in marine environment, especially in intertidal areas including Don Hoi Lot sandbar. It feeds as a benthic scavenger. In addition, the filter respiration and benthic feeding of gastropod can uptake and accumulate heavy metals through gills as well as gastrointestinal tract¹⁴. *N. stolatus* rapidly consumes a large amount of foods compared to its body weight. It was susceptible to heavy metal toxicities, i.e., Cd, Cu and Zn²). Therefore, it could be used as a potential biomonitor for heavy metal contaminated areas.

The aims of this study were to determine the accumulations of Cd, Fe Mn, Ni and Pb in *Nassarius stolatus* collected from Don Hoi Lot sandbar, and to assess the potential use of *N. stolatus* as a biomonitor of heavy metals contaminated water and sediment.

N. stolatus as well as water and sediment were collected from Don Hoi Lot sandbar. Forty sampling points were systematic located by using GPS from N 13°20'02.1" E 100°01'09.9" to 13°21'48.3" E 100°00'32.5". The performances of *N. stolatus* were recorded. Water was filtered through filter paper (Whatman No.4) to remove debris and a few drops of HNO₃ (70%) were added before determining heavy metal concentrations. Sediment and *N. stolatus* tissues were oven dried at 70°C for 48 hours. Then, dried sediment and tissues (0.5 g) were digested with wet acid-digestion using the mixtures of HNO₃ (70%) and H₂O₂ (35%) (1:1 v/v)¹. After complete digestion, all samples were determined for heavy metal

concentrations using a flame atomic absorption spectrophotometer (FAAS; 240B Agilent Technologies). The limits of quantification (LOQ) were 0.01 $\mu\text{g/g}$ for Mn, 0.02 $\mu\text{g/g}$ for Cd and Fe, 0.03 $\mu\text{g/g}$ for Ni and 0.07 $\mu\text{g/g}$ for Pb. The recovery percentages of all heavy metals ranged from 91%–112%. All data were statistically analyzed using SPSS version 16.0 (SPSS Inc., Illinois, USA). The descriptive statistics (mean and standard error of the mean) of heavy metal concentrations in all samples were calculated. The bioaccumulation factor (BAF) was also calculated to compare the heavy metal uptake through all routes of exposure by *N. stolatus* according to the following equation¹¹;

$$\text{BAF} = C_t / C_m$$

Where, C_t is the heavy metal concentration in mollusk tissue and C_m is the heavy metal concentration in water or sediment.

The average shell length and weight of collected *N. stolatus* (n = 109) were 1.28 \pm 0.06 cm and 0.73 \pm 0.05 g, respectively. Their size and weight characteristics indicated that *N. stolatus* is a small to medium sized gastropod similar to other species belonging to the Family Nassariidae. These characteristics were positively correlated with the accumulation of heavy metals found in mollusk tissues^{5,7,10}. The heavy metal accumulation in *N. stolatus* tissues were Fe > Pb > Mn > Ni > Cd (Table 1). Among these heavy metals, the level of Fe was the highest. The finding was similar to other gastropods in previous studies^{6,7,18}. It suggested that Fe is an essential element important for metabolic

Table 2. Comparative bioaccumulation factor (BAF) of *N. stolatus* tissues with other gastropod species

Heavy metal	<i>Nassarius stolatus</i>		<i>Cerathidea cingulata</i> ^a	<i>Cipangopaludina chinensis</i> ^b	<i>Planaxis sulcatus</i> ^a	<i>Thais carinifera</i> ^c
	BAF-water	BAF-sediment	BAF-sediment	BAF-sediment	BAF-sediment	BAF-sediment
Cd	362.77	9.76	0.17–0.30	9.00	–	4.40
Fe	4,164.17	0.04	0.003–0.01	–	247.5	–
Mn	9,517.93	0.27	0.11–0.12	–	100.5	–
Ni	617.35	6.21	0.02–0.03	–	–	–
Pb	890.75	8.02	0.01–0.02	11.20	104.1	0

^aManavi (2013)^bTornimbeni, *et al.* (2013)^cIbrahim and Abu El-Regal (2014)

activities of marine organisms including mollusks^{8,20}. The observed heavy metal concentrations in *N. stolatus* tissue indicated that Don Hoi Lot sandbar was contaminated with these heavy metals.

By using the BAF, *N. stolatus* was classified as a bioaccumulator for heavy metals in water and sediment⁴. The BAF-water and BAF-sediment values in *N. stolatus* were shown in Table 2. The BAF-water of Fe and Mn were greater than 1,000, whereas the BAF-sediment of Cd, Ni and Pb were greater than 2. These results indicated that *N. stolatus* was the macro-concentrator for Fe and Mn as well as Cd, Ni and Pb in water and sediment, respectively⁴. The higher BAF-sediment values of most heavy metals were found in *Planaxis sulcatus* and *Thais carinifera* (Table 2) indicating that these gastropods have high potential to concentrate these heavy metals in their tissues⁴. Because of their life cycle and feeding behaviour, gastropods are exposed to greater pollution including heavy metals and therefore accumulate higher levels of heavy metals than their environments¹³. Mollusks uptake dissolved heavy metals in water during respiration and feeding, whereas benthic feeding gastropods also extensively uptake insoluble heavy metals from sediments^{14,19}. Apart from the environmental or geochemical factors (i.e. sediment texture, rate of sedimentation, resuspension, pH, and organic matter), the biological factors also affect heavy metal bioaccumulation¹⁹. The comparative BAF-sediment values of the other 4 gastropods

(*Cerathidea cingulate*, *P. sulcatus*, *T. carinifera*, and *Cipangopaludina chinensis*) and *N. stolatus* indicated that species variation is one of the important biological factors for heavy metal bioaccumulation. The differences of heavy metal bioaccumulation in gastropod tissues are caused by the variations of heavy metal affinities to the binding proteins, especially metallothioneins¹⁶. In addition, the accumulation can also be different due to gastropod size, sex, age, genotype, phenotype, feeding activity as well as reproductive stage.

In summary, although *N. stolatus* was a small to medium-sized gastropod, it can accumulate 5 heavy metals in their tissues (Fe > Pb > Mn > Ni > Cd). *N. stolatus* also showed a high capacity to bioconcentrate heavy metals from water (Fe and Mn) and sediments (Cd, Ni, and Pb) when compared to other gastropods. Therefore, it can be a suitable heavy metal biomonitor for contaminated water and sediment.

This study was financially supported by Graduate School, Kasetsart University, Thailand.

References

- 1) APHA, AWWA and WEF 2011. *Standard Methods for Examination of Water and Wastewater*, APHA, AWWA, WEF, Washington, DC.
- 2) Cheung, S. G., Tai, K. K., Leung, C. K. and Siu, Y. M. 2002. Effects of heavy metals on the survival and feeding behaviour of the

- sandy shore scavenging gastropod *Nassarius festivus* (Powys). *Mar. Pollut. Bull.*, **45**: 107–113.
- 3) Conti, M. E., Stripeikis, J., Iacobucci, M., Cucina, D., Cecchetti, G. and Tudino, M. B. 2006. Trace metals in molluscs from the Beagle Channel (Argentina): a preliminary study. *WIT Transac. Ecol. Environ.*, **99**: 473–483.
 - 4) Dallinger, R. and Rainbow, P. S. 1993. *Ecotoxicology of metals in invertebrates* (SETAC Special Publications), Lewis, Chelsea, MI.
 - 5) Dumme, V., Kruatrachue, M., Trinachartvanit, W., Tanhan, P., Pokethitiyook, P. and Damrongphol, P. 2012. Bioaccumulation of heavy metals in water, sediments, aquatic plant and histopathological effects on the golden apple snail in Beung Boraphet reservoir, Thailand. *Ecotox. Environ. Safe.*, **86**: 204–212.
 - 6) Ghosh, I., Mitra, A., Rudra, T., Pramanick, P. and Biswas, P. P. 2016. Bioaccumulation pattern of heavy metals by gastropods: a case study from lower Gangetic Delta. *J. Environ. Sci. Computer Sci. Eng. Technol.*, **5**: 99–102.
 - 7) Ibrahim, N. K. and Abu El-Regal, M. A. 2014. Heavy metals accumulation in marine edible molluscs, Timsah lake, Suez canal, Egypt. *ARPN J. Sci. Technol.*, **4**: 282–288.
 - 8) Kesavan, K., Murugan, A., Venkatesan, V. and Kumar, B. S. V. 2013. Heavy metal accumulation in molluscs and sediment from Uppanar estuary, Southeast coast of India. *Thalassas: Int. J. Mar. Sci.*, **29**: 15–21.
 - 9) Khidkhan, K., Tanhan, P., Imsilp, K., Poapolathep, A., Poapolathep, S. and Trinachartvanit, W. 2014. Preliminary Investigation of Toxic Heavy Metals Along Don Hoi Lot Sandbars, Thailand. The 18th Federation of Asian Veterinary Association Congress, Marina Bay Sands, Singapore, 28–30 November, 2014.
 - 10) Kruatrachue, M., Sumritdee, C., Pokethitiyook, P. and Singhakaew, S. 2011. Histopathological effects of contaminated sediments on golden apple snail (*Pomacea canaliculata*, Lamarck 1822). *Bull. Environ. Contam. Toxicol.*, **86**: 610–614.
 - 11) Liao, C. M. and Ling, M. P. 2003. Assessment of human health risks for arsenic bioaccumulation in tilapia (*Oreochromis mossambicus*) and large-scale mullet (*Liza macrolepis*) from blackfoot disease area in Taiwan. *Arch. Environ. Contam. Toxicol.*, **45**: 264–272.
 - 12) Ma, T. and Niyomthai, W. 2010. Contamination of Heavy Metals in Economical Important Bivalve at Don Hoi Lod, Samut Songkhram Province. *Proceedings of 48th Kasetsart University Annual Conference: Natural Resources and Environment*, Kasetsart University, Bangkok, Thailand, 3–5 February, 2010.
 - 13) Manavi, P. N. 2013. Heavy metals in water, sediment and macrobenthos in the intertidal zone of Hormozgan province, Iran. *Mar. Sci.*, **3**: 39–47.
 - 14) Oehlmann, J. and Schulte-Oehlmann, U. 2003. Chapter 17 Molluscs as bioindicators. In: *Trace Metals and other Contaminants in the Environment*, pp. 577–635, A. M. B. B. A. Markert and H. G. Zechmeister, Elsevier, USA.
 - 15) Rermdumri, S., Kareas, S. and Wongsontam, K. 2009. Heavy Metals Contamination in Seawater and Sediment in the Upper Gulf of Thailand. *Proceedings of 47th Kasetsart University Annual Conference: Natural Resources and Environment*, Kasetsart University, Bangkok, Thailand, 17–20 March, 2009.
 - 16) Shirneshan, G. and Bakhtiari, A. R. 2012. Accumulation and distribution of Cd, Cu, Pb and Zn in the soft tissue and shell of oysters collected from the northern coast of Qeshm Island, Persian Gulf, Iran. *Chem. Spec. Bioavailab.*, **24**: 129–138.
 - 17) Taen-on, S. 2007. The Management under Ramsar Convention: Don Hoi Lot Wetland, Samutsongkhram Province. Thesis Master of Science, Thammasat University, Thailand.
 - 18) Tornimbeni, O., Galvez, R., Triffault-Bouchet, G., Dassylva, N. and Roberge, S. 2013. Heavy metal concentrations in *Cipangopaludina chinensis* (Reeve, 1863) and relationships with sediments in Saint-Augustin Lake, Québec City (Qc, Canada). *Ann. Limnol. - Int. J. Lim.*, **49**: 21–29.
 - 19) Wang, W. X. 2009. 9 - Metals and organic contaminants in bivalve molluscs. In: *Shellfish Safety and Quality*, pp. 228–247, Woodhead Publishing.
 - 20) Yusoff, N. A. M. and Long, S. M. 2011. Comparative Bioaccumulation of Heavy Metals (Fe, Zn, Cu, Cd, Cr, Pb) in Different Edible Mollusk Collected from the Estuary Area of Sarawak River. *Universiti Malaysia Terengganu 10th International Annual Symposium (UMTAS 2011)*, Kuala Terengganu, Terengganu, Malaysia, 11–13 July, 2011.