



Title	Identification of novel alternative splicing variants within swine Setd8 gene and their high mRNA expression in testis
Author(s)	Yu, Shuai; Chen, Xiaoxu; Deng, Zhongyuan; Lan, Xianyong; Pan, Chuanying
Citation	Japanese Journal of Veterinary Research, 65(1), 5-15
Issue Date	2017-02
DOI	10.14943/jjvr.65.1.5
Doc URL	<a href="http://hdl.handle.net/2115/64789">http://hdl.handle.net/2115/64789</a>
Type	bulletin (article)
File Information	65-1_005-015.pdf



[Instructions for use](#)

# Identification of novel alternative splicing variants within swine *Setd8* gene and their high mRNA expression in testis

Shuai Yu<sup>1)</sup>, Xiaoxu Chen<sup>1)</sup>, Zhongyuan Deng<sup>1)</sup>, Xianyong Lan<sup>1, 2)</sup> and Chuanying Pan<sup>1,\*)</sup>

<sup>1)</sup> College of Animal Science and Technology, Northwest A&F University, Yangling, Shaanxi 712100, China

<sup>2)</sup> Shaanxi Key Laboratory of Molecular Biology for Agriculture, Northwest A&F University, Yangling, Shaanxi 712100, China

Received for publication, August 26, 2016; accepted, November 28, 2016

## Abstract

SET domain containing (lysine methyltransferase) 8 (*Setd8*), a histone modification enzyme, affects cell cycling, chromosome condensation, high efficient repair of DNA double strand breaks and so on. The objective of this study was to identify novel alternative splicing variants of pig *Setd8* gene and its mRNA expression. Four 180-day-old male Guanzhong Black (GZB) pigs and six male Landrace piglets (including three 30-day-old and three 7-day-old pigs) were collected to study *Setd8* gene. Herein, two novel variants, *Setd8a* and *Setd8b*, were found in pig. The entire sequences of *Setd8a* and *Setd8b* variants were 1,039 bp and 958 bp, respectively. qRT-PCR results showed that *Setd8a* and *Setd8b* were highly expressed in brains and testes of 180-day-old GZB pigs. Moreover, the expressions of the two *Setd8* variants were significantly higher in testis than brain of GZB pig ( $P < 0.05$ ). Further study on testis showed that the mRNA expression of *Setd8a* variant was significantly lower than *Setd8b* variant in 30-day-old and 7-day-old pigs ( $P < 0.05$ ). The mRNA expression of *Setd8a* variant was lower than *Setd8b* variant in GZB pigs ( $P > 0.05$ ). Moreover, the expressions of the two *Setd8* variants were significantly higher along with age enlargement. In conclusion, *Setd8a* and *Setd8b* were firstly identified in pigs and both were expressed in pig testis. *Setd8b* was the major splicing variant of pig *Setd8* gene transcript product. Moreover, the expressions of *Setd8* variants were time-dependent. All these findings would enrich the study of *Setd8* gene in pig testis.

Key Words: pig, testis, SET domain containing (lysine methyltransferase) 8 (*Setd8*), alternative splicing variant, expression patterns.

## Introduction

Pig is not only an important economic large domestic animal for meat industry, but also an ideal mammalian model for human biomedical and diseases studies<sup>13,33,42</sup>. At present, pig industry in China faces severe problems, and one of the

worst is subfertility. One way to solve male subfertility is spermatogonial stem cells (SSCs) transplantation<sup>3,17</sup>. SSCs can transmit genomic information to the offspring by differentiating into spermatozoa<sup>43</sup>. Thus it is worthy to study the biological characteristics and regulation mechanisms of SSCs. Present studies show

\*Corresponding author: Chuanying Pan, College of Animal Science and Technology, Northwest A&F University, Yangling, Shaanxi 712100, P.R. China

Phone: +86-29-87092102. Fax: +86-29-87092164. E-mail: panyu1980@126.com

doi: 10.14943/jjvr.65.1.5

that some critical factors, such as Wnt or proliferating cell nuclear antigen (PCNA), are relevant to regulating the proliferation and stem cell properties maintenance of SSCs<sup>11,12,45,46</sup>. Furthermore, the Wnt and PCNA expression are both mediated by SET domain containing (lysine methyltransferase) 8 (*Setd8*)<sup>24,39,41</sup>. Thus it could be conjectured that *Setd8* might regulate the SSCs activity in mammals. However, little was known about function of *Setd8* gene in pigs.

*Setd8*, also known as *KMT5a*, *SET8* and *PR-Set7/9* gene, is located on pig chromosome 14. The *Setd8* gene in pig is 18,955 bp in length and contains eight exons and seven introns with strictly conserved intron/exon boundaries (NC\_010456.4). As one member of SET domain protein superfamily, *Setd8* has one SET domain, which consists of three elements: Suppressor of variegation 3-9 (Su(var) 3-9), Enhancer of zeste (E(z)), and Trithorax (Trx)<sup>35</sup>. The SET domain has a conserved sequence among the superfamily, and approximate 130 amino acids in length<sup>36</sup>.

*Setd8* has the irreplaceable function in human, mouse and other mammals as a histone modification enzyme<sup>8,24</sup>. Firstly, *Setd8* was the exclusive enzyme catalysing histone H4 monomethylation on Lys 20 (H4K20me1). Secondly, present studies predicted that *Setd8* could regulate cell cycling on S phase progression and promote chromosome condensation in cell cycle progression<sup>15,25,37</sup>. The absence of *Setd8* was shown to be able to induce DNA damage, cell cycle arrest, and sometimes cell apoptosis<sup>34,44</sup>. It was suggested that the mouse embryo could not survive to birth when *Setd8* was deleted<sup>32</sup>. Moreover, latest study demonstrated that *Setd8* took a part in self-renewal of human adult stem cells<sup>8</sup>. Meanwhile, the expression level of DNA methyltransferase influenced the spermatogenesis<sup>40</sup>. Moreover, the expression of *Setd8* was decreased along with spermatogonia differentiating into spermatocyte in mouse<sup>14</sup>. Therefore, it could be hypothesized that *Setd8* was one of the most important protein that support vital movement of pigs and taken effects

on reproduction traits.

Histone modification was found to link with alternative splicing (AS) and AS played a part in vital activity of histone modification in turn<sup>2,18</sup>. *Setd8* had been found to have two certain AS variants and several predicted variants in human, and four predicted variants in mouse and rat<sup>1,10</sup>. However, variants of *Setd8* had not been reported in pig. To better understanding pig *Setd8* gene and its role in SSCs activity, we chose it as a candidate gene to explore whether AS would exist in pig testis or not and analyzed their mRNA expression. The data would contribute to understanding the role of *Setd8* gene in reproduction traits in pig, and further facilitate the development of pig industry.

## Material and methods

**Tissues collection:** Use of all experiments animals and operation programs were permitted by International Animal Care and Use Committee of Northwest A&F University in accordance with the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health.

Four 180-day-old male Guanzhong Black (GZB) pigs and six male Landrace piglets (including three 30-day-old and three 7-day-old pigs) were collected from local farm and Besun agricultural industry group Co., Ltd. in Shaanxi province, China, respectively. A total of seven tissues (including heart, liver, spleen, lung, kidney, brain, and testis) were obtained from four GZB pigs. Only testicular tissues were collected from Landrace pigs. Each tissue was immediately submerged in liquid nitrogen and then stored at  $-80^{\circ}\text{C}$  for subsequent study<sup>22</sup>.

**RNA extraction and cDNA synthesis:** Total RNA of each sample was extracted using RNAiso Plus reagent (TaKaRa, Dalian, China), as well as the RNase-free DNase I (TaKaRa) was used to clean up genomic DNA from the RNA samples. The quantity of RNA was assessed by OD<sub>260/280</sub> value

**Table 1. Primers used for cloning and expression analysis of pig *Setd8* and its variants**

Primer	Primer sequences (5'-3')	Reference sequence	Length of production/bp	Notes
P1	F: ATGACTAAACCTTCCGAG R: TGAGGGCACTTTGTCCAT	XM_013982772.1	1094/1013	mRNA sequence of <i>Setd8a</i> / <i>Setd8b</i> , respectively
P2	F: GTGTCCTTCCAGTGCAACCT R: AGAGCATTTGTTCCGGCTCA	KX254435	97	qRT-PCR for <i>Setd8a</i>
P3	F: GACTAAACCTTCCGAGGCGG R: ACTGAGTTCTTCCGTCGC	KR021363	124	qRT-PCR for <i>Setd8b</i>
GAPDH	F: AACTCACTCTTCTACCTTTG R: CAAATTCATTGTCGTACCAG	NM_001206359.1	90	qRT-PCR for internal control

using the NanoDrop 1000 spectrophotometer (Thermo Fisher Scientific Inc., Wilmington, DE) and agarose electrophoresis<sup>22</sup>. Then reverse transcription PCR was performed to synthesized cDNA using the PrimeSript<sup>TM</sup> RT reagent Kit (TaKaRa) according to the manufacturer's recommended procedure<sup>23</sup>.

*Identification of Setd8 splicing variants:* On the basis of the predicted nucleotide sequence of pig *Setd8* gene (XM\_013982772.1), a pair of primers (P1, Table 1) was designed by Primer Premier 5 software (Premier BioSoft, Palo Alto, CA, USA) to identify novel variants. PCR reactions were performed with Touch-Down PCR in a 25  $\mu$ L volume containing cDNA, 0.5  $\mu$ M of each primers (P2 and P3), 2 $\times$  Eco Taq PCR SuperMix (+dye) Taq DNA polymerase, MgCl<sub>2</sub>, dNTPs, and buffer and rest volumes of distilled water (Beijing TransGen Biotech Co., Ltd)<sup>6</sup>. The PCR products were analyzed by 2% agarose electrophoresis and the target bands were then purified with Gel Extraction Kit (Sangon Biotech, Shanghai, China). After that, the PCR products were sub-cloned into pMD19-T Vector (TaKaRa), and transferred into *Escherichia coli*. Competent cells *DH5 $\alpha$*  (TaKaRa), then verified by sequencing (Gen-Script Co., Ltd, Nanjing, China).

*Bioinformatics analysis:* The sequences alignments of nucleotide were conducted by BioXM 2.6 (Nanjing Agricultural University, Nanjing, China) and NCBI BLAST (<http://blast.ncbi.nlm.nih.gov/>

Blast).


*Measurement of Setd8a and Setd8b mRNA expression:* According to the sequencing results, two pairs of primers (Table 1) were used to detect the mRNA expression levels of the two *Setd8* splicing variants. The house keeping gene glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) (Table 1) was used as internal control<sup>20</sup>. Quantitative real-time PCR (qRT-PCR) was run on a Bio-Rad IQ5 Real-Time PCR system with three repeats for each sample. In addition, a blank control was set in each sample group. The PCR reaction system was 20  $\mu$ L in volume: 10  $\mu$ L SYBR<sup>®</sup> Premix Ex Taq II (TaKaRa) (2 $\times$ ), 1  $\mu$ L cDNA (diluted for 100 times), 0.5  $\mu$ L of each primer (P2 or P3) (10  $\mu$ mol/L) and rest volumes of distilled water.

*Statistical analysis:* The mRNA expressions of *Setd8* variants were computed by 2<sup>- $\Delta\Delta$ CT</sup> method, and normalized by the expression of *GAPDH*<sup>32</sup>. The mRNA expression variation among different samples was calculated by SPSS (version 18.0) (SPSS, Inc., Chicago, IL, USA). Statistical differences of *Setd8* variants expressions in different tissues and different ages were performed by ANOVA<sup>44</sup>.

## Results

### Identification of pig *Setd8* gene variants

Two variants were identified in pig, *Setd8a*



**Fig. 1. The genomic structure diagram, positions of primers and the predictive transcription factors binding sites in swine *Setd8* and its variants.** Note: The schematic used predicted *Sus scrofa Setd8* gene (XM\_013982772.1) as reference sequence. *Setd8a* (KX254435) and *Setd8b* (KR021363) were identified in this study. The arrowhead represented the position of the primers of *Setd8a* and *Setd8b* variants (Primer 2 and 3).

(GenBank Accession number: KX254435) and *Setd8b* (GenBank Accession number: KR021363), and both were novel transcript variants found in swine in this study. The entire coding sequences of *Setd8a* and *Setd8b* were 1039 bp and 958 bp in length, respectively (Fig. 1). After confirming the two alternative splicing variants, these fragments were sub-cloned to pMD19-T Vector and verified by sequencing.


The nucleotide sequencing analysis found that *Setd8* had eight exons, while both *Setd8a* and *Setd8b* had seven exons, lacking of exon 2 (150 nucleotides). The exon 1 of *Setd8a* and *Setd8b* was 37 nucleotides (nt) longer than that of *Setd8*, and the 37 nt was from intron 1 retention (Fig. 1). When the 37 nt sequence was mapped to pig genome sequence, the splicing sites were found to comply with the GT-AG rule for 5' splice donor and 3' splice acceptor sites. The exon 3 of *Setd8b* was 81 nt shorter than those of *Setd8* and *Setd8a*, as well as the exon skipping was fit the GT-AG rule (Fig. 1).

Sequences alignment pointed that pig *Setd8a* and *Setd8b* variants shared the similarity with the predicted *Sus scrofa Setd8* sequence (XM\_013982772.1). They all included the SET domain (Fig. 2).

#### *Expression profiles of pig Setd8 variants*

The RT-PCR results showed that the *Setd8* variants could be identified in testis of all ten male pigs. Histological staining was used to detect the structure of 180-day-old GZB pigs. As shown in Fig. 3, the elongated spermatid cells could be seen in the seminiferous tubules (Fig. 3). Then the GZB pigs in this study were called “puberty”. The mRNA expressions of both *Setd8* variants in puberty GZB pigs were highly expressed in testis and brain tissues (Fig. 4, Fig. 5). Statistics analyses showed that the expressions of *Setd8a* and *Setd8b* in testis were significantly higher than in brain of puberty GZB pigs ( $P < 0.05$ ) (Fig. 5). In addition, the expression levels of *Setd8a* were similar with *Setd8b* in testis of puberty GZB pigs (Fig. 5). Moreover, there was no or rare expression in heart, spleen, lung, kidney and liver of puberty GZB pigs. Then the function of *Setd8* variants in testis was worth to study.

Then the expression of *Setd8* variants in different periods of male pig testis was detected. As shown in Fig. 6, the expression levels of *Setd8* variants were both increased along with age, and the expression of two variants both reached the highest levels in puberty than other childhood periods ( $P < 0.05$ ). When focused on 7-day-old and 30-day-old pigs, the expression level of




**Fig. 2. DNA sequence alignments of *Setd8*, *Setd8a* and *Setd8b* in pig.** Note: *Setd8*: predicted *Sus scrofa Setd8* mRNA (XM\_013982772.1); *Setd8a*: *Sus scrofa Setd8a* transcript variant *Setd8a* mRNA (KX254435); and *Setd8b*: *Sus scrofa Setd8b* transcript variant *Setd8b* mRNA (KR021363). *Setd8a* and *Setd8b* were identified in this study. Gray and black shades indicated regions of two and three sequences identity, respectively. The red box showed the region of SET domain.

*Setd8b* was higher than that of *Setd8a* in testis GZB pigs. ( $P < 0.01$ ) (Fig. 6). However, as mentioned above, there was no significant difference in expression levels of *Setd8a* and *Setd8b* in testis of puberty

## Discussion


Published literatures had revealed that AS



**Fig. 3. H&E staining in 180-day old GZB pig testis** (Scale bar = 50  $\mu$ m). The emergence of elongated spermatid cells illustrated that the 180-day-old GZB pigs were puberty. Note: the white arrowheads indicated cell nucleus of elongated spermatid cells in pig seminiferous tubules.

was a universal phenomenon in animals. As we all know, exons were confirmed with three major points: the 5' splice site, the 3' splice site and the branch point, which was called consensus "GT-AG" splicing rule<sup>18)</sup>. Based on this condition, the mechanism of forming alternative splicing was simply summarized: exon skipping, intron retention, alternative 3' splice site and alternative 5' splice site<sup>23,47)</sup>. AS can not only generate the transcript and protein diversity, but also influence cell differentiation and cell death<sup>2,19,38)</sup>. The epigenetic regulators, such as *Setd8*, could occur the AS in mammalian, suggesting that the AS may be a driving force in regulating the function of these enzymes<sup>21,27,28)</sup>.

Previous study demonstrated that *Setd8* was functionally essential for high efficient repair of DNA double strand breaks<sup>5,9)</sup>. *Setd8* also played a role in human carcinogenesis by aberrant lysine methylation of PCNA, and it was essential for




**Fig. 4. The mRNA expression patterns of *Setd8a* (a), *Setd8b* (b) and *GAPDH* (c) in different tissues of 180-day-old GZB pigs.**


genomic stability<sup>15,41</sup>. Moreover, *Setd8* was a context-dependent GATA-1 corepressor in erythroid cells, thus it was crucial in cell survival and maturation<sup>4,29</sup>. While deficiency of *Setd8* in c-Myc-overexpressing skin blocked cell proliferation and

differentiation as well as caused apoptosis<sup>7</sup>. *Setd8* played a vital role in cell cycle, chromatin modification and organization.

The importance of *Setd8* in life made it important so that its structure should be further



**Fig. 5. Relative expressions of *Setd8a* and *Setd8b* variants in different tissues of 180-day-old GZB pigs.** Note: The *GAPDH* was the housekeeping gene, and the mRNA expression levels of *Setd8a* and *Setd8b* were normalized to that of *GAPDH*. \* and \*\* represented the significant difference ( $P < 0.05$ ) and ( $P < 0.01$ ), respectively.



**Fig. 6. Relative expressions of *Setd8a* and *Setd8b* variants in different periods of testis in male pigs.** Note: The *GAPDH* was the housekeeping gene, and the mRNA expression levels of *Setd8a* and *Setd8b* were normalized to that of *GAPDH*. \* and \*\* represented the significant difference ( $P < 0.05$ ) and ( $P < 0.01$ ), respectively.



examined. *Setd8* was firstly discovered and purified in Hela cells in 2002, while rare studies investigated its AS and mRNA expression in pig<sup>31</sup>. In this study, two variants were identified in testis and brain for the pig *Setd8* gene. Two variants, which named *Setd8a* and *Setd8b*, were found in human *Setd8* gene<sup>1,10</sup>. Sequence alignment of pig *Setd8a* and *Setd8b* indicated that they were similar to their counter-parts of human. Both of them had PIP box2 and SET domain. In addition, sequence analyses showed that the *Setd8* variants of human and pig all experienced the exon skipping. Moreover, *Setd8b* was the main functional splicing variant in human, and the result of qRT-PCR in this study was consistent with that in human<sup>1,10</sup>. Therefore, we presumed that pig *Setd8a* and *Setd8b* should perform the similar expression patterns to human *Setd8a* and *Setd8b*.

There were several points different between pig and human *Setd8* variants. At first, unique additional intron retention and partly missing exon were detected in pig *Setd8* variants, which enhanced the variants complexity in this study. However, the mechanisms how they formed were unknown. In addition, the mRNA expression of pig normal *Setd8* (GenBank Accession number: XM\_013982772.1) was not detected in pig, predicting that the expression of *Setd8* was lowly or rarely in GZB pigs. However, the expression of human *Setd8* gene could be identified in testis.

Compared with other histone methyltransferases, the expression of *Setd8* was low somehow in tissues. Its abnormal expression could be contributed to human cancers<sup>30</sup>. Recent study showed that *Setd8* played an important role in erythroid survival, and deficiency of *Setd8* resulted in loss of H4K20me1<sup>29</sup>. These results predicted that *Setd8* would be expressed in blood, which didn't prove in this study. It was predicted by the expression of *Setd8* variants in the two tissues that *Setd8* might be benefit for reproduction of pigs.

It is said that 76% of all genes were expressed in brain and AS was related with

different developmental stages<sup>47</sup>. In this study, *Setd8* variants were detected in brain and testis of puberty GZB pigs. Testis and brain were the most important organs regulating reproduction traits in pig. The result predicted that *Setd8* might play vital roles in brain, such as regulation of brain development. In addition, the expressions of *Setd8* variants were lower in brain than in testis. There were several studies showed that AS took an important role in species evolution and AS was always found in testis and cancer cell lines<sup>16,18</sup>. AS occurred in testis suggested a natural increase in the variation in expression to allow for evolutionary selection. Then the expressions of the two *Setd8* variants in testis were focused in this study.

In this study, three developmental stages (7-, 30 and 180-day-old) of pig testes were obtained to study the expression patterns of *Setd8* variants in testis. The germ cells were gonocytes in 7 days postnatal pig testis<sup>48</sup>. In addition, the gonocytes were developed into SSCs during 1~2 months in pig<sup>48</sup>. Then the SSCs differentiated in sperm to transferring genetic information to the next generation when pig reached a puberty age<sup>48</sup>. qRT-PCR results predicted the expressions of *Setd8* variants were time-regulated, and they might be related with the differentiation of germ cells. Moreover, *Setd8b* were mainly expressed in testis of young piglets, showed that *Setd8b* would be benefit to testicular development.

## Conclusions

To summarize, two variants (*Setd8a* and *Setd8b*) were firstly identified in pig in this study. In addition, the two variants were expressed in testis and brain of puberty GZB pigs and the expression in testis was significantly higher than in brain. Moreover, the expressions of *Setd8* variants were time-dependent in different age pigs. It could be predicted that *Setd8b* was the major splicing variant of porcine *Setd8* transcript products, and *Setd8a* might play an important

role in testis and brain of puberty pigs.

### Acknowledgments

This work was supported by Fundamental Research Funds for the Central Universities (NWSUAF, No.2452015145), China Postdoctoral Science Foundation funded project (No.2014M560809), Yangling Innovation Major Project on the Integration of Industry, Education and Research (2016CXY-17) and the National Basic Research Program of China (973 program; No.2014CB943100).

### Compliance with ethical standards

### Conflicts of interest

The authors declare no conflict of interest.

### References

- 1) Abbas, T., Shibata, E., Park, J., Jha, S., Karnani, N. and Dutta, A. 2010. CRL4<sup>Cdt2</sup> regulates cell proliferation and histone gene expression by targeting PR-Set7/Set8 for degradation. *Mol. Cell.*, **40**: 9–21.
- 2) Chen, M. and Manley, J. L. 2009. Mechanisms of alternative splicing regulation: insights from molecular and genomics approaches. *Nat. Rev. Mol. Cell Biol.*, **10**: 741–754.
- 3) Ehmcke, J., Wistuba, J. and Schlatt, S. 2006. Spermatogonial stem cells: questions, models and perspectives. *Hum. Reprod. Update*, **12**: 275–282.
- 4) De Vilbiss, A. W., Sanalkumar, R., Hall, B. D., Katsumura, K. R., de Andrade, I. F. and Bresnick, E. H. 2015. Epigenetic determinants of erythropoiesis: role of the histone methyltransferase Setd8 in promoting erythroid cell maturation and survival. *Mol. Cell Biol.*, **35**: 2073–2087.
- 5) Diao, L., Su, H., Wei, G., Li, T., Gao, Y., Zhao, G. and Guo, Z. 2014. Prognostic value of microRNA 502 binding site SNP in the 3'-untranslated region of the SET8 gene in patients with non-Hodgkin's Lymphoma. *Tumori*, **100**: 553–558.
- 6) Don, R. H., Cox, P. T., Wainwright, B. J., Baker, K. and Mattick, J. S. 1991. 'Touchdown' PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Res.*, **19**: 4008.
- 7) Driskell, I., Oda, H., Blanco, S., Nascimento, E., Humphreys, P. and Frye, M. 2012. The histone methyltransferase Setd8 acts in concert with c-Myc and is required to maintain skin. *EMBO J.*, **31**: 616–629.
- 8) Driskell, I., Oeztuerk-Winder, F., Humphreys, P. and Frye, M. 2015. Genetically induced cell death in bulge stem cells reveals their redundancy for hair and epidermal regeneration. *Stem Cells*, **33**: 988–998.
- 9) Dulev, S., Tkach, J., Lin, S. and Batada, N. N. 2014. SET8 methyltransferase activity during the DNA double-strand break response is required for recruitment of 53BP1. *EMBO Rep.*, **15**: 1163–1174.
- 10) Fang, J., Feng, Q., Ketel, C. S., Wang, H., Cao R., Xia L., Erdjument-Bromage H., Tempst P., Simon J. A. and Zhang Y. 2002. Purification and functional characterization of SET8, a nucleosomal Histone H4-lysine 20-specific methyltransferase. *Curr. Biol.*, **12**: 1086–1099.
- 11) Golestaneh, N., Beauchamp, E., Fallen, S., Kokkinaki, M., Uren, A. and Dym, M. 2009. Wnt signaling promotes proliferation and stemness regulation of spermatogonial stem/progenitor Cells. *Reproduction*, **138**: 151–162.
- 12) He, Z., Jiang, J., Kokkinaki, M., Tang, L., Zeng, W., Gallicano, I., Dobrinski, I. and Dym, M. 2013. MiRNA-20 and miRNA -106a regulate spermatogonial stem cell renewal at the post-transcriptional level via targeting STAT3 and Ccnd1. *Stem Cells*, **31**: 2205–2217.
- 13) Jiang, Z. and Rothschild, M. F. 2007. Swine genome science comes of age. *Int. J. Biol. Sci.*, **3**: 129–131.
- 14) Jin, S., Choi, H., Kwon, J. T., Kim, J., Jeong, J., Kim, J., Ham, S., Cho, B. N., Yoo, Y. J. and Cho, C. 2015. Identification and characterization of reproductive KRAB-ZF genes in mice. *Gene*, **565**: 45–55.
- 15) Jørgensen, S., Elvers, I., Trelle, M. B., Menzel, T., Eskildsen, M., Jensen, O. N., Helleday, T., Helin, K. and Sørensen, C. S. 2007. The histone methyltransferase SET8 is required for S-phase progression. *J. Cell*

- Biol.*, **179**: 1337–1345.
- 16) Kan, Z., Garrett-Engele, P. W., Johnson, J. M. and Castle, J. C. 2005. Evolutionarily conserved and diverged alternative splicing events show different expression and functional profiles. *Nucleic Acids Res.*, **33**: 5659–5666.
  - 17) Kanatsu-Shinohara, M. and Shinohara, T. 2013. Spermatogonial stem cell self-renewal and development. *Annu. Rev. Cell Dev. Biol.*, **29**: 163–187.
  - 18) Kelemen, O., Convertini, P., Zhang, Z., Wen, Y., Shen, M., Falaleeva, M. and Stamm, S. 2013. Function of alternative splicing. *Gene*, **514**: 1–30.
  - 19) Kelemen, G. and Márk, M. 2013. E. M. Jellinek's silenced and silencing transgenerational story. *Psychiatr. Hung.*, **28**: 349–369. (in Hungarian).
  - 20) Kerr, C. A., Bunter, K. L., Seymour, R., Shen, B. and Reverter, A. 2005. The heritability of the expression of two stress-regulated gene fragments in pigs. *J. Anim. Sci.*, **83**: 1753–1765.
  - 21) Kolasinska-Zwierz, P., Down, T., Latorre, I., Liu, T., Liu, X. S. and Ahringer, J. 2009. Differential chromatin marking of introns and expressed exons by H3K36me3. *Nat. Genet.*, **41**: 376–381.
  - 22) Lan, X. L., Cretney, E. C., Kropp, J., Khateeb, K., Berg, M. A., Peñagaricano, F., Magness, R., Radunz, A. E. and Khatib, H. 2013. Maternal diet during pregnancy induces gene expression and DNA methylation changes in fetal tissues in sheep. *Front. Genet.*, **4**: 49.
  - 23) Li, M., Sun, X., Hua, L., Huang, Y., Wang, J., Cao, X., Zhan, Z., Lan, X. and Chen, H. 2013. Molecular characterization, alternative splicing and expression analysis of bovine DBC1. *Gene*, **527**: 689–693.
  - 24) Li, Z., Nie, F., Wang, S. and Li, L. 2011. Histone H4 Lys 20 monomethylation by histone methylase SET8 mediates wnt target gene activation. *Proc. Natl. Acad. Sci. U. S. A.*, **108**: 3116–3123.
  - 25) Liu, W., Tanasa, B., Tyurina, O. V., Zhou, T. Y., Gassmann, R., Liu, W. T., Ohgi, K. A., Benner, C., Garcia-Bassets, I., Aggarwal, A. K., Desai, A., Dorrestein, P. C., Glass, C. K. and Rosenfeld, M. G. 2010. PHF8 mediates histone H4 Lysine 20 demethylation events involved in cell cycle progression. *Nature*, **466**: 508–512.
  - 26) Livak, K. J. and Schmittgen, T. D. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, **25**: 402–408.
  - 27) Lois, S., Blanco, N., Martínez-Balbás, M. and de la Cruz, X. 2007. The functional modulation of epigenetic regulators by alternative splicing. *BMC Genomics*, **8**: 252.
  - 28) Luco, R. F., Pan, Q., Tominaga, K., Blencowe, B. J., Pereira-Smith, O. M. and Misteli, T. 2010. Regulation of alternative splicing by histone modifications. *Science.*, **327**: 996–1000.
  - 29) Malik, J., Getman, M. and Steiner, L. A. 2015. Histone methyltransferase Setd8 represses Gata2 expression and regulates erythroid maturation. *Mol. Cell Biol.*, **35**: 2059–2072.
  - 30) Meng, F., Cheng, S., Ding, H., Liu, S., Liu, Y., Zhu, K., Chen, S., Lu, J., Xie, Y., Li, L., Liu, R., Shi, Z., Zhou, Y., Liu, Y. C., Zheng, M., Jiang, H., Lu, W., Liu, H. and Luo, C. 2015. Discovery and optimization of novel, selective histone methyltransferase SET7 inhibitors by pharmacophore- and docking-based virtual screening. *J. Med. Chem.*, **58**: 8166–8181.
  - 31) Nishioka, K., Rice, J. C., Sarma, K. and Erdjument-Bromage, H., Werner, J., Wang, Y., Chuikov, S., Valenzuela, P., Tempst, P., Steward, R., Lis, J. T., Allis, C. D. and Reinberg, D. 2002. PR-Set7 is a nucleosome-specific methyltransferase that modifies Lysine 20 of Histone H4 and is associated with silent chromatin. *Mol. Cell*, **9**: 1201–1213.
  - 32) Oda, H., Okamoto, I., Murphy, N., Chu, J., Price, S. M., Shen, M. M., Torres-Padilla, M. E., Heard, E. and Reinberg, D. 2009. Monomethylation of Histone H4-lysine 20 is involved in chromosome structure and stability and is essential for mouse development. *Mol. Cell Biol.*, **29**: 2278–2295.
  - 33) Park, S. J., Kwon, S. G., Hwang, J. H., Park, D. H., Kim, T. W. and Kim, C. W. 2015. Selection of appropriate reference genes for RT-qPCR analysis in Berkshire, Duroc, Landrace, and Yorkshire pigs. *Gene*, **558**: 152–158.
  - 34) Paucullo, A., Cosenza, G., D'Avino, A., Colimoro, L., Nicodemo, D., Coletta, A., Feligini, M., Marchitelli, C., Di Berardino, D. and Ramunno, L. 2010. Sequence analysis and genetic variability of Stearoyl CoA Desaturase (SCD) gene in the italian mediterranean river buffalo. *Mol. Cell Probes*, **24**: 407–410.
  - 35) Qian, C. and Zhou, M. M. 2006. SET domain protein lysine methyltransferases: structure, specificity and catalysis. *Cell Mol. Life Sci.*,

- 63**: 2755–2763.
- 36) Rea, S., Eisenhaber, F., O’Carroll, D., Strahl, B. D., Sun, Z. W., Schmid, M., Opravil, S., Mechtler, K., Ponting, C. P., Allis, C. D. and Jenuwein, T. 2000. Regulation of chromatin structure by site-specific Histone H3 methyltransferases. *Nature*, **406**: 593–599.
- 37) Rizzardi, L. F., Coleman, K. E., Varma, D., Matson, J. P., Oh, S. and Cook, J. G. 2015. CDK1-dependent inhibition of the E3 ubiquitin ligase CRL4CDT2 ensures robust transition from S phase to mitosis. *J. Biol. Chem.*, **290**: 556–567.
- 38) Stamm, S., Ben-Ari, S., Rafalska, I., Tang, Y., Zhang, Z., Toiber, D., Thanaraj, T. and Soreq, H. 2005. Function of alternative splicing. *Gene*, **344**: 1–20.
- 39) Schotta, G. 2011. H4K20 monomethylation faces the WNT. *Proc. Natl. Acad. Sci. U. S. A.*, **108**: 3097–3098.
- 40) Takashima, S., Takehashi, M., Lee, J., Chuma, S., Okano, M., Hata, K., Suetake, I., Nakatsuji, N., Miyoshi, H., Tajima, S., Tanaka, Y., Toyokuni, S., Sasaki, H., Kanatsu-Shinohara, M. and Shinohara, T. 2009. Abnormal DNA methyltransferase expression in mouse germline stem cells results in spermatogenic defects. *Biol. Reprod.*, **81**: 155–164.
- 41) Takawa, M., Cho, H. S., Hayami, S., Toyokawa, G., Kogure, M., Yamane, Y., Iwai, Y., Maejima, K., Ueda, K., Masuda, A., Dohmae, N., Field, H. I., Tsunoda, T., Kobayashi, T., Akasu, T., Sugiyama, M., Ohnuma, S., Atomi, Y., Ponder, B. A., Nakamura, Y. and Hamamoto, R. 2012. Histone Lysine methyltransferase SETD8 promotes carcinogenesis by deregulating PCNA expression. *Cancer Res.*, **72**: 3217–3227.
- 42) Uddin, M. J., Cinar, M. U., Tesfaye, D., Looft, C., Tholen, E. and Schellander, K. 2011. Age-related changes in relative expression stability of commonly used housekeeping genes in selected porcine tissues. *BMC Res. Notes*, **4**: 441.
- 43) Wang, H., Jiang, M., Bi, H., Chen, X., He, L., Li, X. and Wu, J. 2014. Conversion of female germline stem cells from neonatal and prepubertal mice into pluripotent stem cells. *J. Mol. Cell Biol.*, **6**: 164–171.
- 44) Yang, H. C., Chuang, J. Y., Jeng, W. Y., Liu, C. I., Wang, A. H., Lu, P. J., Chang, W. C. and Hung, J. J. 2014. Pin1-mediated Sp1 phosphorylation by CDK1 increases Sp1 stability and decreases its DNA-binding activity during mitosis. *Nucleic Acids Res.*, **42**: 13573–13587.
- 45) Yeh, J. R., Zhang, X. and Nagano, M. C. 2011. Wnt5a is a cell-extrinsic factor that supports self-renewal of mouse spermatogonial stem cells. *J. Cell Sci.*, **124**: 2357–2366.
- 46) Yeh, J. R., Zhang, X. and Nagano, M. C. 2012. Indirect effects of Wnt3a/ $\beta$ -catenin signalling support mouse spermatogonial stem cells in vitro. *PLoS One*, **7**: e40002.
- 47) Zhang X., Zhou Y., Pan C., Lei C., Dang R., Chen H. and Lan X. 2015. Novel alternative splice variants of NFIX and their diverse mRNA expression patterns in dairy goat. *Gene*, **569**: 250–258.
- 48) Zheng Y., Zhang Y., Qu R., He Y., Tian X. and Zeng W. 2014. Spermatogonial Stem Cells from Domestic Animals: Progress and Prospects. *Reproduction*, **147**: R65–R74.