

Efficient divergent synthesis of new immunostimulant 4''-modified α -galactosylceramide analogues.

Jonas Janssens,^{a,b} Tine Decruy,^{c,d} Koen Venken,^{c,d} Toshiyuki Seki,^{e,f} Simon Krols,^a Johan Van der Eycken,^b Moriya Tsuji,^e Dirk Elewaut,^{c,d} and Serge Van Calenbergh.^{a,*}

^aLaboratory for Medicinal Chemistry, Department of Pharmaceutics (FFW), Ghent University, Ottergemsesteenweg 460, B-9000 Ghent (Belgium)

^bLaboratory for Organic and Bioorganic Synthesis, Department of Organic and Macromolecular Chemistry, Ghent University, Krijgslaan 281 (S4), B-9000 Ghent (Belgium)

^cDepartment of Internal Medicine, Faculty of Medicine and Health Sciences, Ghent University, De Pintelaan 185, B-9000 Ghent (Belgium)

^dVIB Inflammation Research Center, Ghent University, B-9000 Ghent (Belgium)

^eAaron Diamond AIDS Research Center, The Rockefeller University, 1230 York Avenue, New York, NY 10065 (USA)

^fDepartment of Obstetrics and Gynecology, The Jikei University School of Medicine, 3-25-8 Nishi-Shimbashi, Minato-ku, Tokyo 105-8461 (Japan)

Keywords: α -galactosylceramide, KRN7000, CD1d, Th1/Th2, cytokine secretion, immunomodulators.

ABSTRACT: A synthesis strategy for the swift generation of 4''-modified α -galactosylceramide (α -GalCer) analogues is described, establishing a chemical platform to comprehensively investigate the structure-activity relationships (SAR) of this understudied glycolipid part. The strategy relies on a late-stage reductive ring-opening of a *p*-methoxybenzylidene (PMP) acetal to regioselectively liberate the 4''-OH position. The expediency of this methodology is demonstrated by the synthesis of a small yet diverse set of analogues, which were tested for their ability to stimulate invariant natural killer T-cells (iNKT) *in vitro* and *in vivo*. The introduction of a *p*-chlorobenzyl ether yielded an analogue with promising immunostimulating properties, paving the way for further SAR studies.

For over two decades, α -galactosylceramide (α -GalCer or KRN7000; **1**) has been serving as a lead structure for the development of new glycosphingolipids targeting the immune system.¹⁻⁴ α -GalCer is a synthetic glycolipid resulting from the structural optimization of agelasphins, a class of amphiphilic natural products isolated from the marine sponge *Agelas mauritanus*.^{5,6} It is composed of a polar D-galactose unit, α -anomerically linked to a lipophilic ceramide tail. This ceramide, in turn, is built from D-ribo-phytosphingosine, which is *N*-acylated at C2' with cerotic acid (hexacosanoic acid).

α -GalCer binds to the major histocompatibility complex (MHC) class I-like glycoprotein CD1d, associated with the membrane of antigen-presenting cells (APCs).⁷ This binary CD1d-glycolipid complex is presented to the T-cell receptor (TCR) of invariant natural killer T-cells (iNKT cells), evoking simultaneous release of T-helper 1 (Th1) and T-helper 2 (Th2) cytokines.^{8,9} The pro-inflammatory Th1-cytokines, such as interferon- γ (IFN- γ), are involved in anti-tumor, antiviral and antibacterial effects, whereas the anti-inflammatory Th2-cytokines, such as interleukin 4 (IL-4),

counteract the development of autoimmune diseases. However, the antagonizing effect of both cytokine types severely hampers clinical potential of α -GalCer as an immunomodulator. This renders research towards new analogues with an improved immunological profile, slanting towards secretion of either Th1- or Th2-cytokines, highly relevant.





Figure 1. Structural formula of **1** and some reported 4''-analogues.

The list of new α -GalCer analogues continues to grow, with much attention being devoted to modifications of the

hydroxyl groups of the galactose unit, thereby gradually revealing more of the SAR of this key glycolipid part. Modifications at the 2''-position lead to a complete disappearance of antigenicity, due to abrupt disturbance of a vital hydrogen bonding interaction with Gly96 α of the TCR.^{10,11} A sulfate group can be substituted for the 3''-OH-group.¹² The SAR of the 6''-position has been extensively studied, mainly for two reasons. First, the 6''-OH is the only primary hydroxyl group of the sugar part, which from a chemoselective point of view implies that straightforward modification of this position is possible. Indeed, our laboratory has already explored the mild yet powerful regioselective benzylidene ring-opening route to access the 6''-position for the synthesis of fucosyl, galacturonic acid, triazole, carbamate and urea analogues.¹³⁻¹⁵ Second, and more importantly, this hydroxyl group does allow for modifications due to the absence of any major interactions with either CD1d or the TCR,¹⁶ and this has given rise to an array of new immunoreactive analogues.¹⁷⁻²²

In contrast to this, only few 4''-analogues of α -GalCer are known (Figure 1) and, as a consequence, the SAR of this position has been poorly investigated. Crystallographic studies have shown that the 4''-OH forms a hydrogen bond to the main chain carbonyl group of Phe29 α of the TCR, indicating that it acts as a hydrogen bond donor.¹⁶ 4''-Deoxygenation leads to a less active analogue (**2**) as compared to α -GalCer, yet recognition by the TCR is not severely disturbed.²³ Derivatization of the 4''-OH as *O*-methyl ether (**3**) or *O*-ethanol ether (**4**) gives slightly less potent analogues, whereas the *N*-acetyl analogue (**5**) is a significantly weaker antigen.²⁴ Additionally, a couple of active analogues bearing an aromatic ring on the 4''-position have been synthesized (**6-8**).²⁵ Inversion of the 4''-OH gives α -glucosylceramide (α -GlcCer), which is slightly less active as compared to α -GalCer.⁷

Scheme 1. Synthesis of trichloroacetimidate donor **13**.^a



^aReagents and conditions: (a) *p*-thiocresol, $\text{BF}_3 \cdot \text{Et}_2\text{O}$, CH_2Cl_2 , 0 °C (94%); (b) NaOMe, MeOH, rt (quant.); (c) *p*-anisaldehyde dimethylacetal, CSA, CH_2Cl_2 , 4Å MS, rt (87%); (d) NaH, PMBCl, TBAI, DMF, 0 °C to rt (98%); (e) NIS, MeCN, H_2O , 0 °C (99%); (f) Cl_3CCN , DBU, CH_2Cl_2 , 0 °C (94%).

To explore the SAR more thoroughly, a reliable and scalable synthesis route is highly needed. Here, we present the synthesis of two powerful precursors (**16** and **19**) with a free 4''-OH, permitting the fast generation of new 4''- α -GalCer-analogues in a late stage of the synthesis. This divergent


strategy uses PMB ethers as hydroxyl protecting groups, in contrast to most other syntheses, which use benzyl groups. Although they are widely used because of their easy introduction and relative inertness towards a plethora of conditions, benzyl groups suffer from the major drawback that by deprotection *via* catalytic hydrogenolysis some medicinally interesting functionalities, such as alkenes, alkynes, thioethers, naphthalenes, (iso)quinolines, furans, thiophenes, cyclopropanes and chloroarenes, might be (partly) reduced. This unavoidably restricts the structural diversity of the analogues. PMB ethers, on the other hand, can be cleaved under mild and widely tolerated conditions, therefore permitting a broader substrate scope.²⁶

The previously reported trichloroacetimidate donor (**13**)²⁷ was synthesized *via* an improved scalable route from cheap α -D-galactose pentaacetate (**9**) (Scheme 1). Introduction of the *p*-thiitolyl moiety as an anomeric protecting group and subsequent deacetylation under Zemplén conditions was followed by installation of the *p*-methoxybenzylidene acetal and protection of the remaining hydroxyl groups as PMB ethers to yield the fully protected intermediate **11**. Attempts to remove the *p*-thiitolyl group with *N*-iodosuccinimide (NIS) in acetone/ H_2O 10:1 resulted in partial conversion of the *p*-methoxybenzylidene acetal to an isopropylidene acetal. This inconvenience was overcome by performing the reaction in acetonitrile/ H_2O 10:1, giving hemiacetal **12** as a mixture of anomers (α : β ~ 6:4 at 25 °C in CDCl_3) in a nearly quantitative yield. Finally, **12** was converted to the corresponding α -trichloroacetimidate under thermodynamic control, furnishing donor **13** in an overall yield of 75% over 6 steps. All intermediates, except for **13**, can be purified by simple crystallization from a suitable solvent system.

Efforts were undertaken to perform the glycosylation of the known acceptor **14**²⁸ with the thioglycoside donor **11**, as this would shorten the synthesis route. However, the application of neither benzenesulfinyl morpholine (BSM),²⁹ nor NIS or copper(II) triflate³⁰ as thiophilic promoters proved to be successful. These highly electrophilic conditions, typically used in thioglycosylations,³¹ were found to be incompatible with the *p*-methoxybenzylidene acetal. Indeed, *p*-methoxybenzylidene and *p*-thiitolyl cleavage were observed as the main side reactions.

Glycosylation was successful at the trichloroacetimidate stage through application of the “inverse protocol” (Scheme 2).³² By slow addition of donor **13** to a solution of acceptor **14** and $\text{BF}_3 \cdot \text{OEt}_2$ as the promoter at -20 °C, the desired glycoside **15** was obtained as α -anomer only in 85% yield. The use of trimethylsilyl trifluoromethanesulfonate (TMSOTf) as glycosylation promoter led to silylation of the acceptor, while the common protocol with $\text{BF}_3 \cdot \text{OEt}_2$, wherein the promoter is added to a solution of donor and acceptor, gave the glycoside in lower yields (30-50%) due to more extensive decomposition of the donor before glycosylation. The high stereoselectivity observed in this reaction is caused by a well-known conformational effect exhibited by the 4'',6''-*O*-acetal, forming a *cis*-decalin-like system with the galactose ring.³³


Scheme 2. Glycosylation of **13 with **14** and subsequent regioselective ring-opening of glycoside **15**.^a**



^aReagents and conditions: (a) **14**, BF₃·Et₂O, THF, Et₂O, 4 Å MS, -20 °C (85%); (b) DIBALH, toluene, -80 °C (89%); (c) BaO, Ba(OH)₂, MeI, DMF, rt (93%).

Next, attempts were made to regioselectively open the *p*-methoxybenzylidene acetal in order to liberate the 4''-OH function. Experiments involving BH₃·THF/Cu(OTf)₂,³⁴ BH₃·THF/*n*Bu₂BOTf³⁵ or PhBCl₂/TESH³⁶ as reducing agents mainly gave rise to degradation of the *p*-methoxybenzylidene acetal and the glycosidic bond. A more successful approach was the use of diisobutylaluminium hydride (DIBALH) in toluene,³⁷ while keeping the temperature at -80 °C to avoid azide reduction (Scheme 2). This afforded azido alcohol **16** as a single regioisomer in 89% yield. The structure of **16** was unambiguously proven by derivatization as the corresponding methyl ether (**17**) and analysis of the relevant cross peaks in the HSQC and HMBC spectra (see Supporting Information).

Scheme 3. Amide formation of **16.^a**




^aReagents and conditions: (a) (i) H₂S, pyridine, H₂O, rt; (ii) **18**, Et₃N, THF, 70 °C (86% over 2 steps).

Azido alcohol **16** was subjected to azide reduction under the classical Staudinger conditions with PMe₃, followed by EDC-mediated amide formation with hexacosanoic acid (cerotic acid). However, it was observed that the intermediate iminophosphorane was highly stable and even with concentrated sodium hydroxide at elevated temperatures its hydrolysis proceeded sluggishly, providing amide **19** in a rather low yield (43% over 2 steps). To reduce the azide in a more efficient way, we turned our attention to hydrogen sulfide in a pyridine/H₂O mixture. This readily furnished the amine, which was immediately acylated with *N*-succinimidyl hexacosanoate (NSHC, **18**) under basic conditions to deliver amide **19** in 86% yield over 2 steps

(Scheme 3). To drive the reaction to completion within a reasonable amount of time, the amide formation was carried out at 70 °C.

Both **16** and **19** were now suited for late-stage diversification. Alkylation of the 4''-OH had to be performed at the azide stage, since methylation of **19** with methyl iodide and BaO/Ba(OH)₂ in DMF yielded an unseparable mixture of *O*- and *N*-alkylated products in a 4:1 ratio (as determined via ¹H NMR spectroscopy). Thus, alkylation of **16** with *p*-chlorobenzyl bromide gave derivative **22** (Scheme 4). Both **17** and **22** were subjected to H₂S-mediated azide reduction and subsequent amide formation. The final PMB ether cleavage was performed using HCl in 1,4-dioxane, swiftly delivering analogues **21** and **24**. Anisole was added in large excess (10 equiv.) as a scavenger for the highly reactive *p*-methoxybenzyl carbocation.


Scheme 4. Synthesis of *O*-alkylated analogues **21 and **24**.^a**



^aReagents and conditions: (a) (i) H₂S, pyridine, H₂O, rt; (ii) **18**, Et₃N, THF, 70 °C (78% over 2 steps); (b) HCl, 1,4-dioxane, anisole, rt (32%); (c) *p*-chlorobenzyl bromide, BaO, Ba(OH)₂, DMF, rt (92%); (d) (i) H₂S, pyridine, H₂O, rt; (ii) **18**, Et₃N, THF, 70 °C (66% over 2 steps); (e) HCl, 1,4-dioxane, anisole, rt (30%).

Besides alkylation, some other chemistries were explored to create a diverse set of analogues (Scheme 5). Carbamylation of **19** with 1-naphthyl isocyanate delivered, after deprotection, naphthyl carbamate analogue **26**. This analogue structurally resembles the 6''-naphthyl urea, which exerts a strong Th1-bias as a result of the naphthyl moiety occupying an additional binding pocket in CD1d.¹⁸ Oxidation of the alcohol in **16** using a Swern oxidation delivered the corresponding ketone, which, without intermediate purification, smoothly underwent a mild Julia-Kocienski olefination with 1-methyl-2-(methylsulfonyl)-1*H*-benzo[*d*]imidazole (MSBI, **27**)³⁸ to give the exocyclic methylene derivative. Following azide reduction, amide formation and overall deprotection, analogue **29** was obtained. Finally, the OH-group was converted into the corresponding azide **30** via a DPPA-mediated Mitsunobu reaction. The azide was reduced and derivatized as the corresponding naphthyl urea, eventually delivering analogue **32**.

Scheme 5. Synthesis of naphthyl analogues 26 and 32, and alkenyl compound 29.^a



^aReagents and conditions: (a) 1-naphthyl isocyanate, DMF, rt (89%); (b) HCl, 1,4-dioxane, anisole, rt (44%); (c) (i) $(\text{COCl})_2$, DMSO, Et_3N , CH_2Cl_2 , -78°C ; (ii) **27**, KOtBu , DMF, rt; (iii) H_2S , pyridine, H_2O , rt; (iv) **18**, Et_3N , THF, 70°C (61% over 4 steps); (d) HCl, 1,4-dioxane, anisole, rt (31%); (e) DPPA, DEAD, PPh_3 , THF, -20°C to rt (74%); (f) (i) H_2S , pyridine, H_2O , rt; (ii) 1-naphthyl isocyanate, DMF, rt (22% over 2 steps); (g) HCl, 1,4-dioxane, anisole, rt (48%).

The set of new analogues (**21**, **24**, **26**, **29** and **32**) was evaluated for its immunostimulating capacity, both *in vitro* and *in vivo*. C57BL/6 mice were injected intraperitoneally with the glycolipids (5 μg of each analogue: 10 nM for **24**, **26** and **32**, 11 nM for **21** and **29**) and the blood serum levels of the cytokines were measured by ELISA (Figure 2). IFN- γ was quantified 16 h after injection, whereas IL-4 was measured 4 h after injection. All of the analogues were able to stimulate the immune system, albeit at levels comparable to or lower than those elicited by α -GalCer (**1**). Although the 4''-OH acts as a hydrogen bond donor to interact with the TCR, derivatization as a simple ether is allowed. The antigenic effect is most dramatic when introducing a methyl group (**21**). The *p*-chlorobenzyl analogue **24** was found to polarize the cytokine response towards Th₁, although its antigenicity was lower than that of α -GalCer. Yet, we believe that this analogue might be a good lead structure for future optimization towards highly potent Th₁-polarizing α -GalCer analogues.

In the same way, introducing a *gluco*-naphthyl urea on the 4''-position (**32**) yields an analogue that is Th₁-polarizing. Future structural studies might shed light on the mechanism underlying this Th₁-polarization, but enhanced interaction with CD1d may account for this observation. The *galacto*-naphthyl carbamate (**26**), in which the naphthyl group is unlikely to interact with CD1d due to its axial configuration, is significantly less potent.

Remarkably, even when no heteroatom is present at C₄'', as exemplified by the exocyclic alkene (**29**), immunostimulatory capacity is partly preserved. Apparently, the exocyclic alkene, which will partly flatten the galactose ring, induces a pyranose conformation that is still able to bridge between CD1d and the TCR. Further structural studies are required to gain more insight into this phenomenon.




Figure 2. IFN- γ and IL-4 secretion after intraperitoneal injection of 5 μg of each analogues in C57BL/6 mice.

To assess if these analogues also stimulate human *i*NKT cells, the latter were co-cultured with HeLa CD1d cells with varying concentrations of the glycolipids. After 24 h of incubation the IFN- γ levels were determined by ELISA (Figure 3). These results reflect the observations in mice, namely that all of the compounds are able to stimulate *i*NKT cells, with the methyl ether being most antigenic.




Figure 3. IFN- γ levels measured upon incubation of different concentrations of the analogues with human iNKT and HeLa CD1d cells.

In summary, we have developed a concise and scalable synthetic route towards two valuable precursors (**16** and **19**) to perform late-stage diversification of the 4''-OH position of α -GalCer. In this strategy, a triple role has been provided for the 4'',6''-*O*-*p*-methoxybenzylidene acetal: as a protecting group, as a stereocontrolling element during glycosylation and as a structural element to enable a regioselective ring-opening. We have demonstrated the utility of this method by synthesizing a diverse set of analogues, that was tested for the ability to stimulate iNKT cells both *in vitro* and *in vivo*. In both cases, the analogues were able to stimulate iNKT cells, some of them showing a clear pro-inflammatory activity due to Th1-polarization. Even though the 4''-OH is involved in binding to the TCR, we have shown that derivatization is allowed, which should inspire further biological and structural research to comprehensively explore the SAR of this glycolipid part.

ASSOCIATED CONTENT

Supporting Information

The supporting information is available free of charge on the ACS Publications website at DOI: XXX.

Experimental details and characterization data for the reported compounds, NMR spectra, and biological data (PDF).

AUTHOR INFORMATION

Corresponding Author

*E-mail: Serge.VanCalenbergh@UGent.be.

Funding Sources

Ghent University is gratefully acknowledged for financial support to this research, through a Special Research Fund grant to JJ. The work in the laboratory of MT is supported by the NIH (grant A1070258).

ACKNOWLEDGMENTS

Tim Courtin from the NMR & Structural Analysis Unit of Ghent University is thanked for recording the NMR spectra of the final compounds. Izet Karalic is gratefully acknowledged for purity analysis of the final compounds. Joren Guillaume and dr. Martijn D. P. Risseuw are acknowledged for valuable discussions.


ABBREVIATIONS

APC, antigen-presenting cell; CSA, (1*S*)-(+)-10-camphorsulfonic acid; DBU, 1,8-diazabicyclo(5.4.0)undec-7-ene; DEAD, diethyl azodicarboxylate; DIBALH, diisobutylaluminium hydride; DPPA, diphenylphosphoryl azide; EDC, *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide; ELISA, enzyme-linked immunosorbent assay; IFN- γ , interferon- γ ; IL, interleukin; MHC, major histocompatibility complex; MS, molecular sieves; MSBI, 1-methyl-2-(methylsulfonyl)-1*H*-benzo[*d*]imidazole; NIS, *N*-iodosuccinimide; NKT, natural killer T-cell; NSHC, *N*-succinimidyl hexacosanoate; PMB, *p*-methoxybenzyl; PMP, *p*-methoxyphenyl; SAR, structure-activity relationship; TBAl, tetra-*n*-butylammonium iodide; TCR, T-cell receptor; TESH, triethylsilane; Th, T-helper.

REFERENCES

- Murphy, N.; Zhu, X.; Schmidt, R. R. α -Galactosylceramide and analogues – important immunomodulators for use as vaccine adjuvants. *Carbohydr. Chem.* **2010**, *36*, 64-100.
- Banchet-Cadeddu, A.; Hénon, E.; Dauchez, M.; Renault, J.-H.; Monneaux, F.; Haudrechy, A. The stimulating adventure of KRN 7000. *Org. Biomol. Chem.* **2011**, *9*, 3080-3104.
- Birkholz, A. M.; Howell, A. R.; Kronenberg, M. The alpha and omega of galactosylceramides in T cell immune function. *J. Biol. Chem.* **2015**, *290*, 15365-15370.
- Carreño, L. J.; Saavedra-Ávila, N. A.; Porcelli, S. A. Synthetic glycolipid activators of natural killer T cells as immunotherapeutic agents. *Clin. Trans. Immunol.* **2016**, *5*, e69; DOI: 10.1038/cti.2016.14.
- Natori, T.; Morita, M.; Akimoto, K.; Koezuka, Y. Agelasphins, novel antitumor and immunostimulatory cerebroside from the marine sponge *Agelas mauritanus*. *Tetrahedron* **1994**, *50*, 2771-2784.
- Morita, M.; Motoki, K.; Akimoto, K.; Natori, T.; Sakai, T.; Sawa, E.; Yamaji, K.; Koezuka, Y.; Kobayashi, E.; Fukushima, H. Structure-activity relationship of α -galactosylceramides against B16-bearing mice. *J. Med. Chem.* **1995**, *38*, 2176-2187.
- Kawano, T.; Cui, J.; Koezuka, Y.; Toura, I.; Kaneko, Y.; Motoki, K.; Ueno, H.; Nakagawa, R.; Sato, H.; Kondo, E.; Koseki, H.; Taniguchi, M. CD1d-Restricted and TCR-mediated activation of V α 14 NKT cells by glycosylceramides. *Science* **1997**, *278*, 1626-1629.
- Coquet, J. M.; Chakravarti, S.; Kyriassoudis, K.; McNab, F. W.; Pitt, L. A.; McKenzie, B. S.; Berzins, S. P.; Smyth, M. J.; Godfrey, D. I. Diverse cytokine production by NKT cell subsets and identification of an IL-17-producing CD4⁺NK1.1⁺ NKT cell population. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 11287-11292.
- Vartabedian, V. F.; Savage, P. B.; Teyton, L. The processing and presentation of lipids and glycolipids to the immune system. *Immunol. Rev.* **2016**, *272*, 109-119.
- Barbieri, L.; Costantino, V.; Fattorusso, E.; Mangoni, A.; Aru, E.; Parapini, S.; Taramelli, D. Immunostimulatory α -galactoglycosphingolipids: synthesis of a 2'-*O*-methyl- α -Gal-GSL and evaluation of its immunostimulating capacity. *Eur. J. Org. Chem.* **2004**, *2004*, 468-473.
- Barbieri, L.; Costantino, V.; Fattorusso, E.; Mangoni, A.; Basilico, N.; Mondani, M.; Taramelli, D. Immunomodulatory α -galactoglycosphingolipids: synthesis of 2'-fluoro-2'-deoxy- α -galactosylceramide and an evaluation of its immunostimulating properties. *Eur. J. Org. Chem.* **2005**, *2005*, 3279-3285.
- Xing, G.-W.; Wu, D.; Poles, M. A.; Horowitz, A.; Tsuji, M.; Ho, D. D.; Wong, C.-H. Synthesis and human NKT cell

- stimulating properties of 3-O-sulfo- α/β -galactosylceramides. *Bioorg. Med. Chem.* **2005**, *13*, 2907-2916.
13. Pauwels, N.; Aspeslagh, S.; Vanhoenacker, G.; Sandra, K.; Yu, E. D.; Zajonc, D. M.; Elewaut, D.; Linclau, B.; Van Calenbergh, S. Divergent synthetic approach to 6"-modified α -GalCer analogues. *Org. Biomol. Chem.* **2011**, *9*, 8413-8421.
 14. Pauwels, N.; Aspeslagh, S.; Elewaut, D.; Van Calenbergh, S. Synthesis of 6"-triazole-substituted α -GalCer analogues as potent iNKT cell stimulating ligands. *Bioorg. Med. Chem.* **2012**, *20*, 7149-7154.
 15. Guillaume, J.; Pauwels, N.; Aspeslagh, S.; Zajonc, D. M.; Elewaut, D.; Van Calenbergh, S. Synthesis of C-5" and C-6"-modified α -GalCer analogues as iNKT-cell agonists. *Bioorg. Med. Chem.* **2015**, *23*, 3175-3182.
 16. Borg, N. A.; Wun, K. S.; Kjer-Nielsen, L.; Wilce, M. C. J.; Pellicci, D. G.; Koh, R.; Besra, G. S.; Bharadwaj, M.; Godfrey, D. I.; McCluskey, J.; Rossjohn, J. CD1d-lipid-antigen recognition by the semi-invariant NKT T-cell receptor. *Nature* **2007**, *448*, 44-49.
 17. Zhou, X.-T.; Forestier, C.; Goff, R. D.; Li, C.; Teyton, L.; Bendelac, A.; Savage, P. B. Synthesis and NKT cell stimulating properties of fluorophore- and biotin-appended 6"-amino-6"-deoxy-galactosylceramides. *Org. Lett.* **2002**, *4*, 1267-1270.
 18. Trappeniers, M.; Van Beneden, K.; Decruy, T.; Hillaert, U.; Linclau, B.; Elewaut, D.; Van Calenbergh, S. 6'-derivatised α -GalCer analogues capable of inducing strong CD1d-mediated Th1-biased NKT cell responses in mice. *J. Am. Chem. Soc.* **2008**, *130*, 16468-16469.
 19. Hsieh, M.-H.; Hung, J.-T.; Liw, Y.-W.; Lu, Y.-J.; Wong, C.-H.; Yu, A. L.; Liang, P.-H. Synthesis and evaluation of acyl-chain- and galactose-6"-modified analogues of α -GalCer for NKT cell activation. *ChemBioChem* **2012**, *13*, 1689-1697.
 20. Jervis, P. J.; Graham, L. M.; Foster, E. L.; Cox, L. R.; Porcelli, S. A.; Besra, G. S. New CD1d agonists: synthesis and biological activity of 6"-triazole-substituted α -galactosyl ceramides. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 4348-4352.
 21. Hung, J.-T.; Sawant, R. C.; Chen, J.-C.; Yen, Y.-F.; Chen, W.-S.; Yu, A. L.; Luo, S.-Y. Design and synthesis of galactose-6-OH-modified α -galactosyl ceramide analogues with Th2-biased immune responses. *RSC Adv.* **2014**, *4*, 47341-47356.
 22. Compton, B. J.; Tang, C.-w.; Johnston, K. A.; Osmond, T. L.; Hayman, C. M.; Larsen, D. S.; Hermans, I. F.; Painter, G. F. Synthesis and activity of 6"-deoxy-6"-thio- α -GalCer and peptide conjugates. *Org. Lett.* **2015**, *17*, 5954-5957.
 23. Wun, K. S.; Borg, N. A.; Kjer-Nielsen, L.; Beddoe, T.; Koh, R.; Richardson, S. K.; Thakur, M.; Howell, A. R.; Scott-Browne, J. P.; Gapin, L.; Godfrey, D. I.; McCluskey, J.; Rossjohn, J. A minimal binding footprint on CD1d-glycolipid is a basis for selection of the unique human NKT TCR. *J. Exp. Med.* **2008**, *205*, 939-949.
 24. Florence, W. C.; Xia, C.; Gordy, L. E.; Chen, W.; Zhang, Y.; Scott-Browne, J. P.; Kinjo, Y.; Yu, K. O. A.; Keshipeddy, S.; Pellicci, D. G.; Patel, O.; Kjer-Nielsen, L.; McCluskey, J.; Godfrey, D. I.; Rossjohn, J.; Richardson, S. K.; Porcelli, S. A.; Howell, A. R.; Hayakawa, K.; Gapin, L.; Zajonc, D. M.; Wang, P. G.; Joyce, S. Adaptability of the semi-invariant natural killer T-cell receptor towards structurally diverse CD1d-restricted ligands. *EMBO J.* **2009**, *28*, 3579-3590.
 25. Zhang, W.; Xia, C.; Nadas, J.; Chen, W.; Gu, L.; Wang, P. G. Introduction of aromatic group on 4'-OH of α -GalCer manipulated NKT cell cytokine production. *Bioorg. Med. Chem.* **2011**, *19*, 2767-2776.
 26. Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 2nd ed.; John Wiley & Sons: New York, 1991.
 27. Yamamoto, K.; Sato, Y.; Ishimori, A.; Miyairi, K.; Okuno, T.; Nemoto, N.; Shimizu, H.; Kidokoro, S.; Hashimoto, M. Synthesis of D-trigalacturonic acid methylglycoside and conformational comparison with its sulfur analogue. *Biosci. Biotechnol. Biochem.* **2008**, *72*, 2039-2048.
 28. Du, W.; Gervay-Hague, J. Efficient synthesis of α -galactosyl ceramide analogues using glycosyl iodide donors. *Org. Lett.* **2005**, *7*, 2063-2065.
 29. Lu, Y.-S.; Li, Q.; Zhang, L.-H.; Ye, X.-S. Highly direct α -selective glycosylations of 3,4-O-carbonate-protected 2-deoxy- and 2,6-dideoxythioglycosides by preactivation protocol. *Org. Lett.* **2008**, *10*, 3445-3448.
 30. Hou, D.; Lowary, T. L. 2,3-Anhydrosugars in glycoside bond synthesis. Application to 2,6-dideoxypyranosides. *J. Org. Chem.* **2009**, *74*, 2278-2289.
 31. Lian, G.; Zhang, X.; Yu, B. Thioglycosides in *Carbohydrate Research*. *Carbohydr. Res.* **2015**, *403*, 13-22.
 32. Schmidt, R.; Toepfer, A. Glycosylation with highly reactive glycosyl donors: efficiency of the inverse procedure. *Tetrahedron Lett.* **1991**, *32*, 3353-3356.
 33. Yule, J. E.; Wong, T. C.; Gandhi, S. S.; Qiu, D.; Riopel, M. A.; Koganty, R. R. Steric control of N-acetylgalactosamine in glycosidic bond formation. *Tetrahedron Lett.* **1995**, *36*, 6839-6842.
 34. Shie, C.-R.; Tzeng, Z.-H.; Kulkarni, S. S.; Uang, B.-J.; Hsu, C.-Y.; Hung, S.-C. Cu(OTf)₂ as an efficient and dual-purpose catalyst in the regioselective reductive ring opening of benzylidene acetals. *Angew. Chem. Int. Ed.* **2005**, *44*, 1665-1668.
 35. Hernández-Torres, J. M.; Achkar, J.; Wei, A. Temperature-controlled regioselectivity in the reductive cleavage of p-methoxybenzylidene acetals. *J. Org. Chem.* **2004**, *69*, 7206-7211.
 36. Dilhas, A.; Bonnaffé, D. PhBCl₂ promoted reductive opening of 2',4'-O-p-methoxybenzylidene: new regioselective differentiation of position 2' and 4' of α -L-iduronyl moieties in disaccharide building blocks. *Tetrahedron Lett.* **2004**, *45*, 3643-3645.
 37. Sarpe, V. A.; Kulkarni, S. S. Desymmetrization of trehalose via regioselective DIBAL reductive ring opening of benzylidene and substituted benzylidene acetals. *Org. Biomol. Chem.* **2013**, *11*, 6460-6465.
 38. Ando, K.; Kobayashi, T.; Uchida, N. Practical methylenation reaction for aldehydes and ketones using new Julia-type reagents. *Org. Lett.* **2015**, *17*, 2554-2557.



Efficient divergent synthesis of new immunostimulant 4''-modified α -galactosylceramide analogues

Jonas Janssens, Tine Decruy, Koen Venken, Toshiyuki Seki, Simon Krols, Johan Van der Eycken, Moriya Tsuji, Dirk Elewaut and Serge Van Calenbergh
