



## RESEARCH ARTICLE

# *Camponotus fellah* queens are singly mated

Danielle P. Mersch<sup>1,2</sup>  · Christine La Mendola<sup>2</sup> · Laurent Keller<sup>2</sup>Received: 18 November 2016 / Revised: 17 January 2017 / Accepted: 20 January 2017  
© International Union for the Study of Social Insects (IUSI) 2017

**Abstract** The ant *Camponotus fellah* has been used in several behavioral and life history studies. An important factor that affects the genetic structure and division of labour within a colony is whether queens are singly or multiply mated. To determine whether queens are singly mated in *C. fellah*, as is the case in some other *Camponotus* species, we developed nine polymorphic microsatellite markers and sequenced 16 workers each from 20 colonies at six loci. Data in all colonies were compatible with queen monoandry. All the workers of one of the colonies had identical genotypes suggesting that they were clonally produced or that the queen was inbred. We, therefore, genotyped the mother queen as well as 31 more workers of the same colony at the same six loci plus the three remaining loci. These data revealed that the queen was homozygous at eight of the nine loci and that she mated with a male having a shared allele at all but one of the loci. Thus, the queen was apparently not only inbred but also probably mated with a brother.

**Keywords** *Camponotus fellah* · Microsatellites · Monoandry · Ant · Genetic structure

## Introduction

The genetic structure of ant colonies varies widely between species and can impact colony immunity, reproductive output and division of labor (Crozier and Page 1985; Julian and Cahan 1999; Heinze and Keller 2000; Julian and Fewell 2004; Rheindt et al. 2005; Oldroyd and Fewell 2007; Smith et al. 2008). Several studies have shown that in colonies where queens are multiply mated workers from different patriline vary in their tendency to specialize in different tasks, thereby potentially affecting the efficiency of division of labor and colony homeostasis (Robinson 1992; Rheindt et al. 2005; Schwander et al. 2005; Wierasz et al. 2008; Waddington et al. 2010). Increased within-colony genetic diversity due to multiple patriline and/or matriline may also improve resistance to parasites and pathogens (Hughes and Boomsma 2004; Reber et al. 2008; Schmidt et al. 2011) thereby enhancing the lifetime and reproductive output of a colony. Thus, knowledge of the genetic makeup of colonies provides insights into what factors might shape colony function.

Studies in several species of the genus *Camponotus* (*C. herculeanus*, Gertsch et al. 1995; *C. ligniperdus*, Gertsch et al. 1995, Gadau et al. 1998; *C. floridanus*, Gadau et al. 1996; *C. nawai*, Satoh et al. 1997; *C. consobrinus*, Fraser et al. 2000; *C. ocreatus*, Goodisman and Hahn 2004 and *C. festinatus*, Goodisman and Hahn 2005) revealed that queens are generally singly mated with occasional instances of polyandry. A species of the genus *Camponotus* that has not yet been investigated is *C. fellah*. This species has been extensively used to study nestmate recognition (Boulay et al. 2000a, b; Katzav-Gozansky et al. 2004; Katzav-Gozansky et al. 2008), more recently to unravel the impact of social isolation (Boulay et al. 1999; Koto et al. 2015) and to decipher the organizational principles of division

✉ Danielle P. Mersch  
dmersch@mrc-lmb.cam.ac.uk

✉ Laurent Keller  
laurent.keller@unil.ch

<sup>1</sup> MRC-Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge Biomedical Campus, Cambridge CB1 0QH, UK

<sup>2</sup> Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

of labor (Mersch et al. 2013; Greenwald et al. 2015; Kafsi et al. 2016). The colonies exhibit pronounced division of labor with workers organized in distinct social groups that specialize in different tasks (Mersch et al. 2013). The propensity of a worker to join a social group, and preferentially execute some tasks might thus be dependent on a worker's patriline. We developed and tested a set of nine microsatellite loci to determine whether *C. fellah* queens are singly mated, or whether workers come from different fathers.

## Materials and methods

### Collection of ants

Adult workers were collected from 20 lab-reared colonies (A–T) by opening the nest and then collecting workers randomly both near the brood and outside the nest. Each lab colony was established from a single queen collected during a mating flight in 2007 on the campus of the Tel Aviv University, Tel Aviv, Israel, which is within the natural range of the species (Ionescu-Hirsch 2009).

### Genetic analysis

PCR primers were developed for *C. fellah* microsatellite loci from a partial genomic library that was derived from 10 µg of genomic DNA extracted with the BioSprint 96 robot (QIAGEN #900852) from one worker, and sent to Ecogenics (<http://www.ecogenics.ch/>) for 454 sequencing

and primer design. To determine which of the designed primers were polymorphic we tested nine primers on two workers from each of eight colonies (A–H). Six out of the nine microsatellites had five or more alleles in the workers tested (Table 1). We used these six microsatellites to assess queen-mating frequency in 20 colonies using 16 workers per colony. However, because there was a very high level of homozygosity in colony S, we increased the sample size to 47 workers and also genotyped the queen using one of her legs. In this colony we also genotyped all individuals at the three remaining microsatellites (7275, 7207, 7300).

We extracted DNA from each individual using the BioSprint 96 DNA Blood Kit for tissue sample from QIAGEN and eluted the DNA in a final volume of 200 µl. We performed PCR amplifications for each ant sample in a total volume of 20 µl. To reduce the amount of work and sequencing, we multiplexed primers 1284, 6826 and 5134 in one reaction, and primers 3186, 3675, 4380 in the other. Each reaction consisted of 4 µl of genomic DNA, 4 µl of 10×PCR buffer (QIAGEN), 1.92 µl of 25 mM MgCl<sub>2</sub>, 0.32 µl of 25 mM dNTPs, 1 µl of each forward and reverse primer (10 µM), 0.3 µl of 5 U/µl Taq polymerase (QIAGEN), 3.2 µl of 5×enhancers (Qsolution from QIAGEN), and 0.26 µl H<sub>2</sub>O. The PCR consisted of an initial denaturing step of 5 min at 95 °C, followed by 35 cycles that each consisted of 30 s of denaturation at 95 °C, 30 s of annealing at 55 °C, and 60 s elongation at 72 °C. A final elongation step was performed at 72 °C for 10 min. We analyzed PCR products with the ABIprism3100 DNA sequencer and the GeneMapper Software (Applied Biosystems).

**Table 1** Characteristics of the nine microsatellite loci from *Camponotus fellah* colonies

Locus	Primer sequence (5'–3') (F: forward; R: reverse)	T <sub>a</sub> (°C)	Repeat type	Size of cloned amplicon (bp)	No. of alleles	Amplicon size range
1284	F: GTCGGAATGTTGCACCTACG R: CAGACCGCGAGAGGAGATAC	53.8	(AG)	226	5	230–237
3186	F: AATTGAACGTTTCGTCCGCC R: AGTCTTTACGCCCTCCTACG	51.8	(AC)	190	8	194–206
3675	F: TTGTAGAGCGACGACGAGAG R: TGAGAGGTGAGAGCACGAAG	53.8	(CGT)	203	8	221–252
4380	F: TGGCTACAGTTGTTGTGCAG R: TTTGTCGCGGCTCGTAAATC	54.8	(AC)	115	10	126–148
5134	F: TACAATCGAGTGGACGGGAG R: CAGATGAAAGGCGAACGGTG	53.8	(CGA)	247	9	246–299
6826	F: GTACGCGGTCAGAATTACGC R: ACCTGATTCTGGTTGGCAC	53.8	(AG)	176	14	178–207
7207	F: AGGCATTCGATACTGTCCGGC R: TACGACAAGGATGCTACCGC	53.8	(TGC)	163	2	184–187
7275	F: CGCATCGGGAGCATAAACAG R: AAAGCTGGCATGCAGTTACG	53.8	(CA)	102	5	118–123
7300	F: AATTTGCGGAGAATCCGTGG R: CATCGGCAAAGTCGTCCCTC	51.8	(CT)	185	3	201–205

All forward primers are preceded with an 18-bp M13-tag: 5'-TGTA AACGACGGCCAGT-3'

## Data analysis

We estimated linkage disequilibrium between each pair of loci using the Genepop software (Raymond and Rousset 1995; Rousset 2008). For each colony, we determined the number of worker patriline. This was simplified by the fact that all colonies were lab raised with all workers being the offspring of a single queen. Further, in ants, males are haploid because of the haplodiploid sex determination system of Hymenoptera. Thus, at each locus, a worker inherits one allele from her father and one of the two alleles of the queen. If two males share the same alleles at all genotyped loci, it is impossible to distinguish their offspring. The theoretical probability of such non-detection errors was estimated as:

$$\prod \sum q_i^2$$

with  $q_i$  as the frequency of the  $i$  allele (Boomsma and Ratnieks 1996). The sum is over all alleles at a locus, and the multiplication is over all loci.

We discarded from our patriline analyses 8 out of the 351 workers genotyped. Two samples from colony B were contaminated during the PCR and six workers from four different colonies (colony B: 2 workers, colony G: 2 workers, colony L: 1 worker, colony R: 1 worker) had genotypes incompatible with the genotype inferred for their queen. Because colonies were all lab raised and headed by a single queen, we suspect that these workers were intruders that had escaped from other colonies. Aggression toward non-nestmates is often attenuated in the lab for two reasons. First, colonies, which are kept in proximity to each other, habituate to olfactory volatiles of other colonies, which reduces aggression (Katzav-Gozansky et al. 2008). Second, lab colonies are typically fed with the same diet, which can reduce inter-colony differences in hydrocarbon profiles (Richard et al. 2004, Ichinose et al. 2009).

## Results

The six microsatellite loci were very polymorphic with an average of 7.11 alleles per locus (range: 2–14, Table 1), as estimated from all genotyped workers ( $n=365$ ). There was no sign of gametic linkage among the loci ( $p>0.05$  for all pair-wise comparisons of loci after sequential Bonferroni corrections), and all loci were in Hardy–Weinberg equilibrium ( $p>0.4$ ). Overall, the probability of not detecting a second male was extremely low ( $4.89 \times 10^{-4}$ ).

Our patriline analysis showed that for all colonies and all loci, except one (locus 3675 in colony I), worker genotypes were consistent with workers originating from a queen mated to a single male (Table 2). In colony I,

**Table 2** Summary of all worker genotypes and inferred male and queen genotypes

Colony	Locus	Worker genotypes (# workers)	Inferred male	Inferred queen
A	3675	224::230 (7) 221::224 (9)	224	221::230
	1284	230::236 (8) 231::236 (8)	236	230::231
	4380	128::128 (6) 128::132 (10)	128	128::132
	5134	261::261 (9) 246::261 (7)	261	246::261
	6826	188::188 (7) 188::190 (9)	188	188::190
	3186	197::197 (11) 197::203 (5)	197	197::203
B <sup>a</sup>	3675	221::221 (7) 221::228 (5) 221::224 (2) <sup>b</sup>	221	221::228
	1284	232::236 (10) 236::236 (4)	236	232::236
	4380	132::146 (6) 132::132 (7) 128::132 (1) <sup>b</sup> 128::132 (1) <sup>b</sup>	132	132::146
	5134	246::261 (12) 246::246 (1) <sup>b</sup> 246::272 (1) <sup>b</sup>	246 or 261	246::246 or 261::261
	6826	178::188 (8) 178::184 (4) 180::184 (2) <sup>b</sup>	178	184::188
	3186	201::203 (5) 199::201 (7) 203::205(2) <sup>b</sup>	201	199::203
C	3675	221::224 (16)	221 or 224	221::221 or 224::224
	1284	230::230 (8) 230::236 (8)	230	230::236
	4380	130::144 (8) 144::148 (8)	144	130::148
	5134	246::270 (9) 249::270 (5)	270	246::249
	6826 <sup>c</sup> (1)	186::190 (7) 186::196 (8)	186	190::196
	3186	197::203 (9) 203::203 (7)	203	197::203

**Table 2** (continued)

Colony	Locus	Worker genotypes (# workers)	Inferred male	Inferred queen
D	3675 <sup>c</sup> (2)	240::240 (5)	240	224::240
		224::240 (9)		
	1284 <sup>c</sup> (1)	232::237 (10)	237	232::237
		237::237 (5)		
	4380 <sup>c</sup> (1)	130::146 (9)	130	142::146
		130::142 (6)		
	5134 <sup>c</sup> (1)	249::299 (5)	249	249::299
249::249 (10)				
6826	184::194 (12)	194	184::203	
	194::203 (4)			
3186 <sup>c</sup> (1)	196::203 (7)	196	196::203	
	196::196 (8)			
E	3675	228::252 (11)	228	224::252
		224::228 (5)		
	1284	230::232 (12)	232	230::236
		232::236 (4)		
	4380	130::130 (4)	130	130::146
		130::146 (12)		
	5134	246::246 (10)	246	246::249
246::249 (6)				
6826	188::196 (7)	196	188::196	
	196::196 (9)			
3186	197::205 (5)	197	194::205	
	197::197 (11)			
F	3675	221::249 (9)	249	221::224
		224::249 (7)		
	1284	231::231 (4)	231	231::236
		231::236 (12)		
	4380	130::130 (10)	130	126::130
		126::130 (6)		
	5134	246::249 (10)	246	249::261
246::261 (6)				
6826 <sup>c</sup> (1)	194::198 (7)	198	184::194	
	184::198 (8)			
3186	203::205 (7)	203	203::205	
	203::203 (9)			
G	3675	221::228 (14)	221 or 228	221::221 or 228::228
		221::224 (2) <sup>b</sup>		
	1284	231::236 (8)	236	231::232
		232::236 (6)		
		230::230 (2) <sup>b</sup>		
	4380	128::128 (6)	128	128::130
		128::130 (9)		
130::130 (1) <sup>b</sup>				
5134	246::272 (10)	246	249::272	
	246::249 (6)			
6826	186::198 (8)	186	198::207	
	186::207 (6)			
	188::188 (1) <sup>b</sup>			
	188::199 (1) <sup>b</sup>			
3186	201::203 (10)	203	201::203	
	203::203 (4)			
	197::201 (1) <sup>b</sup>			
	201::205 (1) <sup>b</sup>			

**Table 2** (continued)

Colony	Locus	Worker genotypes (# workers)	Inferred male	Inferred queen
H	3675 <sup>c</sup> (1)	221::221 (15)	221	221::221
		231::236 (10)		
	1284	231::231 (6)	231	231::236
		128::130 (6)		
	4380	130::132 (10)	130	128::132
		246::261 (8)		
	5134	246::246 (8)	246	246::261
186::186 (16)				
6826	203::203 (11)	186	186::186	
	203::205 (4)			
I	3675 <sup>d</sup>	240::240 (8)	<sup>d</sup>	<sup>d</sup>
		224::224 (8)		
	1284 <sup>c</sup> (2)	230::236 (12)	236	230::232
		232::236 (2)		
	4380	128::135 (16)	128 or 135	128::128 or 135::135
		246::246 (6)		
	5134 <sup>c</sup> (1)	246::267 (9)	246	246::267
192::192 (7)				
6826	192::199 (9)	192	192::199	
	201::203 (9)			
3186 <sup>c</sup> (1)	203::206 (6)	203	201::206	
	221::224 (16)			
J	3675	230::231 (10)	221 or 224	221::221 or 224::224
		231::236 (6)		
	1284	130::133 (9)	231	230::236
		133::148 (7)		
	4380	246::246 (16)	133	130::148
		190::198 (7)		
	5134	186::198 (9)	198	186::190
197::203 (16)				
6826	197::203 (16)	197 or 203	197::197 or 203::203	
	221::221 (5)			
K	3675 <sup>c</sup> (2)	221::236 (9)	221	221::236
		231::236 (16)		
	1284	231::236 (16)	231 or 236	231::231 236::236
		144::144 (9)		
	4380 <sup>c</sup> (2)	130::144 (5)	144	130::144
		246::261 (16)		
	5134	246::261 (16)	246 or 261	246::246 or 261::261
184::201 (8)				
6826	184::192 (8)	184	192::201	
	203::205 (14)			
3186 <sup>c</sup> (2)	203::205 (14)	203 or 205	203::203 or 205::205	

**Table 2** (continued)

Colony	Locus	Worker genotypes (# workers)	Inferred male	Inferred queen
L	3675 <sup>c</sup> (1)	221::224 (10)	221	221::224
		221::221 (5)		
	1284	236::236 (9)	236	232::236
		232::236 (6)		
		230::236 (1) <sup>b</sup>		
	4380 <sup>c</sup> (5)	128::132 (4)	128	128::132
128::128 (6)				
132::133 (1) <sup>b</sup>				
5134	246::261 (16)	246 or 261	246::246 or 261::261	
6826	184::184 (8)	184	184::188	
	184::188 (7)			
	186::194 (1) <sup>b</sup>			
3186 <sup>c</sup> (1)	203::203 (6)	203	199::203	
	199::203 (8)			
	197::199 (1) <sup>b</sup>			
M	3675	221::221 (7)	221	221::224
		221::224 (8)		
		221::--- (1)		
	1284 <sup>c</sup> (1)	231::236 (8)	231	231::236
		231::231 (7)		
	4380	130::130 (7)	130	130::146
5134	246::275 (10)	246	246::275	
	246::246 (6)			
6826	194::196 (12)	196	194::207	
3186	197::201 (16)	197 or 201	197::197 or 201::201	
N	3675 <sup>c</sup> (1)	221::221 (9)	221	221::224
		221::224 (6)		
	1284	236::237 (6)	236	231::237
		231::236 (9)		
		231::--- (1)		
	4380	128::132 (10)	128	132::146
5134 <sup>c</sup> (1)	246::246 (15)	246	246::246	
	186::194 (16)			
3186 <sup>c</sup> (1)	197::203 (5)	203	197::199	
	199::203 (10)			
	199::203 (10)			
O	3675 <sup>c</sup> (1)	221::221 (15)	221	221::221
		232::232 (9)		
	1284 <sup>c</sup> (1)	231::232 (6) <sup>f</sup>	232	231::232
		231::232 (6) <sup>f</sup>		
	4380 <sup>c</sup> (1)	130::142 (6)	130	128::142
		128::130 (9)		
5134 <sup>c</sup> (1)	246::270 (6)	246	246::270	
	246::246 (9)			
6826 <sup>c</sup> (1)	178::194 (9)	178	188::194	
	178::188 (6)			
3186 <sup>c</sup> (1)	201::203 (3)	203	201::203	
	203::203 (12)			

**Table 2** (continued)

Colony	Locus	Worker genotypes (# workers)	Inferred male	Inferred queen
P	3675	221::221 (16)	221	221::221
		231::236 (7)		
	1284	231::232 (9) <sup>f</sup>	231	232::236
		231::232 (9) <sup>f</sup>		
	4380	128::128 (16)	128	128::128
	5134	261::275 (16)	261 or 275	261::261 or 275::275
6826	180::196 (10)	196	180::207	
	196::207 (6)			
	196::207 (6)			
3186	197::203 (8)	203	197::203	
	203::203 (8)			
	203::203 (8)			
Q	3675	221::221 (16)	221	221::221
		232::232 (6)		
	1284	231::232 (10) <sup>f</sup>	232	231::232
		231::232 (10) <sup>f</sup>		
	4380	128::130 (4)	130	128::130
		130::130 (12)		
5134	249::249 (8)	249	249::272	
	249::272 (8)			
6826	184::203 (16)	184 or 203	184::184 or 203::203	
3186	199::201 (8)	199	199::201	
	199::199 (8)			
	199::199 (8)			
R	3675	224::224 (15)	224	224::224
		221::224 (1) <sup>b</sup>		
	1284	230::236 (9)	230	231::236
		230::231 (7)		
	4380 <sup>c</sup> (1)	128::128 (7)	128	128::146
		128::146 (7)		
5134	126::128 (1) <sup>b</sup>	246	246::263	
	246::246 (8)			
6826	246::263 (7)	196	184::188	
	246::275 (1) <sup>b</sup>			
3186	188::196 (9)	197 or 203	197::197 or 203::203	
	184::196 (6)			
5134	188::207 (1) <sup>b</sup>	246	246::246 <sup>e</sup>	
	197::203 (15)			
S <sup>c</sup>	3675 <sup>c</sup> (3)	221::221 (44)	221	221::221 <sup>e</sup>
		236::236 (47)		
	1284	130::130 (47)	236	236::236 <sup>e</sup>
		130::130 (47)		
	4380	246::246 (47)	130	130::130 <sup>e</sup>
		246::246 (47)		
5134	184::188 (47)	184	188::188 <sup>e</sup>	
	203::203 (47)			
6826	118::120 (23)	120	118::120 <sup>e</sup>	
	120::120 (24)			
7207	184::184 (47)	184	184::184 <sup>e</sup>	
	203::203 (47)			
7300	203::203 (47)	203	203::203 <sup>e</sup>	

**Table 2** (continued)

Colony	Locus	Worker genotypes (# workers)	Inferred male	Inferred queen
T	3675 <sup>c</sup> (1)	221::224 (16)	221 or 224	221::221 or 224::224
	1284	236::236 (10) 230::236 (6)	236	230::236
	4380	128::132 (14) 128::128 (2)	128	128::132
	5134	246::263 (16)	246 or 263	246::246 or 263::263
	6826	180::207 (7) 180::203 (9)	180	203::207
	3186	203::203 (11) 197::203 (5)	203	197::203

<sup>a</sup>Two samples were discarded because they had been contaminated

<sup>b</sup>Worker genotype is incompatible with the queen genotype and thus the worker(s) likely comes from another nest

<sup>c</sup>(X) number of samples that did not amplify

<sup>d</sup>All workers are homozygous for this locus, but there are two alleles in the colony, thus we could not infer the queen and male genotype

<sup>e</sup>We genotyped 47 workers and the queen for all nine primers

<sup>f</sup>Genotype profile is ambiguous, possibly homozygous

eight workers were homozygous for allele 224 at locus 3675 and the remaining worker homozygous for allele 240 (Table 2). Given that workers of this colony had normal genotypes at other loci, this pattern is most likely explained by the queen having a 224/240 genotype, and the male having a null allele at this locus.

In colony S, all 16 workers had identical genotypes. They were homozygous at five of the six loci while at locus 6826 they were all heterozygotes. This pattern could be explained by workers being clonally produced or by the mother being homozygous at the six loci and mated with a male having an identical allele at five of the six loci. To discriminate between these two hypotheses we genotyped all workers at three additional loci, analyzed another 31 workers from the same colony at the nine loci and also genotyped the queen. The 31 additional workers had identical genotypes as the initial 16 workers at the six first loci. For two of the additional loci all 47 workers had again identical homozygous genotypes, but for the third additional locus (locus 7275) 24 workers were homozygotes and 23 were heterozygotes. The queen was homozygote for all loci except locus 7275 for which she was heterozygote. Because workers were heterozygote at locus 6826 where the queen was homozygote, and varied in their genotypes for locus 7275, they were apparently sexually produced. The most parsimonious explanation for the genotypic composition of this colony is that

the queen was inbred and that she mated with a brother that shared the same allele at all but locus 6826.

## Discussion

The genotypic data from a total of 343 workers from 20 colonies suggest that *C. fellah* queens mate with a single male, and thus all workers produced by the queen are full sisters. This, together with previous studies in *C. herculeanus* (Gertsch et al. 1995), *C. ligniperdus* (Gertsch et al. 1995; Gadau et al. 1998), *C. floridanus* (Gadau et al. 1996), *C. nawai* (Satoh et al. 1997), *C. consobrinus* (Fraser et al. 2000), *C. ocreatus* (Goodisman and Hahn 2004) and *C. festinatus* (Goodisman and Hahn 2005) indicate that queens are mostly singly mated in the genus *Camponotus*.

Estimates of patrilineages can be inaccurate due to non-detection errors and small sample size (Pedersen and Boomsma 1999). Given the microsatellites were very polymorphic, the probability of non-detection of a second father was very low (<0.1%). Our sample size of 16 workers per colony enabled us to detect a second patriline with a high probability (>95%) if the second male contributed to about 20% of the workers (Boomsma and Ratnieks, 1996). In colony S we genotyped 47 workers enabling us to detect a second patriline with a probability of more than 90% even if the second male fathered only 5% of the workers (Boomsma and Ratnieks, 1996). However, also in this colony we only detected one patriline.

In one of the 20 colonies analyzed, worker and queen homozygosity was very high with only one heterozygote locus in workers and another one in the queen. Given the high polymorphism of the microsatellites, the most likely explanation for this high homozygosity is that the queen was inbred and that she mated with a brother. In ants, there is usually a low level of inbreeding due to the high synchronization of mating flight among colonies of the same populations (Tabot 1945; McCluskey 1965; Boomsma and Leusink 1981). Queens and males from *C. fellah* colonies typically leave their nest in the evenings of the first warm and dry spring days where individuals of many nests join in a mating flight (Hefetz A, personal communication). However, instances of inbreeding due to sib-matings have been reported in two *Camponotus* species (*C. ocreatus*, Goodisman and Hahn 2004 and *C. yamaokai*, Satoh et al. 1997) as well as several species of other genera (e.g., *Cardiocondyla batesii*, Schrempf et al. 2005; *Cardiocondyla nigra*, Schrempf 2014; *Cardiocondyla shuckardi*, Heinze et al. 2014; and *Formica exsecta*, Sundström et al. 2003; Vitikainen et al. 2015). Interestingly, the inbred queen that we found had apparently mated with a brother. A similar pattern has been uncovered in *Formica exsecta* where more inbred queens are also more likely to have mated with a

brother (Vitikainen et al. 2015). An association between the level of inbreeding of queens and their likelihood of mating with siblings could occur if there is a genetic basis influencing dispersal propensity. Females that are genetically less inclined to disperse may be more likely to mate with males of their nest and, therefore, become more inbred. Alternatively, the propensity to disperse may be directly influenced by the level of inbreeding (Vitikainen et al. 2015). Because more inbred queens may be smaller and less fit than outbred queens (Coltman et al. 1999; Keller and Waller 2002; McQuillan et al. 2012), they might disperse less (Wolf and Seppa 2016), thereby increasing their likelihood to mate with a sibling.

Mating has several costs that may select monoandry over polyandry (Daly 1978; Keller and Reeve 1995). For example, when mating with several males, females increase their risk of contracting sexually transmitted diseases (Knell and Webberley 2004; Peng et al. 2016), and also spent more time outside the nest exposed to predators (Lafaille et al. 2010; Alem et al. 2011). In addition, multiple mating results in sperm competition (Keller and Reeve 1995; den Boer et al. 2010) and associated conflict of interest between the sexes, which can reduce female lifespan and reproductive output (Chapman et al. 1995; Crudginton and Siva-Jothy 2000; Lessells 2005; Alonza and Pizzari 2013). Consequently, if mating with a single male can provide sufficient sperm to cover for the lifetime reproduction of the colony, and the benefits obtained from polyandry fail to compensate the endured mating costs, then queens might benefit more from being monoandrous.

**Acknowledgements** We thank Abraham Hefetz for collecting queens after a mating flight. We thank Mingkwan Nipitwattanaphon for help with the sample preparation for the initial primer design, and for advice on PCRs. We thank Consolée Aletti for sequencing, Raphaël Boulay, Michael Breed, and an anonymous reviewer for valuable comments, and Ben Sutcliffe for proofreading. This work was funded by several grants from the Swiss National Science Foundation, the Human Frontiers Science Program, and an ERC advanced grant.

## References

Alem S, Koselj K, Siemers BM, Greenfield MD (2011) Bat predation and the evolution of leks in acoustic moths. *Behav Ecol Sociobiol* 65:2105–2116. doi:10.1007/s00265-011-1219-x

Alonza SH, Pizzari T (2013) Selection on female remating interval is influenced by male sperm competition strategies and ejaculate characteristics. *Philos T Roy Soc B* 368: 20120044. doi:10.1098/rstb.2012.0044

Boulay R, Hefetz A, Soroker V, Lenoir A (2000a) *Camponotus fellah* colony integration: worker individuality necessitates frequent hydrocarbon exchanges. *Anim Behav* 59:1127–1133

Boulay R, Quagebeur M, Godzinska E, Lenoir A (1999) Social isolation in ants: evidence of its impact on survivorship and behavior in *Camponotus fellah* (Hymenoptera: Formicidae). *Sociobiology* 33:111–124

Boulay R, Soroker V, Godzinska EJ, Hefetz A, Lenoir A (2000b) Octopamine reverses the isolation-induced increase in trophallaxis in the carpenter ant *Camponotus fellah*. *J Exp Biol* 203:513–520

Boomsma JJ, Leusink A (1981) Weather conditions during nuptial flights of four European ant species. *Oecologia* 50:236–241

Boomsma JJ, Ratnieks FLW (1996) Paternity in eusocial Hymenoptera. *Philos T Roy Soc B* 351:947–975.

Chapman T, Liddle LF, Kalb JM, Wolfner MF, Partridge L (1995) Cost of mating in *Drosophila melanogaster* females is mediated by male accessory gland products. *Nature* 373:241–244

Coltman DW, Pilkington JG, Smith JA, Pemberton JM (1999) Parasite-mediated selection against inbred Soay sheep in a free-living, island population. *Evolution* 53:1259–1267

Crozier RH, Page RE (1985) On being the right size: male contributions and multiple mating in social Hymenoptera. *Behav Ecol Sociobiol* 18:105–115

Crudginton HS, Siva-Jothy MT (2000) Genital damage, kicking and early death. *Nature* 407:855–856

Daly M (1978) The cost of mating. *Am Nat* 112:771–774

den Boer SPA, Baer B, Boomsma JJ (2010) Seminal fluid mediates ejaculate competition in social insects. *Science* 327:1506–1509. doi:10.1126/science.1184709

Fraser VS, Kaufmann B, Oldroyd BP, Crozier RH (2000) Genetic influence on caste in the ant *Camponotus consobrinus*. *Behav Ecol Sociobiol* 47:188–194

Gadau J, Gertsch PJ, Heinze J, Pamilo P, Hölldobler B (1998) Oligogyny by unrelated queens in the carpenter ant, *Camponotus ligniperdus*. *Behav Ecol Sociobiol* 44:23–33. doi:10.1007/s002650050511

Gadau J, Heinze J, Hölldobler B, Schmid M (1996) Population and colony structure of the carpenter ant *Camponotus floridanus*. *Mol Ecol* 5:785–792

Gertsch P, Pamilo P, Varvio SL (1995) Microsatellites reveal high genetic diversity within colonies of *Camponotus* ants. *Mol Ecol* 4:257–260

Goodisman MAD, Hahn DA (2004) Colony genetic structure of the ant *Camponotus ocreatus* (Hymenoptera: Formicidae). *Sociobiology* 44:21–33

Goodisman MAD, Hahn DA (2005) Breeding system, colony structure, and genetic differentiation in the *Camponotus festinatus* species complex of Carpenter ants. *Evolution* 59:2185–2199

Greenwal E, Segre E, Feinerman O (2015) Ant trophallactic networks: simultaneous measurements of interaction patterns and food dissemination. *Sci Rep* 5:12496. doi:10.1038/srep12496

Heinze J, Keller L (2000) Alternative reproductive strategies: a queen perspective in ants. *Trends Ecol Evol* 15:508–512

Heinze J, Schrempf A, Wanke T, Rakotondrazafy H, Rakotondranaivo T, Fisher BL (2014) Polygyny, inbreeding and wingless males in the Malagasy ant *Cardiocondyla shuckardi* Forel (Hymenoptera, Formicidae). *Sociobiology* 61:300–306

Hughes WHO, Boomsma JJ (2004) Genetic diversity and disease resistance in leaf-cutting ant societies. *Evolution* 58:1251–1260

Ichinose K, Boulay R, Cerdá X, Lenoir A (2009) Influence of queen and diet on nestmate recognition and cuticular hydrocarbon differentiation in a fission-dispersing ant, *Aphaenogaster senilis*. *Zool Sci* 26:681–685

Ionescu-Hirsch A (2009) An annotated list of *Camponotus* of Israel, with a key and descriptions of new species. *Isr J Entomol* 39:57–98

Julian GE, Cahan S (1999) Undertaking specialization in the desert leaf-cutter ant *Acromyrmex versicolor*. *Anim Behav* 58:437–442

Julian GE, Fewell JH (2004) Genetic variation and task specialization in the desert leaf-cutter ant, *Acromyrmex versicolor*. *Anim Behav* 68:1–8

- Kafsi M, Braunschweig R, Mersch D, Grossglauer M, Keller L, Thiran P (2016) Uncovering latent behaviors in ant colonies. Proceedings of the 2016 SIAM International Conference on Data Mining. doi:[10.1137/1.9781611974348.51](https://doi.org/10.1137/1.9781611974348.51)
- Katzav-Gozansky T, Boulay R, Ionescu-Hirsh A, Hefetz A (2008) Nest volatiles as modulators of nestmate recognition in the ant *Camponotus fellah*. *J Insect Physiol* 54:378–385
- Katzav-Gozansky T, Boulay R, Van der Meer R, Hefetz A (2004) In-nest environment modulates nestmate recognition in the ant *Camponotus fellah*. *Naturwissenschaften* 91:186–190
- Keller L, Reeve HK (1995) Why do females mate with multiple males? The sexually selected sperm hypothesis. *Adv Stud Behav* 24:291–315
- Keller LF, Waller DM (2002) Inbreeding effects in wild populations. *Trends Ecol Evol* 17:230–241
- Knell RJ, Webberley KM (2004) Sexually transmitted diseases of insects: distribution, evolution, ecology and host behavior. *Biol Rev* 79:557–581
- Koto A, Mersch D, Hollis B, Keller L (2015) Social isolation causes mortality by disrupting energy homeostasis in ants. *Behav Ecol Sociobiol* 69:583–591
- Kureck IM, Jongepier E, Nicolai B, Foitzik S (2012) No inbreeding depression but increased sexual investment in highly inbred ant colonies. *Mol Ecol* 21:5613–5623. doi:[10.1111/mec.12060](https://doi.org/10.1111/mec.12060)
- Lafaille M, BimBard G, Greenfield MD (2010) Risk trading in mating behavior: forgoing anti-predator responses reduces the likelihood of missing terminal mating opportunities. *Behav Ecol Sociobiol* 64:1485–1494. doi:[10.1007/s00265-010-0963-7](https://doi.org/10.1007/s00265-010-0963-7)
- Lessells CM (2005) Why are males bad for females? Models for the evolution of damaging male mating behavior. *Am Nat* 165:S46–S63
- McCluskey ES (1965) Circadian rhythms in male ants of five diverse species. *Science* 150:1037–1039
- McQuillan R, Eklund N, Pirastu N, Kuningas M, McEvoy BP, Esko T, Corre T, Davies G, Kaakinen M, Lyytikäinen L-P, Kristiansson K, Havulinna AS, Gögele M et al. (2012) Evidence of inbreeding depression on human height. *Plos Genet* 8: e1002655. doi:[10.1371/journal.pgen.1002655](https://doi.org/10.1371/journal.pgen.1002655)
- Mersch DP, Crespi A, Keller L (2013) Tracking individuals shows spatial fidelity is a key regulator of ant social organization. *Science* 340:1090–1093
- Oldroyd BP, Fewell JH (2007) Genetic diversity promotes homeostasis in insect colonies. *Trend Ecol Evol* 22:408–413
- Peng Y, Grassl J, Millar AH, Baer B (2016) Seminal fluid of honeybees contains multiple mechanisms to combat infections of the sexually transmitted pathogen *Nosema apis*. *Proc R Soc B* 283:20151785
- Raymond M, Rousset F (1995) GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *J Hered* 86:248–249
- Reber A, Castella G, Christe P, Chapuisat M (2008) Experimentally increased group diversity improves disease resistance in an ant species. *Ecol Lett* 11:682–689. doi:[10.1111/j.1461-0248.2008.01177.x](https://doi.org/10.1111/j.1461-0248.2008.01177.x)
- Rheindt FE, Strehl CP, Gadau J (2005) A genetic component in the determination of worker polymorphism in the Florida harvester ant *Pogonomyrmex badius*. *Insect Soc* 52:163–168. doi:[10.1007/s00040-004-0787-4](https://doi.org/10.1007/s00040-004-0787-4)
- Richard FJ, Hefetz A, Christides J-P, Errard C (2004) Food influence on colonial recognition and chemical signature between nestmates in the fungus-growing ant *Acromyrmex subterraneus subterraneus*. *Chemoecology* 14:9–16
- Robinson GE (1992) Regulation of division of labor in insect societies. *Annu Rev Entomol* 37:637–665
- Rousset F (2008) Genepop'007: a complete reimplementation of the Genepop software for Windows and Linux. *Mol Ecol Resour* 8:103–106
- Satoh T, Masuko K, Matsumoto T (1997) Colony genetic structure in the mono- and polygynous sibling species of the ants *Camponotus nawai* and *Camponotus yamaokai*: DNA fingerprint analysis. *Ecol Res* 12:71–76
- Schmidt, AM, Linksvayer TA, Boomsma JJ, Pedersen JS (2011) No benefit in diversity? The effect of genetic variation on survival and disease resistance in a polygynous social insect. *Ecol Entomol* 36:751–759.
- Schrempf, A (2014) Inbreeding, multiple mating and foreign sexuals in the ant *Cardiocondyla nigra* (Hymenoptera: Formicidae). *Myrmecological News* 20:1–5.
- Schrempf A, Reber C, Tinaut A, Heinze J (2005) Inbreeding and local mate competition in the ant *Cardiocondyla batesii*. *Behav Ecol Sociobiol* 57:502–510. doi:[10.1007/s00265-004-0869-3](https://doi.org/10.1007/s00265-004-0869-3)
- Schwander T, Rosset H, Chapuisat M (2005) Division of labour and worker size polymorphism in ant colonies: the impact of social and genetic factors. *Behav Ecol Sociobiol* 59:215–221. doi:[10.1007/s00265-005-0027-6](https://doi.org/10.1007/s00265-005-0027-6)
- Smith CR, Toth AL, Suarez AV, Robinson GE (2008) Genetic and genomic analyses of the division of labour in insect societies. *Nat Rev Genet* 9:735–748. doi:[10.1038/nrg2429](https://doi.org/10.1038/nrg2429)
- Sundström L, Keller L, Chapuisat M (2003) Inbreeding and sex-biased gene flow in the ant *Formica exsecta*. *Evolution* 57:1552–1561. doi:[10.1111/j.0014-3820.2003.tb00363.x](https://doi.org/10.1111/j.0014-3820.2003.tb00363.x)
- Suni SS, Gordon DM (2010) Fine-scale genetic structure and dispersal distance in the harvester ant *Pogonomyrmex barbatus*. *Heredity* 104:168–173. doi:[10.1038/hdy.2009.124](https://doi.org/10.1038/hdy.2009.124)
- Talbot M (1945) A comparison of flights of four species of ants. *Am Midl Nat* 34:504–510
- Vitikainen EIK, Haag-Liautard C, Sundström L (2015) Natal dispersal, mating patterns, and inbreeding in the ant *Formica exsecta*. *Am Nat* 186:716–727. doi:[10.1086/683799](https://doi.org/10.1086/683799)
- Waddington SJ, Santorelli LA, Ryan FR, Hughes WOH (2010) Genetic polyethism in leaf-cutting ants. *Behav Ecol* 21:1165–1169
- Wiernasz DC, Hines J, Parker DG, Cole BJ (2008) Mating for variety increases foraging activity in the harvester ant, *Pogonomyrmex occidentalis*. *Mol Ecol* 17:1137–1144
- Wolf JI, Seppa P (2016) Dispersal and mating in a size-dimorphic ant. *Behav Ecol Sociobiol* 70:1267–1276. doi:[10.1007/s00265-016-2135-x](https://doi.org/10.1007/s00265-016-2135-x)