



Downloaded on 2017-09-05T00:10:56Z



University College Cork, Ireland Coláiste na hOllscoile Corcaigh

## **PROKARYOTES**



# **Complete Genome Sequence of the Gamma-Aminobutyric Acid-Producing Strain Streptococcus thermophilus APC151**

**AMERICAN SOCIETY FOR MICROBIOLOGY**  genome**A**nnouncements<sup>™</sup>

Daniel M. Linares,<sup>a,b</sup> Silvia Arboleya,<sup>a,b</sup> R. Paul Ross,<sup>a,b,c</sup> Catherine Stanton<sup>a,b</sup>

Food Biosciences Department, Teagasc Food Research Centre, Moorepark, Fermoy, County Cork, Ireland<sup>a</sup>; APC, Microbiome Institute, University College Cork, Cork, Irelandb; School of Microbiology, University College Cork, Cork, Ireland<sup>c</sup>

**ABSTRACT** Here is presented the whole-genome sequence of Streptococcus thermophilus APC151, isolated from a marine fish. This bacterium produces gamma-aminobutyric acid (GABA) in high yields and is biotechnologically suitable to produce naturally GABAenriched biofunctional yogurt. Its complete genome comprises 2,097 genes and 1,839,134 nucleotides, with an average  $G+C$  content of 39.1%.

*S*treptococcus thermophilus is a nonpathogenic homofermentative facultative an-aerobic lactic acid bacterium with a long history of use in the artisanal and modern industrial manufacture of fermented dairy products, especially yogurt [\(1\)](#page-2-0). In addition, this commensal bacterium is one of the pioneer colonizers of oral and small intestine mucosal surfaces in newborns [\(2\)](#page-2-1) and remains predominant in the oropharyngeal [\(2,](#page-2-1) [3\)](#page-2-2) and gastrointestinal [\(4\)](#page-2-3) tracts throughout the human life span. Based on the potential beneficial effects on human health, it is sometimes marketed as a probiotic [\(5](#page-2-4)[–](#page-2-5)[9\)](#page-2-6).

We here report the genome of Streptococcus thermophilus APC151, a strain isolated from the digestive tract of a marine fish that produces large amounts of gammaaminobutyric acid (GABA) [\(10\)](#page-2-7). GABA has been classified as a health-promoting bioactive component in foods and pharmaceuticals [\(11\)](#page-2-8) due to its function as an antihypertensive and antidiabetic neurotransmitter [\(11](#page-2-8)[–](#page-2-9)[15\)](#page-2-10).

Genomic DNA was purified using the GenElute bacterial genomic DNA kit (Sigma-Aldrich). The nucleotide sequence of S. thermophilus was resolved utilizing PacBio single-molecule real-time (SMRT) technology on an RS system (GATC Biotech, Konstanz, Germany). Whole-genome sequencing yielded a total of 72,472 reads with a mean length of 12,628 bp. Subsequent de novo assembly utilizing the HGAP3 protocol yielded a single polished contig with 423-fold average reference coverage.

Annotation was performed by the RAST server [\(16\)](#page-2-11). The total genome size was 1,839,134 bp, which included 2,097 predicted open reading frames (ORFs) (2,012 coding genes and 85 RNA genes) and a  $G+C$  content of 39.1%. The genome encompasses one predicted phage packaging cassette, one predicted prophage lysogenic conversion module, three clustered regularly interspaced short palindromic repeat (CRISPR) clusters, and 12 genes related to transposases. Additionally, the genes responsible for GABA biosynthesis, encoding the glutamate decarboxylase (gadB) and the glutamate/GABA antiporter (gadC), have been identified in the genome of this strain. Sequence analysis revealed that the gadB-gadC genes are flanked by transposases, a genetic organization similar to other  $GAD<sup>+</sup>$  genomes available in Gen $B$ ank.

The strain was found to be sensitive to all antibiotics proposed by European Food Safety Authority (EFSA) standards [\(10\)](#page-2-7). In addition, no amino acid decarboxylases

#### **Received** 28 February 2017 **Accepted** 7 March 2017 **Published** 27 April 2017

**Citation** Linares DM, Arboleya S, Ross RP, Stanton C. 2017. Complete genome sequence of the gamma-aminobutyric acid-producing strain Streptococcus thermophilus APC151. Genome Announc 5:e00205-17. [https://doi](https://doi.org/10.1128/genomeA.00205-17) [.org/10.1128/genomeA.00205-17.](https://doi.org/10.1128/genomeA.00205-17)

**Copyright** © 2017 Linares et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0](http://creativecommons.org/licenses/by/4.0/) [International](http://creativecommons.org/licenses/by/4.0/) license.

Address correspondence to Daniel M. Linares, [Daniel.Linares@teagasc.ie.](mailto:Daniel.Linares@teagasc.ie)

associated with biogenic amines biosynthesis or virulence-related genes were identified in this genome.

The S. thermophilus APC151 genome showed a high degree of identity to that of strain S. thermophilus ND03 (GenBank accession no. NC\_017563). The comparison of the two genomes revealed that strain APC151 contains an additional cluster of 13 genes (encoding a 5S RNA, a small subunit rRNA, a large subunit rRNA, and 10 tRNA genes) which are not present in the ND03 strain. In addition, strain APC151 acquired a number of mutations that disrupted or caused a frameshift in seven genes (encoding two transposases, two acyltransferases, the phosphate-transport-ATP-binding protein PstB, the β-carotene 15-15'-monooxygenase, and the substrate-specific component MtsA methionine-regulated ECF transporter).

These data are intended to increase the availability of the genomes of GABAproducing S. thermophilus strains in order to better understand their biodiversity and their potential technological, probiotic, and biofunctional properties.

**Accession number(s).** This whole-genome sequence was deposited at DDBJ/EMBL/ GenBank under the accession number [CP019935.](https://www.ncbi.nlm.nih.gov/nuccore/CP019935)

### **ACKNOWLEDGMENTS**

This work is supported by the APC Microbiome Institute, which is a research center funded by Science Foundation Ireland (SFI). This publication has emanated from research supported by a research grant from Science Foundation Ireland (SFI) under grant number SFI/12/RC/2273.

#### <span id="page-2-0"></span>**REFERENCES**

- 1. Wu Q, Tun HM, Leung FC, Shah NP. 2014. Genomic insights into high exopolysaccharide-producing dairy starter bacterium Streptococcus thermophilus ASCC 1275. Sci Rep 4:4974. [https://doi.org/10.1038/srep04974.](https://doi.org/10.1038/srep04974)
- <span id="page-2-1"></span>2. Seow WK, Lam JH, Tsang AK, Holcombe T, Bird PS. 2009. Oral Streptococcus species in pre-term and full-term children: a longitudinal study. Int J Paediatr Dent 19:406 – 411. [https://doi.org/10.1111/j.1365-263X](https://doi.org/10.1111/j.1365-263X.2009.01003.x) [.2009.01003.x.](https://doi.org/10.1111/j.1365-263X.2009.01003.x)
- <span id="page-2-2"></span>3. Bik EM, Long CD, Armitage GC, Loomer P, Emerson J, Mongodin EF, Nelson KE, Gill SR, Fraser-Liggett CM, Relman DA. 2010. Bacterial diversity in the oral cavity of 10 healthy individuals. ISME J 4:962–974. [https://](https://doi.org/10.1038/ismej.2010.30) [doi.org/10.1038/ismej.2010.30.](https://doi.org/10.1038/ismej.2010.30)
- <span id="page-2-3"></span>4. Wang M, Ahrné S, Jeppsson B, Molin G. 2005. Comparison of bacterial diversity along the human intestinal tract by direct cloning and sequencing of 16S rRNA genes. FEMS Microbiol Ecol 54:219-231. [https://doi.org/](https://doi.org/10.1016/j.femsec.2005.03.012) [10.1016/j.femsec.2005.03.012.](https://doi.org/10.1016/j.femsec.2005.03.012)
- <span id="page-2-4"></span>5. Burton JP, Wescombe PA, Cadieux PA, Tagg JR. 2011. Beneficial microbes for the oral cavity: time to harness the oral streptococci? Benef Microbes 2:93–101. [https://doi.org/10.3920/BM2011.0002.](https://doi.org/10.3920/BM2011.0002)
- 6. Teughels W, Kinder Haake S, Sliepen I, Pauwels M, Van Eldere J, Cassiman JJ, Quirynen M. 2007. Bacteria interfere with A. actinomycetemcomitans colonization. J Dent Res 86:611– 617. [https://doi.org/10.1177/](https://doi.org/10.1177/154405910708600706) [154405910708600706.](https://doi.org/10.1177/154405910708600706)
- 7. Wescombe PA, Heng NC, Burton JP, Chilcott CN, Tagg JR. 2009. Streptococcal bacteriocins and the case for Streptococcus salivarius as model oral probiotics. Future Microbiol 4:819-835. [https://doi.org/10.2217/fmb](https://doi.org/10.2217/fmb.09.61) [.09.61.](https://doi.org/10.2217/fmb.09.61)
- <span id="page-2-5"></span>8. Mignolet J, Fontaine L, Kleerebezem M, Hols P. 2016. Complete genome sequence of Streptococcus salivarius HSISS4, a human commensal bacterium highly prevalent in the digestive tract. Genome Announc 4(1): e01637-15. [https://doi.org/10.1128/genomeA.01637-15.](https://doi.org/10.1128/genomeA.01637-15)
- <span id="page-2-6"></span>9. Jandhyala SM, Talukdar R, Subramanyam C, Vuyyuru H, Sasikala M, Nageshwar Reddy D. 2015. Role of the normal gut microbiota. World J Gastroenterol 21:8787– 8803. [https://doi.org/10.3748/wjg.v21.i29.8787.](https://doi.org/10.3748/wjg.v21.i29.8787)
- <span id="page-2-7"></span>10. Linares DM, O'Callaghan TF, O'Connor PM, Ross RP, Stanton C. 2016. Streptococcus thermophilus APC151 strain is suitable for the manufacture of naturally GABA-enriched bioactive yogurt. Front Microbiol 7:1876. [https://doi.org/10.3389/fmicb.2016.01876.](https://doi.org/10.3389/fmicb.2016.01876)
- <span id="page-2-8"></span>11. Li H, Cao Y. 2010. Lactic acid bacterial cell factories for gammaaminobutyric acid. Amino Acids 39:1107–1116. [https://doi.org/10.1007/](https://doi.org/10.1007/s00726-010-0582-7) [s00726-010-0582-7.](https://doi.org/10.1007/s00726-010-0582-7)
- 12. Shimada M, Hasegawa T, Nishimura C, Kan H, Kanno T, Nakamura T, Matsubayashi T. 2009. Anti-hypertensive effect of gammaaminobutyric acid (GABA)-rich Chlorella on high-normal blood pressure and borderline hypertension in placebo-controlled double blind study. Clin Exp Hypertens 31:342–354. [https://doi.org/10.1080/](https://doi.org/10.1080/10641960902977908) [10641960902977908.](https://doi.org/10.1080/10641960902977908)
- 13. Soltani N, Qiu H, Aleksic M, Glinka Y, Zhao F, Liu R, Li Y, Zhang N, Chakrabarti R, Ng T, Jin T, Zhang H, Lu WY, Feng ZP, Prud'homme GJ, Wang Q. 2011. GABA exerts protective and regenerative effects on islet beta cells and reverses diabetes. Proc Natl Acad Sci U S A 108: 11692–11697. [https://doi.org/10.1073/pnas.1102715108.](https://doi.org/10.1073/pnas.1102715108)
- <span id="page-2-9"></span>14. Wan Y, Wang Q, Prud'homme GJ. 2015. GABAergic system in the endocrine pancreas: a new target for diabetes treatment. Diabetes Metab Syndr Obes 8:79 – 87. [https://doi.org/10.2147/DMSO.S50642.](https://doi.org/10.2147/DMSO.S50642)
- <span id="page-2-10"></span>15. Pouliot-Mathieu K, Gardner-Fortier C, Lemieux S, St-Gelais D, Champagne CP, Vuillemard JC. 2013. Effect of cheese containing gammaaminobutyric acid-producing lactic acid bacteria on blood pressure in men. PharmaNutrition 1:141–148. [https://doi.org/10.1016/j.phanu.2013](https://doi.org/10.1016/j.phanu.2013.06.003) [.06.003.](https://doi.org/10.1016/j.phanu.2013.06.003)
- <span id="page-2-11"></span>16. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. BMC Genomics 9:75. [https://doi.org/10.1186/1471-2164-9-75.](https://doi.org/10.1186/1471-2164-9-75)