



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2016; 5(6): 323-331
Received: 12-09-2016
Accepted: 13-10-2016

Suhail Mohammed Hussain
Department of Microbiology &
Biotechnology, Bangalore
University, Jnana Bharathi
Campus Bangalore - 560 056,
Karnataka, India

Mohammed Sadath Hussain
Department of Biotechnology,
Indian academy of degree
college, Bangalore - 560 043,
Karnataka, India

Arbaaz Ahmed L
Department of Microbiology &
Biotechnology, Bangalore
University, Jnana Bharathi
Campus Bangalore - 560 056,
Karnataka, India

Nahida Arif
Department of Microbiology &
Biotechnology, Bangalore
University, Jnana Bharathi
Campus Bangalore - 560 056,
Karnataka, India

Correspondence

Suhail Mohammed Hussain
Department of Microbiology &
Biotechnology, Bangalore
University, Jnana Bharathi
Campus Bangalore - 560 056,
Karnataka, India

Characterization of isolated bioactive phytoconstituents from *Flacourtia indica* as potential phytopharmaceuticals - An *in silico* perspective

Suhail Mohammed Hussain, Mohammed Sadath Hussain, Arbaaz Ahmed L and Nahida Arif

Abstract

The dependence of mankind for therapeutic applications on plants dates back to the start of the human race. Natural remedies from ethnobotanicals are found to be safe and cost effective. Due to the poor pharmacokinetic profiles and toxicity problems many synthetic drugs often fails to enter the market thus, the pharmacologically active compounds from plants continued to provide an important source of novel drug leads. The early inclusion of pharmacokinetics consideration in the drug discovery process using *in silico* methods is becoming popular due to improved generation of software's. The problems with conventional method required time-consuming multi-step processes against a battery of *in vivo* biological screening and high cost thus, *In silico* prediction of the pharmacokinetic parameters, biological properties and toxicity due to advent of chemo-informatics tools, has reduced the cost dramatically and early application in drug design are realized. The present investigation deals with computational evaluation of six isolated phytocompounds from *Flacourtia indica* for their pharmacological potential and biological activities. These compounds were evaluated for drug likeness properties, bioactivity score, ADME/T profiles, and health affects by using various bioinformatics tools. The result indicated that all the six compounds analyzed were non mutagens, non-carcinogens and having good drug-likeness properties were seen. The ADMET parameters and probability of health effects were analyzed by admet SAR and ACD/I-Lab online tools respectively and results shows the ADMET and probability of health effects values are also in satisfying ranges. Pharmacological activities of these compounds were predicted individually using PASS server many different pharmacological activities and mechanisms of action shown by these compounds were reported. The results of our analysis clearly depict that all six phytocompounds were having good pharmacokinetic profiles with numerous biological activities. These compounds can be further studied *in vitro* and *in vivo* for the discovery of novel preventive and therapeutic drug.

Keywords: *Flacourtia indica*, phytopharmaceuticals, ADMET, PASS, *in silico*, pharmacokinetics, drug discovery

Introduction

Plants have been exploited over the millennia for human welfare in the promotion of health and as therapeutic drugs. Most of the developing countries such as China, India, Sri Lanka and a few others endowed with vast resources of medicinal and aromatic plants. Plants being rich sources of secondary metabolites such as alkaloids, terpenoids, triterpenies, flavanoids, tannins, and phenolic compounds etc. which are responsible for various biological activities have been used as treatment for various ailments (de-Fathima *et al.*, 2006) ^[7]. In the later mediaeval period, Islamic part of the world has flourished the herbalism (David taylor, 2014) and Arab's were encouraged to use plants and fruits as nutraceuticals, the holy Quran mentions many plants as nutraceuticals so that the mankind can enjoy and benefit from their nutritional and health values. Among some of the ethnobotanicals mentioned in the holy Quran and Hadith by the Holy Prophet Mohammed ﷺ (Peace Be Upon Him) are Sweet flag, Myrrh, Memcylon (Tintura), Barley, Sweet basil, Pomegranate, grapes, citrus, melon, squash, Figs, date palm, honey, olive oil, and black seeds etc. (Rahman *et al.*, 2008; Ahmad *et al.*, 2009) ^[19]. The statement of Holy Prophet Mohammed ﷺ (Peace Be upon Him) that "there is no disease that Allah has created, except that He also has created its treatment" encouraged Mankind to engage in medical research and seek out a cure for every disease known to them (Ahmed and Hasan, 2015). Medicinal plants, the backbone of traditional medicine with excessive pharmacological studies are the potential source of lead compounds in drug development (Mathew *et al.*, 2016) ^[15]. Modern pharmacopoeia still contains at least 25% drugs derived from plants (De Silva, 1997) ^[8]. Since many synthetic drugs often fail to enter

the market as a result of poor pharmacokinetic and pharmacodynamics profiles (Ntie-Kang, 2013) [17], therefore phytopharmaceuticals are preferred since Plants have been evolved through biological validation and therefore induce less toxicity and side effects as compared to synthetic drugs (Mathew *et al.*, 2016) [15]. The use of plants in the control and treatment of diseases in recent years has gained considerable importance and major sources of biologically active and high pharmacological active compounds are from plants and fruits (WHO Report, 2002) [22].

The objective of drug design is to find a chemical compound that can fit to a specific cavity called binding pocket cleft on a protein target both geometrically and chemically. After passing the animal tests and human clinical trials, this compound becomes a drug available in market to patients. The conventional drug design methods include random screening of chemicals found in nature or synthesized in laboratories. The problems with conventional method required time-consuming multi-step processes against a battery of *in vivo* biological screening and high cost. Modern approach including structure-based drug design with the help of informatics technologies and computational methods has speeded up the drug discovery process in an efficient manner. Remarkable progress has been made during the past five years in almost all the areas concerned with drug design and discovery. An improved generation of software's with easy operation and superior computational tools to generate chemically stable and worthy compounds with refinement capability has been developed. These tools can tap into cheminformatics to shorten the cycle of drug discovery, and thus make drug discovery in more efficiency, cost effectiveness, time saving, and will provide strategies for combination therapy in addition to overcoming toxic side effect (Mandal *et al.*, 2009; Baldi, 2010; Sharma and Sarkar, 2013) [14, 5, 21].

The present investigation aimed to assess the pharmacokinetics, pharmacological properties and biological activities of the isolated phytocompounds from *Flacourtia indica* (BURM.F.) Merr. By computational approach. *Flacourtia indica* (Burm.f.) Merr, belonging to the family Flacourtiaceae, is a small deciduous thorny shrub indigenous to the Indian Peninsula. *Flacourtia indica* is very popular and used as folklore medicine to treat various diseases, fruits are used in the treatment of jaundice and enlarged spleen, seeds are used as rheumatic pain, bark is applied to the body during intermittent fever and root is used in the nephritic colic (Eramma and Gayathri, 2013; Sashidhara *et al.*, 2013) [9, 20] isolated and purified six phytocompounds viz., (1) 2-(2-benzoyl-b-D-glucopyranosyloxy)-7-(1a,2a,6a-trihydroxy-3-oxocyclohex-4-enoyl)-5-hydroxy benzyl alcohol, (2) poliothryoside, (3) catechin-[5,6-e]-4b-(3,4-dihydroxy phenyl)dihydro-2(3H)-pyranone, (4) 2-(6-benzoyl-b-D-glucopyranosyloxy)-7-(1a,2a,6a-trihydroxy-3-oxocyclohex-4-enoyl)-5-hydroxybenzyl alcohol (5), chrysoeriol-7-O-b-D-glucopyranoside and (6) mururin A" by applying series of chromatographic techniques and Experimentally reported that Compound 6 significantly inhibited the *in vitro* growth of both a chloroquine-sensitive(3D7) and a chloroquine-resistant (K1) strain of Plasmodium falciparum (Sashidhara *et al.*, 2013) [20].

Methods and Implications

Preparation of phyto-ligands

The six phytocompounds isolated from *Flacourtia indica*

(Sashidhara *et al.*, 2013) [20] viz., "(1) 2-(2-benzoyl-b-D-glucopyranosyloxy)-7-(1a,2a,6a-trihydroxy-3-oxocyclohex-4-enoyl)-5-hydroxy benzyl alcohol, (2) poliothryoside, (3) catechin-[5,6-e]-4b-(3,4-dihydroxy phenyl)dihydro-2(3H)-pyranone, (4) 2-(6-benzoyl-b-D-glucopyranosyloxy)-7-(1a,2a,6a-trihydroxy-3-oxocyclohex-4-enoyl)-5-hydroxybenzyl alcohol (5), chrysoeriol-7-O-b-D-glucopyranoside and (6) mururin A" were tested for their pharmacological Potential and biological activity for use as promising therapeutic compounds. The 2D and 3D structures of these isolated phytocompounds were obtained from online server's viz., PubChem (<http://pubchem.ncbi.nlm.nih.gov/>) and ChemSpider (<http://www.chemspider.com/>) and each chemical compound was constructed using ACD/ Chemsketch bioinformatics tool and saved in the '.mol' format.

In silico pharmacokinetics analysis

(a) Prediction Drug-likeness properties

Drug-likeness of a chemical compound is equilibrium amongst the molecular properties of a compound which directly affects biological activity, pharmacodynamics and pharmacokinetics of a drug in human body (Menezes *et al.*, 2011) [16]. The "drug-likeness" test was carried out using Lipinski's "Rule of Five", ro5 (Lipinski *et al.*, 1997). The distributions of the compound molecular weights (MW), calculated lipophilicity (logp), number of hydrogen bond acceptors (HBA) and number of hydrogen bond donors (HBD) were used to assess the "drug-likeness" of Compounds (Ntie-Kang F, 2013) [17]. Depending on these four molecular descriptors, the approach generates a vigilant about apparent absorption trouble; the rule states that most "druglike" molecules must have log P ≤ 5, molecular weight ≤ 500, number of hydrogen bond acceptors ≤ 10, and number of hydrogen bond donors ≤ 5. Molecules violating more than one of these rules may have problems with oral bioavailability (Paramashivam *et al.*, 2015) [18].

(b) Prediction of bioactivity scores of Phytocompounds Molinspiration

Molinspiration tool was used In order to predict bioactivity score, Molinspiration is a free on-line cheminformatics services for calculation of important molecular properties as well as prediction of bioactivity score for the most important phytoconstituents drug targets such as GPCR ligand, Ion channel modulator, Kinase inhibitor, Nuclear receptor ligand, Protease inhibitor, Protease inhibitor and Enzyme inhibitor (Balasundaram *et al.*, 2016) [4].

(c) ADME/Tox Predictions by AdmetSAR

Pharmacokinetics (PK) plays a key role throughout pharmaceutical research and development (Alavijeh *et al.*, 2005) [3]. ADMET is an abbreviation in pharmacokinetics for "absorption, distribution, metabolism, excretion and Toxicity." A set of algorithms are involved in predicting the pharmacological activity of a molecule as a prospective drug (Lydia and Sudarsanam, 2015) [13]. The pharmacokinetic properties such as Absorption, Distribution, Metabolism, Excretion and the Toxicity of the compounds can be predicted using admet SAR (<http://www.admetexp.org>) online database. It provides the latest and most comprehensive manually curated data for diverse chemicals associated with known ADMET profiles.

(d) Computation Lethal Dose LD_{50} and probability of health effects using ACD/I-Lab

ACD/I labs is a free online server which consist of a set of molecular and ADMET descriptors. The probability of health effects and physicochemical properties was predicted using lethal dose LD_{50} value. The comparative analysis of ligand molecules LD_{50} values were studied in mouse and rat by intra peritoneal, oral, intravenous, subcutaneous and the probability of toxic health effects were checked in blood, cardiovascular system, gastro intestinal system, kidney, liver, and lung tissues.

(e) Predictions Biological activity by PASS Server

PASS (Predicted Activity Spectrum for Substances) server (<http://195.178.207.233/PASS/>), this server predicted activity spectrum of a chemical compound as Pa (probable activity) and Pi (probable inactivity). The set of pharmacological effects, mechanisms of action, and specific toxicities, that might be exhibited by a particular compound in its interaction with biological entities, and which is predicted by PASS (Paramashivam *et al.*, 2015) [18], Prediction of this spectrum by PASS was based on structural activity relationship (SAR) analysis of the training set containing 205,000 plus compounds having more than 3750 kinds of Pharmacological effects and biological activities (Goel *et al.*, 2011) [10]. The compounds showing higher Pa value than Pi are the only constituents considered as possible for a particular pharmacological activity (Khurana *et al.*, 2011; Goel *et al.*, 2011) [11, 10].

Results and Discussion

Plants belonging to family Flacourtiaceae are well known for their biological and ethnomedicinal properties and are used as folklore medicine in the treatment of various ailments in India. The structures of the six isolated compounds were retrieved from pubchem and chemspider servers which are represented in table 1. pharmacokinetic and QSAR properties of the compounds were checked for their drug likeness, drug score, bioactivity score, ADME/T profile, and health affect by using various software's mentioned in Section 2. The drug likeliness and drug score of the compounds were predicted by Lipinski's "rule of five" and the results are depicted in Table. 2 and graphical representation of drug likeness score was shown in the figure (fig.1-fig.6). Out of six phytocompounds,



after computation it was found that compound 2 and 6 are significantly follows the Ro5 and other compounds violates one or two parameters viz. acceptable. The drug likeness score of all the compounds are in acceptable ranges but compound 3, 5 and 6 has the highest score when compare to others.

The bioactivity score of the compounds for the drug targets such as GPCR ligand, Ion channel modulator, Kinase inhibitor, Nuclear receptor ligand, Protease inhibitor, Protease inhibitor and Enzyme inhibitor are evaluated using molinspiration tool are listed in the table 3. From the data obtained, one can notice that all the six compounds possess bioactive score in acceptable ranges. The ADMET analysis in early stages of drug discovery is a very crucial since most of the compounds fails due to poor pharmacokinetics properties and toxicity problems. ADMET properties such as blood brain barrier, Caco-2 cell permeability, Human intestinal absorption, P-gp substrate, P-gp inhibitor, Ames mutagenicity and carcinogenicity were analyzed through computational methods and reported in table 4. The result shows that all the six compounds are passes the ADMET filters with no carcinogenicity and no mutagenicity and all the parameters are in acceptable ranges which are the signs of good pharmacokinetics profiles.

The lethal dose 50 (LD_{50}) values of the six compounds were analyzed for the acute toxicity that has administered through oral, intraperitoneal, intravenous and subcutaneous on mouse / rat models and also toxicity with reference to different organs to check adverse effects on organs and their systems are tested and the results are mentioned in table 5. The overall results suggested that all the six phytoconstituents had less toxic effect on internal tissues and no side-effect were observed in the tested dosages.

The pharmacological effects and biological activities of six compounds were analyzed through PASS online server; the values of Pa and Pi vary between 0.000 and 1.000. Only activities with Pa > Pi are considered as possible for a particular compound. If Pa > 0.7, the probability of experimental pharmacological action is high and if $0.5 < Pa < 0.7$, probability of experimental pharmacological action is less. The tested compounds shows many pharmacological activities and mechanism of actions which were represented in table 6.

Table 1: Isolated compounds from *Flacourtia indica* with their 2D and 3D structures.

Sl no	Compound	2D Structure	3D Structure
1	2-(2-benzoyl- b-D-glucopyranosyloxy)-7-(1 a,2 a,6a trihydroxy-3-oxocyclohex-4-enoyl)-5-hydroxy benzyl alcohol		











2	poliothryoside		
3	catechin-[5,6-e]-4b-(3,4dihydroxyphenyl)dihydro-2(3H)-pyranone		
4	2-(6-benzoyl-b-D glucopyranosyloxy)-7-(1a,2 a,6 a-trihydroxy-3-oxocyclohex-4-enoyl)-5-hydroxybenzyl alcohol		
5	chrysoeriol-7-O- b-D-glucopyranoside		
6	mururin A		

Table 2: Lipinski rule of five and Drug-likeness properties prediction using Molsoft online program

Compound	Molecular Formula	Molecular Weight	Hydrogen Bond acceptor (HBA)	Hydrogen Bond Donator (HBD)	Mol LogP	Number of rotatable bonds	Drug-likeness model score
1	C27 H28 O14	576.15	14	7	-2.21	10	-0.10
2	C20 H22 O9	406.13	9	5	0.14	7	-0.10
3	C24 H20 O9	452.11	9	6	2.64	2	1.18
4	C27 H28 O14	576.15	14	7	-2.07	10	-0.19
5	C22 H22 O11	462.12	11	6	0.28	5	0.77
6	C24 H16 O9	448.08	9	5	3.76	1	0.62




Fig 1: Drug-likeness model score: -0.10 **Fig 2:** Drug-likeness model score: -0.10





Fig 3: Drug-likeness model score: 1.18 **Fig 4:** Drug-likeness model score: -0.19





Fig 5: Drug-likeness model score: 0.77 **Fig 6:** Drug-likeness model score: 0.62




Table 3: Bioactivity scores of Phytocompounds predicted by Molinspiration.

Compound	GPCR ligand	Ion channel modulator	Kinase inhibitor	Nuclear receptor ligand	Protease inhibitor	Enzyme inhibitor
1	0.09	-0.29	-0.28	0.11	0.14	0.26
2	0.03	0.01	-0.12	0.12	0.01	0.26
3	0.25	0.03	-0.17	0.33	0.21	0.32
4	0.06	-0.26	-0.27	0.13	0.12	0.24
5	0.05	-0.07	0.14	0.21	-0.06	0.37
6	0.04	-0.32	-0.08	0.42	-0.15	0.27

Table 4: ADME/TOX and pharmacological parameter assessment of phytocompounds predicted using admetSAR toolbox

Compound	PlogBB ^a	PCaco ^b	logHIA ^c	logpGI (substrate) ^d	logpGI (non-inhibitor) ^e	PlogS ^f	AMES Toxicity	Carcinogens
1	0.6121	0.8530	0.7902	0.7055	0.7135	-1.3448	NT	NC
2	0.5000	0.8892	0.7559	0.5911	0.8115	-0.9489	NT	NC
3	0.6208	0.8591	0.9015	0.6008	0.9421	-3.5109	NT	NC
4	0.5993	0.8427	0.8541	0.7230	0.7362	-1.5114	NT	NC
5	0.9247	0.8957	0.7769	0.6724	0.8575	-2.3177	NT	NC
6	0.6553	0.8641	0.9450	0.5638	0.9077	-3.4816	NT	NC

^a-Predicted blood/brain barrier partition coefficient (concern value is -3.0 to 1.0), ^b-predicted Caco-2 cell permeability in nm/s (acceptable range: -1 is poor, 1 is great), ^c-predicted human intestinal absorption in nm/s (acceptable range: 0 poor, >1 great), ^d-predicted P-gp substrate in nm/s (acceptable range of -5 is poor, 1 is great), ^e-predicted P-glycoprotein inhibitor in nm/s (accepted range: 0-1), ^f-predicted aqueous solubility, (concern value is -6.5 to -0.5). P-gp: P-glycoprotein, HIA: Human intestinal absorption, ADME: Absorption, distribution, metabolism, excretion NT: Non Ames toxic, NC: non carcinogens.

Table 5: LD 50 and probability of health effects of Phytocompounds predicted using ACD/I-Lab 2.0

ADME-TOX Parameters	1	2	3	4	5	6
LD ₅₀ mouse ¹ (mg/kg, intraperitoneal)	1100	1000	490	900	1100	470
LD ₅₀ mouse ¹ (mg/kg, oral)	4900	1800	3000	5900	360	2100
LD ₅₀ mouse ¹ (mg/kg, intravenous)	800	500	69	600	67	110
LD ₅₀ mouse ¹ (mg/kg, subcutaneous)	2500	1500	140	4200	190	28
LD ₅₀ Rat ¹ (mg/kg, intraperitoneal)	560	570	1900	440	930	2400
LD ₅₀ Rat ¹ (mg/kg, oral)	11000	2900	3100	10000	930	3100
Probability of blood effect ²	0.98	0.63	1	0.95	1	0.98
Probability of cardiovascular system effect ²	0.97	0.86	0.97	0.96	0.92	0.99
Probability of gastrointestinal system effect ²	0.96	0.07	0.99	0.97	0.94	1
Probability of kidney effect ²	0.62	0.24	0.86	0.63	0.63	0.9
Probability of liver effect ²	0.71	0.19	0.84	0.72	0.8	0.97
Probability of lung effect ²	0.46	0.42	0.98	0.46	0.52	0.97

¹Estimates LD 50 value in mg/kg after intraperitoneal, oral, intravenous and subcutaneous administration to mice and rat, ²Estimates probability of blood, gastrointestinal system, kidney, liver and lung effect at therapeutic dose range, 1-6 represents the phytocompounds and the drugs with moderate effect on reliability index (>0.5), the drugs with border line effect on reliability index (>0.3, <0.5), LD50: Lethal dose50.

Table 6: Predicted Pa and Pi values of phytocompounds using PASS online server.

Compounds	Pa*	Pi#	Predicted Pharmacological activity	pa	pi	Predicted Pharmacological Activity
2-(2-benzoyl- b-D-glucopyranosyloxy)-7-(1 a,2 a,6 a-trihydroxy-3-oxocyclohex-4-enyl)-5-hydroxybenzyl alcohol	0,928	0,006	CDP-glycerol glycerophosphotransferase inhibitor	0,799	0,005	Anticarcinogenic
	0,906	0,010	Membrane integrity agonist	0,747	0,003	Monophenol monooxygenase inhibitor
	0,898	0,006	CYP2H substrate	0,752	0,010	UDP-N-acetylglucosamine 4-epimerase inhibitor
	0,862	0,006	Anaphylatoxin receptor antagonist	0,749	0,008	Mucinaminyserine mucinamidase inhibitor
	0,850	0,007	Antineoplastic	0,751	0,018	Beta-adrenergic receptor kinase inhibitor
	0,850	0,009	Sugar-phosphatase inhibitor	0,751	0,018	G-protein-coupled receptor kinase inhibitor
	0,840	0,004	3-Phytase inhibitor	0,734	0,008	Antiprotozoal (Leishmania)
	0,847	0,011	Benzoate-CoA ligase inhibitor	0,728	0,014	Exoribonuclease II inhibitor
	0,841	0,012	Alkenylglycerophosphocholine hydrolase inhibitor	0,737	0,025	Membrane permeability inhibitor
	0,830	0,003	Lactase inhibitor	0,710	0,005	Fructan beta-fructosidase inhibitor
0,818	0,005	Antiinfective				
poliothyroside	0,973	0,001	Monophenol monooxygenase inhibitor	0,825	0,001	Glucan 1,6-alpha-glucosidase inhibitor
	0,966	0,002	Membrane integrity agonist	0,822	0,001	4-Alpha-glucanotransferase inhibitor
	0,951	0,003	Alkenylglycerophosphocholine hydrolase inhibitor	0,827	0,006	NAD(P)+-arginine ADP-ribosyltransferase inhibitor
	0,949	0,002	Sugar-phosphatase inhibitor	0,822	0,003	Cyclomaltodextrinase inhibitor
	0,946	0,004	CDP-glycerol glycerophosphotransferase inhibitor	0,824	0,006	Glucan endo-1,6-beta-glucosidase inhibitor
	0,932	0,002	3-Phytase inhibitor	0,814	0,001	4-Coumarate-CoA ligase inhibitor
	0,933	0,003	Antiinfective	0,814	0,002	Laminaribiose phosphorylase inhibitor
	0,926	0,003	Membrane permeability inhibitor	0,813	0,005	UDP-glucuronosyltransferase substrate
	0,915	0,002	Beta-mannosidase inhibitor	0,808	0,003	Phenylacetate-CoA ligase inhibitor
	0,915	0,002	Mucinaminyserine mucinamidase inhibitor	0,772	0,009	Membrane integrity antagonist
	0,916	0,003	Anaphylatoxin receptor antagonist	0,771	0,010	Immunostimulant
	0,909	0,002	Fructan beta-fructosidase inhibitor	0,765	0,004	Phosphoinositide 5-phosphatase inhibitor
	0,905	0,003	Exoribonuclease II inhibitor	0,764	0,005	Hepatoprotectant
	0,905	0,005	Benzoate-CoA ligase inhibitor	0,766	0,007	Lipid metabolism regulator

	0,896	0,003	Cholesterol antagonist	0,761	0,004	Phosphatidylglycerophosphatase inhibitor	
	0,894	0,003	UDP-N-acetylglucosamine 4-epimerase inhibitor	0,761	0,007	Vasoprotector	
	0,888	0,003	Antihypercholesterolemic	0,761	0,008	Nucleotide metabolism regulator	
	0,885	0,002	Lactase inhibitor	0,753	0,003	H ⁺ -exporting ATPase inhibitor	
	0,879	0,003	Glucan endo-1,3-beta-D-glucosidase inhibitor	0,751	0,003	Free radical scavenger	
	0,876	0,003	Levanase inhibitor	0,749	0,004	N-acylmannosamine kinase inhibitor	
	0,877	0,006	Beta-adrenergic receptor kinase inhibitor	0,747	0,005	Antithrombotic	
	0,877	0,006	G-protein-coupled receptor kinase inhibitor	0,739	0,004	Mycothioli-S-conjugate amidase inhibitor	
	0,864	0,001	Laxative	0,736	0,003	Lactose synthase inhibitor	
	0,854	0,003	Anthranilate-CoA ligase inhibitor	0,738	0,006	IgA-specific metalloendopeptidase inhibitor	
	0,856	0,005	Fucosterol-epoxide lyase inhibitor	0,731	0,001	Oligo-1,6-glucosidase inhibitor	
	0,850	0,002	Alkenylglycerophosphoethanolamine hydrolase inhibitor	0,734	0,009	Glyceryl-ether monooxygenase inhibitor	
	0,838	0,002	Beta-amylase inhibitor	0,731	0,008	Antiprotozoal (Leishmania)	
	0,838	0,004	Ribulose-phosphate 3-epimerase inhibitor	0,723	0,002	Glucan 1,4-beta-glucosidase inhibitor	
	0,837	0,004	Anticarcinogenic	0,713	0,002	Beta-glucosidase inhibitor	
	0,833	0,004	Manganese peroxidase inhibitor	0,712	0,004	D-threo-aldose 1-dehydrogenase inhibitor	
	0,837	0,012	CYP2H substrate	0,709	0,004	Histamine release stimulant	
	0,827	0,002	Licheninase inhibitor	0,709	0,006	Aspartyltransferase inhibitor	
	catechin-[5,6-e]-4 b-(3,4-dihydroxyphenyl)dihydro-2(3H)-pyranone	0,934	0,005	Membrane integrity agonist	0,774	0,005	Antihypercholesterolemic
		0,924	0,004	CYP1A1 substrate	0,771	0,015	Antineoplastic
0,910		0,005	TP53 expression enhancer	0,748	0,010	Fibrinolytic	
0,892		0,003	UGT1A6 substrate	0,743	0,006	Hepatoprotectant	
0,883		0,016	CYP2C12 substrate	0,731	0,002	Astringent	
0,864		0,003	UGT1A substrate	0,735	0,010	Kinase inhibitor	
0,857		0,005	CYP1A substrate	0,728	0,011	CYP3A4 inducer	
0,836		0,004	UDP-glucuronosyltransferase substrate	0,713	0,004	Free radical scavenger	
0,805		0,003	Pectate lyase inhibitor	0,711	0,005	APOA1 expression enhancer	
0,798		0,004	Antimutagenic	0,713	0,011	CYP3A inducer	
0,798		0,008	CYP2B6 substrate	0,705	0,006	CYP2A11 substrate	
0,787		0,004	Lipid peroxidase inhibitor	0,737	0,040	Mucomembranous protector	
0,780		0,004	HMOX1 expression enhancer	0,702	0,006	CYP2A4 substrate	
0,786		0,013	HIF1A expression inhibitor	0,706	0,014	CYP3A5 substrate	
0,775		0,004	Chemopreventive				
2-(6-benzoyl-bD-glucopyranosyloxy)-7-(1 a,2 a,6 a-trihydroxy-3-oxocyclohex-4-enoyl)-5-hydroxybenzyl alcohol		0,939	0,004	Membrane integrity agonist	0,829	0,014	Alkenylglycerophosphocholine hydrolase inhibitor
	0,933	0,005	CDP-glycerol glycerophosphotransferase inhibitor	0,821	0,009	Antineoplastic	
	0,894	0,004	Anaphylatoxin receptor antagonist	0,810	0,005	Antiinfective	
	0,874	0,004	Membrane permeability inhibitor	0,769	0,006	Anticarcinogenic	
	0,854	0,003	Monophenol monooxygenase inhibitor	0,747	0,004	H ⁺ -exporting ATPase inhibitor	
	0,851	0,010	Benzoate-CoA ligase inhibitor	0,744	0,011	UDP-glucuronosyltransferase substrate	
	0,836	0,003	Lactase inhibitor	0,732	0,009	Mucinamylserine mucinamidase inhibitor	
	0,842	0,011	CYP2H substrate	0,715	0,014	Antiinflammatory	
	0,840	0,011	Sugar-phosphatase inhibitor	0,710	0,016	Exoribonuclease II inhibitor	
0,833	0,004	3-Phytase inhibitor					
Chrysoeriol-7-O- b-D-glucopyranoside	0,982	0,001	Monophenol monooxygenase inhibitor	0,834	0,006	NAD(P) ⁺ -arginine ADP-ribosyltransferase inhibitor	
	0,976	0,001	Membrane permeability inhibitor	0,829	0,005	CYP3A inducer	
	0,976	0,002	Membrane integrity agonist	0,830	0,009	Antineoplastic	
	0,974	0,001	Cardioprotectant	0,825	0,004	UGT1A9 substrate	
	0,973	0,001	Vasoprotector	0,810	0,001	Alpha glucosidase inhibitor	
	0,972	0,001	Free radical scavenger	0,821	0,013	Sugar-phosphatase inhibitor	
	0,971	0,001	Hemostatic	0,818	0,017	Chlordecone reductase inhibitor	
	0,959	0,000	Beta-N-acetylhexosaminidase inhibitor	0,807	0,006	Membrane integrity antagonist	
	0,956	0,003	TP53 expression enhancer	0,803	0,003	Antioxidant	
	0,951	0,002	Chemopreventive	0,795	0,001	Skin whitener	
	0,948	0,002	Hepatoprotectant	0,793	0,002	CYP2E1 inducer	
	0,945	0,002	CYP1A inducer	0,796	0,006	Kinase inhibitor	
	0,943	0,002	Anticarcinogenic	0,792	0,003	Mediator release inhibitor	
	0,943	0,002	Lipid peroxidase inhibitor	0,784	0,004	3-Phytase inhibitor	
	0,940	0,001	NADPH oxidase inhibitor	0,782	0,005	Antiinfective	
	0,941	0,003	Caspase 3 stimulant	0,777	0,004	UGT1A1 substrate	
	0,940	0,003	Anaphylatoxin receptor antagonist	0,767	0,002	Caspase 8 stimulant	
0,940	0,003	UDP-glucuronosyltransferase substrate	0,770	0,010	Apoptosis agonist		

	0,930	0,002	Antiprotozoal (Leishmania)	0,763	0,007	Mucinamylserine mucinamidase inhibitor
	0,929	0,005	CDP-glycerol glycerophosphotransferase inhibitor	0,754	0,002	Sweetener
	0,923	0,003	Antihypercholesterolemic	0,756	0,004	Mycothioliol-S-conjugate amidase inhibitor
	0,920	0,001	CYP1A1 inducer	0,760	0,010	2-Dehydropantoate 2-reductase inhibitor
	0,913	0,001	4-Coumarate-CoA ligase inhibitor	0,772	0,022	Alkenylglycerophosphocholine hydrolase inhibitor
	0,895	0,001	Histamine release stimulant	0,743	0,005	Cyclic AMP phosphodiesterase inhibitor
	0,893	0,001	CYP2C9 inducer	0,745	0,010	Oxidoreductase inhibitor
	0,893	0,001	Laxative	0,736	0,006	Vasodilator
	0,892	0,000	Capillary fragility treatment	0,729	0,002	Glucan 1,6-alpha-glucosidase inhibitor
	0,897	0,006	HIF1A expression inhibitor	0,733	0,008	Radioprotector
	0,887	0,003	UGT1A substrate	0,750	0,026	CYP2H substrate
	0,881	0,001	UGT1A7 substrate	0,727	0,004	Beta glucuronidase inhibitor
	0,881	0,003	Sulfotransferase substrate	0,725	0,010	Nucleotide metabolism regulator
	0,874	0,002	Proliferative diseases treatment	0,719	0,005	Antiviral (Influenza)
	0,871	0,001	Xanthine dehydrogenase inhibitor	0,719	0,007	Histidine kinase inhibitor
	0,863	0,004	CYP3A4 inducer	0,715	0,004	Choleretic
	0,845	0,003	Lactase inhibitor	0,706	0,007	HMOX1 expression enhancer
	0,847	0,005	Cytostatic	0,707	0,009	Antifungal
	0,852	0,010	Benzoate-CoA ligase inhibitor	0,701	0,004	Antitussive
	0,833	0,001	Aryl hydrocarbon receptor agonist			
Mururin A	0,934	0,005	Membrane integrity agonist	0,808	0,006	CYP1A substrate
	0,925	0,004	TP53 expression enhancer	0,781	0,003	Pectate lyase inhibitor
	0,900	0,002	Antimutagenic	0,777	0,004	CYP2A11 substrate
	0,873	0,004	UGT1A6 substrate	0,777	0,004	General pump inhibitor
	0,861	0,003	Histidine kinase inhibitor	0,747	0,004	Histamine release inhibitor
	0,861	0,003	UGT1A substrate	0,745	0,005	HMOX1 expression enhancer
	0,861	0,004	CYP1A1 substrate	0,739	0,006	Hepatoprotectant
	0,843	0,003	Chemopreventive	0,731	0,004	Free radical scavenger
	0,841	0,004	UDP-glucuronosyltransferase substrate	0,730	0,004	CYP1A inhibitor
	0,841	0,008	Antineoplastic	0,726	0,005	UGT1A1 substrate
	0,855	0,023	CYP2C12 substrate	0,714	0,005	Antineoplastic (breast cancer)
	0,821	0,004	Sulfotransferase substrate	0,705	0,006	CYP2A4 substrate
	0,822	0,010	HIF1A expression inhibitor			

*probability of activity, # probability of inactivity

Conclusion

The ethnobotanicals have been regularly used since the ancient civilization as ethnomedicine's for various illness from simple cold to threat full diseases like cancer etc., As recent report of WHO testifies that about 80% of the world's population relies on the phytopharmaceuticals for the treatment of various common diseases. In the era of many emerging and re-emerging diseases phytochemical compounds derived from plant sources have great pharmacological importance. The cure for a particular disease is addressed by a potent lead molecule for their biological activities against the disease therefore; computational approach has led to the identification of drug target and in the prediction of novel leads. In present study we have analyzed six isolated compounds from *Flacourtia indica* through QSAR approach computationally; it is interesting to reveal that all the six compounds were nontoxic and showing good ADMET profiles and high drug likeness properties with many biological activities and mechanism of actions. Screening of bioactive compounds from plants as source, has restored health benefits due to their biological activities, thus in the present study phytochemicals from *Flacourtia indica* provides ethnomedical evidences as potential phytopharmaceuticals, however further *In vitro* and *In vivo* analysis of each compound for various pharmacological benefits can be carried out for the discovery of novel drug compounds.

References

- Ahmad M, Khan MA, Marwat SK, Zafar M, Khan MA, Hassan TU et al. Useful medicinal flora enlisted in Holy Quran and Ahadith. Am Eurasian J Agric Environ Sci. 2009; 5(1):126-40.
- Ahmed S, Hasan M. Muslim heritage in medicine: A concise review on Greco-Arabic contribution. Journal of Pharmacognosy and Phytochemistry. 2016; 5(4):273-83.
- Alavijeh MS, Chishty M, Qaiser MZ, Palmer AM. Drug metabolism and pharmacokinetics, the blood-brain barrier, and central nervous system drug discovery. NeuroRx. 2005; 2(4):554-71.
- Balasundaram A, Ragupathy R, Sankar S, Thiyagarajan M, Ravi L, Karuppasamy R et al. Investigation of Phytocompounds and Computational Approach for the Evaluation of Therapeutic Properties of Ethanolic Leaf Extract of *Callistemon citrinus*., Int. J Pharm. Sci. Rev. Res., 2016; 37(1):20,
- Baldi A. Computational approaches for drug design and discovery: an overview. Systematic Reviews in Pharmacy. 2010; 1(1):99.
- David Taylor, the Pharmaceutical Industry and the Future of Drug Development, in Pharmaceuticals in the Environment, 2015, 1-33.
- De Fátima A, Modolo LV, Conegero LS, Pilli RA, Ferreira CV, Kohn LK et al. Styryl lactones and their derivatives: biological activities, mechanisms of action and potential leads for drug design. Current medicinal chemistry. 2006; 13(28):3371-84.

8. De Silva T. Industrial utilization of medicinal plants in developing countries. Rome: Medicinal plants for forest conservation and health care FAO. 1997, 34-44.
9. Eramma N, Gayathri D. Antibacterial potential and phytochemical analysis of *Flacourtia indica* (Burm. f.) Merr. root extract against human pathogens. Indo Am J Pharm Res. 2013; 3:3832-46.
10. Goel RK, Singh D, Lagunin A, Poroikov V. PASS-assisted exploration of new therapeutic potential of natural products. Medicinal Chemistry Research. 2011; 20(9):1509-14.
11. Khurana N, Ishar MP, Gajbhiye A, Goel RK. PASS assisted prediction and pharmacological evaluation of novel nicotinic analogs for nootropic activity in mice. European journal of pharmacology. 2011; 662(1):22-30.
12. Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. Advanced drug delivery reviews. 2012; 64:4-17.
13. Lydia J, Sudarsanam D. Anticarcinogenic Effects of Selected Alkaloids Identified from *Cyperus rotundus*: An In Silico Perspective. J Appl Bioinform Comput Biol. 2015; 1:2.
14. Mandal S, Moudgil MN, Mandal SK. Rational drug design. European journal of pharmacology. 2009; 625(1):90-100.
15. Mathew S, Sreekumar S, Biju CK. Identification of lead compounds against human hepatitis B viral capsid protein in three medicinal plants through in silico method. IOSR Journal of Pharmacy and Biological Sciences, 2016, 01-06
16. Menezes JC, Kamat SP, Cavaleiro JA, Gaspar A, Garrido J, Borges F. Synthesis and antioxidant activity of long chain alkyl hydroxycinnamates. European journal of medicinal chemistry. 2011; 46(2):773-7.
17. Ntie-Kang F. An in silico evaluation of the ADMET profile of the Streptome DB database. Springer Plus. 2013; 2(1):1.
18. Paramashivam SK, Elayaperumal K, Natarajan BB, Ramamoorthy MD, Balasubramanian S, Dhiraviam KN. In silico pharmacokinetic and molecular docking studies of small molecules derived from *Indigofera aspalathoides* Vahl targeting receptor tyrosine kinases. Bioinformation. 2015; 11(2):73-84.
19. Rahman MT, Nazer R, Brown L, Shogar I, Bouzenita AI. Therapeutic interventions: an Islamic perspective. Journal of the Islamic Medical Association of North America. 2008; 40(2).
20. Sashidhara KV, Singh SP, Singh SV, Srivastava RK, Srivastava K, Saxena JK et al. Isolation and identification of β -hematin inhibitors from *Flacourtia indica* as promising antiplasmodial agents. European journal of medicinal chemistry. 2013; 60:497-502.
21. Sharma V, Sarkar IN. Bioinformatics opportunities for identification and study of medicinal plants. Briefings in bioinformatics. 2013; 14(2):238-50.
22. World Health Organization. WHO traditional medicine strategy 2002-2005.