

Genome Sequence of Avian *Escherichia coli* Strain IHIT25637, an Extraintestinal Pathogenic *E. coli* Strain of ST131 Encoding Colistin Resistance Determinant MCR-1

Christa Ewers,^a Stephan Göttig,^b Maria Bülte,^a Sophie Fiedler,^a Manuela Tietgen,^b Ursula Leidner,^a Carsten Heydel,^a Rolf Bauerfeind,^a Torsten Semmler^c

Institute of Hygiene and Infectious Diseases of Animals, Justus Liebig University, Giessen, Germany^a; Institute of Medical Microbiology and Infection Control, Hospital of the Johann Wolfgang von Goethe-University, Frankfurt am Main, Germany^b; Robert Koch Institute, Berlin, Germany^c

Sequence type 131 (ST131) is one of the predominant *Escherichia coli* lineages among extraintestinal pathogenic *E. coli* (ExPEC) that causes a variety of diseases in humans and animals and frequently shows multidrug resistance. Here, we report the first genome sequence of an ST131-ExPEC strain from poultry carrying the plasmid-encoded colistin resistance gene *mcr-1*.

Received 1 July 2016 Accepted 6 July 2016 Published 1 September 2016

Citation Ewers C, Göttig S, Bülte M, Fiedler S, Tietgen M, Leidner U, Heydel C, Bauerfeind R, Semmler T. 2016. Genome sequence of avian *Escherichia coli* strain IHIT25637, an extraintestinal pathogenic *E. coli* strain of ST131 encoding colistin resistance determinant MCR-1. *Genome Announc* 4(5):e00863-16. doi:10.1128/genomeA.00863-16.

Copyright © 2016 Ewers et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christa Ewers, christa.ewers@vetmed.uni-giessen.de.

Extraintestinal pathogenic *Escherichia coli* (ExPEC) strains are a versatile group of bacteria revealing a complex phylogeny and considerable genomic plasticity. ExPEC strains cause a variety of disease syndromes, ranging from uncomplicated urinary tract infections to life-threatening bloodstream infections in humans and septicemia, peritonitis, and yolk sac infections in poultry (1, 2). Certain subpopulations of avian ExPEC have been suggested to be zoonotic agents (3, 4), which is of great concern, because they also have been associated with multidrug resistant clonal lineages, such as sequence type 131 (ST131) (2, 5).

We report the genome sequence of colistin-resistant avian ExPEC strain IHIT25637 carrying the plasmid-mediated *mcr-1* gene, which was described first in human and food animal samples in China in 2015 (6). IHIT25637 was isolated in Germany in October 2010 from the liver of a broiler chicken that had died of septicemia. Whole-genome sequencing was performed on a MiSeq sequencer using a MiSeq reagent kit v3 (Illumina, USA) resulting in 300 bp paired-end reads and an average coverage of 100×. Filtered 1.75 M clean reads were *de novo* assembled after quality trimming using SPAdes v3.7.1 (7) into scaffolds, and corresponding 90-fold coverage of the genome was generated. The draft genome sequence of *E. coli* IHIT25637 was 5,692,160 bp in size and had a G+C content of 50.4% in 183 contigs (maximum length, 402,413 bp; minimum length, 200 bp; N_{50} contig size 234,455 bp; mean coverage 90). The genome was annotated using the Rapid Annotation using Subsystem Technology (RAST) server (8). Annotation of this assembly identified 5,614 coding sequences (CDSs), 85 tRNAs, and 19 rRNAs.

The sequence type of IHIT25637 was identified as ST131 (<https://cge.cbs.dtu.dk/services/>), and its serotype was O25:H4 as determined by slide agglutination testing. Several virulence-related genes known to be associated with the ExPEC pathotype, like adhesin genes *csgA*, *fim*, *iha*, *tsh*, and *focA*, iron-acquisition related gene *chuA*, *iroN*, *eitA*, *etsB/C*, *fyuA*, *irp2*, *iucD*, and *iutA*, and invasion, serum resistance and capsule synthesis genes *ibeA*, *iss*, *kpsMTIII*, and *neuC*, were present in strain IHIT25637. The allele type of the type I fimbrial adhesin gene was determined as *fimH22*,

which has been associated with a fluoroquinolone-susceptible ST131-subgroup (9). Genome data revealed the presence of colistin resistance gene *mcr-1* which most likely conferred phenotypic resistance to the “last resort” antibiotics colistin and polymyxin B (MIC: 8 mg/L). *Mcr-1* was located on an IncHI2 plasmid (>200 kbp) of plasmid sequence type ST4 (10), which are major carriers of *mcr-1* in *E. coli* isolates from livestock and food and also from humans (11–13). In addition to *mcr-1*, the genome of IHIT25637 carried several resistance genes, such as *aph(3′)-Ic*, *aac(3)-IIa*, *aadA1*, *aadA2*, *strA* and *strB*, *bla_{TEM-1}*, *cmlA1*, *catA1* *sul1/2/3*, *dfrA1*, and *tet(A)*, but lacked genes for known extended-spectrum β -lactamases and carbapenemases. Only two ST131 *E. coli* strains carrying *mcr-1* have yet been described, namely, from chicken meat of European origin imported to Denmark in 2012 and from the urine of a human patient in Taiwan in 2014 (14, 15). However, the appearance of *mcr-1*-harboring, colistin resistant ExPEC belonging to the epidemiological successful clonal lineage ST131 poses a serious public health threat.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number MAIV00000000. The version described in this paper is version MAIV00000000.1.

ACKNOWLEDGMENTS

We thank Petra Krienke, Inga Eichhorn, and Bettina Hopf for excellent technical assistance.

FUNDING INFORMATION

This work, including the efforts of Christa Ewers, was funded by Deutsche Forschungsgemeinschaft (DFG) (Ew-116/2-1). This work, including the efforts of Christa Ewers, Maria Bülte, Sophie Fiedler, and Ursula Leidner, was funded by Bundesministerium für Ernährung und Landwirtschaft (BMEL) (2817701314).

REFERENCES

- Ewers C, Li G, Wilking H, Kiessling S, Alt K, Antão EM, Laturnus C, Diehl I, Glodde S, Homeier T, Böhnke U, Steinrück H, Philipp HC,

- Wieler LH. 2007. Avian pathogenic, uropathogenic, and newborn meningitis-causing *Escherichia coli*: how closely related are they? *Int J Med Microbiol* 297:163–176. <http://dx.doi.org/10.1016/j.ijmm.2007.01.003>.
2. Dale AP, Woodford N. 2015. Extra-intestinal pathogenic *Escherichia coli* (ExPEC): disease, carriage and clones. *J Infect* 71:615–626. <http://dx.doi.org/10.1016/j.jinf.2015.09.009>.
 3. Mellata M. 2013. Human and avian extraintestinal pathogenic *Escherichia coli*: infections, zoonotic risks, and antibiotic resistance trends. *Foodborne Pathog Dis* 10:916–932. <http://dx.doi.org/10.1089/fpd.2013.1533>.
 4. Nandanwar N, Janssen T, Kühl M, Ahmed N, Ewers C, Wieler LH. 2014. Extraintestinal pathogenic *Escherichia coli* (ExPEC) of human and avian origin belonging to sequence type complex 95 (STC95) portray indistinguishable virulence features. *Int J Med Microbiol* 304:835–842. <http://dx.doi.org/10.1016/j.ijmm.2014.06.009>.
 5. Ewers C, Bethe A, Semmler T, Guenther S, Wieler LH. 2012. Extended-spectrum beta-lactamase-producing and AmpC-producing *Escherichia coli* from livestock and companion animals, and their putative impact on public health: a global perspective. *Clin Microbiol Infect* 18:646–655. <http://dx.doi.org/10.1111/j.1469-0691.2012.03850.x>.
 6. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J. 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* 16:161–168. [http://dx.doi.org/10.1016/S1473-3099\(15\)00424-7](http://dx.doi.org/10.1016/S1473-3099(15)00424-7).
 7. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
 8. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
 9. Banerjee R, Johnson JR. 2014. A new clone sweeps clean: the enigmatic emergence of *Escherichia coli* sequence type 131. *Antimicrob Agents Chemother* 58:4997–5004. <http://dx.doi.org/10.1128/AAC.02824-14>.
 10. García-Fernández A, Carattoli A. 2010. Plasmid double locus sequence typing for IncHI2 plasmids, a subtyping scheme for the characterization of IncHI2 plasmids carrying extended-spectrum beta-lactamase and quinolone resistance genes. *J Antimicrob Chemother* 65:1155–1161. <http://dx.doi.org/10.1093/jac/dkq101>.
 11. Veldman K, van Essen-Zandbergen A, Rapallini M, Wit B, Heymans R, van Pelt W, Mevius D. 2016. Location of colistin resistance gene *mcr-1* in *Enterobacteriaceae* from livestock and meat. *J Antimicrob Chemother* 71:2340–2342. <http://dx.doi.org/10.1093/jac/dkw181>.
 12. Haenni M, Poirel L, Kieffer N, Châtre P, Saras E, Métayer V, Dumoulin R, Nordmann P, Mavec J. 2016. Co-occurrence of extended spectrum beta lactamase and MCR-1 encoding genes on plasmids. *Lancet Infect Dis* 16:281–282. [http://dx.doi.org/10.1016/S1473-3099\(16\)00007-4](http://dx.doi.org/10.1016/S1473-3099(16)00007-4).
 13. Falgenhauer L, Waezsada SE, Yao Y, Imirzalioglu C, Kasbohrer A, Roesler U, Michael GB, Schwarz S, Werner G, Kreienbrock L, Chakraborty T. 2016. Colistin resistance gene *mcr-1* in extended-spectrum β -lactamase-producing and carbapenemase-producing Gram-negative bacteria in Germany. *Lancet Infect Dis* 16:282–283. [http://dx.doi.org/10.1016/S1473-3099\(16\)00009-8](http://dx.doi.org/10.1016/S1473-3099(16)00009-8).
 14. Hasman H, Hammerum AM, Hansen F, Hendriksen RS, Olesen B, Agersø Y, Zankari E, Leekitcharoenphon P, Stegger M, Kaas RS, Cavaco LM, Hansen DS, Aarestrup FM, Skov RL. 2015. Detection of *mcr-1* encoding plasmid-mediated colistin-resistant *Escherichia coli* isolates from human bloodstream infection and imported chicken meat, Denmark 2015. *Euro Surveill* 20:49. <http://dx.doi.org/10.2807/1560-7917.ES.2015.20.49.30085>.
 15. Kuo SC, Huang WC, Wang HY, Shiau YR, Cheng MF, Lauderdale TL. 2016. Colistin resistance gene *mcr-1* in *Escherichia coli* isolates from humans and retail meats, Taiwan. *J Antimicrob Chemother* 71:1922–1932. <http://dx.doi.org/10.1093/jac/dkw122>.