



Hridi, Shehla Unaiza and McCann, Lee and McConnell, Gail and Bushell, Trevor J and Jiang, Hui-Rong (2017) Increased expression of IL-16 in the brain of experimental autoimmune encephalomyelitis. Brain and Neuroscience Advances. ISSN 2398-2128 ,

This version is available at <https://strathprints.strath.ac.uk/60998/>

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: strathprints@strath.ac.uk

Multiple Sclerosis (MS) is a demyelinating disease of the CNS, whose pathophysiology involves both inflammatory and neurodegenerative components. CD4⁺ T cells are one of the key mediators of disease initiation and progression; however CD4 is also the receptor for the pro-inflammatory cytokine, interleukin - 16 (IL - 16). IL - 16 has been proposed to play a role in several autoimmune diseases, but the exact role of IL - 16 in the CNS during MS initiation and progression remains unclear. Therefore, the aim of this study was to examine the expression and distribution of IL - 16 in CNS tissue and investigate whether expression levels correlate with neuro-inflammation in experimental autoimmune encephalomyelitis (EAE), a murine model of MS.

EAE was induced in 6 week old C 57BL/6J female mice by immunisation with MOG35 - 55 peptide and adjuvants. Tissue was harvested at onset (day 11), peak (day 16) and resolution (day 26), and immunofluorescence staining carried out to determine CD45, CD4 and IL - 16 expression and localisation in the brain of both control and EAE mice. In addition, co-localisation of IL - 16 with CNS and immune cell subtypes was performed using a Mesolens microscope (McConnell et al., 2016), which allows subcellular detail to be obtained from wide - field epifluorescence images.

Expression of IL - 16 and CD4 was observed primarily within the lesions of cerebellum and hippocampus of the EAE brain, whereas little expression was observed in control brains. IL - 16 expression was highest at onset with $76 \pm 2.8\%$ of cells (n=3) within these lesions expressing IL - 16. This was reduced to $48 \pm 2.4\%$ (n=3) at peak and $16 \pm 1.3\%$ at resolution (n=3). Co-localization studies revealed that IL - 16 was expressed primarily by infiltrating immune cells but not by neurons or astrocytes. Co-localization of IL - 16 with immune cells in brain lesions of EAE mice suggests that infiltrating immune cells are the primary source of IL - 16. Further investigation is required if IL - 16 is pro-inflammatory or anti-inflammatory in the CNS during EAE.

References:

McConnell G et al., (2016). A novel optical microscope for imaging large embryos and tissue volumes with sub-cellular resolution throughout. E-life. <http://dx.doi.org/10.7554/eLife.18659.001>