

Effect of nitrogen source and acclimatization on specific growth rates of microalgae determined by a high throughput in vivo microplate autofluorescence method

Podevin, Michael Paul Ambrose; De Francisci, Davide; Holdt, Susan Løvstad; Fotidis, Ioannis; Angelidaki, Irini

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Podevin, M. P. A., De Francisci, D., Holdt, S. L., Fotidis, I., & Angelidaki, I. (2017). Effect of nitrogen source and acclimatization on specific growth rates of microalgae determined by a high throughput in vivo microplate autofluorescence method.

DTU Library

Technical Information Center of Denmark


General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Effect of nitrogen source and acclimatization on specific growth rates of microalgae determined by a high throughput *in vivo* microplate autofluorescence method



Michael Podevin^{1,3} • Davide De Francisci¹ • Susan L. Holdt^{1,2} • Ioannis Fotidis¹ • Irini Angelidaki¹

¹ Department of Environmental Engineering, Technical University of Denmark, Building 113, DK-2800 Kgs. Lyngby, Denmark

² Current address: Susan L. Holdt: National Food Institute, Technical University of Denmark, Søtofts Plads 221, DK-2800 Kgs. Lyngby, Denmark


³ E-mail: mipp@env.dtu.dk

Abstract


Specific growth rates (SGR) of fresh water algal strains (*Chlorella vulgaris*, *Auxenochlorella protothecoides*, and *Chlorella sorokiniana*) and the marine strain *Nannochloropsis oculata* on various nitrogen sources (ammonium carbonate, ammonium chloride, sodium nitrate, and urea) could be determined by *in vivo* chlorophyll-a autofluorescence. These preferences could be determined before large pH changes occurred in the media, with no significant difference ($P > 0.05$) between buffered and non-buffered media. In all algal species, acclimatization was observed with no significant difference ($P > 0.05$) between SGRs of second and third cultivations. ANOVA of SGRs in the acclimatized second and third cultivation revealed preferences for nitrogen sources among most of the algae; *C. vulgaris* preferred sodium nitrate over other nitrogen sources, *A. protothecoides* adapted to urea after no growth in the first cultivation, and the SGRs of *N. oculata* showed an aversion for sodium nitrate over other nitrogen sources ($P < 0.05$).

Aims

- Developing a microplate screening technique for high throughput screenings of microalgae.
- to determine the preference of nitrogen source for algal growth for four industrially important microalgae
- Determine the acclimatized SGR of each algae species to each nitrogen source.
- Determine the SGR before large changes in pH occur.



Results



Conclusions

- In vivo* microplate batch cultures can be used to compare specific growth rates of microalgae containing chlorophyll-a.
- This method can be used to determine the specific growth rate before significant pH changes in the media occur, making the method useful for comparison of various modes of nitrogen assimilation.
- It has been demonstrated that, at certain conditions, sodium nitrate is the most preferred nitrogen source and urea is the least favored nitrogen source for *Chlorella vulgaris*.
- Auxenochlorella protothecoides* demonstrated an acclimatization to urea, with no growth occurring in nitrate media in autotrophic conditions
- Nannochloropsis oculata* showed a clear aversion for sodium nitrate as a nitrogen source.