

# Metabolic engineering of yeast for fermentative production of flavonoids - DTU Orbit (09/11/2017)

## Metabolic engineering of yeast for fermentative production of flavonoids

Yeast *Saccharomyces cerevisiae* was engineered for de novo production of six different flavonoids (naringenin, liquiritigenin, kaempferol, resokaempferol, quercetin, and fisetin) directly from glucose, without supplementation of expensive intermediates. This required reconstruction of long biosynthetic pathways, comprising up to eight heterologous genes from plants. The obtained titers of kaempferol  $26.57 \pm 2.66 \text{ mgL}^{-1}$  and quercetin  $20.38 \pm 2.57 \text{ mgL}^{-1}$  exceed the previously reported titers in yeast. This is also the first report of de novo biosynthesis of resokaempferol and fisetin in yeast. The work demonstrates the potential of flavonoid-producing yeast cell factories.

### General information

State: Published

Organisations: Novo Nordisk Foundation Center for Biosustainability, Yeast Cell Factories, Research Groups, iLoop, Department of Biotechnology and Biomedicine, Natural Product Discovery, Yeast Metabolic Engineering

Authors: Rodriguez Prado, E. A. (Intern), Strucko, T. (Intern), Stahlhut, S. G. (Intern), Kristensen, M. (Intern), Svenssen, D. K. (Intern), Förster, J. (Intern), Nielsen, J. (Intern), Borodina, I. (Intern)

Pages: 1645-1654

Publication date: 2017

Main Research Area: Technical/natural sciences

### Publication information

Journal: Bioresource Technology

Volume: 245

Issue number: Part B

ISSN (Print): 0960-8524

Ratings:

BFI (2017): BFI-level 2

Web of Science (2017): Indexed yes

BFI (2016): BFI-level 2

Scopus rating (2016): CiteScore 5.94 SJR 2.191 SNIP 1.91

Web of Science (2016): Indexed yes

BFI (2015): BFI-level 2

Scopus rating (2015): SJR 2.255 SNIP 1.908 CiteScore 5.47

Web of Science (2015): Indexed yes

BFI (2014): BFI-level 2

Scopus rating (2014): SJR 2.41 SNIP 2.104 CiteScore 5.3

Web of Science (2014): Indexed yes

BFI (2013): BFI-level 2

Scopus rating (2013): SJR 2.412 SNIP 2.503 CiteScore 5.97

ISI indexed (2013): ISI indexed yes

Web of Science (2013): Indexed yes

BFI (2012): BFI-level 2

Scopus rating (2012): SJR 2.389 SNIP 2.465 CiteScore 5.25

ISI indexed (2012): ISI indexed yes

Web of Science (2012): Indexed yes

BFI (2011): BFI-level 2

Scopus rating (2011): SJR 2.314 SNIP 2.508 CiteScore 5.56

ISI indexed (2011): ISI indexed yes

Web of Science (2011): Indexed yes

BFI (2010): BFI-level 2

Scopus rating (2010): SJR 2.086 SNIP 2.355

Web of Science (2010): Indexed yes

BFI (2009): BFI-level 2

Scopus rating (2009): SJR 1.912 SNIP 2.231

Web of Science (2009): Indexed yes

BFI (2008): BFI-level 2

Scopus rating (2008): SJR 1.734 SNIP 2.732

Web of Science (2008): Indexed yes

Scopus rating (2007): SJR 1.529 SNIP 2.423

Web of Science (2007): Indexed yes

Scopus rating (2006): SJR 1.315 SNIP 1.98

Web of Science (2006): Indexed yes

Scopus rating (2005): SJR 1.269 SNIP 2.006

Web of Science (2005): Indexed yes

Scopus rating (2004): SJR 1.197 SNIP 1.659

Web of Science (2004): Indexed yes

Scopus rating (2003): SJR 0.948 SNIP 1.639

Web of Science (2003): Indexed yes

Scopus rating (2002): SJR 0.882 SNIP 1.3

Web of Science (2002): Indexed yes

Scopus rating (2001): SJR 0.541 SNIP 1.208

Scopus rating (2000): SJR 0.464 SNIP 1.049

Scopus rating (1999): SJR 0.669 SNIP 1.061

Original language: English

Fisetin, Flavonoids, Kaempferol, Resokaempferol, *Saccharomyces cerevisiae*

Electronic versions:

[1\\_s2.0\\_S0960852417309306\\_main.pdf](#)

DOIs:

[10.1016/j.biortech.2017.06.043](#)

Source: FindIt

Source-ID: 2371399339

Publication: Research - peer-review > Journal article – Annual report year: 2017