

## Profiling bacterial kinase activity using a genetic circuit

van der Helm, Eric; Bech, Rasmus; Lehning, Christina Eva; Vazquez-Uribe, Ruben; Sommer, Morten Otto Alexander

*Link to article, DOI:*  
[10.17605/OSF.IO/AKG5X](https://doi.org/10.17605/OSF.IO/AKG5X)

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
van der Helm, E., Bech, R., Lehning, C. E., Vazquez-Uribe, R., & Sommer, M. O. A. (2017). Profiling bacterial kinase activity using a genetic circuit. Novo Nordisk Foundation Center for Biosustainability. DOI: 10.17605/OSF.IO/AKG5X

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



The Novo Nordisk Foundation  
Center for Biosustainability


# PROFILING BACTERIAL KINASE ACTIVITY USING A GENETIC CIRCUIT

Eric van der Helm, Rasmus Bech, Christina Lehning, Ruben Vazquez-Urbe, Morten Sommer


Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark

Phosphorylation is a post-translational modification that regulates the activity of several key proteins in bacteria and eukaryotes. Accordingly, a variety of tools has been developed to measure kinase activity. To couple phosphorylation to an *in vivo* fluorescent readout we used the *Bacillus subtilis* kinase PtkA, transmembrane activator TkmA and the repressor FatR to construct a genetic circuit in *E. coli*. By tuning the repressor and kinase expression level at the same time, we were able to show a 4.2-fold increase in signal upon kinase induction. We furthermore validated that the previously reported FatR Y45E mutation<sup>1</sup> attenuates operator repression. This genetic circuit provides a starting point for computational protein design and a metagenomic library-screening tool.

## 1. Design: sensing phosphorylation




**Figure 1a:** General concept of kinase activity measurement through GFP fluorescence. The repressor binds the operator site upstream of *gfp*, thereby preventing transcription. The kinase phosphorylates the repressor, this attenuates binding of the repressor to the operator site, leading to transcription of *gfp* and thus fluorescence.




**Figure 1b:** An abstraction of the genetic circuit where the first inverter gate represents the phosphorylation of FatR by PtkA-TkmA on the post-translational level. The second inverter gate represents the transcriptional regulation by the binding of FatR to the operator upstream of *gfp*.

## 2. Build: circuit on two plasmids




**Figure 2:** Outline of the implementation with FatR as repressor and PtkA as kinase. The kinase PtkA and kinase-activator TkmA expression is induced by the benzoate/XylS system. The PtkA-TkmA complex can phosphorylate the repressor FatR, which attenuates the DNA binding of FatR to the operator site in front of *gfp*. Derepression of the operator site leads to expression of GFP and thus an increase in GFP/mCherry ratio. Different promoter strengths are used to drive the transcription of the *fatR* repressor. mCherry fluorescence is used to normalize the fluorescent signal of GFP.


## 3. Test: using Fluorescence Activated Cell Sorting (FACS)



**Figure 3a:** Raw FACS counts of 5 different strains with various *fatR* (repressor) genotypes under conditions where the kinase operon is not expressed. These strains each display a different *fatR* activity, which is reflected in the signal in the GFP channel (FITC-A). A strong promoter driving *fatR* results in a low GFP signal (green), whereas the strain containing no *fatR* shows a high GFP signal (red). Only one biological replicate of each strain is shown for clarity. This is the data used to calculate the values in Figure 3b.



**Figure 3b:** GFP/mCherry ratio measured of 5 different strains with various *fatR* (repressor) genotypes (based on data from Figure 3a). The two negative control reporters *fatR*:Y45E and  $\Delta fatR$  show an expected high ratio of GFP/mCherry. A stronger promoter driving *fatR* transcription results in more repression of GFP and thus a lower GFP/mCherry ratio as expected.



**Fig 3c:** Circuit response of *E. coli* containing both the platform plasmid and the kinase plasmid. Increasing amounts of benzoate induces the expression of the kinase PtkA and activation domain TkmA. Error bars represent SD of 3 biological replicates. **d:** The fold increase of GFP/mCherry between 0 and 2 mM benzoate. In the negative control strain ( $\Delta fatR$ , red) the fold increase in GFP/mCherry ratio is  $2.99 \pm 0.05$  upon kinase induction. The strains containing a functional *fatR* gene (green shades), (++) and (+), show a  $3.57 \pm 0.13$  and  $4.17 \pm 0.09$ -fold increase in the GFP/mCherry ratio upon kinase induction, as expected. The strain containing (+++) with the strongest promoter driving *fatR* transcription shows only a  $2.33 \pm 0.04$ -fold increase in GFP/mCherry ratio by inducing the kinase. Error bars denote SD of three biological replicates.

[1] Derouiche A, et al. *Nucleic Acids Res* 2013, 41:9371–81

