



Elst, A. F. M. ter and Gordon, G. and Waurick, M. (2017) The Dirichlet-to-Neumann operator for divergence form problems. Working paper. arXiv.org, Ithaca, NY.. ,

This version is available at <https://strathprints.strath.ac.uk/61320/>

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: strathprints@strath.ac.uk

The Strathprints institutional repository (<https://strathprints.strath.ac.uk>) is a digital archive of University of Strathclyde research outputs. It has been developed to disseminate open access research outputs, expose data about those outputs, and enable the management and persistent access to Strathclyde's intellectual output.

The Dirichlet-to-Neumann operator for divergence form problems

A.F.M. ter Elst, G. Gordon and M. Waurick

July 19, 2017

Abstract

We present a way of defining the Dirichlet-to-Neumann operator on general Hilbert spaces using a pair of operators for which each one's adjoint is formally the negative of the other. In particular, we define an abstract analogue of trace spaces and are able to give meaning to the Dirichlet-to-Neumann operator of divergence form operators perturbed by a bounded potential in cases where the boundary of the underlying domain does not allow for a well-defined trace. Moreover, a representation of the Dirichlet-to-Neumann operator as a first-order system of partial differential operators is provided. Using this representation, we address convergence of the Dirichlet-to-Neumann operators in the case that the appropriate reciprocals of the leading coefficients converge in the weak operator topology. We also provide some extensions to the case where the bounded potential is not coercive and consider resolvent convergence.

Keywords: Dirichlet-to-Neumann operator, resolvent convergence, continuous dependence on the coefficients.

MSC 2010: 35F45, 46E35, 47A07.

1 Introduction

In the theory of elliptic partial differential operators, the Dirichlet-to-Neumann operator is a central object of study. In recent years it attracted a lot of attention and triggered profound research in many directions. In particular, we mention applications of the form method, relations to the extension theory of symmetric operators as well as the intimate connection to the Calderón problem, see, for instance, the references in [BE1].

The Dirichlet-to-Neumann operator relates Dirichlet boundary data to the corresponding Neumann boundary data of solutions to a partial differential equation. As an introduction, we provide a definition for the Dirichlet-to-Neumann operator in its arguably simplest form.

Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with smooth boundary $\Gamma = \partial\Omega$ and where $d \geq 2$. Note that in this case, the trace map Tr from $H^1(\Omega)$ into $H^{1/2}(\Gamma)$ is a well-defined, surjective and

continuous operator. Let $\varphi \in H^{1/2}(\Gamma)$ and let $u \in H^1(\Omega)$ be the solution of the boundary value problem

$$-\Delta u = 0 \text{ weakly on } \Omega \quad \text{and} \quad \text{Tr } u = \varphi.$$

The Dirichlet-to-Neumann operator Λ assigns to φ the normal derivative of u , that is, $\Lambda \varphi = \partial_\nu u \in H^{-1/2}(\Gamma)$.

We can also consider the part of Λ in $L_2(\Gamma)$. If we call this restriction $\Lambda_{L_2(\Gamma)}$, then $\Lambda_{L_2(\Gamma)}$ is an unbounded operator in $L_2(\Gamma)$ such that for all $\varphi, \psi \in L_2(\Gamma)$ it follows that $\varphi \in \text{dom}(\Lambda_{L_2(\Gamma)})$ and $\Lambda_{L_2(\Gamma)}\varphi = \psi$ if and only if there exists a $u \in H^1(\Omega)$ such that $-\Delta u = 0$ weakly on Ω , $\text{Tr } u = \varphi$ and $\psi = \partial_\nu u$. A problem with the above descriptions is that they only make sense if the boundary of Ω is sufficiently smooth. We may also refer to [AE1] for a variant of the Dirichlet-to-Neumann operator for domains with a rough boundary that has finite $(d - 1)$ -dimensional Hausdorff measure. If, however, Ω has for example a fractal boundary with infinite $(d - 1)$ -dimensional Hausdorff measure, then in [AE1] there is no notion of the Dirichlet-to-Neumann operator at hand simply because there is no appropriate notion of a trace. Using the concepts developed in [PTW2] (with extensions in [PTW1] and [Tro]), we are able to provide a substitute for the space $H^{1/2}(\Gamma)$. We note here that this ‘trace-free’ concept has proven to be useful for dealing with boundary value problems on domains with rough boundary, see [PSTW].

The substitute for the space $H^{1/2}(\Gamma)$ is a variant of 1-harmonic functions in Ω . This removes the need for function evaluation at the boundary. For the definition of this substitute of $H^{1/2}(\Gamma)$, the only concept that we use, if we relate our findings to the Laplacian, is that the matrix $\begin{pmatrix} 0 & \text{div} \\ \text{grad} & 0 \end{pmatrix}$ is skew-symmetric on the space of infinite differentiable functions with compact support, see Example 2.3. Thus, without further effort, our results directly apply to similar problems involving the equations of linearized elasticity or the full 3-dimensional system of static Maxwell’s equations. More generally, our methods apply to the covariant derivative defined on suitable L_2 -tensor fields and a formal skew-adjoint.

As our central object of study, we shall deviate from the classical elliptic partial differential operator $-\Delta$ discussed above and treat abstract divergence form operators of the form

$$-DaG + m, \tag{1}$$

where a and m are bounded coercive operators (called coefficients) and D and G are densely defined, closed, unbounded operators in Hilbert spaces H_1 and H_0 with the property $-D^* \subset G$, like div and grad .

If $\text{dom}(G)$, endowed with the graph norm, embeds compactly into H_0 , we will also address the concept of continuous dependence of the Dirichlet-to-Neumann operator associated with (1) on the bounded coefficients a and m under the weak operator topology. This result has applications in homogenization problems, see [Tar] and [Wau] Section 5.5. Moreover, it complements the study of continuous dependence of the Dirichlet-to-Neumann operator on its coefficients in [AEKS], where the authors focus on possible non-coercive cases and convergence of the principal coefficients in $L_\infty(\Omega)$. In order to prove convergence results, we derive a reformulation of the Dirichlet-to-Neumann operator as a system of two first-order partial differential equations, similar to [AKM].

In the present work we also consider removing the coercivity condition on m . That is to say, we define the abstract analogue of the Dirichlet-to-Neumann *graph* with m being possibly not coercive. We note here that these results are the abstract counterpart of results developed in [BE1] and [AEKS]. In the case that the potentials m are not coercive we consider resolvent convergence for Dirichlet-to-Neumann *operators*.

We mention here that a possible non-linear variant of the Dirichlet-to-Neumann operator, where the coercive operator a is replaced by a (strictly) maximal monotone relation, can be discussed using the results of [TW]. This however is beyond the scope of the present manuscript and will be addressed in future work.

We briefly comment on the organization of the paper. In Section 2, we provide the basic functional analytic setting and recall some notions and results of [PTW2], [PTW1] and [Tro]. We then state the definition of the Dirichlet-to-Neumann operator in the abstract setting discussed above. We also provide an extensive example that justifies this abstraction by relating it to the classical formulation of the Dirichlet-to-Neumann operator. In Section 3 we give a representation formula for the Dirichlet-to-Neumann operator as a first-order system and show that this operator is m -sectorial, provided both m and a are coercive. For this we use a representation result for operators given via forms, see [AE2]. In Section 4 we prove resolvent convergence of the Dirichlet-to-Neumann operators when the coefficients converge in an appropriate weak operator topology. Under some additional hypotheses we also obtain in Theorem 4.2 uniform convergence even though the coefficients converge in the weak operator topology only. In Section 5 we consider the non-coercive case and discuss the domain and multi-valued parts of the Dirichlet-to-Neumann *graph* when m is merely assumed to be a bounded operator, that is not necessarily coercive. Moreover, we also prove a convergence theorem for the non-coercive case in Section 6. We conclude with two more examples in Section 7.

2 The Dirichlet-to-Neumann operator and boundary spaces

We start with a description of boundary data spaces as in [PTW2] Subsection 5.2. Throughout this paper fix Hilbert spaces H_0 and H_1 . Further, let G be an operator in H_0 with values in H_1 and let D be an operator in H_1 with values in H_0 . We assume throughout that both G and D are densely defined and closed, and that $-G^* \subset D$. We define $\mathring{D} = -G^*$ and $\mathring{G} = -D^*$.

Note that

$$(\mathring{G}u, q)_{H_1} = -(u, \mathring{D}q)_{H_0}$$

for all $u \in \text{dom}(\mathring{G})$ and $q \in \text{dom}(\mathring{D})$. Equivalently, the matrix

$$\begin{pmatrix} 0 & \mathring{D} \\ \mathring{G} & 0 \end{pmatrix}$$

with dense domain $\text{dom}(\mathring{G}) \times \text{dom}(\mathring{D})$ is skew-symmetric in $H_0 \times H_1$.

Remark 2.1. Note that $\mathring{G} = -D^* \subset -(-G^*)^* = \overline{G} = G$. So one can simultaneously swap H_0 with H_1 and D with G .

Example 2.2. All examples in this paper are of the following type. Let H_0 and H_1 be Hilbert spaces. Consider dense subspaces $\text{dom}(\widehat{G}) \subset H_0$ and $\text{dom}(\widehat{D}) \subset H_1$. Let $\widehat{G}: \text{dom}(\widehat{G}) \rightarrow H_1$ and $\widehat{D}: \text{dom}(\widehat{D}) \rightarrow H_0$ be two operators such that

$$(\widehat{G}u, q)_{H_1} = -(u, \widehat{D}q)_{H_0} \quad (2)$$

for all $u \in \text{dom}(\widehat{G})$ and $q \in \text{dom}(\widehat{D})$. Equivalently, the matrix

$$\begin{pmatrix} 0 & \widehat{D} \\ \widehat{G} & 0 \end{pmatrix}$$

with dense domain $\text{dom}(\widehat{G}) \times \text{dom}(\widehat{D})$ is skew-symmetric in $H_0 \times H_1$.

Then $\widehat{G} \subset -(\widehat{D})^*$ and $\widehat{D} \subset -(\widehat{G})^*$, so both \widehat{G} and \widehat{D} are closable. Let \mathring{G} and \mathring{D} denote the closures. Define $G = -(\mathring{D})^*$ and $D = -(\mathring{G})^*$. Since \mathring{D} and \mathring{G} are closed, therefore closable, it follows that G and D are densely defined. Obviously both G and D are closed. Next $G^* = -(\mathring{D})^{**} = -\mathring{D}$ since \mathring{D} is closed and similarly $D^* = -\mathring{G}$. Also $\mathring{D} \subset -(\widehat{G})^* = -(\mathring{G})^*$, so $\mathring{G} \subset -(\mathring{D})^* = G$. Similarly $\mathring{D} \subset D$. Then $G^* = -\mathring{D} \subset -D$ as required.

The classical example for this paper is as follows. Note that we do not assume any condition on the boundary of Ω .

Example 2.3. Let $\Omega \subset \mathbb{R}^d$ be open. Define $\widehat{G}: C_c^\infty(\Omega) \rightarrow L_2(\Omega)^d$ and $\widehat{D}: C_c^\infty(\Omega)^d \rightarrow L_2(\Omega)$ by

$$\widehat{G}u = (\partial_1 u, \dots, \partial_d u) \quad \text{and} \quad \widehat{D}q = \sum_{k=1}^d \partial_k q_k.$$

Define $H_0 = L_2(\Omega)$ and $H_1 = L_2(\Omega)^d$. Then (2) in Example 2.2 follows from integration by parts. The associated operators are denoted by $G = \text{grad}$, $\mathring{G} = \overset{\circ}{\text{grad}}$, $D = \text{div}$ and $\mathring{D} = \overset{\circ}{\text{div}}$. It is not hard to show that $\text{dom}(\overset{\circ}{\text{grad}}) = H_0^1(\Omega)$, $\text{dom}(\text{grad}) = H^1(\Omega)$ and $\text{dom}(\text{div}) = H_{\text{div}}(\Omega) = \{q \in L_2(\Omega)^d : \text{div } q \in L_2(\Omega)\}$.

We next define an (abstract) variant of the trace spaces $H^{1/2}(\Gamma)$ and $H^{-1/2}(\Gamma)$. Throughout this paper we provide the domain of an operator with the graph norm. Define

$$\text{BD}(G) = \text{dom}(\mathring{G})^{\perp_{\text{dom}(G)}} \quad \text{and} \quad \text{BD}(D) = \text{dom}(\mathring{D})^{\perp_{\text{dom}(D)}}.$$

We provide $\text{BD}(G)$ and $\text{BD}(D)$ with the induced inner products of $\text{dom}(G)$ and $\text{dom}(D)$. We denote by $\pi_{\text{BD}(G)}$ and $\pi_{\text{BD}(D)}$ the corresponding projections onto $\text{BD}(G)$ and $\text{BD}(D)$, respectively.

Example 2.4. Let Ω , G and D be as in Example 2.3. Then $\text{BD}(G) = \{u \in H^1(\Omega) : \Delta u = u \text{ weakly on } \Omega\}$. Indeed, let $u \in \text{BD}(G)$. Then $u \in H^1(\Omega)$ and $0 = (u, v)_{\text{dom}(G)} = (u, v)_{L_2(\Omega)} + (\text{grad } u, \text{grad } v)_{L_2(\Omega)}$ for all $v \in \text{dom}(\mathring{G}) = H_0^1(\Omega)$. So $\Delta u = u$ weakly on Ω . The converse inclusion is similar.

Lemma 2.5. $\text{BD}(G) = \ker(I - DG)$ and $\text{BD}(D) = \ker(I - GD)$.

Proof. By Remark 2.1 it suffices to prove the first equality. Let $u \in \text{BD}(G)$. Then

$$(u, v)_{H_0} + (Gu, \dot{G}v)_{H_1} = (u, v)_{\text{dom}(G)} = 0$$

for all $v \in \text{dom}(\dot{G})$. So $Gu \in \text{dom}((\dot{G})^*) = \text{dom}(D)$ and $DGu = -(\dot{G})^*Gu = u$. Therefore $u \in \ker(I - DG)$. The converse follows similarly. \square

Corollary 2.6. If $u \in \text{BD}(G)$, then $Gu \in \text{BD}(D)$. If $q \in \text{BD}(D)$, then $Dq \in \text{BD}(G)$.

Proof. Let $u \in \text{BD}(G)$. Then $u \in \text{dom}(DG)$ and $DGu = u \in \text{dom}(DG)$. Therefore $u \in \text{dom}(GDG)$ and $(I - GD)Gu = G(I - DG)u = 0$. So $Gu \in \ker(I - GD) = \text{BD}(D)$ by Lemma 2.5. The other statement follows similarly. \square

Define $\dot{G}: \text{BD}(G) \rightarrow \text{BD}(D)$ and $\dot{D}: \text{BD}(D) \rightarrow \text{BD}(G)$ by

$$\dot{G}u = Gu \quad \text{and} \quad \dot{D}q = Dq.$$

Lemma 2.7. The operators \dot{G} and \dot{D} are unitary. Moreover, $(\dot{G})^* = \dot{D}$.

Proof. See [PTW2, Theorem 5.2]. For the convenience of the reader we include the proof. Clearly $\dot{D}\dot{G} = I_{\text{BD}(G)}$ and $\dot{G}\dot{D} = I_{\text{BD}(D)}$ by Lemma 2.5. Moreover,

$$(\dot{G}u, q)_{\text{BD}(D)} = (\dot{G}u, q)_{H_1} + (\dot{D}\dot{G}u, \dot{D}q)_{H_0} = (\dot{G}u, \dot{G}\dot{D}q)_{H_1} + (u, \dot{D}q)_{H_0} = (u, \dot{D}q)_{\text{BD}(G)}$$

for all $u \in \text{BD}(G)$ and $q \in \text{BD}(D)$, from which the lemma follows. \square

In the situation of Example 2.3 the space $\text{BD}(G)$ models the boundary data of an $H^1(\Omega)$ -function if Ω is a bounded Lipschitz domain, as shown in [Tro, Corollary 4.4]. Indeed, let $\Gamma = \partial\Omega$. Since $\text{Tr}: H^1(\Omega) \rightarrow H^{1/2}(\Gamma)$ is continuous, surjective and $\ker \text{Tr} = H_0^1(\Omega) = \text{dom}(\text{grad})$, it follows that

$$\text{Tr}|_{\text{BD}(G)}: \text{BD}(G) \rightarrow H^{1/2}(\Gamma) \tag{3}$$

is bijective and hence a topological isomorphism.

We next consider the space $\text{BD}(D)$. Denote by $\text{BD}(G)'$ the space of all antilinear continuous maps from $\text{BD}(G)$ into \mathbb{C} . There is a natural unitary map from $\text{BD}(D)$ onto $\text{BD}(G)'$.

Proposition 2.8. Define $\Phi: \text{BD}(D) \rightarrow \text{BD}(G)'$ by

$$\left(\Phi(q)\right)(u) = (Dq, u)_{H_0} + (q, Gu)_{H_1}.$$

Then Φ is unitary.

Proof. Let $q \in \text{BD}(D)$ and $u \in \text{BD}(G)$. Then

$$\begin{aligned} \left(\Phi(q)\right)(u) &= (Dq, u)_{H_0} + (q, Gu)_{H_1} \\ &= (q, Gu)_{H_1} + (Dq, DGu)_{H_0} = (q, Gu)_{\text{dom}(D)} = (q, \dot{G}u)_{\text{BD}(D)}. \end{aligned} \tag{4}$$

Then the proposition follows from Lemma 2.7 and the Riesz representation theorem. \square

For clarity and contrast we include the proof of the next proposition. We provide $\text{Tr } H^1(\Omega)$ with the quotient norm.

Proposition 2.9. *Let $\Omega \subset \mathbb{R}^d$ be open, bounded with Lipschitz boundary. Then one has the following.*

(a) *For all $q \in H_{\text{div}}(\Omega)$ there exists a unique $Q \in (\text{Tr } H^1(\Omega))'$ such that*

$$\langle Q, \text{Tr } u \rangle_{(\text{Tr } H^1(\Omega))' \times \text{Tr } H^1(\Omega)} = \int_{\Omega} (\text{div } q) \bar{u} + \int_{\Omega} q \cdot \overline{\nabla u} \quad (5)$$

for all $u \in H^1(\Omega)$.

(b) *If $q \in \text{dom}(\mathring{\text{div}})$, then $Q = 0$, where Q is as in (5).*

(c) *If $q \in H^1(\Omega)^d$, then $Q = \nu \cdot \text{Tr } q$, where ν is the outward normal vector on the boundary Γ of Ω and Q is as in (5).*

Proof. ‘(a)’. Define $F: H^1(\Omega) \rightarrow \mathbb{C}$ by

$$F(u) = \int_{\Omega} (\text{div } q) \bar{u} + \int_{\Omega} q \cdot \overline{\nabla u}.$$

Then $F \in H^1(\Omega)'$. Moreover, if $u \in H_0^1(\Omega)$, then $F(u) = 0$. Hence there exists a unique continuous antilinear map $\tilde{F}: \text{Tr } H^1(\Omega) \rightarrow \mathbb{C}$ such that $\tilde{F}(\text{Tr } u) = F(u)$ for all $u \in H^1(\Omega)$. Then the first statement follows.

‘(b)’. We use the notation as in Example 2.3. Let $q \in \text{dom}(\mathring{D})$. Since $\mathring{D} = -G^*$ one deduces that $F(u) = \int_{\Omega} (\text{div } q) \bar{u} + \int_{\Omega} q \cdot \overline{\nabla u} = (\mathring{D}q, u)_{H_1} + (q, Gu)_{H_0} = 0$ for all $u \in \text{dom}(G)$. So $Q = 0$, because $\text{dom}(G)$ is dense in $H^1(\Omega)$.

‘(c)’. Suppose that $q \in H^1(\Omega)^d$. Let $u \in H^1(\Omega)$. Then $\bar{u}q \in W^{1,1}(\Omega)^d$ and the divergence theorem gives

$$\int_{\Omega} (\text{div } q) \bar{u} + \int_{\Omega} q \cdot \overline{\nabla u} = \int_{\Omega} \text{div}(\bar{u}q) = \int_{\Gamma} \nu \cdot \text{Tr}(\bar{u}q) = \int_{\Gamma} (\nu \cdot \text{Tr } q) \text{Tr } \bar{u}.$$

So $Q = \nu \cdot \text{Tr } q$. □

If $q \in H_{\text{div}}(\Omega)$ and Q is as in Proposition 2.9, then we define $(\nu q) = Q$. So $(\nu q) = \nu \cdot \text{Tr } q$ if $q \in H^1(\Omega)^d$.

Example 2.10. Let Ω be a bounded Lipschitz domain with boundary Γ . Let G and D be as in Example 2.3. Let Φ be as in Proposition 2.8. Then

$$(\Phi(q))(u) = \langle (\nu q), \text{Tr } u \rangle_{(\text{Tr } H^1(\Omega))' \times \text{Tr } H^1(\Omega)}$$

for all $q \in \text{BD}(D)$ and $u \in \text{BD}(G)$.

It follows from (3) and Proposition 2.8 that the spaces $\text{BD}(D)$ and $H^{-1/2}(\Gamma)$ are isomorphic. Hence \mathring{G} is a variant of the Dirichlet-to-Neumann operator.

Next we introduce the (variable) coefficients for our abstract Dirichlet-to-Neumann operator. Recall that a bounded operator M in a Hilbert space H is called **coercive** if there exists a $\mu > 0$ such that $\operatorname{Re} M \geq \mu I$, where $\operatorname{Re} M = \frac{1}{2}(M + M^*)$. That is M is coercive if and only if there exists a $\mu > 0$ such that $\operatorname{Re}(Mx, x) \geq \mu \|x\|_H^2$ for all $x \in H$.

As for the classical Dirichlet-to-Neumann operator, we first show that the Dirichlet problem has a unique solution.

Proposition 2.11. *Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Let $u_0 \in \operatorname{BD}(G)$. Then there exists a unique $u \in \operatorname{dom}(DaG)$ such that $mu - DaGu = 0$ and $u - u_0 \in \operatorname{dom}(\mathring{G})$.*

For the proof of the proposition we need several auxiliary results.

Lemma 2.12. *Let H be a Hilbert space, $M \in \mathcal{L}(H)$ and A a skew-adjoint operator in H . Let $\lambda > 0$ and assume that $\operatorname{Re}(Mx, x)_H \geq \lambda \|x\|_H^2$ for all $x \in H$. Then the operator $M + A$ is invertible. Moreover, the operator $(M + A)^{-1}$ is bounded from H into $\operatorname{dom}(A)$ and $\|(M + A)^{-1}\|_{H \rightarrow \operatorname{dom}(A)} \leq \frac{1+\lambda+\|M\|}{\lambda}$.*

Proof. If $x \in \operatorname{dom}(A)$, then $\operatorname{Re}((M + A)x, x)_H = \operatorname{Re}(Mx, x)_H \geq \lambda \|x\|_H^2$. Hence $M + A$ is one-to-one, its range is closed and $M + A$ is continuously invertible on its range. Since $\operatorname{Re}(Mx, x)_H = \operatorname{Re}(M^*x, x)_H$ for all $x \in H$, we obtain similarly that $(M + A)^* = M^* - A$ is one-to-one. Therefore $M + A$ is onto. So $M + A$ is invertible and $\|(M + A)^{-1}\|_{H \rightarrow H} \leq \frac{1}{\lambda}$.

Since $A(M + A)^{-1} = I - M(M + A)^{-1}$, the operator $A(M + A)^{-1}$ is bounded from H into H and the estimate follows. \square

Next we consider matrix operators.

Lemma 2.13. *Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive.*

- (a) *The operators $\begin{pmatrix} m & -\mathring{D} \\ -G & a^{-1} \end{pmatrix}$ and $\begin{pmatrix} m & -D \\ -\mathring{G} & a^{-1} \end{pmatrix}$ in $H_0 \times H_1$ are invertible.*
- (b) *The operator $\begin{pmatrix} m & -\mathring{D} \\ -G & a^{-1} \end{pmatrix}^{-1}$ is bounded from $H_0 \times H_1$ into $\operatorname{dom}(G) \times \operatorname{dom}(\mathring{D})$.*
- (c) *The operator $\begin{pmatrix} m & -D \\ -\mathring{G} & a^{-1} \end{pmatrix}^{-1}$ is bounded from $H_0 \times H_1$ into $\operatorname{dom}(\mathring{G}) \times \operatorname{dom}(D)$.*

Proof. Let $H = H_0 \times H_1$, $M = \begin{pmatrix} m & 0 \\ 0 & a^{-1} \end{pmatrix}$ and $A = \begin{pmatrix} 0 & -\mathring{D} \\ -G & 0 \end{pmatrix}$ with $\operatorname{dom}(A) = \operatorname{dom}(G) \times \operatorname{dom}(\mathring{D})$. Since $-\mathring{D}^* = G$ and $-G^* = \mathring{D}$, the operator A is skew-adjoint. Also $\operatorname{Re} a^{-1} \geq \|a\|^{-2} \operatorname{Re} a$, so M is coercive. Therefore $M + A$ is invertible and the operator $(M + A)^{-1}$ is bounded from H into $\operatorname{dom}(A)$ by Lemma 2.12. This proves the first part of Statement (a) and Statement (b)

The remaining parts of the lemma follow similarly. \square

Lemma 2.14. *Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Let $u \in \operatorname{dom}(G)$, $q \in \operatorname{dom}(D)$, $u_0 \in \operatorname{BD}(G)$ and $q_0 \in \operatorname{BD}(D)$.*

(a) *The following conditions are equivalent.*

- (i) $Dq = mu$, $q = aGu$ and $u - u_0 \in \text{dom}(\mathring{G})$.
- (ii) $q = aGu$, $u - u_0 \in \text{dom}(\mathring{G})$ and

$$(aGu, \mathring{G}v)_{H_1} = -(mu, v)_{H_0}$$

for all $v \in \text{dom}(\mathring{G})$.

- (iii)
$$\begin{pmatrix} u - u_0 \\ q \end{pmatrix} = \begin{pmatrix} m & -D \\ -\mathring{G} & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} -mu_0 \\ Gu_0 \end{pmatrix}.$$

(b) *The following conditions are equivalent.*

- (i) $Dq = mu$, $q = aGu$ and $q - q_0 \in \text{dom}(\mathring{D})$.
- (ii)
$$\begin{pmatrix} u \\ q - q_0 \end{pmatrix} = \begin{pmatrix} m & -\mathring{D} \\ -G & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} Dq_0 \\ -a^{-1}q_0 \end{pmatrix}.$$

Proof. ‘(a)’. ‘(i) \Leftrightarrow (ii)’. This follows immediately from the equality $D = -(\mathring{G})^*$.

‘(i) \Leftrightarrow (iii)’. By a simple algebraic manipulation Condition (i) is equivalent to

$$u - u_0 \in \text{dom}(\mathring{G}) \quad \text{and} \quad \begin{pmatrix} m & -D \\ -G & a^{-1} \end{pmatrix} \begin{pmatrix} u - u_0 \\ q \end{pmatrix} = \begin{pmatrix} -mu_0 \\ Gu_0 \end{pmatrix}.$$

By Lemma 2.13(a) this is equivalent to Condition (iii).

‘(b)’. The proof is similar. □

Now we are able to prove Proposition 2.11.

Proof of Proposition 2.11. First we show existence. Let $u \in \text{dom}(G)$ and $q \in \text{dom}(D)$ be such that

$$\begin{pmatrix} u - u_0 \\ q \end{pmatrix} = \begin{pmatrix} m & -D \\ -\mathring{G} & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} -mu_0 \\ Gu_0 \end{pmatrix}.$$

Then u satisfies the desired properties by Lemma 2.14(a) (iii) \Rightarrow (i).

It remains to show uniqueness. Let $\tilde{u} \in \text{dom}(DaG)$ and suppose that $m\tilde{u} - DaG\tilde{u} = 0$ and $\tilde{u} - u_0 \in \text{dom}(\mathring{G})$. Set $\tilde{q} = aG\tilde{u}$. Then it follows from Lemma 2.14(a) (i) \Rightarrow (iii) that

$$\begin{pmatrix} \tilde{u} - u_0 \\ \tilde{q} \end{pmatrix} = \begin{pmatrix} m & -D \\ -\mathring{G} & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} -mu_0 \\ Gu_0 \end{pmatrix},$$

which implies that $u = \tilde{u}$. □

There is a similar version of Proposition 2.11 for the Neumann problem.

Proposition 2.15. *Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Let $q_0 \in \text{BD}(D)$. Then there exists a unique $u \in \text{dom}(DaG)$ such that $mu - DaGu = 0$ and $aGu - q_0 \in \text{dom}(\mathring{D})$.*

Proof. This follows similarly to the proof of Proposition 2.11, but now use Lemma 2.14(b) instead of Lemma 2.14(a). □

At this stage we are able to define the Dirichlet-to-Neumann operator with variable coefficients as an operator acting from $\text{BD}(G)$ (the abstract realization of $H^{1/2}(\Gamma)$) to $\text{BD}(D)$ (the abstract realization of $H^{-1/2}(\Gamma)$).

Definition 2.16. Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Define the operator

$$\Lambda: \text{BD}(G) \rightarrow \text{BD}(D)$$

as follows. Let $u_0 \in \text{BD}(G)$. By Proposition 2.11 there exists a unique $u \in \text{dom}(DaG)$ such that $mu - DaGu = 0$ and $u - u_0 \in \text{dom}(\mathring{G})$. Then we define $\Lambda u_0 = \pi_{\text{BD}(D)} aGu$. We call Λ the **Dirichlet-to-Neumann operator associated with $-DaG + m$** .

So the graph of the operator Λ is equal to

$$\{(\pi_{\text{BD}(G)}u, \pi_{\text{BD}(D)}aGu) : u \in \text{dom}(DaG) \text{ and } mu - DaGu = 0\}.$$

Theorem 2.17. Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Then the operator Λ associated with $-DaG + m$ is bounded and invertible. Moreover,

$$\Lambda u_0 = \begin{pmatrix} 0 & \pi_{\text{BD}(D)} \end{pmatrix} \begin{pmatrix} m & -D \\ -\mathring{G} & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} -m \\ G \end{pmatrix} u_0$$

for all $u_0 \in \text{BD}(G)$ and

$$\Lambda^{-1} q_0 = \begin{pmatrix} \pi_{\text{BD}(G)} & 0 \end{pmatrix} \begin{pmatrix} m & -\mathring{D} \\ -G & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} D \\ -a^{-1} \end{pmatrix} q_0$$

for all $q_0 \in \text{BD}(D)$.

Proof. The expression for Λ follows from Lemma 2.14(a), arguing as in the proof of Proposition 2.11. The boundedness of Λ is then a consequence of Lemma 2.13(c).

The proof for Λ^{-1} is similar, using Lemma 2.14(b), Proposition 2.15 and Lemma 2.13(b). \square

3 An intermediate operator and m-sectoriality

In Proposition 2.8 we showed that the space $\text{BD}(D)$ is naturally isomorphic to $\text{BD}(G)'$. In this section we assume that there is a Hilbert space H such that $\text{BD}(G) \hookrightarrow H \hookrightarrow \text{BD}(G)'$ is a Gelfand triple. Then we study the part of the Dirichlet-to-Neumann operator in H . In the model example, Example 2.3, one can take $H = L_2(\Gamma)$.

Throughout this section, we adopt the notation and assumptions as in the beginning of Section 2. In addition, let H be a Hilbert space and $\kappa \in \mathcal{L}(\text{BD}(G), H)$. We assume that κ is one-to-one and has dense range.

Example 3.1. Let Ω be a bounded Lipschitz domain with boundary Γ . Let G and D be as in Example 2.3. Let $\sigma \in (-\infty, \frac{1}{2}]$ and choose $H = H^\sigma(\Gamma)$. Define $\kappa: \text{BD}(G) \rightarrow H$ by

$\kappa(u) = \text{Tr } u$. Then κ is one-to-one and has dense range. Note that κ is compact if and only if $\sigma < \frac{1}{2}$.

Now suppose that $\sigma = 0$, so $H = L_2(\Gamma)$. Let $\psi \in L_2(\Gamma)$ and set $u = \kappa^* \psi$. Then $u \in \text{BD}(G)$, so $u \in H^1(\Omega)$ and $\Delta u = u$ weakly on Ω by Example 2.4. If $v \in \text{BD}(G)$, then

$$\begin{aligned} \int_{\Gamma} \psi \overline{\text{Tr } v} &= (\psi, \kappa(v))_{L_2(\Gamma)} = (\kappa^* \psi, v)_{\text{BD}(G)} = (u, v)_{\text{BD}(G)} \\ &= \int_{\Omega} u \bar{v} + \int_{\Omega} \nabla u \cdot \overline{\nabla v} = \int_{\Omega} (\Delta u) \bar{v} + \int_{\Omega} \nabla u \cdot \overline{\nabla v}. \end{aligned}$$

Alternatively, if $v \in H_0^1(\Omega) = \text{dom}(\mathring{G})$, then

$$\int_{\Gamma} \psi \overline{\text{Tr } v} = 0 = \int_{\Omega} (\Delta u) \bar{v} + \int_{\Omega} \nabla u \cdot \overline{\nabla v}.$$

So by linearity

$$\int_{\Gamma} \psi \overline{\text{Tr } v} = \int_{\Omega} (\Delta u) \bar{v} + \int_{\Omega} \nabla u \cdot \overline{\nabla v}$$

for all $v \in H^1(\Omega)$. Hence u has a weak normal derivative and $\partial_{\nu} u = \psi$.

We consider the Gelfand triple

$$\text{BD}(G) \xrightarrow{\kappa} H \simeq H' \xrightarrow{\kappa'} \text{BD}(G)'$$

with H as pivot space. Recall that $\text{BD}(G)'$ is naturally isomorphic to $\text{BD}(D)$ by Proposition 2.8. We aim to describe the part of the Dirichlet-to-Neumann operator Λ in H . We describe the image of H in $\text{BD}(D)$ under the above maps $H \simeq H' \xrightarrow{\kappa'} \text{BD}(G)' \simeq \text{BD}(D)$.

Lemma 3.2. *Let $\Phi: \text{BD}(D) \rightarrow \text{BD}(G)'$ be as in Proposition 2.8. Define $F: H \rightarrow H'$ by $(F\varphi)(\psi) = (\varphi, \psi)_H$. Then $\Phi^{-1} \circ \kappa' \circ F = G \circ \kappa^*$.*

Proof. Let $\varphi \in H$ and write $q = (\Phi^{-1} \circ \kappa' \circ F)(\varphi)$. Let $u \in \text{BD}(G)$. Then it follows from Lemma 2.7 and (4) that

$$\begin{aligned} (\mathring{D}q, u)_{\text{BD}(G)} &= (q, \mathring{G}u)_{\text{BD}(D)} \\ &= (\Phi(q))(u) = ((\kappa' \circ F)\varphi)(u) = (\varphi, \kappa(u))_H = (\kappa^* \varphi, u)_{\text{BD}(G)}. \end{aligned}$$

So $\mathring{D}q = \kappa^* \varphi$ and $q = \mathring{G} \mathring{D}q = \mathring{G} \kappa^* \varphi$. □

Now we are able to define the part of the Dirichlet-to-Neumann operator in H .

Definition 3.3. Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Define the operator Λ_H in H as follows. Let $\varphi, \psi \in H$. Then we say that $\varphi \in \text{dom}(\Lambda_H)$ and $\Lambda_H \varphi = \psi$ if there exists a $u_0 \in \text{BD}(G)$ such that $\kappa(u_0) = \varphi$ and $\Lambda u_0 = (G \circ \kappa^*)(\psi)$, where Λ is the Dirichlet-to-Neumann operator associated with $-DaG + m$. We call Λ_H the **Dirichlet-to-Neumann operator in H associated with $-DaG + m$** .

Despite the abundance of choice of the space H , see Example 3.1, the operator $-\Lambda_H$ is always a semigroup generator.

Theorem 3.4. *Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Then the Dirichlet-to-Neumann operator Λ_H associated with $-DaG + m$ is m -sectorial. In particular, if both a and m are symmetric, then Λ_H is self-adjoint.*

The proof of this theorem is based on form methods and the next theorem.

Theorem 3.5. *Let \tilde{H}, V be Hilbert spaces and let $j \in \mathcal{L}(V, \tilde{H})$ with dense range. Let $\mathfrak{b}: V \times V \rightarrow \mathbb{C}$ be a continuous coercive sesquilinear form, that is there exists a $\mu > 0$ such that $\operatorname{Re} \mathfrak{b}(v) \geq \mu \|v\|_V^2$ for all $v \in V$. Define the operator A in \tilde{H} as follows. Let $x, f \in \tilde{H}$. Then $x \in \operatorname{dom}(A)$ and $Ax = f$ if there exists a $u \in V$ such that $j(u) = x$ and $\mathfrak{b}(u, v) = (f, j(v))_{\tilde{H}}$ for all $v \in V$. Then A is well-defined and m -sectorial. If, in addition, \mathfrak{b} is symmetric, then A is self-adjoint.*

Proof. See [AE2, Theorem 2.1]. □

In the situation of Theorem 3.5 we call A the **operator associated with (\mathfrak{b}, j)** .

Theorem 3.4 is an immediate consequence of Theorem 3.5 and the next proposition.

Proposition 3.6. *Let $a \in \mathcal{L}(H_1)$ and $m \in \mathcal{L}(H_0)$ be coercive. Define the sesquilinear form $\mathfrak{b}: \operatorname{dom}(G) \times \operatorname{dom}(G) \rightarrow \mathbb{C}$ by*

$$\mathfrak{b}(u, v) = (aGu, Gv)_{H_1} + (mu, v)_{H_0}.$$

Then \mathfrak{b} is coercive and continuous. Further define $j: \operatorname{dom}(G) \rightarrow H$ by $j = \kappa \circ \pi_{\operatorname{BD}(G)}$. Then the Dirichlet-to-Neumann operator Λ_H associated with $-DaG + m$ is equal to the operator associated with (\mathfrak{b}, j) .

Proof. The form \mathfrak{b} is coercive since both a and m are coercive. Obviously \mathfrak{b} is continuous. Let A be the operator associated with (\mathfrak{b}, j) . It remains to prove that $A = \Lambda_H$.

‘ $\Lambda_H \subset A$ ’. Let $\varphi \in \operatorname{dom}(\Lambda_H)$ and set $\psi = \Lambda_H \varphi$. Then there exists a $u_0 \in \operatorname{BD}(G)$ with $\kappa(u_0) = \varphi$ and $\Lambda u_0 = (G \circ \kappa^*)\psi$. By definition there exists a $u \in \operatorname{dom}(DaG)$ such that $mu - DaGu = 0$, $u - u_0 \in \operatorname{dom}(\overset{\circ}{G})$ and $\Lambda u_0 = \pi_{\operatorname{BD}(D)}(aGu)$. Then $(G \circ \kappa^*)\psi = \pi_{\operatorname{BD}(D)}(aGu)$ and $j(u) = \kappa \pi_{\operatorname{BD}(G)} u = \kappa(u_0) = \varphi$.

Next if $v \in \operatorname{dom}(\overset{\circ}{G})$, then

$$\begin{aligned} \mathfrak{b}(u, v) &= (aGu, \overset{\circ}{G}v)_{H_1} + (mu, v)_{H_0} \\ &= -(DaGu, v)_{H_0} + (DaGu, v)_{H_0} = 0 = (\psi, 0)_H = (\psi, j(v))_H. \end{aligned}$$

If $v \in \operatorname{BD}(G)$, then Lemma 2.7 gives

$$\begin{aligned} (\psi, j(v))_H &= (\kappa^* \psi, v)_{\operatorname{BD}(G)} = (G\kappa^* \psi, Gv)_{\operatorname{BD}(D)} = (\pi_{\operatorname{BD}(D)}(aGu), Gv)_{\operatorname{BD}(D)} \\ &= (aGu, Gv)_{\operatorname{dom}(D)} = (aGu, Gv)_{H_1} + (DaGu, DGv)_{H_0} \\ &= (aGu, Gv)_{H_1} + (mu, v)_{H_0} = \mathfrak{b}(u, v). \end{aligned}$$

Since $\operatorname{dom}(G) = \operatorname{BD}(G) \oplus \operatorname{dom}(\overset{\circ}{G})$ it follows that $\mathfrak{b}(u, v) = (\psi, j(v))_H$ for all $v \in \operatorname{dom}(G)$. So $\varphi \in \operatorname{dom}(A)$ and $A\varphi = \psi$.

‘ $A \subset \Lambda_H$ ’. Let $\varphi \in \text{dom}(A)$ and write $\psi = A\varphi$. Then there exists a $u \in \text{dom}(G)$ such that $j(u) = \varphi$ and

$$(aGu, Gv)_{H_1} + (mu, v)_{H_0} = \mathfrak{b}(u, v) = (\psi, j(v))_H \quad (6)$$

for all $v \in \text{dom}(G)$. If $v \in \text{dom}(\mathring{G})$, then

$$(aGu, \mathring{G}v)_{H_1} + (mu, v)_{H_0} = (\psi, j(v))_H = 0.$$

So $aGu \in \text{dom}((\mathring{G})^*) = \text{dom}(D)$ and $DaGu = -(\mathring{G})^*aGu = mu$. Moreover,

$$\Lambda\pi_{\text{BD}(G)}u = \pi_{\text{BD}(D)}(aGu) \quad (7)$$

by the definition of Λ . Note that $\kappa(\pi_{\text{BD}(G)}u) = j(u) = \varphi$.

Now let $v \in \text{BD}(G)$. Then (6) gives

$$\begin{aligned} (\kappa^*\psi, v)_{\text{BD}(G)} &= (\psi, \kappa(v))_H \\ &= (aGu, Gv)_{H_1} + (mu, v)_{H_0} \\ &= (aGu, Gv)_{H_1} + (DaGu, DGv)_{H_0} \\ &= (aGu, Gv)_{\text{dom}(D)} \\ &= (\pi_{\text{BD}(D)}(aGu), Gv)_{\text{BD}(D)} \\ &= (D\pi_{\text{BD}(G)}(aGu), v)_{\text{BD}(G)}, \end{aligned}$$

where we used Lemma 2.7 in the last step. So, $\kappa^*\psi = D\pi_{\text{BD}(D)}(aGu)$. Hence

$$(G \circ \kappa^*)(\psi) = \pi_{\text{BD}(D)}(aGu) = \Lambda\pi_{\text{BD}(G)}u$$

by Lemma 2.7 and (7). Therefore $\varphi \in \text{dom}(\Lambda_H)$ and $\Lambda_H\varphi = \psi$. \square

We next show that the operator Λ_H is invertible and determine its inverse.

Proposition 3.7. *The operator Λ_H is invertible and*

$$\Lambda_H^{-1}\psi = \kappa \left(\begin{array}{cc} \pi_{\text{BD}(G)} & 0 \end{array} \right) \begin{pmatrix} m & -\mathring{D} \\ -G & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ -a^{-1}G \end{pmatrix} \kappa^*\psi$$

for all $\psi \in H$.

Proof. Since the form \mathfrak{b} in Proposition 3.6 is coercive, it follows that the operator Λ_H is invertible. Let $\varphi \in \text{dom}(\Lambda_H)$ and write $\psi = \Lambda_H\varphi$. Then there exists a $u_0 \in \text{BD}(G)$ such that $\kappa(u_0) = \varphi$ and $\Lambda u_0 = G\kappa^*\psi$. By Theorem 2.17 we obtain that

$$\begin{aligned} u_0 = \Lambda^{-1}G\kappa^*\psi &= \left(\begin{array}{cc} \pi_{\text{BD}(G)} & 0 \end{array} \right) \begin{pmatrix} m & -\mathring{D} \\ -G & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} D \\ -a^{-1} \end{pmatrix} G\kappa^*\psi \\ &= \left(\begin{array}{cc} \pi_{\text{BD}(G)} & 0 \end{array} \right) \begin{pmatrix} m & -\mathring{D} \\ -G & a^{-1} \end{pmatrix}^{-1} \begin{pmatrix} I \\ -a^{-1}G \end{pmatrix} \kappa^*\psi, \end{aligned}$$

where we used Lemma 2.7 in the last step. Next apply κ to both sides. Since the inverse matrix maps $H_0 \times H_1$ into $\text{dom}(G) \times \text{dom}(D)$ by Lemma 2.13(b), the proposition follows. \square

4 Resolvent convergence

In this section we consider a sequence of Dirichlet-to-Neumann operators and show resolvent convergence.

Throughout this section we adopt the notation and assumptions as in the beginning of Section 2. Let H be a Hilbert space and $\kappa \in \mathcal{L}(\text{BD}(G), H)$ injective with dense range. Further, we let $m_n, m \in \mathcal{L}(H_0)$ and $a_n, a \in \mathcal{L}(H_1)$ for all $n \in \mathbb{N}$. Let $\mu > 0$ and assume that $\text{Re } m_n, \text{Re } m \geq \mu I_{H_0}$ and $\text{Re } a_n, \text{Re } a \geq \mu I_{H_1}$ for all $n \in \mathbb{N}$. Moreover, assume that $\sup_n \|a_n\|_{\mathcal{L}(H_1)} < \infty$. Let $\Lambda, \Lambda_1, \Lambda_2, \dots$ be the Dirichlet-to-Neumann operators from $\text{BD}(G)$ into $\text{BD}(D)$ associated with $-DaG + m, -Da_1G + m_1, -Da_2G + m_2, \dots$ as in Definition 2.16. Similarly, let $\Lambda_H, \Lambda_H^{(1)}, \Lambda_H^{(2)}, \dots$ be the Dirichlet-to-Neumann operators in H as in Definition 3.3.

Throughout this section we suppose in addition that the inclusion $\text{dom}(G) \hookrightarrow H_0$ is compact.

The compactness assumption is valid in our model case, Example 2.3, if Ω has a continuous boundary or, equivalently, if Ω has the segment property.

We state two well-known consequences of the compactness assumption.

Lemma 4.1.

- (a) *There exists a $c > 0$ such that $\|u\|_{H_0} \leq c\|Gu\|_{H_1}$ for all $u \in \text{dom}(G) \cap \ker(G)^{\perp H_0}$.*
- (b) *The space $\text{ran}(G)$ is closed in H_1 .*

Proof. ‘(a)’. Suppose not. Then there exists a sequence $(u_n)_{n \in \mathbb{N}}$ in $\text{dom}(G) \cap \ker(G)^{\perp H_0}$ such that $\|u_n\|_{H_0} = 1$ and

$$\|u_n\|_{H_0} \geq n\|Gu_n\|_{H_1} \tag{8}$$

for all $n \in \mathbb{N}$. Then $(u_n)_{n \in \mathbb{N}}$ is bounded in $\text{dom}(G)$. We may assume without loss of generality that there exists a $u \in \text{dom}(G)$ such that $\lim u_n = u$ weakly in $\text{dom}(G)$. Since the inclusion $\text{dom}(G) \subset H_0$ is compact we obtain that $\lim u_n = u$ in H_0 . Then $u \in \ker(G)^{\perp H_0}$ since $\ker(G)^{\perp H_0}$ is closed in H_0 . Moreover, $\|u\|_{H_0} = 1$ and in particular $u \neq 0$. Alternatively, (8) implies that $\|Gu\|_{H_1} \leq \liminf_{n \rightarrow \infty} \|Gu_n\|_{H_1} = 0$. So $u \in \ker(G)$. Hence $u \in \ker(G) \cap \ker(G)^{\perp H_0} = \{0\}$ and $u = 0$. This is a contradiction.

‘(b)’. This is a consequence of Statement (a) and the closedness of G . □

We provide $\text{ran}(G)$ with the induced norm of H_1 . Throughout the remainder of this section we denote by $\iota: \text{ran}(G) \hookrightarrow H_1$ the embedding map. Note that ι^* is the orthogonal projection from H_1 onto $\text{ran}(G)$. The main result of this section is the following theorem.

Theorem 4.2. *Suppose that $\lim m_n = m$ in the weak operator topology on $\mathcal{L}(H_0)$ and $\lim_{n \rightarrow \infty} (\iota^* a_n \iota)^{-1} = (\iota^* a \iota)^{-1}$ in the weak operator topology on $\mathcal{L}(\text{ran}(G))$. Then*

$$\lim(\Lambda_H^{(n)})^{-1} = \Lambda_H^{-1}$$

in the weak operator topology on $\mathcal{L}(H)$. Moreover, if in addition the map κ is compact, then the convergence is uniform in $\mathcal{L}(H)$.

For the proof of Theorem 4.2 we need some preliminary results. The first one contains an identity for Λ involving $\text{ran}(G)$.

Lemma 4.3.

- (a) *Let $q \in H_1$. Then $q \in \text{dom}(\mathring{D})$ if and only if $\iota^*q \in \text{dom}(\mathring{D})$. In that case $\mathring{D}q = \mathring{D}\iota^*q$.*
- (b) *The operator $\mathring{D}\iota: \text{ran}(G) \cap \text{dom}(\mathring{D}) \rightarrow H_0$ is a closed and densely defined operator in $\text{ran}(G)$. Moreover, $(\mathring{D}\iota)^* = -\iota^*G$.*
- (c) *The operator $\mathring{D}\iota$ is injective.*
- (d) *The inclusion $\text{dom}(\mathring{D}\iota) \subset H_1$ is compact.*
- (e) *The operator $\begin{pmatrix} m & -\mathring{D}\iota \\ -\iota^*G & (\iota^*a\iota)^{-1} \end{pmatrix}: \text{dom}(G) \times (\text{ran}(G) \cap \text{dom}(\mathring{D})) \rightarrow H_0 \times \text{ran}(G)$ is invertible.*
- (f) *The operator $\begin{pmatrix} m & -\mathring{D}\iota \\ -\iota^*G & (\iota^*a\iota)^{-1} \end{pmatrix}^{-1}$ is bounded from $H_0 \times \text{ran}(G)$ into $\text{dom}(G) \times \text{dom}(\mathring{D})$.*
- (g) *If $q_0 \in \text{BD}(D)$, then*

$$\Lambda^{-1}q_0 = \begin{pmatrix} \pi_{\text{BD}(G)} & 0 \end{pmatrix} \begin{pmatrix} m & -\mathring{D}\iota \\ -\iota^*G & (\iota^*a\iota)^{-1} \end{pmatrix}^{-1} \begin{pmatrix} D \\ -(\iota^*a\iota)^{-1}\iota^* \end{pmatrix} q_0.$$

Proof. ‘(a)’. First $q - \iota^*q \in (\text{ran}(G))^\perp_{H_1} = \ker(G^*) = \ker(\mathring{D}) \subset \text{dom}(\mathring{D})$. This shows the equivalence. Since $\mathring{D}(q - \iota^*q) = 0$, the last statement follows.

‘(b)’. Let $q \in \text{ran}(G)$. Since $\text{dom}(\mathring{D})$ is dense in H_1 there exists a sequence $(q_n)_{n \in \mathbb{N}}$ in $\text{dom}(\mathring{D})$ such that $\lim q_n = q$ in H_1 . Then $\iota^*q_n \in \text{ran}(G) \cap \text{dom}(\mathring{D})$ for all $n \in \mathbb{N}$ by Statement (a) and $\lim \iota^*q_n = \iota^*q = q$ in H_1 . So $\text{ran}(G) \cap \text{dom}(\mathring{D})$ is dense in $\text{ran}(G)$.

Because $\text{ran}(G)$ is closed in H_1 and \mathring{D} is a closed operator one deduces easily that the operator $\mathring{D}\iota$ is closed. It remains to show that $(\mathring{D}\iota)^* = -\iota^*G$.

Let $u \in \text{dom}((\mathring{D}\iota)^*)$. Write $q = (\mathring{D}\iota)^*u$. Note that $q \in \text{ran}(G)$. Let $q' \in \text{dom}(\mathring{D})$. Then Statement (a) implies that

$$(u, \mathring{D}q')_{H_0} = (u, \mathring{D}\iota^*q')_{H_0} = (u, (\mathring{D}\iota)\iota^*q')_{H_0} = ((\mathring{D}\iota)^*u, \iota^*q')_{\text{ran}(G)} = (q, \iota^*q')_{\text{ran}(G)} = (q, q')_{H_1}.$$

So $u \in \text{dom}((\mathring{D}\iota)^*) = \text{dom}(G)$ and $Gu = -(\mathring{D}\iota)^*u = -q$. Therefore, $-\iota^*Gu = q = (\mathring{D}\iota)^*u$. This implies that $(\mathring{D}\iota)^* \subset -\iota^*G$. The converse inclusion is easier and is left to the reader.

‘(c)’. Let $q \in \text{ran}(G) \cap \text{dom}(\mathring{D})$ and suppose that $\mathring{D}\iota q = 0$. There exists a $u \in \text{dom}(G) \cap (\ker G)^\perp_{H_0}$ such that $q = Gu$. Then $\|\iota^*Gu\|_{H_1}^2 = -(q, (\mathring{D}\iota)^*u)_{H_1} = -(\mathring{D}\iota q, u)_{H_0} = 0$. So $u \in \ker G$ and $u = 0$.

‘(d)’. Let $q, q_1, q_2, \dots \in \text{dom}(\mathring{D}\iota)$ and suppose that $\lim q_n = q$ weakly in $\text{dom}(\mathring{D}\iota)$. For all $n \in \mathbb{N}$ there exists a unique $u_n \in \text{dom}(G) \cap (\ker G)^\perp_{H_0}$ such that $q_n = Gu_n$. Since $\lim q_n = q$ weakly in H_1 , the sequence $(q_n)_{n \in \mathbb{N}}$ is bounded in H_1 . Hence the sequence $(u_n)_{n \in \mathbb{N}}$ is bounded in H_0 by Lemma 4.1(a). Passing to a subsequence if necessary, there exists a $u \in H_0$ such that $\lim u_n = u$ weakly in H_0 . Since G is a weakly closed operator, one

deduces that $u \in \text{dom}(G)$ and $Gu = q$. Then $\lim u_n = u$ weakly in $\text{dom}(G)$, so $\lim u_n = u$ strongly in H_0 by the compactness assumption. Note that $G^* = -\mathring{D}$. So

$$\lim_{n \rightarrow \infty} \|q_n\|_{H_1}^2 = \lim_{n \rightarrow \infty} (q_n, Gu_n)_{H_1} = \lim_{n \rightarrow \infty} (-\mathring{D}q_n, u_n)_{H_0} = (-\mathring{D}q, u)_{H_0} = (q, Gu)_{H_0} = \|q\|_{H_1}^2.$$

Hence $\lim q_n = q$ in H_1 .

‘(e)’ and ‘(f)’. This is as in the proof of Lemma 2.13(a) and (b).

‘(g)’. Let $q_0 \in \text{BD}(D)$. By Proposition 2.15 there exists a unique $u \in \text{dom}(DaG)$ such that $mu - DaGu = 0$ and $aGu - q_0 \in \text{dom}(\mathring{D})$. Then $\Lambda^{-1}q_0 = \pi_{\text{BD}(G)}u$. Write $q = aGu$. Then $q - q_0 \in \text{dom}(\mathring{D})$, so $\mathring{D}(q - q_0) = \mathring{D}\iota^*(q - q_0) = (\mathring{D}\iota)\iota^*(q - q_0)$ by Statement (a). Therefore

$$Dq_0 = mu - \mathring{D}(q - q_0) = mu - (\mathring{D}\iota)\iota^*(q - q_0). \quad (9)$$

Also $\iota^*q = \iota^*aGu = (\iota^*a\iota)\iota^*Gu$. Hence $(\iota^*a\iota)^{-1}\iota^*q = \iota^*Gu$ and $-\iota^*Gu + (\iota^*a\iota)^{-1}\iota^*(q - q_0) = -(\iota^*a\iota)^{-1}\iota^*q_0$. Together with (9) this gives

$$\begin{pmatrix} m & -\mathring{D}\iota \\ -\iota^*G & (\iota^*a\iota)^{-1} \end{pmatrix} \begin{pmatrix} u \\ \iota^*(q - q_0) \end{pmatrix} = \begin{pmatrix} D \\ -(\iota^*a\iota)^{-1}\iota^* \end{pmatrix} q_0.$$

Finally use Statement (e). □

Next we need a sequential version of Lemma 2.12.

Lemma 4.4. *Let \tilde{H} be a Hilbert space, $M \in \mathcal{L}(\tilde{H})$ and A a skew-adjoint operator in \tilde{H} . Further let $(M_n)_{n \in \mathbb{N}}$ be a sequence in $\mathcal{L}(\tilde{H})$ and suppose that $\lim M_n = M$ in the weak operator topology on $\mathcal{L}(\tilde{H})$. Assume that the inclusion $\text{dom}(A) \subset \tilde{H}$ is compact and that there exists a $\lambda > 0$ such that $\text{Re } M_n \geq \lambda I_{\tilde{H}}$ for all $n \in \mathbb{N}$. Let $(x_n)_{n \in \mathbb{N}}$ be a sequence in \tilde{H} which converges weakly to $x \in \tilde{H}$. Then $M + A$ is invertible and $\lim_{n \rightarrow \infty} (M_n + A)^{-1}x_n = (M + A)^{-1}x$ weakly in $\text{dom}(A)$.*

Proof. Obviously $\text{Re } M \geq \lambda I_{\tilde{H}}$, so $M + A$ is invertible by Lemma 2.12. Consider $z_n = (M_n + A)^{-1}x_n$ for all $n \in \mathbb{N}$. Then $\|z_n\|_{\text{dom}(A)} \leq \frac{1 + \lambda + \|M_n\|}{\lambda} \|x_n\|_{\tilde{H}}$ for all $n \in \mathbb{N}$ by Lemma 2.12. So the sequence $(z_n)_{n \in \mathbb{N}}$ is bounded in $\text{dom}(A)$. Passing to a subsequence, we may assume without loss of generality that there exists a $z \in \text{dom}(A)$ such that $\lim z_n = z$ weakly in $\text{dom}(A)$. Then $\lim z_n = z$ in \tilde{H} by the compactness assumption. Consequently, $\lim M_n z_n = Mz$ weakly in \tilde{H} . Now $M_n z_n + Az_n = x_n$ for all $n \in \mathbb{N}$. Take the limit $n \rightarrow \infty$ and notice that both sides converge weakly in \tilde{H} . It follows that $Mz + Az = x$, so $z = (M + A)^{-1}x$. Now the lemma follows by a standard subsequence argument. □

We need one more convergence result for the proof of Theorem 4.2. This result is also of independent interest.

Proposition 4.5. *Suppose that $\lim m_n = m$ in the weak operator topology on $\mathcal{L}(H_0)$ and $\lim (\iota^*a_n\iota)^{-1} = (\iota^*a\iota)^{-1}$ in the weak operator topology on $\mathcal{L}(\text{ran}(G))$. Let $q, q_1, q_2, \dots \in \text{BD}(D)$ and assume that $\lim q_n = q$ in $\text{BD}(D)$. Then*

$$\lim_{n \rightarrow \infty} \Lambda_n^{-1}q_n = \Lambda^{-1}q$$

weakly in $\text{BD}(G)$.

Proof. Choose $\tilde{H} = H_0 \times \text{ran}(G)$ and let $A = \begin{pmatrix} 0 & -\mathring{D}\iota \\ -\iota^*G & 0 \end{pmatrix}$ with $\text{dom}(A) = \text{dom}(G) \times (\text{ran}(G) \cap \text{dom}(\mathring{D}))$. Then A is skew-adjoint in \tilde{H} by Lemma 4.3(b). Moreover, the inclusion $\text{dom}(A) \subset \tilde{H}$ is compact by Lemma 4.3(d) and the compactness assumption. Further let

$$M = \begin{pmatrix} m & 0 \\ 0 & (\iota^*a\iota)^{-1} \end{pmatrix} \quad \text{and} \quad M_n = \begin{pmatrix} m_n & 0 \\ 0 & (\iota^*a_n\iota)^{-1} \end{pmatrix}$$

for all $n \in \mathbb{N}$. Then $\lim M_n = M$ in the weak operator topology on $\mathcal{L}(\tilde{H})$. Since

$$\text{Re}(\iota^*a_n\iota)^{-1} \geq \|\iota^*a_n\iota\|_{\mathcal{L}(\text{ran}(G))}^{-2} \text{Re}(\iota^*a_n\iota) \geq \|a_n\|_{\mathcal{L}(H_1)}^{-2} \text{Re}(\iota^*a_n\iota)$$

for all $n \in \mathbb{N}$ and $\sup_n \|a_n\|_{\mathcal{L}(H_1)} < \infty$, it follows that there exists a $\lambda > 0$ such that $\text{Re} M_n \geq \lambda I$ for all $n \in \mathbb{N}$. We use Lemma 4.3(g) for Λ^{-1} and Λ_n^{-1} . Obviously

$$\lim(Dq_n, -(\iota^*a_n\iota)^{-1}\iota^*q_n) = (Dq, -(\iota^*a\iota)^{-1}\iota^*q)$$

weakly in \tilde{H} . Hence

$$\lim_{n \rightarrow \infty} \begin{pmatrix} m_n & -\mathring{D}\iota \\ -\iota^*G & (\iota^*a_n\iota)^{-1} \end{pmatrix}^{-1} \begin{pmatrix} D \\ -(\iota^*a_n\iota)^{-1}\iota^* \end{pmatrix} q_n = \begin{pmatrix} m & -\mathring{D}\iota \\ -\iota^*G & (\iota^*a\iota)^{-1} \end{pmatrix}^{-1} \begin{pmatrix} D \\ -(\iota^*a\iota)^{-1}\iota^* \end{pmatrix} q$$

weakly in $\text{dom}(A)$ by Lemma 4.4. Consequently $\lim \Lambda_n^{-1}q_n = \Lambda^{-1}q$ weakly in $\text{BD}(G)$ by Lemma 4.3(g). \square

Now we are able to prove the main theorem of this section.

Proof of Theorem 4.2. Let $\psi \in H$. Then $\lim \Lambda_n^{-1}G\kappa^*\psi = \Lambda^{-1}G\kappa^*\psi$ weakly in $\text{BD}(G)$ by Proposition 4.5. Hence

$$\lim_{n \rightarrow \infty} (\Lambda_H^{(n)})^{-1}\psi = \lim_{n \rightarrow \infty} \kappa\Lambda_n^{-1}G\kappa^*\psi = \kappa\Lambda^{-1}G\kappa^*\psi = \Lambda_H^{-1}\psi$$

weakly in H . This proves the first statement in Theorem 4.2.

Now suppose that κ is compact. Suppose $\lim(\Lambda_H^{(n)})^{-1} = \Lambda_H^{-1}$ in $\mathcal{L}(H)$ is false. Passing to a subsequence if necessary, there exist $\delta > 0$ and $\psi_1, \psi_2, \dots \in H$ such that

$$\|(\Lambda_H^{(n)})^{-1}\psi_n - \Lambda_H^{-1}\psi_n\|_H > \delta\|\psi_n\|_H \quad (10)$$

for all $n \in \mathbb{N}$. Without loss of generality we may assume that $\|\psi_n\|_H = 1$ for all $n \in \mathbb{N}$. Passing again to a subsequence if necessary, there exists a $\psi \in H$ such that $\lim \psi_n = \psi$ weakly in H . Then $\lim \kappa^*\psi_n = \kappa^*\psi$ in $\text{BD}(G)$ since κ is compact. Therefore $\lim G\kappa^*\psi_n = G\kappa^*\psi$ in $\text{BD}(D)$. Hence $\lim \Lambda_n^{-1}G\kappa^*\psi_n = \Lambda^{-1}G\kappa^*\psi$ weakly in $\text{BD}(G)$ by Proposition 4.5. Using again that κ is compact it follows that $\lim(\Lambda_H^{(n)})^{-1}\psi_n = (\Lambda_H)^{-1}\psi$ in H . Similarly $\lim \Lambda_H^{-1}\psi_n = (\Lambda_H)^{-1}\psi$ in H . So $\lim \|(\Lambda_H^{(n)})^{-1}\psi_n - \Lambda_H^{-1}\psi_n\|_H = 0$. This contradicts (10) for large n . \square

5 The non-coercive case

In this section, we drop the coerciveness condition on m . As a result the Dirichlet-to-Neumann operator can become multi-valued, that is, it is a graph and no longer an operator. The Dirichlet-to-Neumann graph associated with the Schrödinger operator $-\Delta + m$ has been studied in [AEKS] and [BE1].

Throughout this section we adopt the notation and assumptions as in the beginning of Section 2. Further we fix an element $m \in \mathcal{L}(H_0)$ and a coercive $a \in \mathcal{L}(H_1)$. We emphasise that we do not require that m is coercive. The definition of the Dirichlet-to-Neumann *graph*, however, remains the same as in the single-valued case in Definition 2.16.

Definition 5.1. Set

$$\Lambda = \{(\pi_{\text{BD}(G)}u, \pi_{\text{BD}(D)}aGu) \in \text{BD}(G) \times \text{BD}(D) : u \in \text{dom}(DaG) \text{ and } mu - DaGu = 0\}.$$

We call Λ the **Dirichlet-to-Neumann graph associated with $-DaG + m$** .

We briefly recall some definitions in the area of (linear) graphs. Let H, K be Hilbert spaces. Then a **graph** A is a vector subspace of $H \times K$. The **domain**, **multi-valued part** and **inverse** of A are defined by

$$\text{dom}(A) = \{h \in H : \text{there exists a } k \in K \text{ such that } (h, k) \in A\},$$

$$\text{mul}(A) = \{k \in K : (0, k) \in A\} \text{ and}$$

$$A^{-1} = \{(k, h) \in K \times H : (h, k) \in A\}.$$

We say that A is **single-valued** or an **operator** if $\text{mul}(A) = \{0\}$. The next lemma is trivial.

Lemma 5.2.

- (a) $\text{mul}(\Lambda) = \{\pi_{\text{BD}(D)}aGu : u \in \ker(m - Da\mathring{G})\}.$
- (b) *If $\ker(m - Da\mathring{G}) = \{0\}$, then Λ is single-valued.*

As in Proposition 3.6 define the sesquilinear form $\mathfrak{b} : \text{dom}(G) \times \text{dom}(G) \rightarrow \mathbb{C}$ by

$$\mathfrak{b}(u, v) = (aGu, Gv)_{H_1} + (mu, v)_{H_0}.$$

We also need the Dirichlet-version of \mathfrak{b} defined by $\mathring{\mathfrak{b}} = \mathfrak{b}|_{\text{dom}(\mathring{G}) \times \text{dom}(\mathring{G})}$. Then \mathfrak{b} and $\mathring{\mathfrak{b}}$ are continuous. Hence there exist $T \in \mathcal{L}(\text{dom}(G))$ and $\mathring{T} \in \mathcal{L}(\text{dom}(\mathring{G}))$ such that $\mathfrak{b}(u, v) = (Tu, v)_{\text{dom}(G)}$ for all $u, v \in \text{dom}(G)$ and $\mathring{\mathfrak{b}}(u, v) = (\mathring{T}u, v)_{\text{dom}(\mathring{G})}$ for all $u, v \in \text{dom}(\mathring{G})$. Note that $\ker(\mathring{T}) = \ker(m - Da\mathring{G})$, since $(\mathring{G})^* = -D$.

With a condition on $\text{ran}(\mathring{T})$ we can characterise the domain of the Dirichlet-to-Neumann graph Λ .

Proposition 5.3. *Suppose that $\text{ran}(\mathring{T})$ is closed in $\text{dom}(\mathring{G})$. Then*

$$\text{dom}(\Lambda) = \{u_0 \in \text{BD}(G) : (Gu_0, \pi_{\text{BD}(D)}a^*Gv)_{\text{BD}(D)} = 0 \text{ for all } v \in \ker(m^* - Da^*\mathring{G})\}.$$

Proof. ‘ \subset ’. Let $u_0 \in \text{dom}(\Lambda)$. Then there exists a $u \in \text{dom}(G)$ such that $mu - DaGu = 0$ and $u_0 = \pi_{BD(G)}u$. Let $v \in \text{dom}(\mathring{G})$. Then $(mu, v)_{H_0} = (DaGu, v)_{H_0} = -(aGu, \mathring{G}v)_{H_1}$ and

$$\begin{aligned} (\mathring{T}(u - u_0), v)_{\text{dom}(\mathring{G})} &= \mathring{\mathfrak{b}}(u - u_0, v) = (aG(u - u_0), \mathring{G}v)_{H_1} + (m(u - u_0), v)_{H_0} \\ &= -(aGu_0, \mathring{G}v)_{H_1} - (mu_0, v)_{H_0}. \end{aligned}$$

Note that $\mathring{T}(u - u_0) \in \text{ran}(\mathring{T}) = (\ker((\mathring{T})^*))^{\perp_{\text{dom}(\mathring{G})}}$ since $\text{ran}(\mathring{T})$ is closed.

Now let $v \in \ker(m^* - Da^*\mathring{G}) = \ker((\mathring{T})^*)$. Then

$$\begin{aligned} 0 &= -(\mathring{T}(u - u_0), v)_{\text{dom}(\mathring{G})} = (aGu_0, \mathring{G}v)_{H_1} + (mu_0, v)_{H_0} \\ &= (Gu_0, a^*\mathring{G}v)_{H_1} + (u_0, m^*v)_{H_0} \\ &= (Gu_0, a^*\mathring{G}v)_{H_1} + (DGu_0, Da^*\mathring{G}v)_{H_0} \\ &= (Gu_0, a^*\mathring{G}v)_{\text{dom}(D)} = (Gu_0, \pi_{BD(D)}a^*\mathring{G}v)_{BD(D)} \end{aligned}$$

as required.

‘ \supset ’. The proof is similar and for this inclusion it is essential that $\text{ran}(\mathring{T})$ is closed. \square

Corollary 5.4. *Suppose that $\text{ran}(\mathring{T})$ is closed in $\text{dom}(\mathring{G})$. Then*

$$\text{dom}(\Lambda) = \{u_0 \in \text{BD}(G) : (\Phi(\pi_{BD(D)}a^*Gv))(u_0) = 0 \text{ for all } v \in \ker(m^* - Da^*\mathring{G})\},$$

where $\Phi: \text{BD}(D) \rightarrow \text{BD}(G)'$ is the natural unitary map as in Proposition 2.8.

We emphasise that boundary regularity is not needed in Corollary 5.4.

The next lemma gives an easy to verify condition which implies that \mathring{T} has closed range.

Lemma 5.5. *If the inclusion $\tau: \text{dom}(\mathring{G}) \rightarrow H_0$ is compact, then \mathring{T} has closed range.*

Proof. There exist $\mu, \omega > 0$ such that $\mu\|u\|_{\text{dom}(\mathring{G})}^2 \leq \text{Re } \mathring{\mathfrak{b}}(u) + \omega\|\tau u\|_{H_0}^2$ for all $u \in \text{dom}(\mathring{G})$. Then $\mu\|u\|_{\text{dom}(\mathring{G})}^2 \leq \text{Re}(\mathring{T}u, u)_{\text{dom}(\mathring{G})} + \omega(\tau^*\tau u, u)_{\text{dom}(\mathring{G})} = \text{Re}((\mathring{T} + \omega\tau^*\tau)u, u)_{\text{dom}(\mathring{G})}$ for all $u \in \text{dom}(\mathring{G})$. So $\mathring{T} + \omega\tau^*\tau$ is injective and has closed range. Similarly $(\mathring{T})^* + \omega\tau^*\tau$ is injective. So $\mathring{T} + \omega\tau^*\tau$ is invertible. Since $\omega\tau^*\tau$ is compact, the operator \mathring{T} is Fredholm. In particular, the range of \mathring{T} is closed. \square

Note that the operator τ is compact in the situation of Example 2.3.

Example 5.6. Let $\Omega \subset \mathbb{R}^d$ be a bounded Lipschitz domain with boundary Γ . Let G and D be as in Example 2.3. If $u_0 \in \text{BD}(G)$, $v \in H_0$ and $\Phi: \text{BD}(D) \rightarrow \text{BD}(G)'$ is the natural unitary map as in Proposition 2.8, then it follows from Example 2.10 and Proposition 2.9(b) that

$$\begin{aligned} (\Phi(\pi_{BD(D)}a^*Gv))(u_0) &= \langle (\nu\pi_{BD(D)}a^*Gv), \text{Tr } u_0 \rangle_{(\text{Tr } H^1(\Omega))' \times \text{Tr } H^1(\Omega)} \\ &= \langle (\nu a^*Gv), \text{Tr } u_0 \rangle_{(\text{Tr } H^1(\Omega))' \times \text{Tr } H^1(\Omega)} \\ &= \langle (\partial_\nu^{a^*} v), \text{Tr } u_0 \rangle_{H^{-1/2}(\partial\Omega), H^{1/2}(\partial\Omega)}, \end{aligned}$$

where $\partial_\nu^{a^*}$ is the co-normal derivative. So Corollary 5.4 gives

$$\text{dom}(\Lambda) = \{u_0 \in \text{BD}(G) : \langle (\partial_\nu^{a^*} v), \text{Tr } u_0 \rangle_{H^{-1/2}(\partial\Omega), H^{1/2}(\partial\Omega)} = 0 \text{ for all } v \in \ker(m^* - Da^*\mathring{G})\},$$

in agreement with [McL] Proposition 4.10.

Next we turn to the Neumann-to-Dirichlet graph.

Proposition 5.7. *Assume that $\text{ran}(T)$ is closed in $\text{dom}(G)$. Then*

$$\text{dom}(\Lambda^{-1}) = \{q_0 \in \text{BD}(D) : (Dq_0, \pi_{\text{BD}(G)}v)_{\text{BD}(G)} = 0 \text{ for all } v \in \ker(m^* - \mathring{D}a^*G)\}.$$

Before we prove the latter proposition, we need a lemma.

Lemma 5.8. *Let $q_0 \in \text{BD}(D)$. Let $f_0 \in \text{dom}(G)$ be such that*

$$(f_0, v)_{\text{dom}(G)} = (Dq_0, \pi_{\text{BD}(G)}v)_{\text{BD}(G)}$$

for all $v \in \text{dom}(G)$. Let $u \in \text{dom}(G)$. Then the following statements are equivalent.

- (i) $Tu = f_0$.
- (ii) $u \in \text{dom}(DaG)$, $mu - DaGu = 0$ and $q_0 = \pi_{\text{BD}(D)}aGu$.

Proof. ‘(i) \Rightarrow (ii)’. Let $v \in \text{dom}(G)$. Then

$$(mu, v)_{H_0} + (aGu, Gv)_{H_1} = \mathbf{b}(u, v) = (Tu, v)_{\text{dom}(G)} = (f_0, v)_{\text{dom}(G)} = (Dq_0, \pi_{\text{BD}(G)}v)_{\text{BD}(G)}.$$

Hence $(mu, v)_{H_0} + (aGu, \mathring{G}v)_{H_1} = 0$ for all $v \in \text{dom}(\mathring{G})$. So $aGu \in \text{dom}((\mathring{G})^*) = \text{dom}(D)$ and $DaGu = -(\mathring{G})^*aGu = mu$. In particular, $u \in \text{dom}(DaG)$. Alternatively, if $v \in \text{BD}(G)$, then

$$\begin{aligned} (Dq_0, v)_{\text{BD}(G)} &= (Dq_0, \pi_{\text{BD}(G)}v)_{\text{BD}(G)} = (mu, v)_{H_0} + (aGu, Gv)_{H_1} \\ &= (DaGu, DGv)_{H_0} + (aGu, Gv)_{H_1} = (aGu, Gv)_{\text{dom}(D)} \\ &= (\pi_{\text{BD}(D)}aGu, Gv)_{\text{BD}(D)} = (D\pi_{\text{BD}(D)}aGu, v)_{\text{BD}(G)} \end{aligned}$$

by Lemma 2.7. So $q_0 = \pi_{\text{BD}(D)}aGu$.

‘(ii) \Rightarrow (i)’. Let $v \in \text{dom}(\mathring{G})$. Since $(\mathring{G})^* = -D$ one deduces that

$$\begin{aligned} (Tu, v)_{\text{dom}(G)} &= \mathbf{b}(u, v) = (aGu, \mathring{G}v)_{H_1} + (mu, v)_{H_0} \\ &= -(DaGu, v)_{H_0} + (mu, v)_{H_0} = 0 = (Dq_0, \pi_{\text{BD}(G)}v)_{\text{BD}(G)} = (f_0, v)_{\text{dom}(G)}. \end{aligned}$$

Alternatively, if $v \in \text{BD}(G)$, then

$$\begin{aligned} (Tu, v)_{\text{dom}(G)} &= \mathbf{b}(u, v) = (aGu, Gv)_{H_1} + (mu, v)_{H_0} \\ &= (aGu, Gv)_{H_1} + (DaGu, DGv)_{H_0} \\ &= (aGu, Gv)_{\text{dom}(D)} = (\pi_{\text{BD}(D)}aGu, Gv)_{\text{BD}(D)} = (q_0, Gv)_{\text{BD}(D)} \\ &= (Dq_0, v)_{\text{BD}(G)} = (f_0, v)_{\text{dom}(G)}. \end{aligned}$$

So by linearity $(Tu, v)_{\text{dom}(G)} = (f_0, v)_{\text{dom}(G)}$ for all $v \in \text{dom}(G)$ and $Tu = f_0$. \square

Proof of Proposition 5.7. Let $q_0 \in \text{BD}(D)$. Let $f_0 \in \text{dom}(G)$ be as in Lemma 5.8. Then it follows from Lemma 5.8 that $q_0 \in \text{dom}(\Lambda^{-1})$ if and only if $f_0 \in \text{ran}(T)$. But $\text{ran}(T) = (\ker(T^*))^{\perp_{\text{dom}(G)}}$ since $\text{ran}(T)$ is closed in $\text{dom}(G)$. Now $\ker(T^*) = \ker(m^* - \mathring{D}a^*G)$ because $G^* = -\mathring{D}$. Hence $f_0 \in \text{ran}(T)$ if and only if $(Dq_0, \pi_{\text{BD}(G)}v)_{\text{BD}(G)} = 0$ for all $v \in \ker(m^* - \mathring{D}a^*G)$. \square

As in Lemma 5.5 one has the following sufficient condition for the closedness of $\text{ran}(T)$.

Lemma 5.9. *If the inclusion $\text{dom}(G) \subset H_0$ is compact, then $\text{ran}(T)$ is closed in $\text{dom}(G)$.*

In our model case Example 2.3, the inclusion $\text{dom}(G) \subset H_0$ is compact if Ω has a continuous boundary.

We conclude with a variant of the Dirichlet-to-Neumann graph involving an intermediate space as in Section 3. Throughout the remainder of this section let H be a Hilbert space and $\kappa \in \mathcal{L}(\text{BD}(G), H)$ injective with dense range. Define

$$\Lambda_H = \{(\varphi, \psi) \in H \times H : \text{there exists a } u_0 \in \text{BD}(G) \text{ such that } \kappa(u_0) = \varphi \text{ and } (u_0, G\kappa^*\psi) \in \Lambda\}.$$

We call Λ_H the **Dirichlet-to-Neumann graph in H associated with $-DaG + m$** . It follows from Lemma 5.2 that Λ_H is single-valued if $\ker(m - Da\mathring{G}) = \{0\}$.

The graph Λ_H can be described with a form.

Proposition 5.10. *Define $j: \text{dom}(G) \rightarrow H$ by $j = \kappa \circ \pi_{\text{BD}(G)}$. Then*

$$\Lambda_H = \{(\varphi, \psi) \in H \times H : \text{there exists a } u \in \text{dom}(G) \text{ such that } j(u) = \varphi \text{ and } \mathbf{b}(u, v) = (\psi, j(v))_{\text{dom}(G)} \text{ for all } v \in \text{dom}(G)\}.$$

Proof. This follows as in the proof of Proposition 3.6. \square

Corollary 5.11. *If $\ker(-Da\mathring{G} + m) = \{0\}$ and the inclusion $\text{dom}(G) \subset H_0$ is compact, then Λ_H is an m -sectorial operator.*

Proof. Let $j = \kappa \circ \pi_{\text{BD}(G)}: \text{dom}(G) \rightarrow H$ and let $V(\mathbf{b}) = \{u \in \text{dom}(G) : \mathbf{b}(u, v) = 0 \text{ for all } v \in \ker j\}$. Then $V(\mathbf{b}) \cap \ker j = \ker(-Da\mathring{G} + m) = \{0\}$. Then the statement follows from [ACSVV] Theorem 8.11 and Proposition 5.10. \square

Even if the inclusion $\text{dom}(G) \subset H_0$ is compact, then in general Λ_H is not an m -sectorial graph. A counterexample has been given in [BE2] Example 3.7.

6 Resolvent convergence, non-coercive case

In this section we consider resolvent convergence of a sequence of Dirichlet-to-Neumann operators without the coercivity condition on m . Throughout this section, we adopt the notation and assumptions as in the beginning of Section 2. Let H be a Hilbert space and let $\kappa \in \mathcal{L}(\text{BD}(G), H)$ be one-to-one with dense range. Set $j = \kappa \circ \pi_{\text{BD}(G)}: \text{dom}(G) \rightarrow H$.

We need a stronger version of convergence for the leading coefficients, which we next introduce. Let $a, a_1, a_2, \dots \in \mathcal{L}(H_1)$ be coercive. We say that $(a_n)_{n \in \mathbb{N}}$ **converges to a independent of the boundary conditions** if for every strictly increasing sequence $(n_k)_{k \in \mathbb{N}}$ in \mathbb{N} , all $f, f_1, f_2, \dots \in H_0$ and all $u, u_1, u_2, \dots \in \text{dom}(G)$ with

$$\left[\begin{array}{l} \lim_{k \rightarrow \infty} f_k = f \text{ weakly in } H_0, \\ \lim_{k \rightarrow \infty} u_k = u \text{ weakly in } \text{dom}(G), \text{ and} \\ u_k \in \text{dom}(Da_{n_k}G) \text{ and } -Da_{n_k}Gu_k = f_k \text{ for all } k \in \mathbb{N} \end{array} \right. \quad (11)$$

it follows that $\lim_{k \rightarrow \infty} a_{n_k}Gu_k = aGu$ weakly in H_1 .

Note that D is weakly closed and $\lim_{k \rightarrow \infty} D(a_{n_k}Gu_k) = \lim_{k \rightarrow \infty} -f_k = -f$ weakly in H_0 . So $aGu \in \text{dom}(D)$ and $-DaGu = f$. In particular $u \in \text{dom}(DaG)$.

Example 6.1. In this example we show that in the classical situation, convergence of the coefficients independent of the boundary conditions is implied by the already studied notion of H -convergence, see [Tar] and [MT].

Let $\Omega \subset \mathbb{R}^d$ be open and bounded. Further, let H_0, H_1, G and D be as in Example 2.3. We identify an element of $L_\infty(\Omega, \mathbb{C}^{d \times d})$ with an element of $\mathcal{L}(H_1)$ in the natural way. Let $a, a_1, a_2, \dots \in L_\infty(\Omega, \mathbb{C}^{d \times d})$. Suppose that $\text{Re } a_n \geq \mu I$ for all $n \in \mathbb{N}$, $\text{Re } a \geq \mu I$ and $\sup_n \|a_n\|_{\mathcal{L}(H_1)} < \infty$. Further suppose that $(a_n)_{n \in \mathbb{N}}$ is H -convergent to a . Then $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions.

Indeed, let $f, f_1, f_2, \dots \in L_2(\Omega)$, $u, u_1, u_2, \dots \in H^1(\Omega)$ and $(n_k)_{k \in \mathbb{N}}$ satisfy (11). Then every subsequence $(a_{n_k})_{k \in \mathbb{N}}$ is H -convergent to a by the discussion after Definition 6.4 in [Tar]. So without loss of generality we may assume that $n_k = k$ for all $k \in \mathbb{N}$. As $(u_k)_{k \in \mathbb{N}}$ converges to u weakly in $H^1(\Omega)$ it also converges weakly in $H_{\text{loc}}^1(\Omega)$. The inclusion $H_0^1(\Omega) \subset L_2(\Omega)$ is compact since Ω is bounded. Hence also the inclusion $L_2(\Omega) \subset (H_0^1(\Omega))' = H^{-1}(\Omega)$ is compact. Therefore $(f_k)_{k \in \mathbb{N}}$ converges strongly to f in $H^{-1}(\Omega) \subset H_{\text{loc}}^{-1}(\Omega)$. Then the criteria of Lemma 10.3 in [Tar] are fulfilled and we obtain that $(a_k Gu_k)_{k \in \mathbb{N}}$ converges weakly to aGu in $L_{2,\text{loc}}(\Omega)^d$. Since the sequence $(a_k Gu_k)_{k \in \mathbb{N}}$ in $L_2(\Omega)^d$ is bounded in $L_2(\Omega)^d$, there exists a $q \in L_2(\Omega)^d$ and a subsequence of $(a_k Gu_k)_{k \in \mathbb{N}}$ that weakly converges to q in $L_2(\Omega)^d$. By uniqueness of limits in $L_{2,\text{loc}}(\Omega)^d$, we must have that $q = aGu$. So the subsequence converges to aGu in $L_2(\Omega)^d$. Using the standard subsequence argument we deduce that $(a_k Gu_k)_{k \in \mathbb{N}}$ converges weakly to aGu in $L_2(\Omega)^d = H_1$.

The condition $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions, which we use in this section, is stronger than the condition used for the convergence in Theorem 4.2.

Proposition 6.2. *Let $a, a_1, a_2, \dots \in \mathcal{L}(H_1)$ and $\mu > 0$. Suppose that $\text{Re } a_n \geq \mu I$ for all $n \in \mathbb{N}$ and $\text{Re } a \geq \mu I$. Suppose that $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions. Further assume that the inclusion $\text{dom}(G) \subset H_0$ is compact. Let $\iota: \text{ran}(G) \hookrightarrow H_1$ be the embedding map. Then $\lim_{n \rightarrow \infty} (\iota^* a_n \iota)^{-1} = (\iota^* a \iota)^{-1}$ in the weak operator topology on $\mathcal{L}(\text{ran}(G))$.*

Proof. Let $q \in \text{ran } G \cap \text{dom } \mathring{D}$. Let $n \in \mathbb{N}$. Write $r_n = (\iota^* a_n \iota)^{-1} q$. Then $r_n \in \text{ran } G$ and $\|r_n\|_{H_1} \leq \mu^{-1} \|q\|_{H_1}$. There exists a $u_n \in \text{dom } G \cap (\ker G)^\perp$ such that $Gu_n = r_n$. Then

the sequence $(u_n)_{n \in \mathbb{N}}$ is bounded in $\text{dom } G$ by Lemma 4.1(a). Passing to a subsequence if necessary, there exists a $u \in \text{dom } G$ such that $\lim u_n = u$ weakly in $\text{dom } G$. Let $n \in \mathbb{N}$. Then $q = \iota^* a_n \iota r_n = \iota^* a_n G u_n$. Since $q \in \text{dom } \mathring{D}$ it follows from Lemma 4.3(a) that $a_n G u_n \in \text{dom } \mathring{D}$ and $\mathring{D} a_n G u_n = \mathring{D} \iota^* a_n G u_n = \mathring{D} q$. Because $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions, we obtain that $\lim a_n G u_n = a G u$ weakly in H_1 . Since the operator \mathring{D} is closed, we obtain that $a G u \in \text{dom } \mathring{D}$ and $\mathring{D} a G u = \mathring{D} q$. Using again Lemma 4.3(a) one deduces that $\iota^* a G u \in \text{dom } \mathring{D}$ and $\mathring{D} \iota^* a G u = \mathring{D} q$. Hence $(\mathring{D} \iota) \iota^* a \iota G u = (\mathring{D} \iota) q$. Since $\mathring{D} \iota$ is injective by Lemma 4.3(c), it follows that $\iota^* a \iota G u = q$. So $G u = (\iota^* a \iota)^{-1} q$. Then

$$\lim (\iota^* a_n \iota)^{-1} q = \lim r_n = \lim G u_n = G u = (\iota^* a \iota)^{-1} q$$

weakly in $\text{ran } G$.

Finally, since $\sup \|(\iota^* a_n \iota)^{-1}\|_{\mathcal{L}(\text{ran } G)} < \infty$ and $\text{ran } G \cap \text{dom } \mathring{D}$ is dense in $\text{ran } G$ by Lemma 4.3(b), one concludes that $\lim (\iota^* a_n \iota)^{-1} = (\iota^* a \iota)^{-1}$ in the weak operator topology on $\mathcal{L}(\text{ran}(G))$. \square

Remark 6.3. The above proposition is also valid if ι is replaced by the embedding of a closed subspace of $\text{ran } G$ which contains \mathring{G} . This is the motivation for the terminology $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions.

The main theorem of this section is as follows.

Theorem 6.4. *Let $a, a_1, a_2, \dots \in \mathcal{L}(H_1)$, $m, m_1, m_2, \dots \in \mathcal{L}(H_0)$ and $\mu > 0$. Suppose that $\text{Re } a_n \geq \mu I$ for all $n \in \mathbb{N}$, $\text{Re } a \geq \mu I$ and $\sup_n \|a_n\|_{\mathcal{L}(H_1)} < \infty$. Suppose that $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions and $\lim m_n = m$ in the weak operator topology on $\mathcal{L}(H_0)$. Assume that $\ker(m_n - Da_n \mathring{G}) = \{0\}$ for all $n \in \mathbb{N}$ and $\ker(m - Da \mathring{G}) = \{0\}$. Further assume that the inclusion $\text{dom}(G) \subset H_0$ is compact.*

For all $n \in \mathbb{N}$ let $\Lambda_H^{(n)}$ and Λ_H be the Dirichlet-to-Neumann operators in H associated with $-Da_n G + m_n$ and $-Da G + m$, respectively. Then one has the following.

- (a) *The sequence $(\Lambda_H^{(n)})_{n \in \mathbb{N}}$ of operators is uniformly sectorial.*
- (b) *$\lim_{n \rightarrow \infty} (\lambda I + \Lambda_H^{(n)})^{-1} = (\lambda I + \Lambda_H)^{-1}$ in the weak operator topology for all large $\lambda > 0$.*
- (c) *If κ is compact, then*

$$\lim_{n \rightarrow \infty} (\lambda I + \Lambda_H^{(n)})^{-1} = (\lambda I + \Lambda_H)^{-1}$$

uniformly in $\mathcal{L}(H)$ for all large $\lambda > 0$.

The proof requires a lot of preparation. Adopt the notation and assumptions of Theorem 6.4. For all $n \in \mathbb{N}$ define $\mathfrak{b}_n: \text{dom}(G) \times \text{dom}(G) \rightarrow \mathbb{C}$ by

$$\mathfrak{b}_n(u, v) = (a_n G u, G v)_{H_1} + (m_n u, v)_{H_0}$$

and define $V(\mathfrak{b}_n) = \{u \in \text{dom}(G) : \mathfrak{b}_n(u, v) = 0 \text{ for all } v \in \ker j\}$. Define similarly \mathfrak{b} and $V(\mathfrak{b})$.

Lemma 6.5. For all $\varepsilon > 0$ there exists an $\omega > 0$ such that

$$\|u\|_{H_0}^2 \leq \varepsilon \|u\|_{\text{dom}(G)}^2 + \omega \|j(u)\|_H^2$$

for all $n \in \mathbb{N}$ and $u \in V(\mathfrak{b}_n)$.

Proof. Let $n \in \mathbb{N}$. Since $\ker(m_n - Da_n\mathring{G}) = \{0\}$, the restriction $j|_{V(\mathfrak{b}_n)}$ is injective. Because also the inclusion $\text{dom}(G) \subset H_0$ is compact, it follows that for all $\varepsilon > 0$ there exists an $\omega > 0$ such that

$$\|u\|_{H_0}^2 \leq \varepsilon \|u\|_{\text{dom}(G)}^2 + \omega \|j(u)\|_H^2$$

for all $u \in V(\mathfrak{b}_n)$. We next show that one can choose ω uniformly in n .

Suppose the lemma is false. Then without loss of generality and passing to a subsequence if necessary there exist $\varepsilon > 0$ and for all $n \in \mathbb{N}$ there exists a $u_n \in V(\mathfrak{b}_n)$ such that

$$\|u_n\|_{H_0}^2 > \varepsilon \|u_n\|_{\text{dom}(G)}^2 + n \|j(u_n)\|_H^2.$$

Without loss of generality we may assume that $\|u_n\|_{H_0} = 1$ for all $n \in \mathbb{N}$. Then $\varepsilon \|u_n\|_{\text{dom}(G)}^2 \leq 1$ for all $n \in \mathbb{N}$, so the sequence $(u_n)_{n \in \mathbb{N}}$ is bounded in $\text{dom}(G)$. Passing to a subsequence if necessary there exists a $u \in \text{dom}(G)$ such that $\lim u_n = u$ weakly in $\text{dom}(G)$. Since the inclusion $\text{dom}(G) \subset H_0$ is compact it follows that $u = \lim u_n$ in H_0 . In particular $\|u\|_{H_0} = 1$ and $u \neq 0$. Also $j(u) = \lim j(u_n) = 0$ in H , so $u \in \ker j = \text{dom}(\mathring{G})$.

If $n \in \mathbb{N}$, then $(a_n G u_n, \mathring{G} v)_{H_1} = -(m_n u_n, v)_{H_0}$ for all $v \in \text{dom}(\mathring{G}) = \ker j$, since $u_n \in V(\mathfrak{b}_n)$. Therefore $a_n G u_n \in \text{dom}((\mathring{G})^*) = \text{dom}(D)$ and $-Da_n G u_n = -m_n u_n$. Next $\lim m_n u_n = mu$ weakly in H_0 . Since $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions one deduces that $a G u \in \text{dom}(D)$ and $-Da G u = -mu$. Then $u \in \ker(m - Da\mathring{G}) = \{0\}$. So $u = 0$. This is a contradiction. \square

Lemma 6.6. There exist $\tilde{\mu}, \omega > 0$ such that

$$\tilde{\mu} \|u\|_{\text{dom}(G)}^2 \leq \text{Re } \mathfrak{b}_n(u) + \omega \|j(u)\|_H^2$$

for all $n \in \mathbb{N}$ and $u \in V(\mathfrak{b}_n)$.

Proof. Let $\tilde{\omega} = \mu + \sup_n \|m_n\|_{\mathcal{L}(H_0)}$. Then

$$\mu \|u\|_{\text{dom}(G)}^2 \leq \text{Re}(a_n G u, G u)_{H_1} + \mu \|u\|_{H_0}^2 \leq \text{Re } \mathfrak{b}_n(u) + \tilde{\omega} \|u\|_{H_0}^2$$

for all $n \in \mathbb{N}$ and $u \in \text{dom}(G)$.

Choose $\varepsilon = \frac{\mu}{2\tilde{\omega}}$ and let $\omega > 0$ be as in Lemma 6.5. Let $n \in \mathbb{N}$ and $u \in V(\mathfrak{b}_n)$. Then

$$\begin{aligned} \mu \|u\|_{\text{dom}(G)}^2 &\leq \text{Re } \mathfrak{b}_n(u) + \tilde{\omega} \|u\|_{H_0}^2 \\ &\leq \text{Re } \mathfrak{b}_n(u) + \tilde{\omega} \left(\frac{\mu}{2\tilde{\omega}} \|u\|_{\text{dom}(G)}^2 + \omega \|j(u)\|_H^2 \right) \\ &= \text{Re } \mathfrak{b}_n(u) + \frac{\mu}{2} \|u\|_{\text{dom}(G)}^2 + \omega \tilde{\omega} \|j(u)\|_H^2. \end{aligned}$$

So

$$\frac{\mu}{2} \|u\|_{\text{dom}(G)}^2 \leq \text{Re } \mathfrak{b}_n(u) + \omega \tilde{\omega} \|j(u)\|_H^2$$

and the lemma follows. \square

Now we are able to prove Theorem 6.4.

Proof of Theorem 6.4. Let $\tilde{\mu}, \omega > 0$ be as in Lemma 6.6.

‘(a)’. Set $c = \sup_{n \in \mathbb{N}} (\|a_n\|_{\mathcal{L}(H_1)} + \|m_n\|_{\mathcal{L}(H_0)})$. Let $n \in \mathbb{N}$ and $\varphi \in \text{dom}(\Lambda_H^{(n)})$. There exists a $u \in \text{dom}(G)$ such that $j(u) = \varphi$ and $\mathfrak{b}_n(u, v) = (\Lambda_H^{(n)}\varphi, j(v))_H$ for all $v \in \text{dom}(G)$. Then $u \in V(\mathfrak{b}_n)$ and $((\Lambda_H^{(n)} + \omega I)\varphi, \varphi)_H = \mathfrak{b}_n(u) + \omega \|j(u)\|_H^2$, so $\text{Re}((\Lambda_H^{(n)} + \omega I)\varphi, \varphi)_H \geq \tilde{\mu} \|u\|_{\text{dom}(G)}^2$. Therefore

$$|\text{Im}((\Lambda_H^{(n)} + \omega I)\varphi, \varphi)_H| = |\text{Im} \mathfrak{b}_n(u)| \leq c \|u\|_{\text{dom}(G)}^2 \leq \frac{c}{\tilde{\mu}} \text{Re}((\Lambda_H^{(n)} + \omega I)\varphi, \varphi)_H.$$

Hence the operators $\Lambda_H^{(n)}$ are sectorial with vertex $-\omega$ and semi-angle $\arctan \frac{c}{\tilde{\mu}}$, uniformly in n .

‘(b)’. In order not to repeat part of the proof in Statement (c) we first prove something more general. Let $\lambda > \omega$. Let $\psi, \psi_1, \psi_2, \dots \in H$ and suppose that $\lim \psi_n = \psi$ weakly in H . We shall prove that $\lim(\lambda I + \Lambda_H^{(n)})^{-1}\psi_n = (\lambda I + \Lambda_H)^{-1}\psi$ weakly in H .

Let $n \in \mathbb{N}$. Set $\varphi_n = (\lambda I + \Lambda_H^{(n)})^{-1}\psi_n$. There exists a $u_n \in V(\mathfrak{b}_n)$ such that $j(u_n) = \varphi_n$ and

$$\mathfrak{b}_n(u_n, v) + \lambda(j(u_n), j(v))_H = (\psi_n, j(v))_H \quad (12)$$

for all $v \in \text{dom}(G)$. Choose $v = u_n$. Then Lemma 6.6 gives

$$\tilde{\mu} \|u_n\|_{\text{dom}(G)}^2 \leq \text{Re} \mathfrak{b}_n(u_n) + \lambda \|j(u_n)\|_H^2 = \text{Re}(\psi_n, j(u_n))_H \leq \|\psi_n\|_H \|j\|_{\mathcal{L}(\text{dom}(G), H)} \|u_n\|_{\text{dom}(G)}.$$

So $\|u_n\|_{\text{dom}(G)} \leq \tilde{\mu}^{-1} \|\psi_n\|_H \|j\|_{\mathcal{L}(\text{dom}(G), H)}$. Since the sequence $(\psi_n)_{n \in \mathbb{N}}$ is bounded in H , the sequence $(u_n)_{n \in \mathbb{N}}$ is bounded in $\text{dom}(G)$. Passing to a subsequence if necessary, there exists a $u \in \text{dom}(G)$ such that $\lim u_n = u$ weakly in $\text{dom}(G)$. Since the inclusion $\text{dom}(G) \subset H_0$ is compact one deduces that $\lim u_n = u$ in H_0 . Then $\lim m_n u_n = mu$ weakly in H_0 . Moreover, $\lim \varphi_n = \lim j(u_n) = j(u)$ weakly in H . Next we show that $j(u) = (\lambda I + \Lambda_H)^{-1}\psi$.

Let $n \in \mathbb{N}$. If $v \in \ker j = \text{dom}(\overset{\circ}{G})$, then $\mathfrak{b}_n(u_n, v) = 0$, so $(a_n G u_n, \overset{\circ}{G} v)_{H_1} = -(m_n u_n, v)_{H_0}$. Hence $a_n G u_n \in \text{dom}((\overset{\circ}{G})^*) = \text{dom}(D)$ and $-D a_n G u_n = -m_n u_n$. In particular, $u_n \in \text{dom}(D a_n G)$. Moreover, $\lim u_n = u$ weakly in $\text{dom}(G)$ and $\lim m_n u_n = mu$ weakly in H_0 . Since $(a_n)_{n \in \mathbb{N}}$ converges to a independent of the boundary conditions, one deduces that $\lim a_n G u_n = a G u$ weakly in H_1 .

Let $v \in \text{dom}(G)$. If $n \in \mathbb{N}$, then (12) gives

$$(a_n G u_n, G v)_{H_1} + (m_n u_n, v)_{H_0} + \lambda(j(u_n), j(v))_H = (\psi_n, j(v))_H.$$

Taking the limit $n \rightarrow \infty$ one establishes

$$(a G u, G v)_{H_1} + (mu, v)_{H_0} + \lambda(j(u), j(v))_H = (\psi, j(v))_H.$$

So $\mathfrak{b}(u, v) + \lambda(j(u), j(v))_H = (\psi, j(v))_H$. Therefore $j(u) \in \text{dom}(\Lambda_H)$ and $(\lambda I + \Lambda_H)j(u) = \psi$. With the usual subsequence argument we proved that $\lim(\lambda I + \Lambda_H^{(n)})^{-1}\psi_n = (\lambda I + \Lambda_H)^{-1}\psi$ weakly in H . Now Statement (b) follows by choosing $\psi_n = \psi$ for all $n \in \mathbb{N}$.

‘(c)’. Finally suppose that κ is compact. Then also j is compact. Let $\lambda > \omega$. Suppose $\lim(\lambda I + \Lambda_H^{(n)})^{-1} = (\lambda I + \Lambda_H)^{-1}$ in $\mathcal{L}(H)$ is false. Passing to a subsequence if necessary, there exist $\delta > 0$ and $\psi_1, \psi_2, \dots \in H$ such that

$$\|(\lambda I + \Lambda_H^{(n)})^{-1}\psi_n - (\lambda I + \Lambda_H)^{-1}\psi_n\|_H > \delta\|\psi_n\|_H$$

for all $n \in \mathbb{N}$. Without loss of generality we may assume that $\|\psi_n\|_H = 1$ for all $n \in \mathbb{N}$. Passing again to a subsequence if necessary, there exists a $\psi \in H$ such that $\lim \psi_n = \psi$ weakly in H . Let $u_n \in V(\mathbf{b}_n)$ and $u \in \text{dom}(G)$ be as in Part (b) for all $n \in \mathbb{N}$. Then $\lim u_n = u$ weakly in $\text{dom}(G)$, so

$$\lim_{n \rightarrow \infty} (\lambda I + \Lambda_H^{(n)})^{-1}\psi_n = \lim_{n \rightarrow \infty} j(u_n) = j(u) = (\lambda I + \Lambda_H)^{-1}\psi$$

in H by the compactness of j . Similarly $\lim_{n \rightarrow \infty} (\lambda I + \Lambda_H^{(n)})^{-1}\psi = (\lambda I + \Lambda_H)^{-1}\psi$ in H . So

$$\lim_{n \rightarrow \infty} \|(\lambda I + \Lambda_H^{(n)})^{-1}\psi_n - (\lambda I + \Lambda_H)^{-1}\psi_n\|_H = 0.$$

This is a contradiction. □

Note that the limit Dirichlet-to-Neumann graph Λ_H is an operator in Theorem 6.4. In [AEKS] Theorem 5.11 a different condition on the a_n is used to obtain resolvent convergence for symmetric operators/graphs, but possibly multi-valued limit graph Λ_H . Since we do not wish to require symmetry in Theorem 6.4 and we need that the limit graph Λ_H is m -sectorial, we require conveniently that all graphs are single-valued. See also the discussion at the end of Section 5.

7 More examples

The first example is from linearized elasticity.

Example 7.1. Let $\Omega \subset \mathbb{R}^d$ be open. Set

$$L_{2,\text{sym}}(\Omega) = \{S \in L_2(\Omega)^{d \times d} : S^T = S \text{ a.e.}\}.$$

Choose $H_0 = L_2(\Omega)^d$ and $H_1 = L_{2,\text{sym}}(\Omega)$. Define $\widehat{G}: C_c^\infty(\Omega)^d \rightarrow L_{2,\text{sym}}(\Omega)$ by

$$(\widehat{G}u)_{kl} = \frac{1}{2}(\partial_k u_l + \partial_l u_k).$$

Further define $\widehat{D}: C_c^\infty(\Omega)^{d \times d} \cap L_{2,\text{sym}}(\Omega) \rightarrow L_2(\Omega)^d$ by

$$(\widehat{D}q)_k = \sum_{l=1}^d \partial_l q_{kl}.$$

Then $\text{dom}(\widehat{G})$ is dense in H_0 and $\text{dom}(\widehat{D})$ is dense in H_1 . Moreover, using integration by parts one deduces that (2) is valid. Then one can apply Example 2.2.

Korn's first inequality implies that $\|\partial_k u_l\|_{L_2(\Omega)} \leq \sqrt{2}\|\widehat{G}u\|_{H_1}$ for all $u \in C_c^\infty(\Omega)^d$ and $k, l \in \{1, \dots, d\}$. So $\text{dom}(\overset{\circ}{G}) \subset H_0^1(\Omega)$. In particular the inclusion $\text{dom}(\overset{\circ}{G}) \subset H_0$ is compact if Ω is bounded.

Under some regularity conditions on the boundary of Ω , Korn's second inequality states that there exists a $c > 0$ such that $\|\partial_k u_l\|_{L_2(\Omega)} \leq c\|u\|_{\text{dom}(G)}$ for all $u \in \text{dom}(G)$ and $k, l \in \{1, \dots, d\}$. For example, if Ω is bounded with a Lipschitz boundary, then Korn's second inequality is valid. For an easy proof see [Nit] Section 3. If Korn's second inequality is valid, then $\text{dom}(G) \subset H^1(\Omega)^d$. Consequently, if Korn's second inequality is valid and Ω has a continuous boundary, then the inclusion $H^1(\Omega) \subset L_2(\Omega)$ is compact and hence the inclusion $\text{dom}(G) \subset H_0$ is compact. We point out that Korn's second inequality is not a necessary condition for the inclusion $\text{dom}(G) \subset H_0$ to be compact, see [Wec] Theorem 1.

In particular, suppose Ω is bounded with a Lipschitz boundary and write $\Gamma = \partial\Omega$. Let $\sigma \in (-\infty, \frac{1}{2}]$ and set $H = H^\sigma(\Gamma)^d$. Then $\text{Tr } u \in H$ for all $u \in \text{dom}(G)$. Moreover, $\text{Tr}|_{\text{BD}(G)}: \text{BD}(G) \rightarrow H$ is injective and has dense range. So one can consider as in Section 3 a Dirichlet-to-Neumann operator in H . Note that $\text{Tr}|_{\text{BD}(G)}$ is compact if $\sigma < \frac{1}{2}$.

The second example is from electro-magneto statics.

Example 7.2. Let $\Omega \subset \mathbb{R}^3$ be open. Using integration by parts one deduces that

$$(\text{curl } u, v)_{L_2(\Omega)^3} = (u, \text{curl } v)_{L_2(\Omega)^3}$$

for all $u, v \in C_c^\infty(\Omega)^3$. Therefore let $H_0 = H_1 = L_2(\Omega)^3$ and define $\widehat{G} = \widehat{D}: C_c^\infty(\Omega)^3 \rightarrow L_2(\Omega)^3$ by $\widehat{G}u = \widehat{D}u = i \text{curl } u$. Then (2) is satisfied. Using the construction in Example 2.2 one obtains a new example.

Acknowledgements

The third named author expresses his gratitude for the wonderful atmosphere and hospitality extended to him during a two months research visit at the University of Auckland. Part of this work is supported by the Marsden Fund Council from Government funding, administered by the Royal Society of New Zealand. Part of this work is supported by the EU Marie Curie IRSES program, Project AOS, No. 318910. Part of this work was carried out with financial support of the EPSRC grant EP/L018802/2: Mathematical foundations of metamaterials: homogenisation, dissipation and operator theory.

References

- [ACSVV] ARENDT, W., CHILL, R., SEIFERT, C., VOGT, H. and VOIGT, J., Internet Seminar 18, 2015.
- [AE1] ARENDT, W. and ELST, A. F. M. TER, The Dirichlet-to-Neumann operator on rough domains. *J. Differential Equations* **251** (2011), 2100–2124.
- [AE2] ———, Sectorial forms and degenerate differential operators. *J. Operator Theory* **67** (2012), 33–72.

- [AEKS] ARENDT, W., ELST, A. F. M. TER, KENNEDY, J. B. and SAUTER, M., The Dirichlet-to-Neumann operator via hidden compactness. *J. Funct. Anal.* **266** (2014), 1757–1786.
- [AKM] AXELSSON, A., KEITH, S. and M^cINTOSH, A., Quadratic estimates and functional calculi of perturbed Dirac operators. *Invent. Math.* **163** (2006), 455–497.
- [BE1] BEHRNDT, J. and ELST, A. F. M. TER, Dirichlet-to-Neumann maps on bounded Lipschitz domains. *J. Differential Equations* **259** (2015), 5903–5926.
- [BE2] ———, The Dirichlet-to-Neumann map for Schrödinger operators with complex potentials. *Discrete Contin. Dyn. Syst. Ser. S* **10** (2017), 661–671.
- [McL] MCLEAN, W., *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, 2000.
- [MT] MURAT, F. and TARTAR, L., *H*-convergence. In CHERKAEV, A. and KOHN, R., eds., *Topics in the mathematical modelling of composite materials*, Progr. Nonlinear Differential Equations Appl. 31, 21–43. Birkhäuser, Boston, 1997.
- [Nit] NITSCHKE, J. A., On Korn’s second inequality. *RAIRO Anal. Numér.* **15** (1981), 237–248.
- [PSTW] PICARD, R., SEIDLER, S., TROSTORFF, S. and WAURICK, M., On abstract div–grad systems. *J. Differential Equations* **260** (2016), 4888–4917.
- [PTW1] PICARD, R., TROSTORFF, S. and WAURICK, M., On a class of boundary control problems. *Oper. Matrices* **8** (2014), 185–204.
- [PTW2] ———, On a comprehensive class of linear control problems. *IMA J. Math. Control Inform.* **33** (2016), 257–291.
- [Tar] TARTAR, L., *The general theory of homogenization. A personalized introduction*. Lecture Notes of the Unione Matematica Italiana 7. Springer-Verlag, Berlin, 2009.
- [Tro] TROSTORFF, S., A characterization of boundary conditions yielding maximal monotone operators. *J. Funct. Anal.* **267** (2014), 2787–2822.
- [TW] TROSTORFF, S. and WAURICK, M., A note on elliptic type boundary value problems with maximal monotone relations. *Math. Nachr.* **287** (2014), 1545–1558.
- [Wau] WAURICK, M., On the continuous dependence on the coefficients of evolutionary equations, 2016. Habilitation thesis, TU Dresden.
- [Wec] WECK, N., Local compactness for linear elasticity in irregular domains. *Math. Methods Appl. Sci.* **17** (1994), 107–113.

A.F.M. ter Elst
University of Auckland
Department of Mathematics
Auckland 1142
New Zealand
`terelst@math.auckland.ac.nz`

G. Gordon
University of Auckland
Department of Mathematics
Auckland 1142
New Zealand
`g.gordon@auckland.ac.nz`

Marcus Waurick
Department of Mathematics and Statistics
University of Strathclyde
Livingstone Tower
26 Richmond Street
Glasgow G1 1XH
UK
`marcus.waurick@strath.ac.uk`