



Methods of probing the interactions between small molecules and disordered proteins

Gabriella T. Heller¹ · Francesco A. Aprile¹ · Michele Vendruscolo¹

Received: 18 May 2017 / Accepted: 1 June 2017
© The Author(s) 2017. This article is an open access publication

Abstract It is generally recognized that a large fraction of the human proteome is made up of proteins that remain disordered in their native states. Despite the fact that such proteins play key biological roles and are involved in many major human diseases, they still represent challenging targets for drug discovery. A major bottleneck for the identification of compounds capable of interacting with these proteins and modulating their disease-promoting behaviour is the development of effective techniques to probe such interactions. The difficulties in carrying out binding measurements have resulted in a poor understanding of the mechanisms underlying these interactions. In order to facilitate further methodological advances, here we review the most commonly used techniques to probe three types of interactions involving small molecules: (1) those that disrupt functional interactions between disordered proteins; (2) those that inhibit the aberrant aggregation of disordered proteins, and (3) those that lead to binding disordered proteins in their monomeric states. In discussing these techniques, we also point out directions for future developments.

Keywords Disordered proteins · Small molecules · Drugs · Binding · Molecular interactions

Introduction

Disordered proteins do not adopt well-defined secondary and tertiary structures under native conditions [1–7]. These proteins can be represented as ensembles of many

conformationally distinct states, each with its own statistical weight (i.e. its probability of being occupied) [1–7]. Quite generally, all proteins exhibit some level of disorder, ranging from those that have just short dynamic terminal regions to those that are almost completely unstructured [1–7]. In many cases, the conformational heterogeneity of the latter proteins is believed to play important biological roles, as it enables them to interact with myriad partners. This multifunctionality is further enhanced by structural variations from post-translational modifications, as well as by the presence of multiple isoforms as a result of alternative splicing or pre-translational modifications [8]. Consequently, disordered proteins and proteins with disordered regions can act as central hubs in protein interaction networks for crucial regulation and signalling processes [9, 10]. Thus, it is not surprising that the dysregulation of disordered proteins is often correlated with biochemical pathways involved in cancer, cardiovascular diseases, diabetes, autoimmune disorders, and neurodegenerative conditions [10–12]. Illustrative examples of the involvement of disordered proteins in disease include the Cip/Kip cell cycle inhibitors, breast cancer type 1 susceptibility protein, and securin in the case of cancer, amyloid β , tau, α -synuclein, and huntingtin in the case of neurodegenerative disorders, and amylin (IAPP) in the case of type II diabetes [13].

For both structured and disordered proteins, the molecular mechanisms underlying protein-associated diseases can be divided into two broad categories, as a pathological condition can be triggered by either the total or partial inactivation of a protein (loss of function) or the acquisition of a new aberrant activity (gain of function). A well-known example of loss-of-function mechanism involving a disordered protein is that of the tumour suppressor protein, p53. p53 is a multi-domain protein with extended unfolded

✉ Michele Vendruscolo
mv245@cam.ac.uk

¹ Department of Chemistry, University of Cambridge,
Cambridge CB2 1EW, UK

regions under native conditions, including its N-terminal and C-terminal domains. Several cancer-related mutations of p53 are localized in these regions and alter the interactome of this protein, thereby inhibiting its regulatory activity [8, 14]. On the other hand, an example of a disordered protein that exhibits gain of toxic function in disease is α -synuclein. Several missense mutations and genomic multiplications of α -synuclein affect its native state, solubility and cellular interactions, eventually prompting the protein to form amyloid aggregates associated with Parkinson's disease [15–22].

Despite the high prevalence of disordered proteins in diseases, it is still very challenging to target these proteins using therapeutic compounds [11, 12, 23–29]. Two major obstacles in the drug discovery process for disordered proteins are: (1) the limited number of fully quantitative experimental techniques that can accurately probe disordered protein interactions with candidate therapeutic molecules compared to those available for ordered proteins, and (2) a lack of understanding of the molecular mechanisms underlying such interactions.

For the past six decades, techniques such as X-ray crystallography have yielded highly accurate structural insights about ordered proteins, which have paved the way for drug discovery and potency-optimization efforts. Such techniques, however, are generally poorly informative in the case of disordered proteins as a result of their highly dynamical nature. Progress has been made when the disordered proteins are not fully disordered and have highly ordered regions or domains for which crystal structures can be obtained, as was done to identify small-molecule inhibitors of the cancer-associated p53-murine double minute 2 (MDM2) interaction [30, 31]. However, protein crystallization is largely inapplicable to the vast majority of disordered proteins due to their conformational heterogeneity. While ordered proteins usually have a single, well-defined conformation representing a global free-energy minimum with the potential to bind small molecules with high affinity, the free-energy landscape of disordered proteins is characterized by a large number of local minima (Fig. 1). These minima correspond to the many conformations within the structural ensemble populated by disordered proteins, which can transiently bind small molecules with very weak affinity. In fact, disordered proteins can remain disordered in their bound states, and characterization of these bound states is only recently becoming possible due to experimental and computational advances [32–35]. There are also examples of disordered proteins interacting with partner proteins in which they undergo disorder-to-order transitions through coupled folding upon binding mechanisms [36, 37] or templated-folding [38], resulting in high specificity, but low affinity complexes with usually large surface areas [37]. While




Fig. 1 Schematic representation of the free-energy landscapes of ordered and disordered proteins. Structured or ordered proteins (*red*) have a free-energy landscape with a well-defined global minimal conformation, which can bind small molecules with high affinity. In contrast, disordered proteins have multiple minima within their free-energy landscape, which represent the many conformations capable of interacting with small molecules with lower affinities

these folded complexes may be crystallized, the identification of potential binding pockets from the crystal structures of the bound states is far from trivial [23]. Despite these challenges, several small molecules have been identified to modulate the behaviour of disordered proteins including the neurodegeneration-associated α -synuclein [39] and amyloid β [40], and the cancer-associated p27^{Kip1} [41], c-Myc [35, 42–44], and EWS–FLI1 [45].

To make further progress, effective techniques to probe small-molecule binding to disordered proteins must be further developed. Indeed, the lack of such methods has been a major bottleneck for the identification of molecules able to interact with disordered proteins and modulate their disease-promoting behaviour. The current situation has resulted in a poor understanding of the mechanisms underlying these interactions, which has, in turn, hindered the development of drugs active against disordered proteins. In this review, we highlight the most prominent techniques that have enabled so far major contributions to be made to the understanding of whether and how small molecules can alter the disease-promoting behaviour of disordered proteins.

Methods of identifying inhibitors of interactions between disordered proteins

The lack of well-structured binding sites within disordered proteins makes it challenging to target them directly using well-established drug discovery techniques developed for

ordered proteins such as enzymes and receptors. Instead, some approaches involve targeting disordered proteins indirectly, by blocking their binding interfaces with other proteins [46] and lipid membranes [47]. In the cases where the binding surfaces are structured and well characterized, this approach can be highly specific, as it may be amenable to standard affinity-optimization techniques. However, a thorough understanding of the binding partners involved, as well as the contact sites of interest, must usually already be well established. In this section, we discuss the state of art of this approach and present some notable examples of what we define here as ‘interface blockers’.

Perhaps one of the most well studied systems in this context is the interaction between the two disordered proteins, c-Myc and Max, which have been probed by a wide variety of techniques. c-Myc is a transcription factor associated with many types of cancer, whose interaction with its regulator Max is associated with cellular growth, metabolism, apoptosis and differentiation [48, 49]. A basic helix–loop–helix–leucine zipper (bHLHZip) in each of these two proteins facilitates their coupled folding and binding upon dimerization, creating an interface of approximately 3200 Å² in the coiled-coil dimer [35, 43, 46, 50]. The identification and characterization of small-molecule inhibitors of this interaction has represented a major milestone in demonstrating the feasibility of therapeutic targeting of disordered proteins.

Fluorescence resonance energy transfer

Inhibitors of the c-Myc/Max association were found via high-throughput screening assays of combinatorial small-molecule or peptidomimetic libraries based on fluorescence resonance energy transfer (FRET, Table 1) measurements of the bHLHZip domains of c-Myc and Max fused to cyan fluorescent protein (CFP) and yellow fluorescent protein (YFP), respectively [51–53].

FRET signals arise upon the interaction of two chromophores, whereby an excited donor chromophore transfers its excitation energy to a nearby acceptor chromophore through nonradiative dipole–dipole coupling. This transfer of energy results in both a quenching of the fluorescence of the donor and an appearance of a fluorescence emission spectra of the acceptor. Importantly, the efficiency of the energy transfer is strongly dependent on the distance between the donor and acceptor (in the range between 1 and 10 nm), thereby enabling FRET to effectively quantify molecular associations [54].

In a seminal study, c-Myc/Max FRET experiments identified two compounds, called IIA4B20 and IIA6B17, capable of inhibiting c-Myc-dependent cell growth [51]. These molecules, however, also showed activity against another oncogenic transcription factor, c-Jun, suggesting

poor specificity. Nevertheless, a follow-up study using a related combinatorial library with members assembled from a racemic, *trans*-3,4 dicarboxylic acid template yielded c-Myc/Max dimerization inhibitors that did not affect c-Jun [52], suggesting that specificity is potentially achievable in targeting disordered proteins.

Additional libraries have been screened against c-Myc/Max interfaces based on FRET experiments. One library consisted of 285 so-called ‘credit card’ compounds, designed to insert themselves into a shallow protein–protein interface hotspot of about 600 Å² rich in hydrophobic and aromatic residues, and force the protein partners to remain in their monomeric forms [53]. After FRET screening, the initial hits were further characterized by an electrophoretic mobility shift assay (EMSA) to confirm their activity. Based on the observation that the electrophoretic mobility of a bound system is less than that of the unbound system, EMSA provides quantitative information about modifiers of DNA-binding protein complexes [55]. In general, compounds in the ‘credit card’ library tend to be planar, with varying chemical diversity, designed to have favourable enthalpic contributions from van der Waals interactions, π -stacking, and favourable entropy gains from desolvation [53]. Two compounds, NY2267 and NY2280, were identified to disrupt c-Myc–Max dimerization, and inhibit both specific DNA binding and its associated oncogenic transformation. As in the case of IIA4B20 and IIA6B17, however, these molecules also showed an inhibition of c-Jun [53].

FRET experiments have also been used to characterize the conformational changes induced when the natural product trodusquemine (also known as MSI-1436) allosterically inhibits protein-tyrosine phosphatase 1B (PTP1B) by interacting with its disordered region. By labelling the N- and C-termini of PTP1B with CFP and YFP, respectively, conformational changes upon binding could be detected, suggesting that the presence of trodusquemine induced a more compact structure upon binding [56]. This binding-induced conformational change was further characterized by nuclear magnetic resonance (NMR) spectroscopy and small-angle X-ray scattering (SAXS), discussed in the “[Methods of characterizing ligand interactions with monomeric disordered proteins](#)”.

Yeast two-hybrid system

Another powerful high-throughput screening technique involves using a yeast two-hybrid system to identify small molecules capable of disrupting protein–protein interaction. In a yeast two-hybrid system (Table 1), two candidate interacting proteins are fused to the DNA-binding domain (BD) and the activation domain (AD) of a transcription

Table 1 Summary of techniques discussed in this review

Technique	Applicability	Limitations	Throughput	Selected examples	Further reading
Fluorescence resonance energy transfer (FRET)	Detection of modulators of protein–protein interactions; detection of protein–ligand interactions	Fluorescent labels required	High	c-Myc/Max and ligands [51–53], protein-tyrosine phosphatase 1B and MSI-1436 [56]	[54]
Yeast two-hybrid system	Detection of modulators of protein–protein interactions	Indirectly quantitative	High	c-Myc/Max and ligands [44, 58]	[200]
Fluorescence polarization	Detection of modulators of protein–protein interactions; detection of protein–ligand interactions	Fluorescent labels required	High	c-Myc/Max and ligands [43], c-Myc-Max complex/DNA and ligands [59, 60]	[61–63]
Circular dichroism spectroscopy (CD)	Determination of the changes in secondary structure upon binding	Low sensitivity	Low	c-Myc/Max and ligands [35, 43]	[65, 201]
Fluorescence-based aggregation kinetic assays	Identification of inhibitors of protein aggregation	Fluorescent dyes required	High	A β [40], α -synuclein [47]	[89, 92, 97]
Surface-plasmon resonance/other surface-based techniques	Real-time detection of modulators of protein–protein/interactions; detection of protein–ligand interactions	Non-specific interactions may yield false positives	Medium	EWS-FLI1 and YK-4-279 [45]	[116, 117, 202]
Small-angle X-ray scattering (SAXS)	Detection of large conformational changes upon binding at nanometer resolution	Low resolution	Variable	Protein-tyrosine phosphatase 1B and trodusquemine [56]	[114, 115]
Thermal denaturation screening	Detection of monomeric binders	Non-quantitative	High	Nuclear protein 1 and ligands [119]	[203]
Isothermal titration calorimetry (ITC)	Label-free measurement of the heat associated with binding events	Significant heat change required upon binding	Medium–low	Nuclear protein 1 and ligands [119]	[121]
Single-molecule techniques	Determination of the structure and dynamics of disordered proteins in presence of ligands	Labels required	Medium–low	α -Synuclein [125, 129]	[125]
Mass spectrometry	Localization of noncovalent interactions	May miss ligand interactions, gas-phase dissociation constants may differ from solution	Low	Polycationic spermine and α -synuclein [141]	[133, 134, 204]
Nuclear magnetic resonance (NMR) spectroscopy	Detection of protein–ligand interactions at atomic resolution	Ligand monitoring: fast, protein monitoring: time intensive, isotopic labelling may be required	Medium–low	Osteopontin/heparin [169], protein-tyrosine phosphatase 1B and trodusquemine [56]	[145, 147]
Integrative structural biology methods	Modelling of unbound/bound structural ensembles	Time intensive, computationally expensive	Low	c-Myc and ligands [28, 34]	[6]

factor, which hence functions only when a complex between the two proteins is formed. Using this setup the HLHZip domains of c-Myc and Max were fused to the BD and AD domains, respectively, of the yeast transcription factor Gal4. Only upon c-Myc/Max dimerization a fully functional transcriptional activator is produced, which is

able to induce the expression of β -galactosidase from the corresponding gene containing a Gal4-binding site within its promoter [44] (Fig. 2). As small molecules that alter this association prevent the induction of β -galactosidase in a quantitative manner, yeast two-hybrid systems represent effective screening tools to identify interface blockers. We




Fig. 2 Schematic representation of the yeast two-hybrid system. In the type of yeast two-hybrid system used to identify inhibitors of c-Myc/Max dimerization [51–53], recombinant genes encoding the HLHZip domain of c-Myc fused to the DNA-binding domain and

HLHZip domain of Max fused to the transcriptional activation domain are introduced into a yeast cell (a). Upon c-Myc/Max association, the transcriptional activation domain induces expression of β-galactosidase in a quantitative manner (b)

should add, however, that although high-throughput screening approaches of this type identified a number of c-Myc/Max dimerization inhibitors, screening against other similar dimers suggested that the majority of the resulting hits lacked specificity. Many of these hits, including the small molecule 10058-F4, were confirmed to directly interact with monomeric HLHZip region of c-Myc [35, 43]. The optimization of the 10058-F4 structure led to analogues with increased potency [42], and the development of a pharmacophore model [57]. A further discussion of the interaction of 10058-F4 with its c-Myc is continued

below in the “Methods of characterizing ligand interactions with monomeric disordered proteins”. A yeast two-hybrid system was also used to identify dihydroxycapnellene, a coral-derived sesquiterpene capable of preventing c-Myc–Max dimerization and effective against the proliferation of cancer cells [58].

Fluorescence polarization

Further studies focused on targeting the interface between the c-Myc/Max dimer and DNA by screening molecular

libraries based on fluorescence polarization experiments [59, 60] (Table 1). Fluorescence polarization is a physical phenomenon that occurs when fluorescent small molecules are excited with polarized light. The resulting emitted light is largely depolarized due to the rapid tumbling of the small molecules in solution. However, when the small molecules are bound to other species, this tumbling is quantitatively slowed as their effective hydrodynamic radii are increased, thereby better maintaining the polarization of the emitted light (Fig. 3). Because the measured polarization value is a weighted average of the free and bound states, fluorescence polarization is an important biophysical tool for drug discovery to measure the fraction of a fluorescent ligand bound to a receptor [61–63].

Fluorescence polarization screens identified two compounds, Mycro1 and Mycro2, capable of preventing DNA binding of the c-Myc/Max dimer and transcription, measured by a c-Myc reporter gene. This approach was implemented by labelling the DNA target sequences with fluorophores. However, these molecules showed signs of non-specificity as they also inhibited Max/Max DNA binding in addition to transcription from an AP-1 dependent reporter [59]. Upon further screening of Mycro1 and

Mycro2 derivatives, Mycro3 was identified to strongly inhibit c-Myc transcription while leaving AP-1 unaffected [60]. Furthermore, once the binding sites of 10058-F4, as well as that of the compound 10074-G5, within the c-Myc monomer were established using deletion and mutagenesis studies of the bHLHZip domain of c-Myc, fluorescence polarization competition affinity experiments were performed to determine the binding sites of seven other inhibitors, taking advantage of the intrinsic fluorescence of the drug-like molecules. Six of these seven compounds bound one of the binding sites already established, whereas one, 10074-A4, bound a region adjacent to the site of 10075-G5. It is notable that these three binding sites are all within a span of 85 residues, which suggests that drug-binding regions may fall within specific disordered sequences [43].

Circular dichroism spectroscopy

The case of targeting monomeric c-Myc demonstrated the feasibility of targeting a disordered protein in its monomeric state as a therapeutic strategy [25, 35, 44, 64]. Many different optical techniques contributed to the




Fig. 3 Schematic representation of a fluorescence polarization experiment. As a result of rapid tumbling of molecules in solution, when a fluorescently labelled ligand is excited with plane-polarized light, the resulting emitted light is largely depolarized (a). Upon

binding another species, a larger proportion of the emitted light remains in the same plane as the excitation energy, because the rotation is slowed as the effective molecular size increases, whether it is an ordered molecular structure (b) or one that is disordered (c)

characterization of this type of binding interaction, including in particular circular dichroism (CD) experiments (Table 1), which are based on the differential absorption of left- and right-handed circularly polarized light and can be used to determine the secondary-structure content of proteins. With this approach it was demonstrated that 10058-F4 and 10074-G5 caused an unfolding of the c-Myc/Max coiled-coil dimer into disordered monomeric states. Furthermore, CD was used to confirm the two distinct 11 and 19-residue binding regions identified for 10058-F4 and 10074-G5, respectively, by deletion and mutagenesis studies. This binding was further characterized by performing fluorescence polarization titrations, which take advantage of the intrinsic fluorescence of these compounds [35, 43]. In addition to conventional CD experiments, CD spectroscopy obtained using beamline synchrotron radiation offers improved sensitivity at a wider range of wavelengths to detect subtle changes upon complex formation [65].

Byproducts of screenings to identify enzyme inhibitors

Small molecules have also been identified to interact with the Alzheimer's-related disordered amyloid- β peptide ($A\beta$, discussed more in detail below). Serendipitously, some of these small molecules were not identified by direct screening against the peptide itself, but rather during a search for modulators of γ -secretase, which, together with β -secretase, cleaves the amyloid precursor protein (APP) to produce toxic $A\beta$. Derivatives of two modulators (tarenflurbil and fenofibrate) were created to contain a benzophenone group (a UV-active moiety used for labeling) and a biotin tag. It was thus found that these derivatives bind directly to APP within the $A\beta$ region, and act as a 'molecular clamps' or substrate-targeted inhibitors preventing the cleavage of $A\beta$ [66].

Chemical kinetics approaches to identify protein aggregation inhibitors

Under certain conditions, some disordered peptides and proteins, such as $A\beta$, α -synuclein and amylin, undergo a self-assembly process, which leads to the formation of fibrillar aggregates known as amyloid fibrils. This aggregation process is typically associated with pathological conditions such as Alzheimer's and Parkinson's diseases, and type II diabetes [15, 67–69]. Given the clinical relevance of the aggregation phenomenon, efforts have been put forward to inhibit the aggregation process from occurring, many of which have been carried out via *in vitro* assays [23, 70–77].

The kinetics of formation of these aggregates can be monitored experimentally via the use of amyloid-specific fluorescent dyes (Table 1), such as the thioflavin T (ThT). Complementary biophysical techniques to monitor this process include transmission electron microscopy (TEM), atomic force microscopy (AFM), and Fourier transform infrared spectroscopy (FTIR). Such experiments highlight the presence of three typical macroscopic phases of aggregation *in vitro*, namely, the lag phase, growth phase, and plateau phase. The molecular pathways that control this aggregation process, however, have been extremely difficult to characterize, mainly because of the challenges in establishing accurate and highly reproducible *in vitro* assays for monitoring fibril formation and in formulating an overall kinetic theory to analyse the resulting measurements. For example, as the aggregation of $A\beta$ has emerged as a key feature of the onset and progression of Alzheimer's disease [78, 79], various compounds [80–87] have been reported to interfere with the aggregation process of $A\beta$, but none of these molecules has yet found a therapeutic application because of the poor understanding of their mechanism of action.

Recently, this situation has begun to change due to advances in defining a chemical kinetics theory of aggregation [15, 88, 89]. It is now understood that the overall aggregation process is the result of complex non-linear combinations of microscopic events, including: (1) primary nucleation, in which initial aggregates form from monomeric species; (2) elongation, in which existing fibrils increase in length by monomer addition, (3) secondary nucleation, whereby the surfaces of existing aggregates catalyse the formation of new aggregates and (4) fragmentation in which existing fibrils break apart, increasing the total number of fibrils [15, 88]. The contributions of each of these microscopic events to the lag, growth, and plateau phases are highly protein and condition specific. It has thus become possible to obtain microscopic rates from macroscopic measurements, thereby revealing the mechanisms of aggregation of specific proteins and the effects of small molecules on such mechanisms [15, 88, 89].

Furthermore, reproducible protocols to measure the kinetics of $A\beta$ aggregation have also been established [88, 90–92], thus providing accurate data that could be fitted with the chemical kinetics theory. These advances have helped in elucidate the crucial mechanisms in the aggregation process of $A\beta_{42}$, the 42-residue form of $A\beta$, which forms the most toxic species associated with Alzheimer's disease. In particular, it has been found that once a critical concentration of amyloid fibrils has formed, secondary nucleation overtakes primary nucleation in becoming the major source of toxic oligomers [93]. Further developments of this chemical kinetics framework have shown that therapeutic strategies against amyloid

aggregation should not simply aim at a complete inhibition of fibril formation, but rather at specifically targeting toxic oligomeric species, as generic and non-specific effects could lead to the increase in the concentration of these oligomers and hence result in a negative outcome in terms of suppressing pathogenicity [89].

Recently, the small molecule bexarotene was discovered to target the primary nucleation step in the aggregation of A β 42. Its presence delays the formation of oligomers of A β 42 and suppresses the toxicity in neuroblastoma cells and in a *Caenorhabditis elegans* model of A β -mediated dysfunction. While this small molecule suggests that compounds may be found that act as 'neurostatins' to delay Alzheimer's disease if taken before the onset of disease, research efforts are now also focused on developing a strategy for specifically targeting secondary nucleation processes which may yield a therapeutic capable of inhibiting toxicity after the onset of symptoms [40]. As a proof-of-principle, it has been demonstrated that the molecular chaperone Brichos is able to block the formation of toxic oligomers of A β 42 by specifically inhibiting the secondary nucleation [94, 95]. To this end, kinetic analysis applied to a range of derivatives of bexarotene has been recently employed in order to evolve systematically potential inhibitors and to obtain libraries of compounds with increased anti-aggregation activity [96] (Fig. 4).

In contrast to monitoring aggregation with fibril-specific dyes, an alternative in cell high-throughput screening method for detecting A β inhibitors has been proposed which involves the expression of a fusion of A β 42 to the green fluorescent protein (GFP) in *Escherichia coli* cells. In the absence of inhibition, the aggregation of A β 42 results in a quenching of the GFP fluorescence. However, in the

presence of an aggregation inhibitor, the fluorescence of GFP is preserved, thus enabling the identification of molecules based on a triazine scaffold that inhibit A β aggregation [97].

Furthermore, in addition to small-molecule compounds, protein-like compounds capable of specifically suppressing protein aggregation have inspired new technological advances aimed to produce peptides, such as β -hairpins [98] and β -breakers [99, 100], antibodies [101], antibody fragments [102, 103], or other biomolecules, including molecular chaperones [104], to act as highly effective and specific protein aggregation inhibitors. Specifically, antibody fragments, particularly single-domain and single-chain antibodies, are becoming highly explored molecules for the inhibition of amyloid aggregation. Since the first production of conformationally distinct antibodies able to uniquely target fibrillar and oligomeric species of various amyloidogenic proteins [105], many other amyloid-specific antibodies have been generated by means of direct immunization or using hybridoma technology [101], phage display [106] or, more recently by rational design [99, 103].

In addition to directly modulating homogeneous aggregation processes, as illustrated above in the case of bexarotene for A β aggregation, small molecules have also been shown to also impact heterogeneous nucleation processes associated with aggregation. For example, the antimicrobial aminosterol, squalamine, alters the heterogeneous aggregation of α -synuclein [47]. The primary nucleation of α -synuclein is an intrinsically slow process, whose rate increases by a thousand fold as a consequence of the interaction of α -synuclein monomers with lipid membranes [107]. Squalamine has been proved to inhibit




Fig. 4 Schematic representation of a fluorescence-based kinetic aggregation assay. Aggregation assays to monitor the kinetics of formation of fibrillar aggregates are performed using a fluorescence dye molecule, in this case thioflavin T (ThT). Binding can be fitted with a kinetic model from which microscopic aggregation parameters

can be derived [88, 91, 92]. Monitoring how these microscopic parameters change in the presence of small molecules is a powerful approach for screening molecules capable of inhibiting the aggregation process [40, 89]




Fig. 5 Schematic representation of mass spectrometry with electron capture dissociation (ECD). This is a technique that enables the identification of local binding regions within disordered proteins.

the lipid-induced primary nucleation of α -synuclein by displacing monomers from the membranes [47].

In summary, as the cases of the A β and α -synuclein have shown, reproducibility of high-throughput fluorescence aggregation assays and a chemical kinetic framework underlying these complex aggregation processes have emerged as essential tools to identify molecules as modulators of these toxic aggregation processes. Furthermore, these tools enable the quantification of the effects of such therapeutics on various microscopic aggregation steps, thus creating novel opportunities in drug discovery against neurodegenerative diseases.

Methods of characterizing ligand interactions with monomeric disordered proteins

Experimental methods to characterize the binding of molecules to disordered proteins in their monomeric forms

In contrast to targeting disordered proteins in their aggregated or bound forms, it is often desirable to target them in their monomeric forms, upstream of any biological effect. Small molecule binding to a monomeric disordered protein, however, may come at a high entropic cost due to restraining a conformationally heterogeneous protein into a bound state [11]. Consequently, disordered protein interactions with small molecules are not readily amenable to the traditional ‘binding site docking’, which is generally exploited in the case of designing and optimizing small-molecule binders of structured proteins. Even some mechanisms used to describe protein–protein interactions involving at least one disordered partner tend to not be applicable because generally in these cases, since the enthalpic contributions over large interaction surface areas outweigh the entropic costs. In the case of small molecules, which lack these large surface areas, the entropic cost of restraining a disordered protein can be too high. Instead,

ECD breaks covalent backbone bonds of the disordered protein, while leaving noncovalent interactions intact, thus preserving the disordered protein–ligand interaction [141]

the currently reported interactions between small molecules and monomeric disordered proteins are relatively weaker than traditional drug–protein interactions [2], may involve multiple binding sites, and the protein may remain disordered in its bound state [108].

X-ray crystallography is the gold standard for determining small-molecule binding sites within ordered proteins for which an average conformation is well defined at the atomic level by mapping corresponding electron densities to atomic coordinates [109]. In the case of disordered proteins, however, dynamical regions generally appear as missing electron density [110–113]. Therefore, solution-state techniques that do not require crystallization, such as nuclear magnetic resonance (NMR) spectroscopy and other techniques described here (Table 1), coupled within integrative structural biology methods (Table 1) are better suited to probe disordered proteins, as they can directly characterize their conformational heterogeneity.

Small-angle X-ray scattering

Small-angle X-ray scattering (SAXS, Table 1) is a label-free biophysical technique that is particularly well suited to quantitatively analyse heterogeneous and flexible systems such as disordered proteins in solution [114]. Based on the scattering of X-rays upon exposure to a sample, it is a useful technique to quantify conformational changes upon ligand binding [115]. As previously mentioned, SAXS, in combination with FRET and NMR experiments, was employed to demonstrate the compaction of PTP1B upon binding trodusquemine, which alters the allosteric communication of the disordered C-terminal region of PTP1B and the folded catalytic domain, thereby inhibiting its phosphatase activity [56].

Surface plasmon resonance and other surface-based techniques

Surface plasmon resonance (SPR, Table 1) is a sensitive, label-free, optical method based on the detection of the

changes upon binding of the refractive index at the surface of a bio-functionalized gold-coated prism. At certain angles of incidence, electrons at the gold surface absorb some photons of the incident light, giving rise to surface plasmons. Because this phenomenon is extremely sensitive to changes in the surface of the biochip due to changes in mass, SPR is particularly sensitive for monitoring association and dissociation of biomolecules immobilized on a surface. SPR was used to screen a 3000-molecule library for small molecules able to bind EWS–FLI1, a predominantly disordered oncogenic fusion protein associated with Ewing’s sarcoma family tumours [45]. An initial hit was optimized to produce the small molecule YK-4-279, with a reported affinity of 10 μ M, which showed in vitro and in vivo inhibition of the RNA helicase A binding ability of EWS–FLI1. Like SPR, other surface-based techniques including bio-layer interferometry (BLI) [116] and quartz crystal microbalance (QCM) [117] are extremely sensitive and well suited to study disordered protein interactions with small molecules [25]. We also point out, however, that with any surface-based technique, one should carefully minimize any non-specific interaction with the sensor or the tip, or to account for them appropriately in the analysis [118].

Thermal denaturation screening

By comparing temperature-dependent denaturation patterns of proteins in the presence and absence of small molecules, one can identify potential hits, because interacting ligands may induce structural rearrangements and changes in stability (Table 1). These effects can also be monitored extrinsically using dyes, such as 8-anilino-1-naphthalene sulfonic acid (ANS), whose fluorescence increases upon binding to hydrophobic protein regions. This screening method was recently exploited to identify several binders, including trifluoperazine, of nuclear protein 1 (NUPR1) [119], which is of great therapeutic interest due to its association with pancreatic adenocarcinoma, and many other diseases [120]. However, one shortcoming of this approach is that there is no direct correlation between the stabilization effect and the affinity, making it difficult to rank hits.

Isothermal titration calorimetry

Isothermal titration calorimetry (ITC, Table 1) is an experimental technique that measures the heat exchanged during binding events between molecules in solution [121]. In this experiment, direct measurements of the absorbed or released heat are taken as one binding partner (either the protein or ligand) is titrated into a solution containing the second binding partner, offering invaluable information

that cannot be readily observed by other means. In one single experiment, one can obtain the binding constant (K_d), Gibbs free energy of binding (ΔG), enthalpy (ΔH), entropy (ΔS), and stoichiometry of the interaction. Furthermore, ITC has many advantages over other techniques; measurements can be carried out in a physiologically relevant buffer, no surface effects need to be taken into account, and the species of interest do not need to be immobilized or labelled [121]. In a standard setup, one binding partner, whose concentration is known, is titrated into a solution of the second binding partner, whose concentration is also known, while changes in the heat of the system are monitored. Over time, the protein–ligand system reaches equilibrium while the differences between heat changes diminish. Plotting the heats of the titration as a function of the molar ratio of ligand and protein inside the cell yields a curve that can be analysed with a binding model to determine the thermodynamic parameters [121].

In the case of disordered proteins, ITC can be particularly useful when a protein adopts a rigid conformation upon binding a partner, such that the contributions of enthalpy to the Gibbs free energy are significant. Such contributions can arise from the formation and breaking of noncovalent bonds, namely protein–solvent hydrogen bonds, protein–ligand bonds, van der Waals interactions, salt bridges, reorganization of atoms and solvent molecules near the binding site, and many more. ITC enabled a validation and quantitative ranking of the binders of NURP1 (introduced above in the “[Thermal denaturation screening](#)”) in terms of affinity, and suggested that this binding is largely entropically driven [119].

Single-molecule methods

Major technological advances have recently created exciting opportunities to probe disordered protein interactions with ligands at the single-molecule level. Single-molecule techniques (Table 1) are particularly promising to probe the structure and function of disordered proteins, because measurements are not ensemble-averaged as in the case of the vast majority of other available experimental techniques. Generally, two types of these experiments can be performed to elucidate the interactions of disordered proteins with binding partners: fluorescence-based techniques [122, 123] and force-probe methods [124].

Single-molecule FRET measurements are one of the several fluorescence experiments that can be performed at the single-molecule level. Similarly to the bulk-phase FRET experiments (described above), single-molecule FRET techniques require labelling with donor and acceptor dyes, but both the dyes are generally on the same protein. Experiments can either be performed on surface-immobilized samples using a total internal reflection fluorescence

(TIRF) setup or performed on freely diffusing molecules. While TIRF may enable the collection of long measurements of the fluctuations of a single molecule, interactions with the surface can perturb the native ensemble of the disordered protein. Consequently, it is more common to perform experiments on freely diffusing disordered proteins in which a laser is focused at a dilute solution (usually 50–100 pM) of labelled protein. The resulting fluorescence from both the donor and acceptor is measured and related to the distance between the two fluorophores, thereby reflecting the conformation of that molecule in the presence or absence of a ligand. Unlike bulk FRET measurements, this value is not ensemble-averaged, and many measurements enable one to construct the distribution of conformations within a given sample [125]. For example, single-molecule FRET was applied to study the conformations and dynamics of monomeric α -synuclein in the presence of sodium dodecyl sulphate (SDS) as a lipid mimetic. This technique enabled a detailed thermodynamic characterization of the multi-state conformational changes of α -synuclein folding in the presence of SDS [126].

Single-molecule force-probe microscopy also offers intriguing complementary approaches to the single-molecule fluorescence-based methods. These techniques involve the use of optical tweezers, magnetic tweezers, or atomic force microscopy by which the ends of individual protein molecules are constrained in order to apply and measure forces which yield information about their extensions and resulting conformational transitions [127]. This type of approach has been widely employed for characterizing the conformational and dynamic behaviour of disordered proteins, including α -synuclein [128, 129] and A β [130]. Furthermore, these techniques have characterized disordered and unfolded proteins in the presence of binding partners, including molecular chaperones [131] and ions [132].

Mass spectrometry methods

The development of soft ionization methods, such as electrospray ionization (ESI) and matrix-assisted laser desorption/ionization (MALDI), has facilitated the application of mass spectrometry (MS) to protein characterization and protein binding, offering insight on stoichiometry, reversibility, specificity, and binding affinities [133, 134]. Furthermore, MS is a particularly powerful probe of protein behaviour due to its ability to monitor discrete conformers in a mixture [135–137]. While there are a number of MS-based techniques to probe protein behaviour, ranging from those that monitor how variance in buffer conditions affect the distribution of charge states into the gas phase [138, 139], to those that trap protein ions in the gas phase and observe conformational changes on a μ s–s timescale [140], here we highlight

one particular advance that has enabled the localization of a ligand binding site within a disordered protein. By combining ESI-MS with electron capture dissociation (EDC), a technique to fragment gas-phase ions (Fig. 5), the polycationic compound spermine was found to bind α -synuclein in the region of residues 106–138. It was shown that EDC breaks certain covalent backbone bonds of α -synuclein, while leaving noncovalent interactions intact, thus preserving the spermine– α -synuclein interaction [141]. This technique is highly complementary to nuclear magnetic resonance spectroscopy (see below) and other biophysical measurements, although it should be noted that the parameters observed in the gas phase may differ from those in solution, as the hydrophobic effect is essentially lost in the gas phase, while electrostatic interactions are strengthened due to the fact that the dielectric constant is lower than water [142, 143]. We also mention that MS methods have been applied to identify inhibitors of aggregation (introduced above) [134, 144].

Nuclear magnetic resonance spectroscopy

Nuclear magnetic resonance spectroscopy (NMR, Table 1) can be employed in two complementary ways to monitor the binding between a disordered protein and a small molecule. Changes in the one-dimensional hydrogen spectrum of the ligand in the presence of a disordered protein offer a fast and sensitive indication of binding, but offers little insight regarding the binding site and mode of interaction. In addition, monitoring the protein (which usually requires ^{15}N or ^{13}C isotopic labelling) is a powerful method that can yield informative structural and dynamical binding information about disordered proteins, as a result of a systematic series of advances within the past decade [145–148]. In particular, the sensitivity of the latter technique offers highly quantitative insights into how the properties of disordered proteins change in the presence of small molecules. Quite generally, in NMR structural information is derived by exploiting the conformational dependence of the transitions between different energy levels of atomic nuclear spins, which can be made to split in an external magnetic field and resonate using electromagnetic radiation. While, in contrast with structured proteins, nuclear Overhauser effects (NOEs) [148] cannot always be readily exploited to obtain inter-proton distances for disordered proteins due to their conformational heterogeneity, other NMR parameters, including chemical shifts, hydrogen exchange rates, residual dipolar couplings (RDCs) and paramagnetic relaxation enhancements (PREs), can provide atomic-resolution structural information [145, 147, 149–151].

In contrast to the high-resolution assignments for globular proteins, which can be obtained using triple resonance

coherence transfer experiments on isotopically labelled proteins, equivalent measurements of disordered proteins often yield overlapping peaks within collapsed spectra. This is a result of a combination of structural disorder and solvent exposure, which creates similar environments for many residues. This problem is often worsened by the low sequence complexity found within disordered proteins [145, 147, 152, 153], especially as they are enriched in proline residues, which are invisible to hydrogen-detected NMR spectra [154, 155]. Furthermore, such high solvent exposure also contributes to decreasing the signal-to-noise ratios for disordered proteins, as significant chemical exchange with bulk solvent reduces the intensities of amide hydrogen signals. While signal overlap of disordered proteins can be partially ameliorated by sample preparation at low pH and by taking measurements at low temperatures, the largest improvements have been a result of technological advances. Such advances include increased instrumental sensitivity, faster sampling rates exploiting longitudinal relaxation enhancements [156] and the use of non-uniform sampling for high-dimensionality experiments [145, 152, 153, 157, 158]. Additionally, by replacing hydrogen detection with carbon detection and by exploiting cryoprobe technology, it is possible to separate peaks accurately, while remaining insensitive to broadening and salt concentrations [147, 152, 153, 159]. Despite their poor spectral resolution, disordered proteins produce particularly sharp peaks, making them ideal for relaxation experiments, and as such, additional improvements include relaxation-optimized detection schemes [145, 160]. Furthermore, the structural properties of the aggregates formed by some disordered proteins can be studied by other NMR techniques such as solid-state magic-angle spinning which is discussed in detail elsewhere [161, 162].

Among the most useful NMR parameters for characterizing disordered proteins that we discuss here are chemical shifts, which report on the population-weighted average across the conformations sampled within a millisecond time scale. By calculating deviations from random coil values, one can describe the local geometry and quantify local secondary structure propensity in disordered proteins [163–165] and quantify changes in the absence and presence of therapeutic molecules. Furthermore, 2D NMR ‘fingerprint’ spectra, most commonly obtained with either ^1H detected ^1H – ^{15}N (HN) [166] and ^{13}C detected ^{13}C – ^{15}N (CON) [167] for disordered proteins, are simple indicators of monomeric binding, and provide highly detailed site-specific information if protein assignments have already been established. While the HN experiment has higher sensitivity and requires less time to record, the CON experiment displays better spectral resolution, can detect proline residues, and is not prone to hydrogen-exchange-induced line broadening, and thus spectra can be




Fig. 6 Schematic illustration of the chemical shift perturbation mapping method. By identifying and quantifying changes in two-dimensional spectra (in this case ^1H – ^{15}N HSQC) in the absence (*red*) and presence (*blue*) of ligands, chemical shift perturbation mapping is a powerful technique to identify whether ligands interact with disordered proteins, and identify binding sites or locations of conformational change

recorded at higher pH and temperatures [159]. Chemical shift perturbations are a sensitive technique that can simultaneously provide binding affinity (K_d) values and insight about a binding site or the location of conformational changes induced upon binding (Fig. 6). Finally, within integrative methods (see below), chemical shifts can be used to determine structural ensembles of disordered proteins [168].

Two-dimensional (2D) ^1H – ^{15}N heteronuclear single quantum coherence spectroscopy (HSQC) experiments were used to confirm the binding of heparin to the intrinsically disordered osteopontin [169], an extracellular structural protein associated with many pathological conditions, including autoimmune diseases [170], cancer metastasis [171], Crohn’s disease and ulcerative colitis [172], allergy and asthma [173], and muscle disease [174]. Chemical shift differences only at certain residues between the free and bound forms of osteopontin suggested a specific interaction, and enabled mapping of the binding site [169]. Similarly, 2D ^1H – ^{15}N HSQC experiments were used to characterize the specific binding of hits from ‘fragment-like’ small-molecule hits against p27, a disordered cell cycle regulator protein. These hits were identified from 1D ^1H WaterLOGSY [175] and standard transfer difference (STD) [176] NMR screening methods, and one molecule in particular, was shown to inhibit the Cdk2/cyclin A binding function of p27 by fluorescence anisotropy and 2D ^1H – ^{15}N TROSY [41]. A similar

Fig. 7 Schematic representation of integrative methods for protein ensemble generation. Integrative (or hybrid) methods, such as metainference [191, 199], combine the strengths of experimental techniques and computational methods to overcome the challenges associated with each technique alone [6]





Fig. 8 Summary of approaches for modulating the behaviour of disordered proteins using small molecules. Small molecules can be used to: **a** disrupt functional interactions, **b** modify the properties of

approach based on 2D ^1H - ^{15}N TROSY [160] measurements was used to characterize the binding site of trodusquemine to the disordered C-terminal region of PTP1B [56]. Modifications to the HN and CON spectra enable the detection of other observables including RDCs, PREs, cross-relaxation and cross-correlation rates, in addition to solvent exchange rates. All these observables describe the structure and dynamics of disordered proteins at atomic resolution and are sensitive to changes in the presence of small molecules.

As mentioned above, RDCs are additional sensitive NMR observables that are particularly well suited to study disordered proteins in their monomeric states. These

native states, or **c** inhibit aberrant aggregation. Modifying the properties of monomeric disordered proteins (**b**) has the potential to also inhibit (**a**, **c**)

observables arise when disordered protein samples are partially aligned in a magnetic field by preparing samples in anisotropic media, for example, in a liquid crystal [177], polyacrylamide gels [178], filamentous phages [179], or bicelles [180]. As a result of restricted overall reorientation in the presence of the anisotropic media and dynamic conformational averaging, non-zero RDCs are observed which reflect the weighted average conformation of the ensemble [180]. Additionally, chemically modifying the disordered protein of interest with covalently attached paramagnetic spin labels, one can observe PREs, which report on tertiary structure, and the distances and orientations with respect to the principal axes frame of the

paramagnetic centre. As for chemical shifts, RDCs and PREs can be implemented as structural restraints for ensemble generation [181], which is discussed in the next section.

Integrative methods to characterize the effects of small molecules on protein ensembles

It is becoming increasingly clear that disordered proteins often bind ligands in transient and delocalized manners, in which the disordered protein remains in a disordered state upon association [32–34, 108]. In this context, high-resolution characterizations of conformational ensembles of disordered proteins, and of the ways in which such ensembles change in the presence of therapeutic molecules, have the potential to yield both functional mechanistic details and insights towards drug optimization.

Unfortunately, however, such detailed descriptions are currently difficult to obtain because the dynamic nature of disordered proteins makes it challenging to acquire accurate experimental measurements, as well as to interpret them in terms of structural models [5]. For example, as noted above, while NMR spectroscopy and other solution-state methods can provide valuable information on structural ensembles, these techniques alone are insufficient to provide all the conformational restraints needed to fully characterize the conformations within such ensembles. This is because experimental techniques, in addition to being inevitably affected by systematic and random errors, often measure sparse and sometimes ambiguous time- and ensemble-averages over the many heterogeneous conformations of the disordered proteins [6, 182].

To overcome these problems, computational techniques such as molecular dynamics simulations can provide accurate descriptions of protein ensembles [6]. In these simulations, the conformational space of a protein is sampled via the integration of the equations of motion over a sufficiently long time interval to ensure the exploration of the most relevant states and corresponding estimates of their populations. Such approaches have been used to investigate many small-molecule interactions with disordered proteins, particularly amyloidogenic ones [41, 83, 183–185] in addition to identifying potential binding pockets within disordered monomers [186]. Unfortunately, however, despite continuous advances, the force fields used to represent the interatomic forces needed to solve the equations of motion are still approximate [187–189], which leads to the need of validating the results through the comparison with experimental data [184, 186].

We should also remark that as most proteins of interest are large macromolecules in a complex environment, they are at the limit of what can be simulated. Conformational sampling as a result of limited computational resources is

in fact often a major issue. While this problem can be partially alleviated through the use of enhanced sampling techniques [190, 191], the resulting ensembles may still be dependent on the simulation time, which is an approximation that requires careful control.

We believe that an effective way forward is to bring together the advantages of the experimental and computational approaches (Fig. 7). A series of recent results indicate that by combining sparse experimental data on disordered proteins with a priori information from force fields [192–195] in molecular dynamics simulations, it is possible to generate descriptions of conformational ensembles with corresponding equilibrium probabilities for each state, such that the ensembles are consistent with the overall theoretical understanding of disordered proteins and with the available experimental data. There are many available methods for integrative structural ensemble determination of heterogeneous systems [6]. Because of the limited space that we have here, however, we only briefly highlight recent advances which enable one both to incorporate experimental data directly as structural restraints and to account for systematic and random errors by employing Bayesian inference techniques. These include ‘multi-state Bayesian modelling [196, 197], the ‘Bayesian ensemble refinement [198], and ‘metadynamic meta-inference’ [191, 199].

Integrative structural biology methods were used to characterize the binding interactions between binding sites within c-Myc and small molecules. Metadynamics simulations using NMR chemical shift data [35] as restraints were employed to show that these interactions are highly delocalized, the binding sites remain disordered, and the conformational space of the binding regions are slightly altered [28, 34].

In summary, integrative computational methods for determining ensembles of disordered proteins that incorporate experimental measurements and account for different sources of error represent a powerfully detailed and increasingly accurate approach to study the behaviour of ensembles in the presence of candidate therapeutic molecules.

Conclusions and outlook

We have discussed three strategies to use small molecules to modify the behaviour of disordered proteins (Fig. 8). We have begun with the strategy of modulating the functional interactions involving disordered proteins using small-molecule inhibitors. We then reviewed recent advances in using chemical kinetics to identify compounds capable of blocking the aggregation of disordered proteins, and finally discussed methods of finding small molecules capable of binding disordered proteins, emphasizing the importance of

more closely integrating experimental and computational techniques. Overall, we believe that upon further developments, the methods that we have reviewed will lead to a progressive ability to identify compounds of therapeutic interest for disordered proteins. We anticipate that an area of research of crucial importance will be to understand the role of specificity in these interactions, which will likely require the development of new assays, as well as possibly innovative conceptual tools.

Acknowledgements Gabriella T. Heller is supported by the Gates Cambridge Trust Scholarship. Francesco A. Aprile is supported by a Senior Research Fellowship award from the Alzheimer's Society, UK (Grant Number 317, AS-SF-16-003).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Habchi J, Tompa P, Longhi S, Uversky VN (2014) Introducing protein intrinsic disorder. *Chem Rev* 114:6561–6588
- Csizmok V, Follis AV, Kriwacki RW, Forman-Kay JD (2016) Dynamic protein interaction networks and new structural paradigms in signaling. *Chem Rev* 116:6424–6462
- Tompa P (2012) Intrinsically disordered proteins: a 10-year recap. *Trends Biochem Sci* 37:509–516
- Bhowmick A et al (2016) Finding our way in the dark proteome. *J Am Chem Soc* 138:9730–9742
- Sormanni P et al (2017) Simultaneous quantification of order and disorder in proteins. *Nat Chem Biol* 13:339–342
- Bonomi M, Heller GT, Camilloni C, Vendruscolo M (2017) Principles of protein structural ensemble determination. *Curr Opin Struct Biol* 42:106–116
- Wright PE, Dyson HJ (2015) Intrinsically disordered proteins in cellular signalling and regulation. *Nat Rev Mol Cell Biol* 16:18–29
- Uversky VN (2016) P53 proteoforms and intrinsic disorder: an illustration of the protein structure–function continuum concept. *Int J Mol Sci* 17:1874
- Ward JJ, Sodhi JS, McGuffin LJ, Buxton BF, Jones DT (2004) Prediction and functional analysis of native disorder in proteins from the three kingdoms of life. *J Mol Biol* 337:635–645
- Babu MM, van der Lee R, de Groot NS, Gsponer J (2011) Intrinsically disordered proteins: regulation and disease. *Curr Opin Struct Biol* 21:432–440
- Heller GT, Sormanni P, Vendruscolo M (2015) Targeting disordered proteins with small molecules using entropy. *Trends Biochem Sci* 40:491–496
- Uversky VN, Oldfield CJ, Dunker AK (2008) Intrinsically disordered proteins in human diseases: introducing the d2 concept. *Annu Rev Biophys* 37:215–246
- Tompa P (2009) Structure and function of intrinsically disordered proteins. Chapman & Hall/CRC Press, Boca Raton, p 331
- Uversky VN et al (2014) Pathological unfoldomics of uncontrolled chaos: intrinsically disordered proteins and human diseases. *Chem Rev* 114:6844–6879
- Knowles TPJ, Vendruscolo M, Dobson CM (2014) The amyloid state and its association with protein misfolding diseases. *Nat Rev Mol Cell Biol* 15:384–396
- Krüger R et al (1998) Ala30pro mutation in the gene encoding α -synuclein in Parkinson's disease. *Nat Gen* 18:106–108
- Zarranz JJ et al (2004) The new mutation, e46k, of α -synuclein causes Parkinson and Lewy body dementia. *Ann Neurol* 55:164–173
- Polymeropoulos MH et al (1997) Mutation in the α -synuclein gene identified in families with Parkinson's disease. *Science* 276:2045–2047
- Singleton AB et al (2003) α -Synuclein locus triplication causes Parkinson's disease. *Science* 302:841
- Chartier-Harlin MC et al (2004) α -Synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet* 364:1167–1169
- Ibáñez P et al (2004) Causal relation between α -synuclein gene duplication and familial Parkinson's disease. *Lancet* 364:1169–1171
- Cookson MR (2009) α -Synuclein and neuronal cell death. *Mol Neurodegener* 4:9
- Cuchillo R, Michel J (2012) Mechanisms of small-molecule binding to intrinsically disordered proteins. *Biochem Soc Trans* 40:1004–1008
- Joshi P, Vendruscolo M (2015) Druggability of intrinsically disordered proteins. *Adv Exp Med Biol* 870:383–400
- Metallo SJ (2010) Intrinsically disordered proteins are potential drug targets. *Curr Opin Chem Biol* 14:481–488
- Dunker AK, Uversky VN (2010) Drugs for 'protein clouds': targeting intrinsically disordered transcription factors. *Curr Opin Pharmacol* 10:782–788
- Cheng Y et al (2006) Rational drug design via intrinsically disordered protein. *Trends Biotechnol* 24:435–442
- Jin F, Yu C, Lai L, Liu Z (2013) Ligand clouds around protein clouds: a scenario of ligand binding with intrinsically disordered proteins. *PLoS Comput Biol* 9:e1003249
- Uversky VN (2012) Intrinsically disordered proteins and novel strategies for drug discovery. *Expert Opin Drug Discov* 7:475–488
- Shangary S, Wang S (2009) Small-molecule inhibitors of the mdm2–p53 protein–protein interaction to reactivate p53 function: a novel approach for cancer therapy. *Annu Rev Pharmacol Toxicol* 49:223–241
- Vassilev LT (2004) Small-molecule antagonists of p53–mdm2 binding: research tools and potential therapeutics. *Cell Cycle* 3:419–421
- Tompa P, Fuxreiter M (2008) Fuzzy complexes: polymorphism and structural disorder in protein–protein interactions. *Trends Biochem Sci* 33:2–8
- Fuxreiter M, Tompa P (2012) Fuzzy complexes: a more stochastic view of protein function. *Adv Exp Med Biol* 725:1–14
- Michel J, Cuchillo R (2012) The impact of small molecule binding on the energy landscape of the intrinsically disordered protein c-myc. *PLoS One* 7:e41070
- Follis AV, Hammoudeh DI, Wang H, Prochownik EV, Metallo SJ (2008) Structural rationale for the coupled binding and unfolding of the c-myc oncoprotein by small molecules. *Chem Biol* 15:1149–1155
- Dogan J, Gianni S, Jemth P (2014) The binding mechanisms of intrinsically disordered proteins. *Phys Chem Chem Phys* 16:6323–6331
- Dyson HJ, Wright PE (2002) Coupling of folding and binding for unstructured proteins. *Curr Opin Struct Biol* 12:54–60
- Toto A et al (2016) Molecular recognition by templated folding of an intrinsically disordered protein. *Sci Rep* 6:21994

39. Tóth G et al (2014) Targeting the intrinsically disordered structural ensemble of α -synuclein by small molecules as a potential therapeutic strategy for Parkinson's disease. *PLoS One* 9:e87133
40. Habchi J et al (2016) An anticancer drug suppresses the primary nucleation reaction that initiates the production of the toxic $a\beta_{42}$ aggregates linked with Alzheimer's disease. *Sci Adv* 2:e1501244–e1501244
41. Iconaru LI et al (2015) Discovery of small molecules that inhibit the disordered protein, p27kip1. *Sci Rep* 5:15686
42. Wang H et al (2007) Improved low molecular weight myc–max inhibitors. *Mol Cancer Ther* 6:2399–2408
43. Hammoudeh DI, Follis AV, Prochownik EV, Metallo SJ (2009) Multiple independent binding sites for small-molecule inhibitors on the oncoprotein c-myc. *J Am Chem Soc* 131:7390–7401
44. Yin X, Giap C, Lazo JS, Prochownik EV (2003) Low molecular weight inhibitors of myc–max interaction and function. *Oncogene* 22:6151–6159
45. Erkizan HV et al (2009) A small molecule blocking oncogenic protein ews–fli1 interaction with rna helicase a inhibits growth of Ewing's sarcoma. *Nat Med* 15:750–756
46. Follis AV, Hammoudeh DI, Daab AT, Metallo SJ (2009) Small-molecule perturbation of competing interactions between c-myc and max. *Bioorg Med Chem Lett* 19:807–810
47. Perni M et al (2017) A natural product inhibits the initiation of α -synuclein aggregation by displacing it from lipid membranes and suppresses its toxicity. *Proc Natl Acad Sci USA* 114:E1009–E1017
48. Iakoucheva LM, Brown CJ, Lawson JD, Obradović Z, Dunker AK (2002) Intrinsic disorder in cell-signaling and cancer-associated proteins. *J Mol Biol* 323:573–584
49. Lebel R, McDuff FO, Lavigne P, Grandbois M (2007) Direct visualization of the binding of c-myc/max heterodimeric b-hlhlz to e-box sequences on the htert promoter. *Biochemistry* 46:10279–10286
50. Nair SK, Burley SK (2003) X-ray structures of myc–max and mad–max recognizing DNA: molecular bases of regulation by proto-oncogenic transcription factors. *Cell* 112:193–205
51. Berg T et al (2002) Small-molecule antagonists of myc/max dimerization inhibit myc-induced transformation of chicken embryo fibroblasts. *Proc Natl Acad Sci USA* 99:3830–3835
52. Shi J, Stover JS, Whitby LR, Vogt PK, Boger DL (2009) Small molecule inhibitors of myc/max dimerization and myc-induced cell transformation. *Bioorg Med Chem Lett* 19:6038–6041
53. Xu Y et al (2006) A credit-card library approach for disrupting protein–protein interactions. *Bioorg Med Chem* 14:2660–2673
54. Hovius R, Vallotton P, Wohland T, Vogel H (2000) Fluorescence techniques: shedding light on ligand–receptor interactions. *Trends Pharmacol Sci* 21:266–273
55. Fried MG (1989) Measurement of protein–DNA interaction parameters by electrophoresis mobility shift assay. *Electrophoresis* 10:366–376
56. Krishnan N et al (2014) Targeting the disordered c terminus of ptp1b with an allosteric inhibitor. *Nat Chem Biol* 10:558–566
57. Mustata G et al (2009) Discovery of novel myc–max heterodimer disruptors with a three-dimensional pharmacophore model. *J Med Chem* 52:1247–1250
58. Grote D, Hänel F, Dahse HM, Seifert K (2008) Capnellenes from the soft coral dendronephthya rubeola. *Chem Biodiver* 5:1683–1693
59. Kiessling A, Sperl B, Hollis A, Eick D, Berg T (2006) Selective inhibition of c-myc/max dimerization and DNA binding by small molecules. *Chem Biol* 13:745–751
60. Kiessling A, Wiesinger R, Sperl B, Berg T (2007) Selective inhibition of c-myc/max dimerization by a pyrazolo [1, 5-a] pyrimidine. *ChemMedChem* 2:627–630
61. Nasir MS, Jolley ME (1999) Fluorescence polarization: an analytical tool for immunoassay and drug discovery. *Comb Chem High Throughput Screen* 2:177–190
62. Burke TJ, Loniello KR, Ja Beebe, Ervin KM (2003) Development and application of fluorescence polarization assays in drug discovery. *Comb Chem High Throughput Screen* 6:183–194
63. Lea WA, Simeonov A (2011) Fluorescence polarization assays in small molecule screening. *Expert Opin Drug Discov* 6:17–32
64. Jeong K-C, Ahn K-O, Yang C-H (2010) Small-molecule inhibitors of c-myc transcriptional factor suppress proliferation and induce apoptosis of promyelocytic leukemia cell via cell cycle arrest. *Mol Biosyst* 6:1503–1509
65. Cowieson NP et al (2008) Evaluating protein: protein complex formation using synchrotron radiation circular dichroism spectroscopy. *Proteins* 70:1142–1146
66. Kukar TL et al (2008) Substrate-targeting γ -secretase modulators. *Nature* 453:925–929
67. Uversky VN, Eliezer D (2009) Biophysics of Parkinson's disease: structure and aggregation of α -synuclein. *Curr Protein Pept Sci* 10:483–499
68. Ma Q-L, Chan P, Yoshii M, Uéda K (2003) A-synuclein aggregation and neurodegenerative diseases. *J Alzheimer's Dis* 5:139–148
69. Dobson CM (2004) Principles of protein folding, misfolding and aggregation. *Semin Cell Dev Biol* 15:3–16
70. Pickhardt M et al (2005) Anthraquinones inhibit tau aggregation and dissolve Alzheimer's paired helical filaments in vitro and in cells. *J Biol Chem* 280:3628–3635
71. Cohen T, Frydman-Marom A, Rechter M, Gazit E (2006) Inhibition of amyloid fibril formation and cytotoxicity by hydroxyindole derivatives. *Biochemistry* 45:4727–4735
72. Carter MD, Ga Simms, Weaver DF (2010) The development of new therapeutics for Alzheimer's disease. *Clin Pharmacol Ther* 88:475–486
73. McLaurin JA, Golomb R, Jurewicz A, Antel JP, Fraser PE (2000) Inositol stereoisomers stabilize an oligomeric aggregate of alzheimer amyloid β peptide and inhibit A β -induced toxicity. *J Biol Chem* 275:18495–18502
74. Conway KA et al (2001) Kinetic stabilization of the α -synuclein protofibril by a dopamine– α -synuclein adduct. *Science* 294:1346–1349
75. Li J, Zhu M, Manning-Bog AB, Da Di Monte, Fink AL (2004) Dopamine and l-dopa disaggregate amyloid fibrils: implications for Parkinson's and Alzheimer's disease. *FASEB J* 18:962–964
76. Harrington C et al (2008) O1-06-04: methylthioninium chloride (mtc) acts as a tau aggregation inhibitor (TAI) in a cellular model and reverses Tau pathology in transgenic mouse models of Alzheimer's disease. *Alzheimer's Dement* 4:T120–T121
77. Bulic B, Pickhardt M, Mandelkow EM, Mandelkow E (2010) Tau protein and tau aggregation inhibitors. *Neuropharmacology* 59:276–289
78. Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297:353–356
79. Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid β -peptide. *Nat Rev Mol Cell Biol* 8:101–112
80. Shimmyo Y, Kihara T, Akaike A, Niidome T, Sugimoto H (2008) Multifunction of myricetin on A β : neuroprotection via a conformational change of A β and reduction of A β via the interference of secretases. *J Neurosci Res* 86:368–377
81. Zheng X et al (2012) Z-phe-ala-diazomethylketone (padk) disrupts and remodels early oligomer states of the Alzheimer disease $a\beta_{42}$ protein. *J Biol Chem* 287:6084–6088

82. Ge JF, Qiao JP, Qi CC, Wang CW, Zhou JN (2012) The binding of resveratrol to monomer and fibril amyloid β . *Neurochem Int* 61:1192–1201
83. Li G, Rauscher S, Baud S, Pomès R (2012) Binding of inositol stereoisomers to model amyloidogenic peptides. *J Phys Chem B* 116:1111–1119
84. Connors CR et al (2013) Tranilast binds to ab monomers and promotes ab fibrillation. *Biochemistry* 52:3995–4002
85. Groenning M (2010) Binding mode of thioflavin t and other molecular probes in the context of amyloid fibrils—current status. *J Chem Biol* 3:1–18
86. Lendel C, Bolognesi B, Wahlström A, Dobson CM, Gräslund A (2010) Detergent-like interaction of congo red with the amyloid β peptide. *Biochemistry* 49:1358–1360
87. Caesar I, Jonson M, Nilsson KPR, Thor S, Hammarström P (2012) Curcumin promotes A β fibrillation and reduces neurotoxicity in transgenic drosophila. *PLoS One* 7:e31424
88. Knowles TPJ et al (2009) An analytical solution to the kinetics of breakable filament assembly. *Science* 326:1533–1537
89. Arosio P, Vendruscolo M, Dobson CM, Knowles TPJ (2014) Chemical kinetics for drug discovery to combat protein aggregation diseases. *Trends Pharmacol Sci* 35:127–135
90. Hellstrand E, Boland B, Walsh DM, Linse S (2010) Amyloid β -protein aggregation produces highly reproducible kinetic data and occurs by a two-phase process. *ACS Chem Neurosci* 1:13–18
91. Meisl G et al (2016) Molecular mechanisms of protein aggregation from global fitting of kinetic models. *Nat Protoc* 11:252–272
92. Cohen SIA, Vendruscolo M, Dobson CM, Knowles TPJ (2012) From macroscopic measurements to microscopic mechanisms of protein aggregation. *J Mol Biol* 421:160–171
93. Cohen SIA et al (2013) Proliferation of amyloid- β 42 aggregates occurs through a secondary nucleation mechanism. *Proc Natl Acad Sci USA* 110:9758–9763
94. Arosio P et al (2016) Kinetic analysis reveals the diversity of microscopic mechanisms through which molecular chaperones suppress amyloid formation. *Nat Commun* 7:10948
95. Cohen SIA et al (2015) A molecular chaperone breaks the catalytic cycle that generates toxic A β oligomers. *Nat Struct Mol Biol* 22:207–213
96. Habchi J et al (2017) Systematic development of small molecules to inhibit specific microscopic steps of a β 42 aggregation in Alzheimer's disease. *Proc Natl Acad Sci USA* 114:E200–E208
97. Kim W et al (2006) A high-throughput screen for compounds that inhibit aggregation of the Alzheimer's peptide. *ACS Chem Biol* 1(7):461–469. doi:10.1021/cb600135w
98. Mirecka EA et al (2016) β -hairpin of islet amyloid polypeptide bound to an aggregation inhibitor. *Sci Rep* 6:33474. doi:10.1038/srep33474
99. Sievers SA et al (2011) Structure-based design of non-natural amino-acid inhibitors of amyloid fibril formation. *Nature* 475:96–100
100. Gazit E (2005) Mechanisms of amyloid fibril self-assembly and inhibition: model short peptides as a key research tool. *FEBS J* 272:5971–5978
101. Sevigny J et al (2016) The antibody aducanumab reduces A β plaques in Alzheimer's disease. *Nature* 537:50–56
102. Perchiacca JM, Ladiwala ARA, Bhattacharya M, Tessier PM (2012) Structure-based design of conformation- and sequence-specific antibodies against amyloid β . *Proc Natl Acad Sci USA* 109:84–89
103. Sormanni P, Aprile FA, Vendruscolo M (2015) Rational design of antibodies targeting specific epitopes within intrinsically disordered proteins. *Proc Natl Acad Sci USA* 112:9902–9907
104. Aprile FA, Sormanni P, Vendruscolo M (2015) A rational design strategy for the selective activity enhancement of a molecular chaperone toward a target substrate. *Biochemistry* 54:5103–5112
105. Necula M, Kaye R, Milton S, Glabe CG (2007) Small molecule inhibitors of aggregation indicate that amyloid β oligomerization and fibrillization pathways are independent and distinct. *J Biol Chem* 282:10311–10324
106. Lee CC et al (2016) Design and optimization of anti-amyloid domain antibodies specific for β -amyloid and islet amyloid polypeptide. *J Biol Chem* 291:2858–2873
107. Galvagnion C et al (2015) Lipid vesicles trigger α -synuclein aggregation by stimulating primary nucleation. *Nat Chem Biol* 11:229–234
108. Fuxreiter M (2012) Fuzziness: linking regulation to protein dynamics. *Mol Biosyst* 8:168
109. Blundell TL, Johnson LN (1976) Protein crystallography. Academic Press, New York
110. Bode W, Fehlhämmer H, Huber R (1976) Crystal structure of bovine trypsinogen at 1.8 Å resolution. I. Data collection, application of Patterson search techniques and preliminary structural interpretation. *J Mol Biol* 106:325–335
111. Bloomer AC, Champness JN, Bricogne G, Staden R, Klug A (1978) Protein disk of tobacco mosaic virus at 2.8 Å resolution showing the interactions within and between subunits. *Nature* 276:362–368
112. Ringe D, Petsko GA (1985) Mapping protein dynamics by X-ray diffraction. *Prog Biophys Mol Biol* 45:197–235
113. Dunker AK, Oldfield CJ (2015) Back to the future: nuclear magnetic resonance and bioinformatics studies on intrinsically disordered proteins. *Adv Exp Med Biol* 870:1–34
114. Bernadó P, Mylonas E, Petoukhov MV, Blackledge M, Svergun DI (2007) Structural characterization of flexible proteins using small-angle X-ray scattering. *J Am Chem Soc* 129:5656–5664
115. Bernadó P, Svergun DI (2012) Structural analysis of intrinsically disordered proteins by small-angle X-ray scattering. *Mol Biosyst* 8:151–167
116. Concepcion J et al (2009) Label-free detection of biomolecular interactions using bilayer interferometry for kinetic characterization. *Comb Chem High Throughput Screen* 12:791–800
117. Heller GT, Mercer-Smith AR, Johal MS (2015) Quartz microbalance technology for probing biomolecular interactions. Protein–protein interactions: methods and applications, 2nd edn. Springer, New York, NY, pp 153–164
118. Heller GT et al (2014) Accounting for unintended binding events in the analysis of quartz crystal microbalance kinetic data. *Colloids Surf B* 117:425–431
119. Neira JL et al (2017) Identification of a drug targeting an intrinsically disordered protein involved in pancreatic adenocarcinoma. *Sci Rep* 7:39732
120. Chowdhury UR, Samant RS, Fodstad O, La Shevde (2009) Emerging role of nuclear protein 1 (nupr1) in cancer biology. *Cancer Metastasis Rev* 28:225–232
121. Bronowska AK (2011) Thermodynamics of ligand–protein interactions: implications for molecular design. INTECH Open Access Publisher
122. Schuler B, Eaton WA (2008) Protein folding studied by single-molecule fret. *Curr Opin Struct Biol* 18:16–26
123. Cremades N et al (2012) Direct observation of the interconversion of normal and toxic forms of α -synuclein. *Cell* 149:1048–1059
124. Forman JR, Clarke J (2007) Mechanical unfolding of proteins: insights into biology, structure and folding. *Curr Opin Struct Biol* 17:58–66
125. Ferreón ACM, Moran CR, Gambin Y, Deniz AA (2010) Single-molecule fluorescence studies of intrinsically disordered proteins. *Methods Enzymol* 472:179–204

126. Ferreon ACM, Deniz AA (2007) α -Synuclein multistate folding thermodynamics: implications for protein misfolding and aggregation. *Biochemistry* 46:4499–4509
127. Neuman KC, Nagy A (2008) Single-molecule force spectroscopy: optical tweezers, magnetic tweezers and atomic force microscopy. *Nat Methods* 5:491
128. Sandal M et al (2008) Conformational equilibria in monomeric α -synuclein at the single-molecule level. *PLoS Biol* 6:e6
129. Solanki A, Neupane K, Woodside MT (2014) Single-molecule force spectroscopy of rapidly fluctuating, marginally stable structures in the intrinsically disordered protein α -synuclein. *Phys Rev Lett* 112:158103
130. Lv Z, Roychoudhuri R, Condran MM, Teplow DB, Lyubchenko YL (2013) Mechanism of amyloid β -protein dimerization determined using single-molecule afm force spectroscopy. *Sci Rep* 3:2880
131. Mashaghi A, Gn Kramer, Lamb DC, Mayer MP, Tans SJ (2013) Chaperone action at the single-molecule level. *Chem Rev* 114:660–676
132. Hane FT, Hayes R, Lee BY, Leonenko Z (2016) Effect of copper and zinc on the single molecule self-affinity of Alzheimer's amyloid- β peptides. *PLoS One* 11:e0147488
133. Liko I, Allison TM, Hopper JT, Robinson CV (2016) Mass spectrometry guided structural biology. *Curr Opin Struct Biol* 40:136–144
134. Young LM et al (2015) Screening and classifying small-molecule inhibitors of amyloid formation using ion mobility spectrometry–mass spectrometry. *Nat Chem* 7:73–81
135. Uetrecht C, Rose RJ, van Duijn E, Lorenzen K, Heck AJ (2010) Ion mobility mass spectrometry of proteins and protein assemblies. *Chem Soc Rev* 39:1633–1655
136. Schwartz BL et al (1995) Dissociation of tetrameric ions of noncovalent streptavidin complexes formed by electrospray ionization. *J Am Soc Mass Spectrom* 6:459–465
137. Cohen SL, Chait BT, Ferré-D'Amaré AR, Burley SK (1995) Probing the solution structure of the DNA-binding protein max by a combination of proteolysis and mass spectrometry. *Protein Sci* 4:1088–1099
138. Eyles SJ, Speir JP, Kruppa GH, Gierasch LM, Kaltashov IA (2000) Protein conformational stability probed by Fourier transform ion cyclotron resonance mass spectrometry. *J Am Chem Soc* 122:495–500
139. Konermann L, Douglas D (1998) Equilibrium unfolding of proteins monitored by electrospray ionization mass spectrometry: distinguishing two-state from multi-state transitions. *Rapid Commun Mass Spectrom* 12:435–442
140. Freitas MA, Hendrickson CL, Emmett MR, Marshall AG (1999) Gas-phase bovine ubiquitin cation conformations resolved by gas-phase hydrogen/deuterium exchange rate and extent. *Int J Mass Spectrom* 185:565–575
141. Xie Y, Zhang J, Yin S, Loo JA (2006) Top-down ESI–ECD–FT-ICR mass spectrometry localizes noncovalent protein–ligand binding sites. *J Am Chem Soc* 128:14432–14433
142. Sanglier S, Atmanene C, Chevreux G, Van Dorsselaer A (2008) Nondenaturing mass spectrometry to study noncovalent protein/protein and protein/ligand complexes: technical aspects and application to the determination of binding stoichiometries. *Funct Proteomics: Methods Protoc* 484:217–243
143. Yin S, Xie Y, Loo JA (2008) Mass spectrometry of protein–ligand complexes: enhanced gas-phase stability of ribonuclease–nucleotide complexes. *J Am Soc Mass Spectrom* 19:1199–1208
144. Young L et al (2013) Monitoring oligomer formation from self-aggregating amylin peptides using ESI–IMS–MS. *Int J Ion Mobility Spectrom* 16:29–39
145. Konrat R (2014) NMR contributions to structural dynamics studies of intrinsically disordered proteins. *J Magn Reson* 241:74–85
146. Brutscher B et al (2015) NMR methods for the study of intrinsically disordered proteins structure, dynamics, and interactions: general overview and practical guidelines. *Adv Exp Med Biol* 870:49–122
147. Felli IC, Pierattelli R (2015) Intrinsically disordered proteins studied by NMR spectroscopy. Springer, Berlin
148. Wüthrich K (1986) NMR of proteins and nucleic acids. Wiley, New York
149. Dyson HJ, Wright PE (2004) Unfolded proteins and protein folding studied by NMR. *Chem Rev* 104:3607–3622
150. Larsson G, Martinez G, Schleucher J, Wijmenga SS (2003) Detection of nano-second internal motion and determination of overall tumbling times independent of the time scale of internal motion in proteins from NMR relaxation data. *J Biomol NMR* 27:19–312
151. Pawley NH, Wang C, Koide S, Nicholson LK (2001) An improved method for distinguishing between anisotropic tumbling and chemical exchange in analysis of ^{15}N relaxation parameters. *J Biomol NMR* 20:149–165
152. Felli IC, Pierattelli R (2014) Novel methods based on ^{13}C detection to study intrinsically disordered proteins. *J Magn Reson* 241:115–125
153. Nováček J, Židek L, Sklenář V (2014) Toward optimal-resolution NMR of intrinsically disordered proteins. *J Magn Reson* 241:41–52
154. Tompa P, Tompa P (2002) Intrinsically unstructured proteins. *Trends Biochem Sci* 27:527–533
155. Theillet F-X et al (2013) The alphabet of intrinsic disorder. *Intrinsically Disord Proteins* 1(1):e24360. doi:10.4161/idp.24360
156. Schanda P, Van Melckebeke H, Brutscher B (2006) Speeding up three-dimensional protein NMR experiments to a few minutes. *J Am Chem Soc* 128:9042–9043
157. Mobli M, Maciejewski MW, Schuyler AD, Stern AS, Hoch JC (2012) Sparse sampling methods in multidimensional NMR. *Phys Chem Chem Phys* 14:10835–10843
158. Kazimierczuk K, Stanek J, Zawadzka-Kazimierczuk A, Koźmiński W (2010) Random sampling in multidimensional NMR spectroscopy. *Prog Nucl Magn Reson Spectrosc* 57:420–434
159. Gil S et al (2013) NMR spectroscopic studies of intrinsically disordered proteins at near-physiological conditions. *Angew Chem Int Ed* 52:11808–11812
160. Solyom Z et al (2013) Best-trosy experiments for time-efficient sequential resonance assignment of large disordered proteins. *J Biomol NMR* 55:311–321
161. Tycko R (2006) Solid-state NMR as a probe of amyloid structure. *Prot Pept Lett* 13:229–234
162. Bertini I et al (2011) Solid-state NMR of proteins sedimented by ultracentrifugation. *Proc Natl Acad Sci USA* 108:10396–10399
163. Ja Marsh, Singh VK, Jia Z, Forman-Kay JD (2006) Sensitivity of secondary structure propensities to sequence differences between α - and gamma-synuclein: implications for fibrillation. *Protein Sci* 15:2795–2804
164. Camilloni C, De Simone A, Vranken WF, Vendruscolo M (2012) Determination of secondary structure populations in disordered states of proteins using nuclear magnetic resonance chemical shifts. *Biochemistry* 51:2224–2231
165. Tamiola K, Mulder FAA (2012) Using NMR chemical shifts to calculate the propensity for structural order and disorder in proteins. *Biochem Soc Trans* 40:1014–1020
166. Favier A, Brutscher B (2011) Recovering lost magnetization: polarization enhancement in biomolecular NMR. *J Biomol NMR* 49:9–15
167. Bermel W, Bertini I, Felli IC, Kümmerle R, Pierattelli R (2006) Novel ^{13}C direct detection experiments, including extension to the third dimension, to perform the complete assignment of proteins. *J Magn Reson* 178:56–64

168. Camilloni C, Vendruscolo M (2014) Statistical mechanics of the denatured state of a protein using replica-averaged metadynamics. *J Am Chem Soc* 136:8982–8991
169. Platzner G et al (2011) The metastasis-associated extracellular matrix protein osteopontin forms transient structure in ligand interaction sites. *Biochemistry* 50:6113–6124
170. Chabas D et al (2001) The influence of the proinflammatory cytokine, osteopontin, on autoimmune demyelinating disease. *Science* 294:1731–1735
171. Wang KX, Denhardt DT (2008) Osteopontin: role in immune regulation and stress responses. *Cytokine Growth Factor Rev* 19:333–345
172. Gassler N et al (2002) Expression of osteopontin (eta-1) in crohn disease of the terminal ileum. *Scand J Gastroenterol* 37:1286–1295
173. Xanthou G et al (2007) Osteopontin has a crucial role in allergic airway disease through regulation of dendritic cell subsets. *Nat Med* 13:570–578
174. Uaesoontrachoon K et al (2008) Osteopontin and skeletal muscle myoblasts: association with muscle regeneration and regulation of myoblast function in vitro. *Int J Biochem Cell Biol* 40:2303–2314
175. Dalvit C, Fogliatto G, Stewart A, Veronesi M, Stockman B (2001) Waterlogsy as a method for primary NMR screening: practical aspects and range of applicability. *J Biomol NMR* 21:349–359
176. Mayer M, Meyer B (1999) Characterization of ligand binding by saturation transfer difference NMR spectroscopy. *Angew Chem Int Ed* 38:1784–1788
177. Rückert M, Otting G (2000) Alignment of biological macromolecules in novel nonionic liquid crystalline media for NMR experiments. *J Am Chem Soc* 122:7793–7797
178. Sass HJ, Musco G, Stahl SJ, Wingfield PT, Grzesiek S (2000) Solution NMR of proteins within polyacrylamide gels: diffusional properties and residual alignment by mechanical stress or embedding of oriented purple membranes. *J Biomol NMR* 18:303–309
179. Hansen MR, Mueller L, Pardi A (1998) Tunable alignment of macromolecules by filamentous phage yields dipolar coupling interactions. *Nat Struct Biol* 5:1065–1074
180. Tjandra N, Bax A (1997) Direct measurement of distances and angles in biomolecules by NMR in a dilute liquid crystalline medium. *Science* 278:1111–1114
181. Camilloni C, Vendruscolo M (2014) A tensor-free method for the structural and dynamical refinement of proteins using residual dipolar couplings. *J Phys Chem B* 119:653–661
182. Schneidman-Duhovny D, Pellarin R, Sali A (2014) Uncertainty in integrative structural modeling. *Curr Opin Struct Biol* 28:96–104
183. Herrera FE et al (2008) Inhibition of α -synuclein fibrillization by dopamine is mediated by interactions with five c-terminal residues and with e83 in the nac region. *PLoS One* 3:e3394
184. Convertino M, Vitalis A, Caffisch A (2011) Disordered binding of small molecules to A β (12–28). *J Biol Chem* 286:41578–41588
185. Xu L, Gao K, Bao C, Wang X (2012) Combining conformational sampling and selection to identify the binding mode of zinc-bound amyloid peptides with bifunctional molecules. *J Comput-Aided Mol Des* 26:963–976
186. Zhu M et al (2013) Identification of small-molecule binding pockets in the soluble monomeric form of the A β 42 peptide. *J Chem Phys* 139:07B609_1
187. Rauscher S et al (2015) Structural ensembles of intrinsically disordered proteins depend strongly on force field: a comparison to experiment. *J Chem Theory Comput* 11:5513–5524
188. Palazzesi F, Prakash MK, Bonomi M, Barducci A (2015) Accuracy of current all-atom force-fields in modeling protein disordered states. *J Chem Theory Comput* 11:2–7
189. Henriques J, Cragnell C, Skepö M (2015) Molecular dynamics simulations of intrinsically disordered proteins: force field evaluation and comparison with experiment. *J Chem Theory Comput* 11:3420–3431
190. Abrams C, Bussi G (2014) Enhanced sampling in molecular dynamics using metadynamics, replica-exchange, and temperature-acceleration. *Entropy* 16:163–199
191. Bonomi M, Camilloni C, Vendruscolo M (2016) Metadynamic meta-inference: enhanced sampling of the meta-inference ensemble using metadynamics. *Sci Rep* 6:31232
192. Brooks BR et al (1983) CHARMM: a program for macromolecular energy, minimization, and dynamics calculations. *J Comput Chem* 4:187–217
193. Hornak V et al (2006) Comparison of multiple amber force fields and development of improved protein backbone parameters. *Proteins* 65:712–725
194. Lindorff-Larsen K et al (2012) Systematic validation of protein force fields against experimental data. *PLoS One* 7:e32131
195. Wang J, Wolf RM, Caldwell JW, Kollman PA, Case DA (2004) Development and testing of a general amber force field. *J Comput Chem* 25:1157–1174
196. Molnar KS et al (2014) Cys-scanning disulfide crosslinking and bayesian modeling probe the transmembrane signaling mechanism of the histidine kinase, phoQ. *Structure* 22:1239–1251
197. Street TO et al (2014) Elucidating the mechanism of substrate recognition by the bacterial hsp90 molecular chaperone. *J Mol Biol* 426:2393–2404
198. Hummer G, Kofinger J (2015) Bayesian ensemble refinement by replica simulations and reweighting. *J Chem Phys* 143:243150
199. Bonomi M, Camilloni C, Cavalli A, Vendruscolo M (2016) Meta-inference: a Bayesian inference method for heterogeneous systems. *Sci Adv* 2:e1501177
200. Vidal M, Endoh H (1999) Prospects for drug screening using the reverse two-hybrid system. *Trends Biotechnol* 17:374–381
201. Greenfield NJ (2006) Using circular dichroism spectra to estimate protein secondary structure. *Nat Protoc* 1:2876–2890
202. Myszka DG, Rich RL (2000) Implementing surface plasmon resonance biosensors in drug discovery. *Pharm Sci Technol Today* 3:310–317
203. Lo M-C et al (2004) Evaluation of fluorescence-based thermal shift assays for hit identification in drug discovery. *Anal Biochem* 332:153–159
204. Beveridge R, Chappuis Q, Macphee C, Barran P (2013) Mass spectrometry methods for intrinsically disordered proteins. *Analyst* 138:32–42