



## **A spatiotemporal model for snow crab (*Chionoecetes opilio*) stock size in the southern Gulf of St. Lawrence**

**Cadigan, Noel G.; Wade, Elmer; Nielsen, Anders**

*Published in:*  
Canadian Journal of Fisheries and Aquatic Sciences

*Link to article, DOI:*  
[10.1139/cjfas-2016-0260](https://doi.org/10.1139/cjfas-2016-0260)

*Publication date:*  
2017

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Cadigan, N. G., Wade, E., & Nielsen, A. (2017). A spatiotemporal model for snow crab (*Chionoecetes opilio*) stock size in the southern Gulf of St. Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences*, 74(11), 1808-1820. DOI: 10.1139/cjfas-2016-0260

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



**A spatial-temporal model for snow crab (*Chionoecetes opilio*) stock size in the Southern Gulf of St. Lawrence**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2016-0260.R2
Manuscript Type:	Article
Date Submitted by the Author:	28-Oct-2016
Complete List of Authors:	Cadigan, Noel; Memorial University of Newfoundland, Marine Institute Wade, Elmer; Fisheries and Oceans Canada Gulf Region Nielsen, Anders; Technical University of Denmark (DTU Aqua), Marine living resources
Keyword:	Catch per unit of effort, depletion model, density-dependence catchability, STOCK ASSESSMENT < General, Gaussian markov random field



1 A spatial-temporal model for snow crab (*Chionoecetes opilio*) stock size in the Southern Gulf of  
2 St. Lawrence.

3  
4 Noel G. Cadigan<sup>1</sup>, Elmer Wade<sup>2</sup>, Anders Nielsen<sup>3</sup>

5  
6 <sup>1</sup>Centre for Fisheries Ecosystems Research  
7 Marine Institute of Memorial University of Newfoundland  
8 P.O. Box 4920  
9 St. John's, NL  
10 Canada A1C 5R3

11  
12 <sup>2</sup>Aquatic Resources Division  
13 Oceans and Science Branch  
14 Department of Fisheries and Oceans  
15 343 Université Avenue  
16 Moncton, NB  
17 Canada E1C 5K4

18  
19 <sup>3</sup>National Institute of Aquatic Resources  
20 Technical University of Denmark  
21 Charlottenlund Slot Jægersborg Allé 1  
22 2920 Charlottenlund

23  
24 Corresponding author: Noel Cadigan, tel:1 709 778 0603; fax: 1 709 778 0669; email:  
25 [noel.cadigan@mi.mun.ca](mailto:noel.cadigan@mi.mun.ca)

26

## 27 **Abstract**

28 We develop a high resolution spatio-temporal model of stock size and harvest rates for snow  
29 crab (*Chionoecetes opilio*) in the Southern Gulf of St. Lawrence, which supports an  
30 economically important fishery off the east coast of Canada. It is a spatial and weekly model  
31 during 1997-2014 that utilizes within-season depletion based on catch per unit of effort (CPUE;  
32 kg per pot), and also biomass values from a survey designed specifically for this stock. The  
33 model is formulated in a state-space framework. The main contribution of the model is to  
34 provide a better understanding of fishery-dependent factors that affect CPUE. There is strong  
35 evidence of density-dependence in the relationship with CPUE and stock biomass, in addition to  
36 a general increase in CPUE catchability over time that may be related to changes in gear soak  
37 time, and spatial variation in catchability. We also find that a natural mortality rate of 0.4  
38 provides a better fit to survey results. Model results suggest that there is no evidence of effort  
39 saturation in the fishery.

40

41 Keywords: Catch per unit of effort; depletion model; Gaussian markov random field; density-  
42 dependence catchability; stock assessment.

43

## 44 **Introduction**

45 Snow crab (*Chionoecetes opilio*) have been fished commercially in the southern Gulf of St.  
46 Lawrence (sGSL; Figures 1 and 2) since the mid-1960's using baited traps (i.e. pots). Monitoring  
47 of this fishery is based on logbook data, dockside monitoring of the catch, and at-sea sampling  
48 by observers. Snow crab normally molt every year until maturity (Conan and Comeau 1986)

49 prior to the fishery (e.g. Hébert et al. 2002). After molting, crabs have a soft shell for 8 to 10  
50 months. Males live approximately 5 years (Sainte-Marie et al. 1995) to 7.7 years (Fonseca et al.  
51 2008) after their terminal molt. The fishery targets hard-shelled adult male animals with a  
52 minimum landing size of 95 mm carapace width (CW). The management of this fishery is based  
53 on total allowable catch (TAC), effort controls (number of licenses, trap limits, and seasons), and  
54 a limit on the percentage of the low commercial value soft-shelled crabs in the catch.

55 This stock is assessed using size and/or stage-based survey indices of biomass and abundance  
56 from a post-season trawl survey designed specifically for this stock (Moriyasu et al. 2016;  
57 Hébert et al. 2016b). A grid-based sampling design has been used in all survey years, although  
58 the survey coverage and grid cell size has changed over time. Total survey biomass for various  
59 life stages of crab is estimated using kriging with external drift using depth as a secondary  
60 variable (Hébert et al. 2016b). These stages include residual biomass of commercial sized males  
61 remaining after the fishery, and recruitment biomass (soft-shelled adult males  $\geq 95$  mm CW)  
62 that will be available to the fishery in the following fishing season. The exploitable biomass of  
63 commercial males is inferred from the residual and recruited biomass from the previous year's  
64 survey. A harvest decision rule including precautionary approach reference points based on  
65 commercial biomass is used to provide total allowable catch (TAC) advice (Chaput et al. 2014).  
66 A simple model has been used to project biomass in the next several years based on survey  
67 estimates of pre-recruit stages (Surette and Wade, 2006). The only other model published for this  
68 stock was an age-, sex- and stage-structured model by Siddeek et al. (2009), and the purpose of  
69 this model was for developing limit and target harvest control rules, but not directly for stock  
70 assessment.

71 Commercial catch rates are considered in sGSL snow crab stock assessments but the relationship  
72 between commercial-sized adult male biomass estimated from the snow crab survey and CPUE  
73 calculated from logbooks is weak;  $r^2 = 0.22$  (Hébert et al. 2016a) and varies across different  
74 locations. This results in differences in perception of stock abundance between the fishing  
75 industry observations and those from the survey. Fishing industry perceptions not only involve  
76 the overall level of CPUE but also the rate of decline in CPUE as the fishing season progresses.  
77 They consider that if CPUE does not decline much then this is an indication of a larger stock size  
78 than when CPUE declines substantially. This is also related to the sedentary nature of snow crabs  
79 (Biron et al. 2008) during the sGSL fishery.

80 It is possible to estimate absolute biomass based on the within-season rate of decline of a  
81 biomass index in response to the fishery removals. The Leslie method (Leslie 1952) has been  
82 used for many crustacean species (see review by Smith and Addison 2003) and others (e.g.  
83 Bishir and Lancia 1996). Its popularity is partly related to its simplicity. If  $C_t$  and  $E_t$  are the catch  
84 and effort of a fishery at time  $t$ , and CPUE is  $X_t = C_t/E_t$  and is proportional to stock biomass  $B_t$ ,  
85 then the Leslie method involves fitting the linear model  $X_t = qB_o - qK_t$ , where  $q$  is the  
86 catchability coefficient,  $B_o$  is the biomass at the start of the fishing season, and  $K_t$  is the  
87 cumulative catch at time  $t$ . These parameters can be estimated using a linear regression of  $X_t$  on  
88  $K_t$ . Alternatively, the DeLury method (DeLury 1947) involves fitting to cumulative effort rather  
89 than cumulative catch. Both methods use the rate of decline of CPUE throughout the fishing  
90 season, as a function of either catch or effort, to estimate initial stock size. These are two  
91 examples of depletion models.

92 In practise there are many problems with assumptions for these simple depletion estimators.  
93 Miller and Mohn (1993) examined issues with the Leslie method for species like crabs and

94 lobsters. An important assumption is that catchability is the same for all individuals throughout  
95 the fishing season. Of particular relevance to sGSL commercial snow crab is the potential for 1)  
96 a change (decrease or increase) in capture probability depending on how many crabs are in a  
97 trap, which will result in density-dependent CPUE catchability, and 2) increased activity in  
98 warmer waters could lead to higher catchability in nearshore areas during the spring and  
99 summer. The physical habitat and temperature may affect crab walking rates, and current  
100 direction and speed may affect the bait “plume” and trap detection. CPUE catchability may also  
101 change for many other reasons (also see Miller 1990, Bacheler et al. 2013). Varying  
102 characteristics of fishing effort such as soak time, bait type and amount, and trap design and  
103 spacing can affect catchability. Trap saturation may happen with longer soak times for many  
104 reasons including space limitation of the gear, inter-species or intra-species interactions, or the  
105 loss and degradation of bait (Shertzer et al. 2015). However, Charles et al. (2014) found soak  
106 time had little effect in a study of sGSL snow crab. Temporal, spatial, and density-dependent  
107 variations in CPUE catchability have been examined extensively in the fisheries literature.  
108 Density-dependence relationships between CPUE and stock size (i.e. catchability) are commonly  
109 reported (e.g. Harley 2001, Ward et al. 2013) but are not ubiquitous (e.g. Zhou et al. 2007).  
110 Many factors may produce a power-law relationship between CPUE and stock size (e.g. Cooke  
111 and Beddington 1984). If such factors change in a systematic way, bias can be introduced to  
112 depletion estimators.

113 Another important assumption is that fishing effort is uniformly distributed over the area  
114 occupied by the stock. This is not the case for sGSL snow crab. Fishing fleets may adjust their  
115 fishing locations when catch rates decline and this can mask the decline in stock size due to  
116 fishing (e.g. Hilborn and Walters 1987). Depletion methods also require landings and effort to be

117 accurately reported. Fisheries monitoring for sGSL snow crab is rigorous so reported catch  
118 should be reliable. However, effort is not evaluated in dockside monitoring and may be  
119 inaccurate. The Leslie method assumes a closed population, with no emigration, natural  
120 mortality, or recruitment. The last two issues are relevant for snow crab because of the length of  
121 the fishing season. Concerns about within-season recruitment and fleet movements led scientists  
122 to conclude that the Leslie method was inappropriate for stock assessment of sGSL snow crab  
123 (Loch et al. 1995). Corrections for these problems, when they exist, require additional  
124 information (Miller and Mohn 1993).

125

126 Various extensions of depletion models have been proposed to address some of the above  
127 problems. Gedamke et al. (2004) used information from vessel monitoring systems to  
128 disentangle the effects of nonrandom fishing patterns for scallops so that a depletion model could  
129 be applied. They used spatial analyses to define regions where assumptions of the depletion  
130 method were met. Walter et al. (2007) extended this stratification approach and conducted a  
131 geostatistical analysis of their depletion model results. Robert et al. (2010) formulated the  
132 DeLury model in a Bayesian framework and included natural mortality and within-season  
133 recruitment for an octopus fishery, with simulation testing of various model formulations and  
134 data configurations. They also modelled a range of years in a hierarchical framework in which  
135 initial abundance and catchability were related across years. Roa-Ureta (2012) proposed  
136 generalized depletion models for open populations and a nonlinear relationship between catch  
137 and effort. His model allowed for waves of within-season immigration to the fishing grounds. He  
138 applied his model to a squid fishery. Roa-Ureta (2015) extended the generalized depletion model  
139 to be multi-annual and to include multiple fleets with different catchabilities for a Spanish



140 mackerel fishery. Zhou et al. (2011, 2015) presented a generalized Leslie-type model to estimate  
141 natural mortality rates as well as stock size and harvest rates in a Bayesian framework. Their  
142 multi-annual biomass model included within-season somatic growth and a parametric year-  
143 varying catchability function.

144 There is often little evidence of depletion in total fishery CPUE for sGSL snow crab. In the 2014  
145 fishery, total CPUE (total catch divided by total effort) increased overall and only decreased for  
146 three short periods (Figure 3). Average soak times rapidly increased from about 50 hours at the  
147 start of the fishery to about 120 hours by week 25. This may partly explain the stability of CPUE  
148 during this period. However, another factor is a change in the distribution of the fishery (Figure  
149 4). There is a pronounced change in the fleet distribution starting in August (week 29). This is a  
150 consistent feature throughout the time-series. Also, in some weeks there may be a large  
151 concentration of effort in just a few grids cells and pots may compete for crab in these cases. If  
152 multiple pots are placed within their effective fishing area then this will eventually produce a  
153 saturation effect in which catch is no longer proportional to effort. Not surprisingly, in  
154 preliminary analyses we found poor correspondence between stock size and harvest rates from a  
155 sGSL spatial depletion model and survey results. Hence, in this paper we also explore changes in  
156 model catchability assumptions, both temporal and spatial, to account for changes in the fishery  
157 to improve this correspondence.

158 We propose a novel CPUE and survey integrated spatial biomass depletion model that provides  
159 spatio-temporal estimates of snow crab stock size and also density-dependent and within-season  
160 spatio-temporal changes in CPUE catchability. We provide more precision on how to interpret  
161 CPUE catchability for snow crab, or more generally for stocks in which it is reasonable to  
162 assume that the gear catches all fish in the area the gear fishes. We use high resolution spatio-

163 temporal (i.e. weeks and grids) catch and effort data, and also spatial survey data, for model  
164 estimation. Extracting more information about stock size using data at the highest resolution  
165 possible can produce more reliable stock assessments and more relevant information for fisheries  
166 managers, but an important cost of doing this is the need for a highly parameterized model that is  
167 potentially computationally complex. Another contribution of this paper is a high resolution  
168 spatial biomass depletion model that is restricted in certain ways to make the model  
169 computationally efficient. We also include a novel effort saturation effect that is spatial because  
170 of differences in the fraction of crab “grounds” within grid cells. Our model is implemented in a  
171 state-space modelling framework which has become a favored approach in modeling time  
172 varying population dynamics (e.g. Nielsen and Berg, 2014; Cadigan, 2016).

173

## 174 **Materials and Methods**

175 Model notation and parameters are described in the text and also Table 1. The data we used are  
176 weekly fisherman logbook observations of catch and effort for 314 spatial grid cells in sGSL  
177 during 1997-2014. Also, we use the post fishery snow crab survey trawl catches of hard-shelled  
178 adult males (i.e. exploitable crab). These data are displayed in Supplementary figures, for catch  
179 weight (Figures SC.1-20), effort (Figures SE.1-20), CPUE (Figures SX.1-20), and the snow crab  
180 survey results for exploitable biomass (Figures SSEB.1-19) and total biomass (Figures SSTB.1-  
181 19) although the latter information is not used in our model but is used in the DFO stock  
182 assessment.

183 The biomass ( $B$ ) depletion model for grid cell  $i$  and week  $t$  in year  $y$  is

$$B_{y,t,i} = B_{y,t-1,i} \exp(-Z_{y,t-1,i}), \quad (1)$$

184 where the weekly total mortality rate is  $Z_{y,t,i} = M/52 + F_{y,t,i}$ ,  $F_{y,t,i}$  is the fishing mortality rate,  
 185 and  $M$  is an assumed constant annual value for the natural mortality rate which we prorate  
 186 equally to weeks. We examine  $M = 0.2, 0.3,$  and  $0.4$  which are values that are consistent with the  
 187 survival times in Sainte-Marie et al. (1995) and Fonseca et al. (2008). The depletion model starts  
 188 at an initial biomass  $B_{y,o,i}$  in year  $y$  and we project this to the start of the fishery using Equation  
 189 (1) with no  $F$ . Our model does not include movement of crab between grid cells which provides  
 190 substantial computational efficiencies. Snow crab in the sGSL are relatively sedentary during the  
 191 fishery (Biron et al. 2008) but they may move somewhat between grid cells during the winter  
 192 months before the fishery so  $B_{y,o,i}$  is only a nominal initial biomass for a grid cell and the actual  
 193 biomass may be different. However, we assume that  $\sum_i B_{y,o,i}$  is the total stock biomass at the  
 194 beginning of the year because errors due to movements will cancel out in the total biomass. The  
 195 catch equation is

$$C_{y,t,i} = B_{y,t,i} \{1 - \exp(-Z_{y,t,i})\} \frac{F_{y,t,i}}{Z_{y,t,i}}. \quad (2)$$

196

197 A basic assumption underlying our spatial depletion model is that  $CPUE = C/E = qB$  where  $E$  is  
 198 the effort which we define as the number of pots fished. This implies that the harvest rate is  $H \equiv$   
 199  $C/B = qE$  and  $q$  represent the harvest rate of a single pot. Some boundary grid cells are small so  
 200 an area-adjustment is useful. Let  $a_{fi}$  be the area fished by a pot which we assume varies spatially  
 201 ( $i$ ) and temporally ( $t$ ) because of a variety of factors related to the fishery (e.g. soak time);  
 202 however, we assume the area fished by a pot is constant within a grid cell and a week. Let  $A_i$  be

203 the area of the grid cell and  $p_i$  be the fraction of area that are crab grounds. If crab are  
 204 homogeneously distributed within the crab grounds and if a pot catches all crab within the area it  
 205 fishes then the expected catch from a single randomly placed pot in cell  $i$  with total biomass  $B_{y,t,i}$   
 206 is  $q_{y,t,i}B_{y,t,i}/A_i$  where  $q_{y,t,i} = a_{fy,t,i}/p_i$  is the potential area fished; that is,  $a_{fy,t,i} \leq q_{y,t,i}$ . The total catch  
 207 from  $E_{y,t,i}$  pots is  $C_{y,t,i} = q_{y,t,i}E_{y,t,i}D_{y,t,i}$  where  $D_{y,t,i} = B_{y,t,i}/A_i$  is the biomass density in grid cell  $i$ .  
 208 Here we assumed that  $E_{y,t,i} \ll q_{y,t,i}$  so that pots fish independently. Hence the expected CPUE is  
 209  $X_{y,t,i} = q_{y,t,i}D_{y,t,i}$ . The CPUE biomass catchability is proportional to grid cell size and varies across  
 210 years, weeks and grids.

211 We have incomplete information on the factors that may cause catchability to change over time  
 212 but we expect that these factors will be auto-correlated. Hence, we assume that

$$q_{y,t,i} = q_{y,i} \exp(\delta_{y,t,i}), \quad (3a)$$

213 where  $\delta_{y,t,i}$  are weekly  $q$  process errors that we assume for simplicity are AR(1) auto-correlated  
 214 normal random variables with mean zero, standard deviation  $\sigma_\delta$ , and auto-correlation parameter  
 215  $\varphi_\delta$ . These errors are assumed to be correlated across weeks within grid cells and years but  
 216 otherwise are independent. Including process error is typical of contemporary stock assessment  
 217 models (Maunder and Piner, 2014). The  $q_{y,i}$  annual spatial effects are further separated into  
 218 independent year and space effects

$$q_{y,i} = q_y q_i, \quad (3b)$$

219 The  $q_y$  year-effects account for long-term changes in fishing practices and are modelled as a  
 220 random walk,

$$\log(q_{y+1}/q_y) \sim N(0, \sigma_{q_y}^2). \quad (4)$$

221 The  $q_i$  spatial effects are described below. Hence, we model catchability at high resolution using  
 222 correlated spatial effects, a random walk year effect, and auto-correlated weekly process errors  
 223 within grid cells.

224 The idealized maximum effective effort that can be fished without pot competition is  $E_{max,y,t,i} =$   
 225  $A_i/q_{y,t,i}$ . This is the total number of distinct pot fishing areas in grid  $i$  with total area  $A_i$ . The  
 226 model-predicted effort corresponding to  $C_{y,t,i}$  is  $E_{y,t,i} = C_{y,t,i}/q_{y,t,i}D_{y,t,i}$  and

$$E_{y,t,i} = \frac{C_{y,t,i}A_{y,t,i}}{q_{y,t,i}B_{y,t,i}} = H_{y,t,i}E_{max,y,t,i} \quad (5)$$

227 We will use Equation (5) to model observed effort. Note that unlike much of the fisheries  
 228 literature we do not assume effort is a covariate that is known without error and we are therefore  
 229 better able to account for the reliability of this data. However, if there is pot saturation then pots  
 230 may compete for crabs and the effective (i.e. independent) effort is less than the model  
 231 prediction.

232 If fishermen randomly chose fishing locations regardless of gear that may already be present then  
 233 this behavior can be approximated as a Poisson process. We assume that each grid cell is  
 234 comprised of  $E_{max,y,t,i}$  distinct fishing locations and that all pots within one location exploit the  
 235 same as a single pot. Let  $E_{o,y,t,i}$  be the nominal total effort and let  $Y$  be the number of pots in a  
 236 fishing location which has an approximate Poisson distribution with mean  
 237  $\lambda_{y,t,i} = E_{o,y,t,i}/E_{max,y,t,i}$ . The probability that a location has at least one pot is  $Prob(Y > 0) =$   
 238  $1 - \exp(-\lambda_{y,t,i})$ . The number of fishing locations with at least one pot, which is the effective  
 239 effort  $E_{y,t,i}$ , is

$$E_{y,t,i} = E_{max,y,t,i} \{1 - \exp(-E_{o,y,t,i}/E_{max,y,t,i})\}. \quad (6a)$$

240 However, crab fishermen are aware of gear already present at a location and they will usually  
 241 avoid setting new gear close to existing gear. In this case Equation (6a) will under-estimate  
 242 effective effort. We propose an alternative smooth “hockey-stick” formulation (e.g. Mesnil and  
 243 Rochet, 2010) to account for this behavior,

$$E_{y,t,i} = \frac{1}{2}E_{o,y,t,i} + \frac{1}{2}E_{max,y,t,i} \left\{ \sqrt{1 + \gamma^2/4} - \sqrt{(E_{max,y,t,i}^{-1}E_{o,y,t,i} - 1)^2 + \gamma^2/4} \right\}. \quad (6b)$$

244 Note that we modified the model formulation of Mesnil and Rochet (2010) so that  $\gamma$  is a fraction  
 245 of  $E_{max,y,t,i}$ .

246 Equations (6a) and (6b) are illustrated in Figure 5 for  $E_{max,y,t,i} = 2000$ . We set  $\gamma = 0.3$  in  
 247 Equation (6b). Effective effort based on Equation (6b) and nominal effort are almost the same  
 248 until effort is close to  $E_{max}$  but effective effort based on Equation (6a) is usually much less than  
 249 nominal effort unless effort is low. Both Equations (6a) and (6b) imply that  $E_{y,t,i} \leq E_{o,y,t,i}$ . If  
 250  $E_{o,y,t,i} \ll E_{max,y,t,i}$  then  $E_{y,t,i} \approx E_{o,y,t,i}$  but as  $E_{o,y,t,i} \rightarrow \infty$  then  $E_{y,t,i} \rightarrow E_{max,y,t,i}$ . We can use  
 251 the inverse of these equations as functions of  $E_{o,y,t,i}$  to get the model predicted nominal fishing  
 252 effort. For Equation (6a) this is

$$E_{o,y,t,i} = -E_{max,y,t,i} \log(1 - H_{y,t,i}). \quad (7a)$$

253 Let  $U_{y,t,i} = 2H_{y,t,i} - 1 - \sqrt{1 + \gamma^2/4}$ . The inverse of Equation (6b) is

$$E_{o,y,t,i} = E_{max,y,t,i} \left\{ 1 + \frac{U_{y,t,i}^2 - \gamma^2/4}{2U_{y,t,i}} \right\}. \quad (7b)$$

254 If there is negligible effort saturation then

$$E_{o,y,t,i} = H_{y,t,i} E_{max,y,t,i}. \quad (7c)$$

255

256 We also include a density dependence effect in  $q$ . Our basic generating equation is  $CPUE = qD^\beta$   
 257 and  $\beta = 1$  means no density dependence. However,  $\beta$  changes the interpretation of  $q$  so we use a  
 258 slightly modified version,  $CPUE = qDD_s^{\beta-1}$  where  $D_s = D/\bar{D}$  is the density relative to the  
 259 average over space and time. We computed an average value of  $D$  for all years, weeks, and grid  
 260 cells in which fishing occurred, using a specific model formulation (M1; see Results) but kept  
 261 this same value (i.e. 867 kg/km<sup>2</sup> or 49 Kt for the sGSL total area) in other model formulations.  
 262 The re-scaling of density only affects estimation of  $q$  and not  $\beta$ . When  $D = \bar{D}$  then  $CPUE = qD$ ,  
 263 so  $q$  represents catchability at average stock density. Our density-dependence model only  
 264 requires modifying Equation (3),

$$q_{y,t,i} = D_{s,y,t,i}^{\beta-1} q_{y,i} \exp(\delta_{y,t,i}). \quad (8)$$

265 There is otherwise no change in the way annual and spatial effects ( $q_{y,i}$ ) or weekly effects ( $\delta_{y,t,i}$ )  
 266 are modelled. However, this density dependence may be confounded with effort saturation.

267 There are two groups of spatial effects in our model, the yearly initial biomass densities  $D_{y,o,i} =$   
 268  $B_{y,o,i}/A_i$  and the catchabilities  $q_i$ . We have insufficient data to estimate these effects freely, nor  
 269 would this be a good idea because we expect spatial correlation in these effects. Hence, we also  
 270 model these spatial effects as latent random variables on the snow crab survey grid. Let  $G$  be the  
 271 total number of grid cells. The covariance structure we use for these random variables is the  
 272 same as that described in Kristensen et al. (2013) and is based on a precision matrix (i.e. inverse

273 covariance matrix) that implies that a grid cell is, conditional on the cell's neighbors,  
 274 independent of all other cells. The precision matrix  $\Psi_{G \times G}$  for the logs of  $D_{y,o,i}$  or  $q_i$  has  $i,j$ 'th  
 275 element

$$\Psi_{i,j} = \begin{cases} -\tau, & \text{if cell } i \text{ neighbors cell } j, \\ \tau(m_i + \vartheta), & \text{if } i = j, \\ 0, & \text{otherwise,} \end{cases} \quad (9)$$

276 where  $m_i$  is the number of neighbors of grid cell  $i$  which will typically be four but will be less  
 277 for boundary cells. The  $\tau$  and  $\vartheta$  parameters are estimated separately for each spatial effect. We  
 278 denote these as  $\tau_D$  and  $\vartheta_D$ , etc. The precision matrix approach is typical of those used in  
 279 Gaussian Markov random fields (GMRF) and implies that the correlation decreases with distance  
 280 traveled through water and not simply the straight-line distance between two grid cells. This is  
 281 particularly relevant for the sGSL because of irregular coast lines (i.e. Chaleur Bay) and island  
 282 obstacles (i.e. the Magdalen Islands; see Figure 1). Let  $\underline{D}_{y,o}$  be a vector of spatial density effects  
 283 for year  $y$ . We also expect that the spatial distribution of biomass will be similar from year to  
 284 year. We accommodate this behavior by modelling the density in the first year,  $\log(\underline{D}_{1,o})$ , as a  
 285 GMRF with parameters  $\tau_{D1}$  and  $\vartheta_{D1}$ , and the relative difference in densities in successive years,  
 286  $\log(\underline{D}_{y+1,o}/\underline{D}_{y,o})$ , as independent GMRFs with parameters  $\tau_{D2}$  and  $\vartheta_{D2}$ . These latter parameters  
 287 are assumed to be the same for all years. Hence, our model involves a spatially correlated  
 288 random walk for biomass density from year to year.

289 The data available to estimate model parameters ( $\theta$ , see next Section) are spatial catch and effort  
 290 measurements each week during 1997-2014, and the snow crab survey catches for the same  
 291 years (see Supplementary figures). Our model is for male commercial biomass so we only use  
 292 survey catch information for hard-shelled legal size males that were exploited in the same year of



293 the survey (i.e. residual exploitable biomass). We assume a small and fixed value for log-catch  
 294 measurement error (ME) standard deviation (SD; i.e.  $\sigma_C = 0.05$ ) because this data is obtained  
 295 from dockside monitoring with reportedly high accuracy. This is approximately the catch  
 296 measurement error CV. The state-space model log-likelihood (i.e. observation) equation for  
 297 catch is

$$l(\theta|\{C_{obs}\}) = \sum_y \sum_t \sum_i \log \left[ \sigma_C^{-1} \varphi_N \left\{ \frac{\log(C_{obs,y,t,i}) - \log(C_{y,t,i})}{\sigma_C} \right\} \right], \quad (10)$$

298 where  $\varphi_N$  is the standard normal probability density function and  $\{C_{obs}\}$  indicates the set of all  
 299 catch observations. The model values for catches,  $C_{y,t,i}$ , come from Equation (2). Effort is not  
 300 evaluated in dockside monitoring and is reported less accurately than catches. We assume  
 301 multiplicative ME,  $E_{obs,y,t,i} = E_{o,y,t,i} e^{\varepsilon_{y,t,i}}$  where the model values for reported effort,  $E_{o,y,t,i}$ ,  
 302 come from Equations (7a), (7b), or (7c) depending on whether a saturation effect is included or  
 303 not, and the type of effect, and  $\varepsilon$  is the effort ME. This ME may be auto-correlated across weeks  
 304 within grid cells and years because it is likely that the same fishing crews are involved and they  
 305 are likely to report effort with similar error. If there is no effort saturation then using Equations  
 306 (3) and (5) we can show that

$$\log(E_{obs,y,t,i}) = \log(H_{y,t,i}) + \log(A_{y,t,i}) - \log(q_{y,i}) + \varepsilon_{y,t,i} - \delta_{y,t,i}. \quad (11)$$

307 The  $\varepsilon$  effort ME's and the  $\delta$  catchability process errors are confounded. We can only estimate  
 308 their total effect. We still refer to these errors as  $\delta$  process errors although they are a mixture of  
 309 changes in catchability and ME in effort. Let  $t_{f,y,i}$  denote the first week of the fishery in year  $y$  and  
 310 grid cell  $i$ . We re-define  $\delta_{y,t,i} = \log(E_{obs,y,t,i}) - \log(E_{o,y,t,i})$  and the likelihood equation for  
 311 effort is

$$l(\theta|\{E_{obs}\}) = \sum_y \sum_i \sum_{t>t_{f,y,i}} \log \left[ \sigma_\delta^{-1} \varphi_N \left\{ \frac{\delta_{y,t,i} - \varphi_\delta \delta_{y,t-1,i}}{\sigma_\delta} \right\} \right]. \quad (12)$$

312 For the first week of the fishery, when  $t = t_{f,y,i}$  then  $\delta_{y,t-1,i} = 0$ .

313 The snow crab survey biomass values,  $S_{obs,y,i}$ , are assumed to be absolute estimates of the  
 314 biomass in the trawl swept area. We re-scaled these biomass values to be equivalent to a tow  
 315 with a standard swept area of 2700 m<sup>2</sup> which is typical in the survey design. The model predicted  
 316 survey biomass values are  $S_{y,i} = 2700B_{y,34,i}/A_i$  where the grid cell area is in m<sup>2</sup>. We assumed  
 317 this survey occurred in week 34 which is about the end of August. This is considered further in  
 318 the Discussion. In preliminary analysis we found that the ME SD ( $\sigma_S$ ) did not depend much on  
 319 predicted values. Also, there are a large number of zero's in the survey time-series so log-  
 320 transformation was not an option and did not seem necessary. Hence, the survey likelihood  
 321 equation is

$$l(\theta|\{S_{obs}\}) = \sum_y \sum_i \log \left[ \sigma_S^{-1} \varphi_N \left\{ \frac{S_{obs,y,i} - S_{y,i}}{\sigma_S} \right\} \right]. \quad (12)$$

322

323 The  $F$ 's in Equation (2) are also treated as unstructured random effects with a common mean and  
 324 variance. Let  $\mu_F = E(F)$  and  $\sigma_F^2 = Var\{\log(F)\}$ . The likelihood equation for  $F$ 's is

$$l(\{F\}|\theta) = \sum_y \sum_t \sum_i \log \left[ \sigma_F^{-1} \varphi_N \left\{ \frac{\log(F_{y,t,i}) - \log(\mu_F)}{\sigma_F} \right\} \right]. \quad (13)$$

325 We will show that the estimate of  $\sigma_F$  is fairly large and so the  $F$ 's are estimated almost freely.

326 **Estimation**

327 The model is high dimensional, based on 314 spatial grid cells and 936 time steps for the 18  
 328 years and 52 weeks per year, resulting in slightly over 290 000 space-time cells. However, there  
 329 are only a small number of parameters to estimate. Most of the model effects are random and are  
 330 not freely estimated. Fixed effect parameters, denoted collectively as the parameter vector  $\theta$ , are  
 331 estimated via maximum likelihood (MLE) based on the total marginal likelihood,  $L(\theta)$ , in which  
 332 random effects are “integrated out”. Let  $\Gamma$  denote a vector of all random effects. There are 13 or  
 333 14  $\theta$ 's, depending on the model formulation, but there are almost 27 000  $\Gamma$ 's. Note that there are  
 334 far fewer  $\Gamma$ 's than the total number of space-time model cells because  $\delta$ 's and  $F$ 's are only  
 335 estimated for cells that had reported catches, and in many weeks there is no fishing in a grid cell  
 336 (see Supplementary figures).

337 Let  $S$  denote the set of all survey, catch, and effort data used in the model. The marginal  
 338 likelihood is

$$L(\theta) = \iiint_{\Gamma} f_{\theta}(S|\Gamma)g_{\theta}(\Gamma)\partial\Gamma, \quad (14)$$

339 where  $f_{\theta}(S|\Gamma)$  is the pdf of the data and  $g_{\theta}(\Gamma)$  is the pdf for the  $\Gamma$  random effects. The  
 340 template model builder (TMB; Kristensen, 2015) package within R (R Core Team, 2016) was  
 341 used to implement the model. The MLE's of  $\theta$  maximize  $L(\theta)$ . The user has to provide C++  
 342 computer code to calculate  $f_{\theta}(S|\Gamma)$  and  $g_{\theta}(\Gamma)$  but the integration in Equation (14) and  
 343 calculation of  $\theta$  to maximize this equation is then provided by TMB. The high dimensional  
 344 integral is numerically evaluated in TMB using the Laplace approximation. The random effects  
 345  $\Gamma$  can be predicted by maximizing the joint likelihood,  $f_{\theta}(S|\Gamma)g_{\theta}(\Gamma)$ . Additional information  
 346 on these procedures is provided by Skaug and Fournier (2006). TMB uses automatic  
 347 differentiation to evaluate the gradient function of Equation (14) and in the Laplace

348 approximation. The gradient function is produced automatically from  $f_{\theta}(S|\Gamma)$  and  $g_{\theta}(\Gamma)$ . This  
349 greatly improves parameter estimation using a derivative-based optimizer. We use the *nlminb*  
350 function within R (R Core Team, 2016) to find the MLE for  $\theta$ .

351 The reliability of model estimates was assessed through detailed examination of model estimates  
352 and residuals, and also retrospective analyses in which the model was fit to subsets of data with  
353 recent years left out. Some covariance parameters are shared across years in the model so it is  
354 possible that there may be retrospective patterns in model estimates. This was done for  
355 retrospective years 2007 to 2014. Sensitivity analyses to model assumptions about  $M$  (0.2, 0.3, or  
356 0.4), saturation (none or hockey-stick type), and density-dependence (none or estimated) were  
357 also performed, leading to a total of  $3 \times 2 \times 2 = 12$  model formulations that were fitted.

358

## 359 **Results**

360 Including a saturation effect did not make much difference to estimates and model fits because  
361 the saturation level was usually estimated to be much greater than typical levels of effort. For  
362 example, two models described in Table 2 (i.e. M1 and M2), that differed only in whether a  
363 “hockey-stick” saturation effect was included or not, produced nearly identical parameter  
364 estimates and the model without effort saturation (i.e. M2) produced a better fit. This was a  
365 consistent feature in other model formulations we compared for different values of  $M$  and the  
366 density dependence parameter  $\beta$ . There is also no evidence of saturation in plots of catch versus  
367 effort (unreported results) for various regions in sGSL.

368 The statistical evidence for density dependence was very strong (e.g. compare M2 and M3, and  
369 M4 and M5) and the estimate of  $\beta \ll 1$  indicating high hyper-stability in CPUE. This effect  
370 resulted in a much improved fit to the snow crab survey residual biomass values, as evidenced by  
371 the lower value of  $\sigma_S$  for M3 and M4. Density dependence is evident in plots of CPUE versus  
372 mean biomass for each grid cell and year (Figure 6) even from a model (i.e. M2 in Table 2) with  
373 no density dependence. The slope of log CPUE versus log biomass should be close to one if  
374 there is no density dependence but this is clearly not the case. Note that to simplify this figure we  
375 only plotted average CPUE and biomass for all weeks within each grid cell and year. Results  
376 based on weekly values were very similar. Including density dependence also resulted in  
377 substantially different estimates of spatial effects. For example, the total variation,  $\text{trace}(\Psi^{-1})$ , of  
378 the spatial effects for CPUE catchability (i.e.  $q_i$ ) was 303.0 for M2 but only 51.8 for M3,  
379 indicating more between grid variation in  $q$  for M2. Conversely, the total variation of the D2  
380 spatial random walk for between year changes in biomass density was 152.1 for M3 but only  
381 69.9 for M2 indicating larger between year changes in the spatial distribution of biomass for M3  
382 compared to M2. The total variation of the D1 spatial distribution of biomass in the first year (i.e.  
383 1997) was 16 221 for M2 and 17 671 for M3. The D1 total variance is much larger than the D2  
384 total variance which indicates relatively small changes in the spatial distribution of biomass from  
385 year to year for both models. The  $\vartheta$  parameters control the decorrelation range in the spatial  
386 covariance matrix. When these parameters are small then the decorrelation range is large (see  
387 Kristensen et al. 2013); that is, the spatial correlation is high. The relatively large CV's for  $\vartheta$ 's  
388 estimates for models M1-M5 indicate that the spatial correlations are estimated with less  
389 precision than other parameters.

390 Another model formulation issue was the choice of  $M$ . We investigated three choices ( $M = 0.2$ ,  
391  $0.3$ , and  $0.4$ ). All values resulted in fairly similar fits (see Table 2 for  $M = 0.3$  or  $0.4$ ). Models in  
392 which  $\beta = 1$  indicated  $M = 0.4$  was the best choice. When  $\beta$  was estimated and no effort  
393 saturation was assumed then the best fit was obtained with  $M = 0.2$ . The fit was 2.6 AIC units  
394 lower than the fit for  $M = 0.3$ . However, models with  $M=0.4$  produced estimates of beginning of  
395 year commercial biomass that agreed more closely with the stock assessment kriging biomass  
396 estimates (DFO, 2016) scaled by  $\exp(-18M/52)$  to account for natural mortality between the  
397 time of the post-season survey and the end of the year (Figure 7). This scaling is described in  
398 more detail later. We fit the model only to survey catches of residual biomass so the improved fit  
399 to DFO kriging estimates of residual plus recruited biomass provides additional and partially  
400 independent support for  $M = 0.4$ . Hence, our preferred model formulation (i.e. M4 in Table 2) is  
401  $M = 0.4$ , with density dependence in CPUE, but no effort saturation effect. More specific results  
402 from this model are presented in Table 3. Detailed spatial models results are provided in  
403 Supplementary Figures S5-S22.

404 On a tow-by-tow basis the M4 fit to the snow crab survey catches was highly variable (Figure 8;  
405 Figures S1, S5-S22). The mean predicted survey catch was 1.0799 and  $\hat{\sigma}_S = 1.153$  (Table 2)  
406 indicating that the survey CV was 108%. The snow crab survey catches have substantial spatial  
407 variability (see Figures SSEB.1 – SSEB.19) and many zero catches that the model did fit well.  
408 Of particular concern is the tendency of the model to under-estimate high values (Figure 8)  
409 which suggests potential model mis-specification. The model is constructed to fit catches almost  
410 exactly. Effort observations were fit less closely but without apparent bias (Figures S2-S4).  
411 There was very little retrospective variation in model results (unreported results).

412 All models indicated significant changes in catchability over year. The standard deviation ( $\sigma_{qy}$ )  
413 of the random walk for the  $q_y$  year-effects was around 0.1 (Table 2). There was also significant  
414 spatial variability in  $q$ 's (Figure 9). We computed the effort-weighted average of  $q_{y,i}$  in Equation  
415 (3) for all grid cells each year, which is proportional to  $q_y$ , and divided this by the average grid  
416 area to estimate the catchability of a pot which is also the exploitation rate of a single pot (Figure  
417 10). We compared this with average soak times and the correlation was high (0.75) suggesting  
418 that changes in soak times may be a major factor related to change in trap catchability. The  
419 effort-weighted average of  $q_{y,i}$  is also an estimate of the average potential pot fishing area and  
420 the model results indicate that this has ranged from 46 000 to 79 000 m<sup>2</sup> and has increased over  
421 time (unreported results). Recall that these values are upper bounds because they are based on  
422 the assumption that an entire grid is snow crab "grounds". We computed the average  $E_{max}$  over  
423 years for each grid cell,  $\bar{E}_{max,i} = A_i/q_i\bar{q}_y$ , and  $\bar{q}_y$  is the average of  $q_{1997}, \dots, q_{2014}$  (Figure 11).  
424 Regions where the area fished by a pot is higher, as indicated in Figure 9, have lower values for  
425  $E_{max}$ . Typical levels of effort (i.e. hundreds of pots per grid cell per week) are far less than these  
426 saturation levels.

427 The M4 model harvest rates and those from the stock assessment were usually similar (Figure  
428 12). When computing harvest rates from stock assessment estimates of commercial male  
429 biomass, we also projected post-season biomass from year  $y-1$  to beginning of year biomass in  
430 year  $y$  using  $\exp(-18M/52)$  to account for natural mortality between the time of the survey and  
431 the beginning of the following year. For  $M = 0.4$  this results in a 13% reduction in biomass and a  
432 15% increase in harvest rates compared to stock assessment values (see Figure 9 in DFO, 2016).  
433 Although similar in magnitude to values from the stock assessment, the M4 harvest rates were  
434 more variable. If the goal of snow crab fisheries management is a constant harvest rate then the

435 M4 model indicates that past quotas have resulted in landings and harvest rates that were more  
436 variable than required, especially the large decreases in 2010 and 2011 (Figure 2). Landings were  
437 reduced from around 25 Kt in 2008-2009 to 10 Kt in 2010-2011 which resulted in a sharp drop in  
438 M4 exploitation rates, well below potential targets. The M4 model suggests that quotas around  
439 15 Kt in 2010 and 2011 would have resulted in more constant harvest rates.

440

## 441 **Discussion**

442 We developed a spatial depletion model for sGSL snow crab based on high resolution (weekly  
443 and spatial) catch and effort data, and spatial survey data, to provide more precise estimates of  
444 snow crab biomass and population dynamics. Additional information on population size is  
445 obtained from the high resolution temporal information by using a within-season biomass  
446 depletion model based on the change in fishery CPUE. Such information is often not utilized in  
447 stock assessments. However, there are many problems when interpreting CPUE as an index of  
448 stock size and we had to incorporate spatio-temporal changes in CPUE catchability. Nonetheless  
449 our model did provide somewhat different trends in a few years compared to the snow crab  
450 survey that the stock assessment is based on. In particular, our model suggested that some  
451 historic reductions in fishery quotas may not have been completely warranted if the objective of  
452 snow crab fisheries management was a constant harvest rate near some target level. Also, our  
453 model results provide additional information about how CPUE catchability may change and  
454 problems with interpreting CPUE as an index of stock size. Some aspects of the model we  
455 proposed are specific to sGSL snow crab, or to cases where the fishing gear catches most or all  
456 of the fish available to the gear. However, we suggest the basic approach can be used for other



457 stocks where: 1) fish movement within the time-frame of the fishery is expected to be low, 2)  
458 high resolution spatio-temporal catch and effort data are available, and 3) high resolution spatial  
459 data from a research survey are available.

460 The original motivation for this research was to validate a spatial CPUE depletion model for  
461 application with Newfoundland and Labrador (NL) snow crab. The depletion model was  
462 originally formulated with a spatially and temporally constant  $q$  and without density-dependence  
463 in  $q$  or saturation effects. It was felt that sGSL snow crab was a good test case for that model  
464 because there were independent estimates of exploitable biomass from the snow crab survey to  
465 compare with depletion model estimates. Research surveys in NL have low catchability for snow  
466 crab and did not provide a useful comparison. However, this original model based only on  
467 seasonal CPUE did not produce plausible values for sGSL snow crab because of the general lack  
468 of a CPUE depletion signal for that fishery. This is very different from NL snow crab that often  
469 does show a decline in CPUE throughout the fishing season (e.g. see Figures 47, 61, and 78 in  
470 Mullaney et al 2016). We then extended the sGSL model to integrate the snow crab survey  
471 catches which provided a basis to estimate various changes (i.e. seasonal, spatial, annual,  
472 density-dependent) in CPUE catchability.

473 Our results indicate that there have been substantial changes in sGSL snow crab CPUE  
474 catchability and these changes seem to be linked with changes in soak time, among other  
475 reasons. There is strong evidence of density-dependence in the relationship with CPUE and stock  
476 biomass, in addition to a general increase in CPUE catchability over time, very consistent with  
477 changes in fishery soak times across years, and spatial variation in catchability. However, we did  
478 not find evidence of CPUE saturation. Hence, our model has contributed additional insights into  
479 problems with inferring sGSL snow crab resource status based on CPUE. There are many factors

480 influencing CPUE in addition to stock size. This has been a source of uncertainty in the  
481 assessment of this stock (DFO 2016). In addition, the model also indicated potential snow crab  
482 survey year effects in 2009 for exploitable biomass and 2011 for total biomass, a problem that  
483 has been identified in snow crab assessments (e.g. DFO 2012).

484 We used our model and additional information from the snow crab survey on soft-shelled male  
485 biomass that will recruit to the fishery in the following year to infer that  $M = 0.4$  was a more  
486 plausible value. This implies 5% survival at the 7.7 year male longevity estimate of Fonseca et  
487 al. (2008) which is reasonably consistent. At  $M = 0.3$  the survival at 7.7 years is 10%. We  
488 showed that with  $M = 0.4$  our model provided a good fit to both M-projected beginning of year  
489 commercial biomass and post-season residual biomass from the DFO kriging analyses of the  
490 snow crab survey. It has often been noted in sGSL snow crab assessments (e.g. DFO, 2016) that  
491 the estimated commercial biomass from the survey tends to be higher than the sum of the  
492 residual biomass and the landings of the following year, and the difference is attributed to a  
493 number of factors. Our results indicate that additional natural mortality at  $M = 0.4$  between the  
494 end of the post-season survey and the start of the fishery in the following year can explain this  
495 difference, but  $M = 0.3$  does not completely explain the discrepancy.

496 The standard deviation of the process error ( $\sigma_\delta$ ) was estimated to be large for all model  
497 formulations we investigated. This reflects measurement error in effort and also within-season  
498 changes in catchability. Reducing this source of error will require additional information from  
499 the snow crab fishing fleet, such as accurate measurements of effort and soak time, pot spacing,  
500 and other characteristics that may affect catch rates. Such information could be used as  
501 covariates to model and explain some of the variation in catchability, and this seems like a useful  
502 area for future research. Nonetheless, the model uncertainty in estimates of stock size and harvest

503 rates is low, with CV's in the 2-3% range, which is consistent with the low retrospective  
504 variation we found in model M4. Hence, reducing the process error variation may not lead to  
505 more precise estimates of stock size or harvest rates.

506 The low CV's on stock size and harvest rates may seem unrealistic given the highly flexible  
507 model with a CPUE  $q$  that may vary across years, spatial locations, weeks, and stock density.  
508 However, the model used a large amount of data, nearly 21 000 catch and effort measurements  
509 and 4200 survey catch values, and the model was constrained to fit the catches very closely,  
510 which will result in low CV's. There is some evidence of model mis-specification in the lack of  
511 fit to the larger catches from the snow crab survey which leads to an overall average positive  
512 residual in most years. This suggests that the CV's may be too small because they do not reflect  
513 uncertainty due to model mis-specification. There are many zero catches in the snow crab survey  
514 that the model also does not fit well. This could be due to site effects in the fixed-station survey  
515 sampling design or other operational issues. There have been changes in snow crab survey  
516 vessels during 1997 to 2015, including a change in 2013 from the vessel which had been used  
517 during 2003 to 2012. Comparative fishing experiments were not conducted to assess if there  
518 were changes in catchability between vessels; however, Benoît and Cadigan (2016) found some  
519 evidence of a change in vessel catchability by comparing snow crab survey catches from those  
520 from the DFO multi-species survey. Hence, changes in survey catch rates over time may not  
521 accurately reflect changes in stock size if survey vessels have different catchabilities and this is  
522 another source of unaccounted variability.

523 We assumed that all snow crab survey sets occurred in a single week. In reality the survey takes  
524 around 80 days to complete. It would be fairly simple to use the week a grid cell was sampled  
525 because the model predicts weekly and spatial biomass.

526 To simplify the model we treated stock dynamics separately each year and no connection is  
527 made between biomass from year to year. Roa-Ureta (2012) took a similar approach. However,  
528 population dynamics are of course linked across years, and the biomass remaining at the end of a  
529 year is part of the biomass at the start of the next year in addition to recruitment. Linking across  
530 years creates additional challenges related to movement between grid cells which may be  
531 ignored within a single fishing season because of the low mobility of sGSL snow crab (Biron et  
532 al. 2008), but movement will become more of an issue if crab dynamics are modelled over longer  
533 time-scales. Linking across years will also create other computational problems related to  
534 reduced sparseness of the covariance matrix of the random effects. This may significantly  
535 increase model run times, which are about 6-7 minutes for model formulations we investigated.

536 Best stock assessment practise should involve an assessment model that integrates all of the  
537 relevant productivity data for sGSL snow crab, including fishery CPUE, both the snow crab and  
538 DFO multi-species trawl survey catch rates, and their stage (maturity/length) composition  
539 information. Shell-stage (new/intermediate/old) compositions for mature males can provide  
540 important information on mortality rates. An assessment model should include a growth  
541 component between immature and newly mature soft-shell crabs and legal-sized hard-shell  
542 males. This will provide some capability to provide short-term projections for tactical harvest  
543 advice. A spatial assessment model can better integrate data collected at varying spatial scales  
544 and provide spatial management advice which is an issue for sGSL snow crab (e.g. DFO, 2016).  
545 The results of this paper provide a good basis for further development of such an integrated  
546 spatial stock assessment model.

547

548 **References**

549 Note that most DFO documents cited in this paper are freely available online, at [http://www.dfo-](http://www.dfo-mpo.gc.ca/csas-sccs/index.htm)  
550 [mpo.gc.ca/csas-sccs/index.htm](http://www.dfo-mpo.gc.ca/csas-sccs/index.htm).

551

552 Bacheler, N.M., Bartolino, V. and Reichert, M.J., 2013. Influence of soak time and fish  
553 accumulation on catches of reef fishes in a multispecies trap survey. *Fishery Bulletin*, 111: 218-  
554 232.

555

556 Benoît, H.P., and Cadigan, N. 2016. Trends in the biomass, distribution, size composition and  
557 model-based estimates of commercial abundance of snow crab (*Chionoecetes opilio*) based on  
558 the multi-species bottom trawl survey of the southern Gulf of St. Lawrence, 1980-2014. DFO  
559 Can. Sci. Advis. Sec. Res. Doc. 2015/084. v + 25 p.

560

561 Biron, M., Ferron, C. and Moriyasu, M. 2008. Movement of adult male snow crab, *Chionoecetes*  
562 *opilio*, in the southern Gulf of St. Lawrence and eastern Nova Scotia, Canada. *Fisheries*  
563 *Research*, 91: 260-270.

564

565 Bishir, J. and Lancia, R.A., 1996. On catch-effort methods of estimating animal abundance.  
566 *Biometrics* 52: 1457-1466.

567

568 Cadigan, N.G. 2016. A state-space stock assessment model for northern cod, including under-  
569 reported catches and variable natural mortality rates. *Canadian Journal of Fisheries and Aquatic*  
570 *Sciences* 73: 296-308.

571

572 Chaput, G., and DFO/Industry Joint Working Group. 2014. Evaluation of compliance to the  
573 Precautionary Approach of harvest decision rules for the snow crab (*Chionoecetes opilio*) stock  
574 of the southern Gulf of St. Lawrence. DFO Can. Sci. Advis. Sec. Res. Doc. 2014/083. v + 57 p.

575

576 Charles, Colin, Darren Gillis, and Elmer Wade. Using hidden Markov models to infer vessel  
577 activities in the snow crab (*Chionoecetes opilio*) fixed gear fishery and their application to catch  
578 standardization. *Canadian Journal of Fisheries and Aquatic Sciences* 71: 1817-1829.

- 579
- 580 Conan, G.Y., and Comeau, M. 1986. Functional maturity of male snow crab, (*Chionoecetes*  
581 *opilio*). Canadian Journal of Fisheries and Aquatic Sciences, 43: 1710-1719.
- 582
- 583 Cooke, J.G. and Beddington, J.R., 1984. The relationship between catch rates and abundance in  
584 fisheries. Mathematical Medicine and Biology, 1: 391-405.
- 585
- 586 DeLury, D.B., 1947. On the estimation of biological populations. Biometrics, 3: 145-167.
- 587
- 588 DFO. 2012. Assessment of snow crab in the southern Gulf of St. Lawrence (Areas 12, 19, 12E  
589 and 12F) and advice for the 2012 fishery. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep.  
590 2012/003.
- 591
- 592 DFO. 2016. Assessment of snow crab (*Chionoecetes opilio*) in the southern Gulf of St. Lawrence  
593 (Areas 12, 19, 12E and 12F) and advice for the 2016 fishery. DFO Can. Sci. Advis. Sec. Sci.  
594 Advis. Rep. 2016/010.
- 595
- 596 Fonseca, D.B., Sainte-Marie, B., and Hazel, F. 2008. Longevity and change in shell condition of  
597 adult male snow crab *Chionoecetes opilio* inferred from dactyl wear and mark-recapture data.  
598 Transactions of the American Fisheries Society, 137: 1029-1043.
- 599
- 600 Gedamke, T., DuPaul, W.D. and Hoenig, J.M., 2004. A spatially explicit open-ocean DeLury  
601 analysis to estimate gear efficiency in the dredge fishery for sea scallop *Placopecten*  
602 *magellanicus*. North American Journal of Fisheries Management, 24: 335-351.
- 603
- 604
- 605
- 606

- 607 Harley, S.J., Myers, R.A. and Dunn, A., 2001. Is catch-per-unit-effort proportional to  
608 abundance? Canadian Journal of Fisheries and Aquatic Sciences, 58: 1760-1772.
- 609
- 610 Hébert, M., Benhalima, K., Miron, G., and Moriyasu, M. 2002. Molting and growth of male  
611 snow crab, *Chionoecetes opilio*, (O. Fabricius, 1788) (Crustacea: Majidae) in the southern Gulf  
612 of St. Lawrence. Crustaceana 75: 671-702.
- 613
- 614 Hébert, M., Wade, E., DeGrâce, P., and Moriyasu, M. 2016a. Review of the 2014 snow crab  
615 (*Chionoecetes opilio*) fishery in the southern Gulf of St. Lawrence (Areas 12, 19, 12E and 12F).  
616 DFO Can. Sci. Advis. Sec. Res. Doc. 2015/081. v + 43 p.
- 617
- 618 Hébert, M., Wade, E., DeGrâce, P., and Moriyasu, M. 2016b. The 2014 assessment of the snow  
619 crab (*Chionoecetes opilio*) stock in the southern Gulf of St. Lawrence (Areas 12, 19, 12E  
620 and 12F). DFO Can. Sci. Advis. Sec. Res. Doc. 2015/083. v + 43 p.
- 621
- 622 Hilborn, R. and Walters, C.J., 1987. A general model for simulation of stock and fleet dynamics  
623 in spatially heterogeneous fisheries. Canadian Journal of Fisheries and Aquatic Sciences, 44:  
624 1366-1369.
- 625
- 626 Kristensen, K., Thygesen, U.H., Andersen, K.H. and Beyer, J.E., 2013. Estimating spatio-  
627 temporal dynamics of size-structured populations. Canadian Journal of Fisheries and Aquatic  
628 Sciences: 326-336.
- 629
- 630 Kristensen, K., Nielsen, A., Berg, C.W., Skaug, H. and Bell, B., 2015. Template model builder  
631 TMB. Journal of Statistical Software, 70: 1-21.
- 632
- 633 Leslie, P.H., 1952. The estimation of population parameters from data obtained by means of the  
634 capture-recapture method. II. The estimation of total numbers. Biometrika, 39: 363-388.
- 635

- 636 Loch, J.S., Moriyasu, M. and Jones, J.B., 1995. An improved link between industry,  
637 management and science: review of case history of the Southwestern Gulf of St. Lawrence snow  
638 crab fishery. *Aquatic Living Resources*, 8: 253-265.
- 639
- 640 Maunder, M. N., and Piner, K. R. 2014. Contemporary fisheries stock assessment: many issues  
641 still remain. *ICES Journal of Marine Science*, doi:10.1093/icesjms/fsu015.
- 642
- 643 Mesnil, B. and Rochet, M.J., 2010. A continuous hockey stick stock–recruit model for estimating  
644 MSY reference points. *ICES Journal of Marine Science: Journal du Conseil*, 67: 1780-1784.
- 645
- 646 Mullaney, D. Dawe, E., Coffey, W., Quilty, S., Colbourne, E., and Maddock Parsons, D. 2016.  
647 An Assessment of Newfoundland and Labrador Snow Crab (*Chionoecetes opilio*) in 2014. DFO  
648 Can. Sci. Advis. Sec. Res. Doc. 2016/026. v+179 p.
- 649
- 650 Nielsen, A., Berg, C. W. 2014. Estimation of time-varying selectivity in stock assessments using  
651 state-space models. *Fisheries Research*, 158, 96-101..
- 652
- 653 Miller, R.J., 1990. Effectiveness of crab and lobster traps. *Canadian Journal of Fisheries and*  
654 *Aquatic Sciences*, 47: 1228-1251.
- 655
- 656 Miller, R.J. and Mohn, R.K., 1993. Critique of the Leslie method for estimating sizes of crab and  
657 lobster populations. *North American Journal of Fisheries Management*, 13: 676-685.
- 658
- 659 Moriyasu, M., Wade, E., Landry, J.F., DeGrâce, P., Surette, T., and Hébert, M. 2016. Summary  
660 of 2014 snow crab trawl survey activities in the southern Gulf of St. Lawrence. DFO Can. Sci.  
661 Advis. Sec. Res. Doc. 2015/082. v + 39 p.
- 662
- 663 R Core Team (2016). R: A language and environment for statistical computing. R Foundation for  
664 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 665



- 666 Roa-Ureta, R.H., 2012. Modelling in-season pulses of recruitment and hyperstability-  
667 hyperdepletion in the *Loligo gahi* fishery around the Falkland Islands with generalized depletion  
668 models. ICES Journal of Marine Science: Journal du Conseil, 69:1403-1415.
- 669
- 670 Roa-Ureta, R.H., 2015. Stock assessment of the Spanish mackerel (*Scomberomorus commerson*)  
671 in Saudi waters of the Arabian Gulf with generalized depletion models under data-limited  
672 conditions. Fisheries Research, 171: 68-77.
- 673
- 674 Robert, M., Faraj, A., McAllister, M.K. and Rivot, E., 2010. Bayesian state-space modelling of  
675 the De Lury depletion model: strengths and limitations of the method, and application to the  
676 Moroccan octopus fishery. ICES Journal of Marine Science: Journal du Conseil, 67: 1272-1290.
- 677
- 678 Sainte-Marie, B., Raymond, S., and Brêthes, J.-C. 1995. Growth and maturation of the benthic  
679 stages of male snow crab, *Chionoecetes opilio* (Brachyura: Majidae). Canadian Journal of  
680 Fisheries and Aquatic Sciences, 52: 903-924.
- 681
- 682
- 683 Siddeek, M.S.M., Wade, E. and Moriyasu, M., 2009. Proposed harvest control rule for managing  
684 the snow crab stock in the southwestern Gulf of St. Lawrence, Canada. Fisheries Research: 268-  
685 279.
- 686
- 687 Skaug H.J., and Fournier D.A. 2006. Automatic approximation of the marginal likelihood in  
688 non-Gaussian hierarchical models. Computational Statistics & Data Analysis, 51: 699-709.
- 689
- 690 Smith, M.T. and Addison, J.T., 2003. Methods for stock assessment of crustacean fisheries.  
691 Fisheries Research, 65: 231-256.
- 692
- 693
- 694 Surette, T.J. and Wade, E., 2006. Bayesian serial linear regression models for forecasting  
695 the short-term abundance of commercial snow crab (*Chionoecetes opilio*). Canadian Technical  
696 Report of Fisheries and Aquatic Sciences, 2672: vi + 33 p.

- 697
- 698 Ward, H.G., Askey, P.J. and Post, J.R., 2013. A mechanistic understanding of hyperstability in  
699 catch per unit effort and density-dependent catchability in a multistock recreational fishery.  
700 Canadian Journal of Fisheries and Aquatic Sciences, 70: 1542-1550.
- 701
- 702 Walter, J.F., Hoenig, J.M. and Gedamke, T., 2007. Correcting for effective area fished in fishery-  
703 dependent depletion estimates of abundance and capture efficiency. ICES Journal of Marine  
704 Science: Journal du Conseil, 64: 1760-1771.
- 705
- 706 Zhou, S., Dichmont, C., Burridge, C.Y., Venables, W.N., Toscas, P.J. and Vance, D., 2007. Is  
707 catchability density-dependent for schooling prawns?. Fisheries Research, 85: 23-36.
- 708
- 709 Zhou, S., Punt, A.E., Deng, R. and Bishop, J., 2011. Estimating multifleet catchability  
710 coefficients and natural mortality from fishery catch and effort data: comparison of Bayesian  
711 state-space and observation error models. Canadian Journal of Fisheries and Aquatic Sciences,  
712 68: 1171-1181.
- 713
- 714 Zhou, S., Buckworth, R.C., Ellis, N., Deng, R.A. and Pascoe, S., 2015. Getting all information  
715 out of logbooks: estimating banana prawn fishable biomass, catchability, and fishing power  
716 increase, with a focus on natural mortality. ICES Journal of Marine Science: Journal du Conseil,  
717 72: 54-61.
- 718

719 **Figure Captions**

720 Figure 1. Locations of snow crab (*Chionoecetes opilio*) fishing grounds and management areas in  
721 the southern Gulf of St. Lawrence. PEI indicates the province of Prince Edward Island. 20 and  
722 200 meter depth contours are shown with heavy grey lines.

723 Figure 2. Landings (tonnes) by fishery management areas in the southern Gulf of St. Lawrence  
724 snow crab fishery, 1969 to 2015.

725 Figure 3. CPUE and average soak time versus week for the 2014 snow crab fishery.

726 Figure 4. Weekly catch per unit of effort (CPUE) of snow crab in the southern Gulf of St.  
727 Lawrence in 2014. Colors of spatial grids correspond to CPUE levels, as indicated in the legend  
728 on the right-hand side. Weeks are indicated in the top-left corner of each panel. Darkest red grids  
729 indicate  $CPUE > 110$  Kg/pot which is the 95th percentile.

730 Figure 5. Illustration of two models for effort saturation. Effective effort is the number of  
731 independent fishing locations, with a maximum of  $E_{max} = 2000$ . PP (Poisson Process) indicates  
732 randomly chosen fishing locations and HS (hockey-stick) indicates locations chosen to avoid  
733 existing gear. Dashed grey lines are references for  $E_{max}$  and  $E_{max}/2$ , and the 1:1 line. Vertical  
734 lines indicate the nominal effort corresponding to  $E_{max}/2$  effective effort.

735 Figure 6. Log CPUE (total catch divided by total effort) versus log mean biomass for all weeks  
736 in each grid cell. Panels correspond to years. Biomass values were obtained from model M1 (see  
737 Table 2). The dashed reference lines indicate average log CPUE. The grey lines are linear  
738 regressions of log CPUE versus log biomass. Red lines are linear regressions with the slope fixed  
739 at one (i.e. no density dependence).

740 Figure 7. DFO estimates of biomass from the snow crab survey (SCS; green and blue lines) and  
741 M4 model estimates (red and black lines). Colors correspond to sources and shaded regions  
742 indicate 95% confidence intervals. CPUE is shown using dashed lines. Horizontal lines indicate  
743 the series average. SCS biomass values were obtained using kriging estimation (see Hébert et al.  
744 2016b

745 Figure 8. Snow crab survey (SCS) M4 residuals versus predicted values. The red curve is  
746 smoother of residuals versus biomass.

747 Figure 9. Spatial effects in snow crab catchability. Colors of spatial grids correspond catchability  
748 effects, as indicated in the legend on the right hand side and directly on the figure. Darkest red  
749 grids indicate effects > 95th percentile.

750 Figure 10. Pot catchability (solid lines) and soak time (dashed lines) versus year. Catchability is  
751 effort-weighted across grid cells and standardized by grid area.

752 Figure 11. Annual average maximum effective effort (pots x 1000). Colors of spatial grids  
753 correspond to effort, as indicated in the legend on the right hand side and directly on the figure.  
754 Darkest red grids indicate effort > 5 000 pots.

755 Figure 12. Harvest rate estimates from the M4 model and values based on projected stock  
756 assessment estimates of snow crab survey (SCS) commercial biomass. Colors correspond to  
757 sources and shaded regions indicate 95% confidence intervals. Horizontal lines indicate the  
758 series average.

Table 1. Acronyms, model notation and parameters.

<i>Acronyms and notation</i>	
sGSL	Southern Gulf of St. Lawrence
CW	Carapace width
TAC	Total allowable catch
ME	Measurement error
SD	Standard deviation
CV	Coefficient of Variation
CPUE, X	Catch per unit of effort
$C_{\text{obs}}$	Reported fishery Catch
$E_{\text{obs}}$	Reported fishery effort
$q$	CPUE Catchability/potential area fished
$E_{\text{max}}$	Maximum effective effort
$E_o$	Nominal fishing effort without adjustment for saturation
$S_{\text{obs}}$	Snow crab survey trawl catch (biomass)
$A$	Grid cell area
$p$	Fraction of A that is crab grounds
$a_f$	Area fished by a pot
$y$	year
$i$	Grid cell, $i=1, \dots, G$
$G$	Total number of grid cells.
$t$	week
<i>GMRF</i>	Gaussian Markov random field
$\varphi_N$	Standard normal probability density function
$\theta$	Fixed effect model parameters
$\Gamma$	Model random effects
<i>model parameters (P) and random effects (RE)</i>	
$F$	Fishing mortality rate (RE)
$\mu_F$	mean $F$ (P)
$\sigma_F$	CV $F$ (P)
$q_y$	Catchability year effects (RE)
$\sigma_{qy}$	CV for $q_y$ (P)
$\delta$	Weekly process errors in catchability and ME in effort (RE)
$\sigma_\delta$	SD for $\delta$ (P)
$\varphi_\delta$	Auto-correlation for $\delta$ (P)
$q_i$	Catchability spatial effects (RE)
$\tau_{q_i}, \vartheta_{q_i}$	Precision matrix parameters for $q_i$ (P)
$D_{y,o,i}$	Initial biomass density (RE)
$\tau_{D1}, \vartheta_{D1}$	$\Psi$ parameters for $D_{1,o,i}$ in first year (P)
$\tau_{D2}, \vartheta_{D2}$	$\Psi$ parameters for log-deviations in $D_{y,o,i}$ in other years (P)
$\beta$	Density-dependence catchability parameter (P)
$\sigma_C$	Catch measurement error CV (P)
$\sigma_S$	Standard deviation of snow crab survey catch (P)

*model derived quantities*

$Z$	Total mortality rate, $Z = F + M$
$\Psi$	Spatial precision matrix
$q_{y,t,i}$	CPUE catchability in year $y$ , week $t$ and grid cell $i$
$B_{y,o,i}$	Initial annual biomass, $B_{y,o,i} = D_{y,o,i}A_i$
$C_{y,t,i}$	Catch
$H_{y,t,i}$	Harvest rate, $H_{y,t,i} = C_{y,t,i}/B_{y,t,i}$
$E_{y,t,i}$	Model predicted effort
$D_{y,o,i}$	Initial biomass density,

*fixed model inputs*

$M$	Natural mortality rate
$\gamma$	Fixed parameter controlling the transition of effective effort to $E_{\max}$
$m_i$	The number of neighbors of grid cell $i$

---

Draft

Table 2. Estimates (Est) of model parameters (see Table 1) and some population quantities, with percent coefficients of variation (CVx100), for five model formulations. nll denotes the negative loglikelihood and  $\Delta$ AIC is the difference in Akaike information criterion from the minimum of the models.

	M1		M2		M3		M4		M5	
$\Delta$ AIC:	3442.1		3429.4		0		2.7		3409.7	
nll:	56899		56893		55177		55178		56883	
Quantity	Est	CV%	Est	CV%	Est	CV%	Est	CV%	Est	CV%
$q_{1997} (x10^{-4})$	6.609	18.7	6.568	18.8	6.474	13.8	6.439	13.8	6.454	18.7
$\beta$	-	-	-	-	0.387	2.0	0.388	1.9	-	-
$\mu_F$	0.053	1.3	0.053	1.3	0.078	2.1	0.076	2.0	0.052	1.3
$\sigma_F$	1.149	0.5	1.149	0.5	1.130	0.7	1.130	0.7	1.149	0.5
$\sigma_{qy}$	0.090	23.0	0.090	23.0	0.106	18.3	0.106	18.3	0.089	23.1
$\sigma_\delta$	0.474	0.6	0.474	0.6	0.463	0.6	0.464	0.6	0.474	0.6
$\sigma_S$	1.409	1.4	1.409	1.4	1.145	4.4	1.153	3.6	1.409	1.5
$\tau_{D1}$	0.614	10.2	0.610	10.2	0.511	11.0	0.514	11.0	0.611	10.2
$\vartheta_{D1}$	0.000	141.8	0.000	141.8	0.000	141.9	0.000	141.9	0.000	141.8
$\tau_{D2}$	3.936	4.8	3.932	4.8	1.440	5.4	1.467	5.1	4.016	5.0
$\vartheta_{D2}$	0.042	24.5	0.042	24.5	0.090	22.1	0.087	21.9	0.041	24.8
$\tau_q$	0.543	11.4	0.535	11.4	3.913	13.3	3.911	13.3	0.538	11.4
$\vartheta_q$	0.227	52.9	0.229	52.8	0.110	66.6	0.109	66.7	0.229	52.8
$\varphi_q$	0.359	2.8	0.360	2.8	0.289	3.2	0.288	3.1	0.359	2.8
$B_{2014}$ (Kt)	66.815	1.9	66.802	1.9	56.498	2.3	59.792	2.3	70.799	1.9
$B_{2014}/\bar{B}$	1.118	1.8	1.118	1.8	1.116	2.1	1.119	2.1	1.119	1.8
$H_{2014}$ (%)	34.803	1.9	34.810	1.9	41.155	2.3	38.887	2.3	32.844	1.9
$H_{2014}/\bar{H}$	1.047	1.8	1.047	1.8	1.049	2.1	1.047	2.1	1.046	1.8
$a_{f,2014} (x10^{-4})$	7.315	3.4	7.269	3.4	7.566	2.3	7.521	2.3	7.146	3.4
$a_{f,2014}/\bar{a}_f$	1.052	2.8	1.052	2.8	1.128	2.0	1.128	2.0	1.053	2.8

M1: No density dependence ( $\beta = \mathbf{1}$ ), effort saturation,  $M = 0.3$ .

M2: No density dependence ( $\beta = \mathbf{1}$ ), no effort saturation,  $M = 0.3$ .

M3: Density dependence ( $\beta$  estimated), no effort saturation,  $M = 0.3$ .


M4: Density dependence, no effort saturation,  $M = 0.4$ .

M5: No density dependence, no effort saturation,  $M = 0.4$ .


Table 3. Southern Gulf of St. Lawrence snow crab M4 model estimates (Est) and coefficients of variation (CV, %) of 1) beginning of year (i.e. initial) biomass, 2) post-season biomass, 3) harvest rates, and 4) area fished. See Table 2 footnotes for M4 formulation.

Year	Initial Biomass (Kt)		Post-Season Biomass (Kt)		Harvest Rate (%)		Area Fished ( $\times 10^4$ m <sup>2</sup> )	
	Est	CV%	Est	CV%	Est	CV%	Est	CV%
1997	57.3	2.9	28.3	4.5	29.9	2.9	6.3	2.3
1998	53.1	2.5	28.3	3.7	25.7	2.5	5.8	2.4
1999	57.4	2.7	30.2	3.9	26.6	2.7	6.0	2.5
2000	37.7	2.6	12.3	6.1	48.2	2.6	5.2	2.2
2001	40.7	2.5	14.7	5.3	44.8	2.5	5.9	2.3
2002	46.7	1.9	12.4	5.5	55.5	1.9	5.5	2.2
2003	57.7	2.6	25.6	4.4	35.1	2.6	6.1	2.4
2004	66.8	1.9	22.5	4.4	47.0	1.9	6.6	2.1
2005	75.1	1.7	26.0	3.7	46.1	1.7	7.4	2.0
2006	57.8	1.9	20.5	4.0	46.2	1.9	7.5	2.1
2007	61.3	2.1	25.1	4.0	39.6	2.1	7.3	2.3
2008	53.9	2.5	21.5	4.7	40.1	2.5	6.7	2.3
2009	47.4	1.9	16.4	4.1	46.2	1.9	6.5	2.1
2010	30.7	3.0	15.4	4.7	29.9	3.0	7.0	2.7
2011	54.4	3.2	32.4	4.2	19.1	3.2	6.7	2.9
2012	59.6	2.2	26.9	3.8	35.5	2.2	8.3	2.3
2013	65.2	2.0	27.5	3.7	38.1	2.0	9.0	2.3
2014	59.8	2.3	24.5	4.3	39.0	2.3	7.5	2.3






140x125mm (150 x 150 DPI)




Landings (tonnes) by fishery management areas in the southern Gulf of St. Lawrence snow crab fishery, 1969 to 2015.

275x179mm (150 x 150 DPI)



CPUE and average soak time versus week for the 2014 snow crab fishery.



Weekly catch per unit of effort (CPUE) of snow crab in the southern Gulf of St. Lawrence in 2014. Colors of spatial grids correspond to CPUE levels, as indicated in the legend on the right-hand side. Weeks are indicated in the top-left corner of each panel. Darkest red grids indicate CPUE > 110 Kg/pot which is the 95th percentile.







Illustration of two models for effort saturation. Effective effort is the number of independent fishing locations, with a maximum of  $E_{max} = 2000$ . PP (Poisson Process) indicates randomly chosen fishing locations and HS (hockey-stick) indicates locations chosen to avoid existing gear. Dashed grey lines are references for  $E_{max}$  and  $E_{max}/2$ , and the 1:1 line. Vertical lines indicate the nominal effort corresponding to  $E_{max}/2$  effective effort.






DFO estimates of biomass from the snow crab survey (SCS; green and blue lines) and M4 model estimates (red and black lines). Colors correspond to sources and shaded regions indicate 95% confidence intervals. CPUE is shown using dashed lines. Horizontal lines indicate the series average. SCS biomass values were obtained using kriging estimation (see Hébert et al. 2016b).




Snow crab survey (SCS) M4 residuals versus predicted values. The red curve is smoother of residuals versus biomass.







Spatial effects in snow crab catchability. Colors of spatial grids correspond catchability effects, as indicated in the legend on the right hand side and directly on the figure. Darkest red grids indicate effects > 95<sup>th</sup> percentile.



Pot catchability (solid lines) and soak time (dashed lines) versus year. Catchability is effort-weighted across grid cells and standardized by grid area.



Annual average maximum effective effort (pots x 1000). Colors of spatial grids correspond to effort, as indicated in the legend on the right hand side and directly on the figure. Darkest red grids indicate effort > 5 000 pots.



Harvest rate estimates from the M4 model and values based on projected stock assessment estimates of snow crab survey (SCS) commercial biomass. Colors correspond to sources and shaded regions indicate 95% confidence intervals. Horizontal lines indicate the series average.




Figure S1. Observed versus predicted snow crab survey exploitable biomass catch per tow (grey '+'s). Panels correspond to years. Predicted biomass values were obtained from model M4 (see Table 2). Thin solid lines indicates 1:1 lines, and thick solid lines indicate smoothers.




Figure S2. Observed and predicted total annual effort. Shaded regions indicate 95% confidence intervals. Predicted values were obtained from model M4 (see Table 2).

Draft




Figure S3. Observed versus predicted total weekly effort. Panels correspond to years. Predicted values were obtained from model M4 (see Table 2).




Figure S4. Observed versus predicted effort. Predicted values were obtained from model M4 (see Table 2). The 1:1 line is shown in red.

Draft






Figure S5. Model M4 (see Table 2) spatial results for 1997. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S6. Model M4 (see Table 2) spatial results for 1998. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S7. Model M4 (see Table 2) spatial results for 1999. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S8. Model M4 (see Table 2) spatial results for 2000. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S9. Model M4 (see Table 2) spatial results for 2001. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S10. Model M4 (see Table 2) spatial results for 2002. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S11. Model M4 (see Table 2) spatial results for 2003. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S12. Model M4 (see Table 2) spatial results for 2004. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.






Figure S13. Model M4 (see Table 2) spatial results for 2005. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S14. Model M4 (see Table 2) spatial results for 2006. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S15. Model M4 (see Table 2) spatial results for 2007. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S16. Model M4 (see Table 2) spatial results for 2008. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S17. Model M4 (see Table 2) spatial results for 2009. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S18. Model M4 (see Table 2) spatial results for 2010. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S19. Model M4 (see Table 2) spatial results for 2011. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S20. Model M4 (see Table 2) spatial results for 2012. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.






Figure S21. Model M4 (see Table 2) spatial results for 2013. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure S22. Model M4 (see Table 2) spatial results for 2014. Catch and effort are aggregated across weeks, and CPUE is the ratio of aggregated catch divided by aggregated effort. Q process errors are averaged over weeks. White cells indicate no observations.




Figure SC.1. Top panel: Total catch (tonnes) of snow crab each week (rows) and year (columns). Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile. Bottom panel: Total catch (000 tonnes) each year for all weeks.




Figure SC.2. Total annual catch (tonnes) of snow crab in each grid cell, averaged for 1997-2014. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.3a. Total annual catch (tonnes) of snow crab in each grid cell in 1997. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.3b. Catch (tonnes) of snow crab in each week and grid cell in 1997. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.3c. Catch (tonnes) of snow crab in each week and grid cell in 1997. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.4a. Total annual catch (tonnes) of snow crab in each grid cell in 1998. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.






Figure SC.4b. Catch (tonnes) of snow crab in each week and grid cell in 1998. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.4c. Catch (tonnes) of snow crab in each week and grid cell in 1998. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.5a. Total annual catch (tonnes) of snow crab in each grid cell in 1999. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.5b. Catch (tonnes) of snow crab in each week and grid cell in 1999. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.5c. Catch (tonnes) of snow crab in each week and grid cell in 1999. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.6a. Total annual catch (tonnes) of snow crab in each grid cell in 2000. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.6b. Catch (tonnes) of snow crab in each week and grid cell in 2000. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.6c. Catch (tonnes) of snow crab in each week and grid cell in 2000. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.






Figure SC.7a. Total annual catch (tonnes) of snow crab in each grid cell in 2001. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.7b. Catch (tonnes) of snow crab in each week and grid cell in 2001. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.7c. Catch (tonnes) of snow crab in each week and grid cell in 2001. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.8a. Total annual catch (tonnes) of snow crab in each grid cell in 2002. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.8b. Catch (tonnes) of snow crab in each week and grid cell in 2002. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.8c. Catch (tonnes) of snow crab in each week and grid cell in 2002. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.9a. Total annual catch (tonnes) of snow crab in each grid cell in 2003. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.9b. Catch (tonnes) of snow crab in each week and grid cell in 2003. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.






Figure SC.9c. Catch (tonnes) of snow crab in each week and grid cell in 2003. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.10a. Total annual catch (tonnes) of snow crab in each grid cell in 2004. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.10b. Catch (tonnes) of snow crab in each week and grid cell in 2004. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.10c. Catch (tonnes) of snow crab in each week and grid cell in 2004. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.11a. Total annual catch (tonnes) of snow crab in each grid cell in 2005. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.11b. Catch (tonnes) of snow crab in each week and grid cell in 2005. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.11c. Catch (tonnes) of snow crab in each week and grid cell in 2005. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.12a. Total annual catch (tonnes) of snow crab in each grid cell in 2006. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.






Figure SC.12b. Catch (tonnes) of snow crab in each week and grid cell in 2006. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.12c. Catch (tonnes) of snow crab in each week and grid cell in 2006. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.13a. Total annual catch (tonnes) of snow crab in each grid cell in 2007. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.13b. Catch (tonnes) of snow crab in each week and grid cell in 2007. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.13c. Catch (tonnes) of snow crab in each week and grid cell in 2007. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.14a. Total annual catch (tonnes) of snow crab in each grid cell in 2008. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.14b. Catch (tonnes) of snow crab in each week and grid cell in 2008. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.14c. Catch (tonnes) of snow crab in each week and grid cell in 2008. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.






Figure SC.15a. Total annual catch (tonnes) of snow crab in each grid cell in 2009. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.15b. Catch (tonnes) of snow crab in each week and grid cell in 2009. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.15c. Catch (tonnes) of snow crab in each week and grid cell in 2009. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.16a. Total annual catch (tonnes) of snow crab in each grid cell in 2010. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.16b. Catch (tonnes) of snow crab in each week and grid cell in 2010. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.16c. Catch (tonnes) of snow crab in each week and grid cell in 2010. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.17a. Total annual catch (tonnes) of snow crab in each grid cell in 2011. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.17b. Catch (tonnes) of snow crab in each week and grid cell in 2011. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.






Figure SC.17c. Catch (tonnes) of snow crab in each week and grid cell in 2011. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.18a. Total annual catch (tonnes) of snow crab in each grid cell in 2012. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.18b. Catch (tonnes) of snow crab in each week and grid cell in 2012. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.18c. Catch (tonnes) of snow crab in each week and grid cell in 2012. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.19a. Total annual catch (tonnes) of snow crab in each grid cell in 2013. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.19b. Catch (tonnes) of snow crab in each week and grid cell in 2013. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.19c. Catch (tonnes) of snow crab in each week and grid cell in 2013. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.20a. Total annual catch (tonnes) of snow crab in each grid cell in 2014. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.






Figure SC.20b. Catch (tonnes) of snow crab in each week and grid cell in 2014. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SC.20c. Catch (tonnes) of snow crab in each week and grid cell in 2014. The week number is indicated in the top left-hand corner. Colors correspond to catch levels, as indicated in the legend on the right-hand side. Darkest red grids indicate catch > 98th percentile.




Figure SE.1. Top panel: Total effort (number of pots) of snow crab each week (rows) and year (columns). Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile. Bottom panel: Total effort each year for all weeks.




Figure SE.2. Total annual effort (number of pots) for snow crab in each grid cell, averaged for 1997-2014. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.3a. Total annual effort (number of pots) for snow crab in each grid cell in 1997. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.3b. Effort (number of pots) for snow crab in each week and grid cell in 1997. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.3c. Effort (number of pots) for snow crab in each week and grid cell in 1997. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.4a. Total annual effort (number of pots) for snow crab in each grid cell in 1998. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.






Figure SE.4b. Effort (number of pots) for snow crab in each week and grid cell in 1998. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.4c. Effort (number of pots) for snow crab in each week and grid cell in 1998. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.5a. Total annual effort (number of pots) for snow crab in each grid cell in 1999. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.5b. Effort (number of pots) for snow crab in each week and grid cell in 1999. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.5c. Effort (number of pots) for snow crab in each week and grid cell in 1999. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.6a. Total annual effort (number of pots) for snow crab in each grid cell in 2000. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.6b. Effort (number of pots) for snow crab in each week and grid cell in 2000. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.6c. Effort (number of pots) for snow crab in each week and grid cell in 2000. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.






Figure SE.7a. Total annual effort (number of pots) for snow crab in each grid cell in 2001. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.7b. Effort (number of pots) for snow crab in each week and grid cell in 2001. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.7c. Effort (number of pots) for snow crab in each week and grid cell in 2001. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.8a. Total annual effort (number of pots) for snow crab in each grid cell in 2002. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.8b. Effort (number of pots) for snow crab in each week and grid cell in 2002. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.8c. Effort (number of pots) for snow crab in each week and grid cell in 2002. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.9a. Total annual effort (number of pots) for snow crab in each grid cell in 2003. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.9b. Effort (number of pots) for snow crab in each week and grid cell in 2003. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.






Figure SE.9c. Effort (number of pots) for snow crab in each week and grid cell in 2003. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.10a. Total annual effort (number of pots) for snow crab in each grid cell in 2004. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.10b. Effort (number of pots) for snow crab in each week and grid cell in 2004. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.10c. Effort (number of pots) for snow crab in each week and grid cell in 2004. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.11a. Total annual effort (number of pots) for snow crab in each grid cell in 2005. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.11b. Effort (number of pots) for snow crab in each week and grid cell in 2005. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.11c. Effort (number of pots) for snow crab in each week and grid cell in 2005. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.12a. Total annual effort (number of pots) for snow crab in each grid cell in 2006. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.






Figure SE.12b. Effort (number of pots) for snow crab in each week and grid cell in 2006. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.12c. Effort (number of pots) for snow crab in each week and grid cell in 2006. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.13a. Total annual effort (number of pots) for snow crab in each grid cell in 2007. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.13b. Effort (number of pots) for snow crab in each week and grid cell in 2007. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.13c. Effort (number of pots) for snow crab in each week and grid cell in 2007. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.14a. Total annual effort (number of pots) for snow crab in each grid cell in 2008. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.14b. Effort (number of pots) for snow crab in each week and grid cell in 2008. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.14c. Effort (number of pots) for snow crab in each week and grid cell in 2008. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.






Figure SE.15a. Total annual effort (number of pots) for snow crab in each grid cell in 2009. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.15b. Effort (number of pots) for snow crab in each week and grid cell in 2009. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.15c. Effort (number of pots) for snow crab in each week and grid cell in 2009. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.16a. Total annual effort (number of pots) for snow crab in each grid cell in 2010. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.16b. Effort (number of pots) for snow crab in each week and grid cell in 2010. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.16c. Effort (number of pots) for snow crab in each week and grid cell in 2010. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.17a. Total annual effort (number of pots) for snow crab in each grid cell in 2011. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.17b. Effort (number of pots) for snow crab in each week and grid cell in 2011. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.






Figure SE.17c. Effort (number of pots) for snow crab in each week and grid cell in 2011. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.18a. Total annual effort (number of pots) for snow crab in each grid cell in 2012. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.18b. Effort (number of pots) for snow crab in each week and grid cell in 2012. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.18c. Effort (number of pots) for snow crab in each week and grid cell in 2012. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.19a. Total annual effort (number of pots) for snow crab in each grid cell in 2013. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.19b. Effort (number of pots) for snow crab in each week and grid cell in 2013. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.19c. Effort (number of pots) for snow crab in each week and grid cell in 2013. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.20a. Total annual effort (number of pots) for snow crab in each grid cell in 2014. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.






Figure SE.20b. Effort (number of pots) for snow crab in each week and grid cell in 2014. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.




Figure SE.20c. Effort (number of pots) for snow crab in each week and grid cell in 2014. The week number is indicated in the top left-hand corner. Colors correspond to effort levels, as indicated in the legend on the right-hand side. Darkest red grids indicate effort > 98th percentile.

Total annual average = 55 (kg/pot),  
346 cells; cell average = 56.96 (kg/pot),




Figure SX.1. Top panel: Total CPUE of snow crab each week (rows) and year (columns). Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile. Bottom panel: Total CPUE each year for all weeks.




Figure SX.2. Total annual CPUE of snow crab in each grid cell, averaged for 1997-2014. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.3a. Total annual CPUE of snow crab in each grid cell in 1997. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.3b. CPUE of snow crab in each week and grid cell in 1997. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.3c. CPUE of snow crab in each week and grid cell in 1997. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.4a. Total annual CPUE of snow crab in each grid cell in 1998. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.






Figure SX.4b. CPUE of snow crab in each week and grid cell in 1998. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.4c. CPUE of snow crab in each week and grid cell in 1998. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.5a. Total annual CPUE of snow crab in each grid cell in 1999. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.5b. CPUE of snow crab in each week and grid cell in 1999. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.5c. CPUE of snow crab in each week and grid cell in 1999. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.6a. Total annual CPUE of snow crab in each grid cell in 2000. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.6b. CPUE of snow crab in each week and grid cell in 2000. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.6c. CPUE of snow crab in each week and grid cell in 2000. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.






Figure SX.7a. Total annual CPUE of snow crab in each grid cell in 2001. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.7b. CPUE of snow crab in each week and grid cell in 2001. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.7c. CPUE of snow crab in each week and grid cell in 2001. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.8a. Total annual CPUE of snow crab in each grid cell in 2002. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.8b. CPUE of snow crab in each week and grid cell in 2002. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile




Figure SX.8c. CPUE of snow crab in each week and grid cell in 2002. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.9a. Total annual CPUE of snow crab in each grid cell in 2003. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.9b. CPUE of snow crab in each week and grid cell in 2003. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.






Figure SX.9c. CPUE of snow crab in each week and grid cell in 2003. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.10a. Total annual CPUE of snow crab in each grid cell in 2004. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.10b. CPUE of snow crab in each week and grid cell in 2004. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.10c. CPUE of snow crab in each week and grid cell in 2004. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.11a. Total annual CPUE of snow crab in each grid cell in 2005. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.11b. CPUE of snow crab in each week and grid cell in 2005. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.11c. CPUE of snow crab in each week and grid cell in 2005. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.12a. Total annual CPUE of snow crab in each grid cell in 2006. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.






Figure SX.12b. CPUE of snow crab in each week and grid cell in 2006. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.12c. CPUE of snow crab in each week and grid cell in 2006. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.13a. Total annual CPUE of snow crab in each grid cell in 2007. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.13b. CPUE of snow crab in each week and grid cell in 2007. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.13c. CPUE of snow crab in each week and grid cell in 2007. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.14a. Total annual CPUE of snow crab in each grid cell in 2008. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.14b. CPUE of snow crab in each week and grid cell in 2008. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.14c. CPUE of snow crab in each week and grid cell in 2008. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.






Figure SX.15a. Total annual CPUE of snow crab in each grid cell in 2009. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.15b. CPUE of snow crab in each week and grid cell in 2009. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.15c. CPUE of snow crab in each week and grid cell in 2009. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.16a. Total annual CPUE of snow crab in each grid cell in 2010. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.16b. CPUE of snow crab in each week and grid cell in 2010. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.16c. CPUE of snow crab in each week and grid cell in 2010. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.17a. Total annual CPUE of snow crab in each grid cell in 2011. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.17b. CPUE of snow crab in each week and grid cell in 2011. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.






Figure SX.17c. CPUE of snow crab in each week and grid cell in 2011. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.18a. Total annual CPUE of snow crab in each grid cell in 2012. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.18b. CPUE of snow crab in each week and grid cell in 2012. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.18c. CPUE of snow crab in each week and grid cell in 2012. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.19a. Total annual CPUE of snow crab in each grid cell in 2013. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.19b. CPUE of snow crab in each week and grid cell in 2013. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.19c. CPUE of snow crab in each week and grid cell in 2013. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.20a. Total annual CPUE of snow crab in each grid cell in 2014. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.






Figure SX.20b. CPUE of snow crab in each week and grid cell in 2014. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SX.20c. CPUE of snow crab in each week and grid cell in 2014. The week number is indicated in the top left-hand corner. Colors correspond to CPUE levels, as indicated in the legend on the right-hand side. Darkest red grids indicate CPUE > 98th percentile.




Figure SSEB.1. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 1997. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.2. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 1998. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.3. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 1999. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.4. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2000. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.5. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2001. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.6. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2002. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.






Figure SSEB.7. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2003. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.8. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2004. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.9. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2005. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.10. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2006. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.11. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2007. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.12. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2008. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.13. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2009. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.14. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2010. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.






Figure SSEB.15. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2011. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.16. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2012. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.17. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2013. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.18. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2014. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSEB.19. Snow crab survey exploitable biomass (shell conditions 3-5; tonnes per grid) results for 2015. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.1. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 1997. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.2. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 1998. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.3. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 1999. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.






Figure SSTB.4. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2000. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.5. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2001. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.6. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2002. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.7. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2003. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.8. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2004. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.9. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2005. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.10. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2006. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.11. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2007. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.






Figure SSTB.12. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2008. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.13. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2009. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.14. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2010. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.15. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2011. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.16. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2012. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.17. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2013. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.18. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2014. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.




Figure SSTB.19. Snow crab survey total biomass (shell conditions 1-5; tonnes per grid) results for 2015. Colors correspond to biomass levels, as indicated in the legend on the right-hand side. Darkest red grids indicate biomass > 98th percentile.