

Technical University of Denmark



## Data regarding the growth of *Lactobacillus acidophilus* NCFM on different carbohydrates and recombinant production of elongation factor G and pyruvate kinase

Celebioglu, Hasan Ufuk; Olesen, Sita Vaag; Prehn, Kennie; Lahtinen, Sampo J; Pedersen, Susanne Brix; Abou Hachem, Maher ; Svensson, Birte

*Published in:*  
Data in Brief

*Link to article, DOI:*  
[10.1016/j.dib.2017.07.021](https://doi.org/10.1016/j.dib.2017.07.021)

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Celebioglu, H. U., Olesen, S. V., Prehn, K., Lahtinen, S. J., Pedersen, S. B., Abou Hachem, M., & Svensson, B. (2017). Data regarding the growth of *Lactobacillus acidophilus* NCFM on different carbohydrates and recombinant production of elongation factor G and pyruvate kinase. *Data in Brief*, 14, 118-122. DOI: 10.1016/j.dib.2017.07.021

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)

## Data Article

# Data regarding the growth of *Lactobacillus acidophilus* NCFM on different carbohydrates and recombinant production of elongation factor G and pyruvate kinase



Hasan Ufuk Celebioglu<sup>a</sup>, Sita Vaag Olesen<sup>a</sup>, Kennie Prehn<sup>a</sup>, Sampo J. Lahtinen<sup>b</sup>, Susanne Brix<sup>c</sup>, Maher Abou Hachem<sup>d</sup>, Birte Svensson<sup>a,\*</sup>

<sup>a</sup> Enzyme and Protein Chemistry, Department of Biotechnology and Biomedicine, Technical University of Denmark, Denmark

<sup>b</sup> Active Nutrition, DuPont, Nutrition & Health, Finland

<sup>c</sup> Disease Systems Immunology, Department of Biotechnology and Biomedicine, Technical University of Denmark, Denmark

<sup>d</sup> Protein Glycoscience and Biotechnology, Department of Biotechnology and Biomedicine, Technical University of Denmark, Denmark

## ARTICLE INFO

## Article history:

Received 17 May 2017

Accepted 11 July 2017

Available online 14 July 2017

## ABSTRACT

The present study describes the growth of the very well-known probiotic bacterium *Lactobacillus acidophilus* NCFM on different carbohydrates. Furthermore, recombinant production of putative moonlighting proteins elongation factor G and pyruvate kinase from this bacterium is described. For further and detailed interpretation of the data presented here, please see the research article “Mucin- and carbohydrate-stimulated adhesion and subproteome changes of the probiotic bacterium *Lactobacillus acidophilus* NCFM” (Celebioglu et al., 2017) [1].

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.jprot.2017.05.015>

\* Corresponding author: Enzyme and Protein Chemistry, Department of Biotechnology and Biomedicine, Technical University of Denmark, Elektrovej, Building 375, DK-2800 Kgs. Lyngby, Denmark. Phone: +45 45252740.

E-mail address: [bis@bio.dtu.dk](mailto:bis@bio.dtu.dk) (B. Svensson).

<http://dx.doi.org/10.1016/j.dib.2017.07.021>

2352-3409/© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Specifications Table

Subject area	Biology
More specific subject area	Microbiology, Biochemistry
Type of data	Table, graph, figure
How data was acquired	Bacterial growth in batch cultures, Heterologous production of recombinant proteins
Data format	Raw and analyzed
Experimental factors	Bacterial cells were grown on different carbohydrates until stationary phase (24 h) in batch cultures. Genes encoding elongation factor G and pyruvate kinase were cloned and recombinantly produced in <i>Escherichia coli</i> cells.
Experimental features	Growth was measured at 600 nm at the stationary phase (24 h). Recombinant elongation factor G and pyruvate kinase were purified using HisPur™ Cobalt Purification Kit (Thermo Scientific), followed by SDS-PAGE to visualize the purified proteins.
Data source location	Technical University of Denmark, Kgs. Lyngby, Denmark
Data accessibility	All data are presented in this article

## Value of the data


- Growth data presented here shows growth potential of the probiotic bacterium *Lactobacillus acidophilus* NCFM on different carbon sources.
- The growth data may be used by researchers to understand the ability of the bacterium to grow on different carbon sources.
- Data regarding recombinant production of elongation factor G and pyruvate kinase from *Lactobacillus acidophilus* NCFM may be used by researchers to apply the same procedure.
- Data shown here and in Ref. [1] are useful for the researchers who are working on gut microbiota, probiotic bacteria, carbohydrate-microbe interactions, and moonlighting proteins.

## 1. Data

The extensively used probiotic bacterium *Lactobacillus acidophilus* NCFM was grown on nine different carbohydrates including growth on glucose supplemented with mucin (Fig. 1). The putative moonlighting proteins elongation factor G and pyruvate kinase also identified from this bacterium by differential proteomics [1] were recombinantly produced using the primers shown in Table 1 and purified by his-tag affinity chromatography. The purified proteins were analysed by SDS-PAGE (Fig. 2). Table 2 reports on the mass spectrometric identification of the two purified proteins without any identification of *E. coli* proteins.

## 2. Experimental design, materials and methods

*L. acidophilus* NCFM (NCFM) ( $1.50 \times 10^{10}$  CFU/g DuPont, USA Inc., Madison, US) was grown aerobically without shaking at 37 °C in batch cultures (50 mL) in semisynthetic lactic acid bacteria medium (LABSEM) [2] containing 1% FOS (Sigma-Aldrich); GOS; polydextrose (both DuPont); melibiose (Fluka); lactulose (Sigma-Aldrich); cellobiose (Sigma-Aldrich); isomaltulose (palatinose; Sigma-Aldrich); or trehalose (Sigma-Aldrich); and the reference glucose (Sigma-Aldrich). Porcine gastric mucin to 0.1% (Sigma-Aldrich) was included in cultures with 1% glucose. The bacterium was sub-



**Fig. 1.** *in vitro* evaluation of growth of *Lactobacillus acidophilus* NCFM (early stationary phase, 24 h) on different carbon sources (1%) or supplemented with mucin (0.1%). Asterisk (\*) indicates that the difference in growth of the bacteria are statistically significant compared to growth on glucose ( $p \leq 0.05$ ).

**Table 1**

Primers used for cloning of *lba0289* encoding elongation factor G and *lba0957* encoding pyruvate kinase.


Gene	Primers
<i>lba0289</i> (Elongation factor G)	<b>F:</b> CGCGGGCAGCCATATGAGGAGAGACTAATTTATGGCTAACA <b>R:</b> GCTCGAATTCGGATCCTTATTACGCGTCGCCG
<i>lba0957</i> (Pyruvate kinase)	<b>F:</b> CGCGGGCAGCCATATGGAGAGGATTATTAATAATGAAGAAAAC <b>R:</b> GCTCGAATTCGGATCCTTAAAGGTTTGAGATTTCACCTTG

**F:** Forward Primer

**R:** Reverse Primer

cultured for three cycles in LABSEM and growth was monitored at early stationary phase (24 h) by measuring absorbance at 600 nm.

Gene-specific primers for *lba0289* (elongation factor G, EF-G) and *lba0957* (pyruvate kinase, PK) with extra 15 bp complementary to the pET28a(+) vector linearized with BamHI and NdeI (Table 1) were designed using CLC Main Workbench software (Qiagen), primer blasted (NCBI), and used to amplify genes by PCR. Cloning was performed with In-Fusion Cloning kit (Clontech) per the user manual. The resulting plasmids were transformed into competent *E. coli* DH5 $\alpha$  and positive colonies were selected using kanamycin LB agars. Inserted genes were confirmed by sequencing (GATC Biotech). Soluble recombinant proteins were obtained in *E. coli* BL21 (DE3) induced with 0.1 mM isopropyl- $\beta$ -D-thiogalactoside. Cells were disrupted using a high-pressure cell homogenizer (Stanstead), followed by centrifugation (10,000  $\times$  g, 20 min). Recombinant proteins (rEF-G and rPK) were purified (HisPur™ Cobalt Purification Kit; Thermo Scientific) according to the manufacturer's instructions and verified by SDS-PAGE (Fig. 2) [3].



**Fig. 2.** SDS-PAGE of purified recombinant elongation factor G (rEF-G) and pyruvate kinase (rPK). M, molecular weight marker; L, lysate of *E. coli* BL21(DE3); FT, flow-through; Elutions, eluted proteins from HisPur Cobalt resin.

**Table 2**

MALDI-TOF MS results of recombinantly produced elongation factor G and pyruvate kinase of *Lactobacillus acidophilus* NCFM. No *E. coli* proteins were identified.

Protein name	Database	Accession number	Score	Expect	Mw/pI	Peptides matched/identified	Protein sequence coverage
rEF-G	NCBIprot	YP_193213.1	173	7.7e-11	76,806/4.94	42/169	55%
rPK	NCBIprot	YP_193840.1	231	1.2e-16	63,136/5.23	39/145	62%

Bands corresponding to elutions in Fig. 2 were excised manually, subjected to in-gel degradation by trypsin and MS protein identification. Briefly, gel pieces were washed with 40% ethanol (200  $\mu$ L, 10 min) followed by acetonitrile (ACN) (50  $\mu$ L), reduced with DTT (10 mM in 100 mM  $\text{NH}_4\text{HCO}$ ) and alkylated with iodoacetamide (55 mM in 100 mM  $\text{NH}_4\text{HCO}$ ), incubated with 12.5 ng/mL trypsin (Promega) in 10 mM ammonium bicarbonate (5  $\mu$ L, on ice, 45 min), added 10 mM ammonium bicarbonate (10  $\mu$ L), and incubated (37  $^\circ\text{C}$ , overnight). Supernatant (1  $\mu$ L) was applied onto an Anchor Chip target (Bruker-Daltonics), added matrix (1  $\mu$ L 0.5 mg/mL CHCA in 90% ACN, 0.1% TFA) and washed (2  $\mu$ L 0.02% TFA). MS spectra were obtained using an Ultraflex II MALDI-TOF MS mass spectrometer (Bruker-Daltonics) in auto-mode with Flex Control v3.0 (Bruker-Daltonics) and processed by Flex Analysis v3.0 (Bruker-Daltonics). Spectra were externally calibrated by trypsin-generated  $\beta$ -lactoglobulin peptides (5 pmol/mL). MS spectra were searched against the NCBIprot database (ver. 20170215) or SwissProt for bacteria using the MASCOT 2.0 software (<http://www.matrixscience.com>) integrated with BioTools v3.1 (Bruker-Daltonics). Protein identifications by Peptide Mass Fingerprinting (PMF) were confirmed with a MASCOT score of 80 (60 for SwissProt),  $p \leq 0.05$  and a minimum of six matched peptides.

## Acknowledgements

Karina Jansen, Anne Blicher, and Lisbeth Buus Rosholm are thanked for technical assistance. This work was supported by the Danish Strategic Research Council's Program Committee on Health, Food and Welfare (FøSu)(2101-07-105), the Danish Council for Independent Research | Natural Sciences and the Danish Center for Advanced Food Studies (LMC). HUC is grateful to the Republic of Turkey, Ministry of National Education for a PhD scholarship.

## Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2017.07.021>.

## References

- [1] H.U. Celebioglu, S.V. Olesen, K. Prehn, S.J. Lahtinen, S. Brix, M. Abou Hachem, B. Svensson, Mucin- and carbohydrate-stimulated adhesion and subproteome changes of the probiotic bacterium *Lactobacillus acidophilus* NCFM, *J. Proteom.* 163 (2017) 102–110.
- [2] A. Majumder, A. Sultan, R.R. Jersie-Christensen, M. Ejby, B.G. Schmidt, S.J. Lahtinen, S. Jacobsen, B. Svensson, Proteome reference map of *Lactobacillus acidophilus* NCFM and quantitative proteomics towards understanding the prebiotic action of lactitol, *Proteomics* 11 (2011) 3470–3481.
- [3] D.K. Patel, K.R. Shah, A. Pappachan, S. Gupta, D.D. Singh, Cloning, expression and characterization of a mucin-binding GAPDH from *Lactobacillus acidophilus*, *Int. J. Biol. Macromol.* 91 (2016) 338–346.