

Technical University of Denmark



## Synchronizing legume residue nutrient release with Kale (*Brassica oleracea* var. *acephala*) uptake in a Nitrosol of Kabete, Kenya

Onwonga, Richard N.; Chepkoech, Caroline; Wahome, R.G.; Jensen, Henning Høgh; Freyer, Bernhard

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Onwonga, R. N., Chepkoech, C., Wahome, R. G., Jensen, H. H., & Freyer, B. (2017). Synchronizing legume residue nutrient release with Kale (*Brassica oleracea* var. *acephala*) uptake in a Nitrosol of Kabete, Kenya. Poster session presented at Tropentag 2017, Bonn, Germany.

**DTU Library**  
Technical Information Center of Denmark

---

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Tropentag 2017, Bonn, Germany  
September 20-22, 2017

Conference on International Research on Food Security, Natural Resource  
Management and Rural Development  
organised by the University of Bonn, Bonn, Germany

---

## **Synchronizing Legume Residue Nutrient Release with Kale (*Brassica oleracea var. acephala*) Uptake in a Nitisol of Kabete, Kenya**

Richard N. Onwonga<sup>a</sup>, Caroline Chepkoech<sup>a</sup>, R.G. Wahome<sup>b</sup>, Henning Høgh Jensen<sup>c</sup>, Bernhard Freyer<sup>d</sup>

<sup>a</sup> University of Nairobi, Department of Land Resource Management and Agricultural Technology, Kenya

<sup>b</sup> University of Nairobi, Department of Animal Production, Kenya

<sup>c</sup> Technical University of Denmark, National Food Institute, Denmark

<sup>d</sup> University of Natural Resources and Life Sciences (BOKU), Division of Organic Farming, Austria

### **Introduction**

Nitrogen (N) and phosphorus (P) are identified as major limiting nutrients for many cropping systems (Kwambiah et al., 2003) and their application from organic and inorganic sources is essential to maximize and sustain crop yield potentials (Hartermink et al., 2000). N is commonly considered a key factor limiting crop growth in organic systems (Möller et al., 2008) and, unlike conventional farming systems, rely on management of soil organic matter to optimize crop production (Watson et al., 2002). Addition of plant residues, quantity wise and timing, has thus become a pivotal strategy in soil fertility improvement for crop production e.g. kales (*Brassica oleracea var. acephala*) under organic farming systems.

Chickpea (*Cicer arietinum*) and white lupin (*Lupinus albus L.*) are leguminous crops commonly intercropped with kales (Genga, 2014) and their influence on crop yield and soil nutrient status has been widely studied (Nduku 2014, Genga 2014; Onwonga et al., 2015). There is however a dearth of information with respect to synchronization of nutrient released by legume residues with pattern of nutrient uptake by kales to match their demand. The objective of the current study was therefore to assess decomposition and nutrient release rates of chickpea and lupin residues and kale nutrient uptake patterns for better synchrony of nutrient supply and demand.

### **Material and Methods**

**Site description:** Kabete field station (latitude 1°15'S, longitude 36° 41'E; 1940 m above sea level; mean temperature 24.3°C - maximum and 13.7°C -minimum; mean annual precipitation 1000 mm) of the University of Nairobi, is located 10 km north of Nairobi and falls in agroecological zone III (Jaetzold and Schmidt, 2006). Soils are deep red nitisols containing 70% clay particles (FAO, 1990).

**Agronomic practices, sampling and analysis:** Kale seedlings (4 week old) were grown, parallel to legume residue decomposition plots, in pure stands with application of farm yard manure (FYM) (10t/ha). All recommended agronomic practices were applied (Chepkoech, 2015). Legumes, to provide residue after harvest, were previously planted and managed as described by Lelei et al. (2014) in a neighbouring field with 10t/ha of FYM applied. Kale leaves for N and P analyses were sampled from three middle rows at 30, 60 and 90 days after planting.

**Decomposition studies:** Chickpea and white lupin residues were weighed (proportional to their dry matter production per hectare) and put in litter bags (20 × 20 × 20 cm) without chopping

them, and buried (in triplicates) in soil to a depth of between 10 and 15 cm. The litter bags were retrieved at 0, 15, 30, 45, 60, 75, 90, 105 and 120 days of incubation, residues washed with distilled water to remove soil particles and carefully transferred to khaki bags. The residues were then oven dried, grounded to pass through a 2mm sieve and analyzed for N and P content using standard laboratory procedures.


**Determination of weight loss and nutrient release rates:** The percent of dry weight remaining after each retrieval was calculated as;  $XR (\%) = (XR_t/X_0) * 100$ . Where XR = percent weight remaining,  $X_t$  = weight content at each sampling time and  $X_0$  = the starting weight values. The decomposition rates were calculated using a first-order exponential decay function,  $Y = y_0 e^{-kt}$  (Wielder and Lang 1982). Where Y = is the weight or nutrients remaining at time t,  $y_0$  = the initial weight or nutrients available before the decomposition, k = the decomposition constants, t = time of decomposition. The half-life ( $t_{1/2}$ ) was calculated using the equation;  $T_{1/2} = \ln(2)/k$ . Where  $t_{1/2}$  = time when half of the nutrients or weight is lost,  $\ln(2)$  = natural logarithm and k = decomposition rate.

## Results and Discussion



**Residue Weight loss:** Legume residue weight loss (days), with days of incubation, was rapid (0-30), moderately rapid (30-60), moderate (60-90) and gradual (90-120). About 30 (first 15 days), 50 (after 45 days) and 86.5% (at 120 days) weight loss was registered for both residues (Figure 1).

The drastic weight loss in the first 30 days could be attributed to high content of fast decomposable components such as sugars, amino acids and proteins and with progress of incubation, recalcitrant materials increase (Berg et al. (2007)). The calculated half-life for both legumes was 52 days.

**Nutrient release rates:** The N release was rapid (first 30 days), moderate (30 and 75 days) and gradual (75 to 120 days) (Figure 2A). There was a 10-33, 50 and 80 - 86.7% loss of N by chickpea and lupin within the first 15, 30-60 and 120 days, respectively. The N half-life (N release) was 20 days.



**Figure 1: Weight losses of chickpea and white lupin within 120 days of soil incorporation**




**Figure 2: Nitrogen (A) and Phosphorus (B) release of chickpea and lupin residues in 120 days of incubation**

The rapid release of N by chickpea (C/N=24) and white lupin (C/N=26) residues can be attributed to high initial N contents and lower C/N ratios (see also Danga et al., 2010). Shi (2013), reported that C (%) of decomposing residues decreased with increase of incubation days.

The P release (days) was rapid (0-15), moderately rapid (15 and 45), moderate (45 and 75) and gradual (75 and 120) for chickpea and lupin, respectively. About 50 % loss of P occurred between 15 and 30 days corresponding to 33.3 and 38.8% and 3.5 and 1.7% of initial P in lupin and chickpea (Figure 2B), (see also Mburu et al. (2013). The P half-life was 30 days.

### Kale nutrient uptake

During kale growth (Figure 3), nutrient uptake was rapid (0-30 days), followed by a moderate phase (30-60) and progressive decline (60-120 days). Nutrient uptake was significantly correlated with kale growth (Figure 4). Rapid nutrient uptake in first 30 days of growth is a factor of more




**Figure 3: Nutrient uptake by kale in 120 days of growth**


uptake.

leaves being produced and hence more nutrients needed. Highest concentration of nutrients in kale leaves was recorded between day 30 and 60 (see also Salisbury and Ross 1992).

**Legume N and P release compared with kale uptake:** The N and P release by legume residues curves (Figures 2A and B) were superimposed on kale nutrient uptake curves (Figure 3) with a view of determining the point at which they intersect, herein interpreted as optimum synchrony of nutrient release with




A



B

**Figure 4: Kale nitrogen (A) and phosphorus (B) uptake in relation with nutrient release by white lupin**

The N release by chickpea and white lupin, and uptake by kale intersected at day 25 and 30, respectively (Figure 4A and B) with the intersection of P for both legumes being at day 30. The N and P release rate were significantly correlated ( $R^2=0.970$ ) to uptake by kales (Figure 4).



**Figure 4: Kale N (A) and P (B) uptake in relation to nutrients release by chickpea**

The nutrient release and kale nutrient uptake curve superimposition results, closely mirrored the findings from the calculated half-life of N (20) and P (30). As described by Palm et al. (2001) and Rowe et al. (2004), strategies in legume-based systems to potentially reduce the rate of N supply around periods of potential asynchrony include changing the timing and placement of legume residues, manipulating residue quality through choice of legume tissue or species.

### Conclusions and Outlook

Weight loss of legume residues was rapid (0-30 days), moderately rapid (30-60), moderate (60-90) and gradual (90-120). About 10-33, 50 and 80-86.7% loss of N by chickpea and lupin was realized in 10-15, 30-60 and 120 days, respectively. Approximately 50% of P released occurred between day 15 and 30 for both legumes. Kale leaves had higher N and P concentrations/uptake from day 30 to 60. Half-life for N and P was at day 20 and 30, respectively. Similarly, the N and P release by chickpea and uptake by kale intersected at day 25 and 30, respectively. Chickpea and white lupin residues have the potential to provide nutrients upon mineralization but for maximum nutrient synchrony between nutrient release and kale nutrient uptake, the legume residues should be incorporated in soil at kale seedling (4 weeks old) transplanting to optimize on residue benefits and minimize loss of available nutrients.

### Acknowledgement

This study was funded by DANIDA through productivity and growth of organic value chains (ProGroV) project.

### References

- Chepkoech Caroline 2015. Effect of legume integration and application of organic fertilizers on soil nutrient status and kale yield in Kabete, Kenya. MSc. Thesis, University of Nairobi
- Nduku Haggai 2014. Crop Performance and Soil Nutrient Dynamics under Different Organic Based Cropping Systems and Modeling the Effect of Climate Change on Maize (*Zea May*) Growth in Central Kenya. PhD. Thesis, University of Nairobi
- Genga, Quintar A. 2014. Organic based soil fertility management strategies for production of marketable quality tomatoes. MSc. Thesis, University of Nairobi
- Onwonga Richard, Noel A. Templer, Joyce J. Lelei, Faith J. Toroitich 2015. Combined Effects of Legumes with Phosphorus Fertilizer on Nutrient Balances and Gross Margins in Maize (*Zea mays* L.) systems of Kabete sub-County, Kenya. *Journal of Biology, Agriculture and Healthcare* 5, (16) 65-77.