

Vitamin E May Protect Against Contrast-Induced Acute Kidney Injury

Harri Hemilä and Yousef Rezaei

This is a manuscript version of a paper published in its final version in:

Journal of the American College of Cardiology 2017;69(14):1878

<https://doi.org/10.1016/j.jacc.2016.11.091>

<https://www.ncbi.nlm.nih.gov/pubmed/28385322>

A meta-analysis of the three RCTs was published in another report:

Rezaei Y, Hemilä H. Vitamin E may decrease the incidence of contrast-induced acute kidney injury.

American Journal of Kidney Diseases 2017

<https://doi.org/10.1053/j.ajkd.2016.12.022>

<https://www.ncbi.nlm.nih.gov/pubmed/28279510>

Correspondence:

Harri Hemilä

Department of Public Health, POB 20, Tukholmankatu 8B,

University of Helsinki, Helsinki

FI-00014, Finland

harri.hemila@helsinki.fi

Please note: Both authors have reported that they have no relationships relevant to the contents of this paper to disclose.

McCullough et al. (1) reviewed the pathophysiology and treatment options for contrast-induced acute kidney injury (CI-AKI). They stated that no effective adjunctive pharmaceutical had been demonstrated that either prevented or treated CI-AKI. However, they also suggested that of the agents being investigated, statins were the most promising. We would like to point out that strong evidence has also emerged regarding the effect of vitamin E against CI-AKI, which was not mentioned in their review.

Three randomized placebo-controlled trials found that vitamin E significantly prevented CI-AKI, with point estimates ranging from 52% to 75% for the decrease in the incidence of CI-AKI (2–4). All participants had chronic kidney disease and had been subjected to coronary catheterization or angiography. The latest trial reported CI-AKI cases in 14.1% of the placebo group, but in only 6.7% of the vitamin E group, which corresponded to 7.4% of participants benefitting from the vitamin, with a number needed to treat (NNT) of 13.5 (4). The 2 earlier studies found NNTs of 5.8 (2) and 10.6 (3). In each study, approximately one-half of the patients were on statin therapy; therefore, the effects of vitamin E might have also been beneficial in addition to statins.

Two of the vitamin E trials were carried out in Thailand (2,3), and 1 was carried out in Iran (4). Thus, it is not known whether the findings can be directly generalized to Western countries. Even if the positive findings might only be applicable to less developed countries, the findings are important for the populations of such countries.

Vitamin E is an essential nutrient, and therefore, its potential benefit in preventing CI-AKI is interesting. Furthermore, vitamins E and C may interact. Vitamin E decreased total mortality in male smokers aged older than 65 years if their dietary vitamin C intake level was high, but not if their vitamin C intake was low (5). Thus, a large factorial trial seems warranted to examine the effect of statins and vitamins E and C, and their combinations to discover the optimal protocol to prevent CI-AKI.

References

1. McCullough PA, Choi JP, Feghali GA, et al. Contrast-induced acute kidney injury. *J Am Coll Cardiol* 2016;68:1465–73.
<https://doi.org/10.1016/j.jacc.2016.05.099>
<https://www.ncbi.nlm.nih.gov/pubmed/27659469>
2. Tasanarong A, Piyayotai D, Thitiarchakul S. Protection of radiocontrast induced nephropathy by vitamin E (alpha tocopherol): a randomized controlled pilot study. *J Med Assoc Thai*. 2009;92(10):1273-1281.
<https://www.ncbi.nlm.nih.gov/pubmed/19845233>
3. Tasanarong A, Vohakiat A, Hutayanon P, Piyayotai D. New strategy of alpha- and gamma-tocopherol to prevent contrast-induced acute kidney injury in chronic kidney disease patients undergoing elective coronary procedures. *Nephrol Dial Transplant*. 2013;28(2):337-344.
<https://www.ncbi.nlm.nih.gov/pubmed/23314316>
<https://doi.org/10.1093/ndt/gfs525>
4. Rezaei Y, Khademvatani K, Rahimi B, Khoshfetrat M, Arjmand N, Seyyed-Mohammadzad MH. Short-term high-dose vitamin E to prevent contrast medium-induced acute kidney injury in patients with chronic kidney disease undergoing elective coronary angiography: a randomized placebo-controlled trial. *J Am Heart Assoc*. 2016;5(3):e002919.
<https://www.ncbi.nlm.nih.gov/pubmed/27068631>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4943266>
<https://doi.org/10.1161/JAHA.115.002919>
5. Hemilä H, Kaprio J. Modification of the effect of vitamin E supplementation vitamin E group on the mortality of male smokers by age and dietary vitamin C. *Am J Epidemiol* 2009;169:946–53.
<https://www.ncbi.nlm.nih.gov/pubmed/19218294>
<https://doi.org/10.1093/aje/kwn413>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2661323>