

Deltex-1 mutations predict poor survival in diffuse large B-cell lymphoma

Diffuse large B-cell lymphoma (DLBCL) is a group of clinically aggressive and heterogeneous malignancies, of which approximately 60% can be cured with anthracycline-based chemoimmunotherapy. Based on gene expression profiling, DLBCL can be classified into two molecularly distinct subgroups showing germinal center B-cell (GCB) and activated B-cell (ABC) lymphoma signatures,^{1,2} which differ in their genotypic, phenotypic and clinical features. Recently, studies applying next generation sequencing (NGS) techniques have provided further insights into the heterogeneity and pathogenesis of DLBCL.³ Many recurrent mutations have been identified as drivers of lymphoma pathogenesis,³ yet their association with survival remains to be established. It is also likely that other mutations, which do not contribute to lymphomagenesis *per se*, but rather predispose to a more drug-resistant disease and poor survival, will be recognized.

To identify mutations that contribute to the clinical outcome of DLBCL patients, we performed whole exome sequencing for 8 primary lymphoma samples and matched normal DNA. The patients were clinically at high-risk of relapse [age-adjusted International Prognostic Index (aaIPI)]. A total of 922 somatic point mutations and insertion-deletions in 838 different genes were identified (*Online Supplementary Figure S1*). Several of them were detected in established oncogenes recurrently mutated in DLBCL, including *MYD88*, *MEF2B* and *B2M*.³ Fifty-one genes were mutated in at least 2 patients including lymphoma related genes *CD79B*, *P TEN* and *DTX1*.

For validation and correlation of the findings with survival we used a Cancer Genome Characterization Initiative (CGCI) cohort of 92 DLBCL patients (*Figure 1A*).⁴ The clinical characteristics of the cohort are shown in *Table 1*. Of the 838 genes, which were mutated in our initial screen, 456 genes were also mutated in the CGCI cohort. Using three different filtering approaches to exclude germline polymorphisms and to recognize disease-causing alterations, mutations in 41 genes were found to correlate with survival (*Online Supplementary Figure S1*). When the variants were filtered with CADD scoring predicting altered protein function, survival-associated mutations were found only in 4 genes, including *DTX1*, *PIM1*, *GNA13* and *TMSB4X* (*Online Supplementary Figure S1*). Of these, only the mutations in the *DTX1* gene encoding an E3 ubiquitin ligase and a regulator of the Notch pathway⁵⁻⁷ were significantly associated both with time to progression (TTP) and overall survival (OS) with all the filtering criteria used, and were thus selected for further analysis.

In our initial exome sequencing screen 3 patients harbored mutations in their *DTX1* genes, and a total of 8 different non-synonymous mutations were identified (*Online Supplementary Table S2*). All the mutations in the *DTX1* gene were validated using targeted capillary sequencing and confirmed to be somatic. In the CGCI study population, 14% (*Table 1*) of the patients were carrying non-synonymous *DTX1* mutations (*Online Supplementary Table S3*). Of these, 2 had 2 mutations in their *DTX1* genes and 1 had 3. Mutations were equally distributed between GCB and ABC subtypes and clinical risk groups (*Table 1*).

For the CGCI cohort, the median follow-up time was 59 months, with TTP, PFS (progression-free survival) and OS at 5 years of 74%, 72% and 79%, respectively. The

Table 1. Patient characteristics according to *DTX1* mutations in CGCI and Nordic cohorts of DLBCL patients.

Patient Characteristics	CGCI				Nordic			
	All	<i>DTX1</i> wt	<i>DTX1</i> mut	<i>P</i>	All	<i>DTX1</i> WWE1 wt	<i>DTX1</i> WWE1 mut	<i>P</i>
Sex								
Male	61 (66)	51 (65)	10 (77)	0.532	83 (57)	66 (57)	17 (59)	1.000
Female	31 (34)	28 (35)	3 (23)		62 (43)	50 (43)	12 (41)	
Age								
<60 y	39 (42)	35 (44)	4 (31)	0.546	82 (57)	68 (59)	14 (48)	0.403
>60 y	53 (58)	44 (56)	9 (69)		63 (43)	48 (41)	15 (52)	
Stage								
I-II	44 (48)	36 (46)	8 (62)	0.373	52 (36)	42 (36)	10 (34)	1.000
III-IV	48 (52)	43 (54)	5 (38)		93 (64)	74 (64)	19 (66)	
IPI								
Low (0-2)	40 (43)	35 (44)	5 (38)	0.770	87 (60)	69 (59)	18 (62)	0.836
High (3-5)	52 (57)	44 (56)	8 (62)		58 (40)	47 (41)	11 (38)	
Subtype*								
GCB	51 (55)	46 (58)	5 (38)		68 (47)	57 (49)	11 (38)	
ABC/Non-GCB	32 (35)	25 (32)	7 (54)	0.234	62 (43)	46 (40)	16 (55)	0.199
Unclassified/NA	9 (10)	8 (10)	1 (8)		15 (10)	13 (11)	2 (7)	

Median age in the CGCI and the Nordic cohorts is 61 (range 17-75) and 58 (range, 16-84) years. *In the CGCI cohort, molecular subtyping is based on gene expression profiling allowing for the classification of GCB, ABC and unclassified subtypes, whereas in the Nordic cohort, the classification is based on the Hans algorithm (GCB vs. non-GCB). NA (not available) refers to the Nordic cohort. IPI: International Prognostic Index; GCB: germinal center B cell; ABC: activated B cell; CGCI: Cancer Genome Characterization Initiative; wt: wild-type; mut: mutated.

patients with non-synonymous *DTX1* mutations had significantly worse survival in comparison to the patients with wild-type *DTX1* (Figure 1B). When *DTX1* mutations were analyzed according to different functional protein domains, a substantial enrichment of mutations (11 out of 16, 65%) was observed in the WWE1-domain (Figure 1C). In Kaplan-Meier and Cox regression analyses, decreases in TTP, PFS and OS rates were observed, if a mutation was located in the WWE1 domain (TTP, RR=4.533, 95% CI 1.684-12.207, $P=0.003$; PFS, RR=3.831, 95% CI 1.443-10.170, $P=0.007$; OS, RR=3.551, 95% CI 1.192-10.574, $P=0.023$; Figure 1D).

To validate the survival association of the non-synonymous mutations in the *DTX1* WWE1 domain, we expanded our initial screening cohort to a 'Nordic cohort' of 145 DLBCL patients (Figure 1A). Non-synonymous

mutations in the *DTX1* WWE1 domain were detected in 20% of the samples (29/145) (Figure 1C, *Online Supplementary Table S4*). Consistent with the CGCI cohort, WWE1 mutations were equally distributed between molecular and clinical risk groups (Table 1). The median follow-up time for the whole cohort was 62 months (range 3-133 months). 5-year TTP, PFS and OS rates were 73%, 69%, and 77%, respectively. However, for the patients with non-synonymous WWE1 domain mutations of *DTX1*, survival rates at 5 years were significantly worse than for the patients without mutations (Figure 1E). In Cox multivariate analysis with an IPI score, *DTX1* WWE1 mutations remained independent prognostic factors for TTP and OS (Table 2). Collectively, the results in our two independent cohorts show that the mutational status of the *DTX1* WWE1 domain is a novel




Figure 1. *DTX1* mutations are enriched in the first exon encoding the WWE 1 domain and predict survival in DLBCL. (A) Flow of the study. (B) OS in the CGCI cohort according to *DTX1* mutations. (C) Structure of the *DTX1* gene and localization of the *DTX1* mutations in the Nordic and CGCI cohorts. (D) OS in the CGCI cohort according to mutations in the *DTX1* WWE1 domain. (E) OS in the Nordic validation cohort according to mutations in the *DTX1* WWE1 domain. DLBCL: diffuse large B-cell lymphoma; OS: overall survival; CGCI: Cancer Genome Characterization Initiative.

Table 2. Cox multivariate models for *DTX1* mutations in the Nordic cohort.

Survival	TTP			OS		
	RR	95% CI	P	RR	95% CI	P
Variable						
<i>DTX1</i> (wt vs. WWE1 mutated)	2.140	1.102-4.154	0.025	2.114	1.050-4.255	0.036
IPI	1.434	1.095-1.878	0.009	1.482	1.110-1.980	0.008

DTX1 mutational status, mutated worse; IPI, higher IPI worse. RR: relative risk; CI: confidence interval; IPI: International Prognostic Index; TTP: time to progression; OS: overall survival; wt: wild-type.

negative predictor of survival in DLBCL.

To examine *DTX1* expression in the lymphoma tissue and localization in different cellular compartments, we extended our analyses to the protein level. In the normal lymph node, *DTX1* immunoreactivity was cytoplasmic and primarily localized to germinal center B-cells with centroblast morphology (Online Supplementary Figure S2A). In DLBCL tissue, the staining intensity varied between the samples, but otherwise resembled that of normal centroblasts (Online Supplementary Figure S2B). Interestingly, the samples carrying *DTX1* mutations showed significantly lower protein levels (*t*-test, $P=0.007$; Online Supplementary Figure S2C-D).

Further examination of *DTX1* mutations and gene expression in the CGCI repository confirmed that the samples carrying *DTX1* mutations showed significantly lower *DTX1* expression (*t*-test, $P=0.011$; Online Supplementary Figure S2E). In addition, *DTX1* expression was significantly higher in the GCB than the ABC type DLBCLs (*t*-test, $P=0.009$). The association was further validated in the Lymphoma/Leukemia Molecular Profiling Project (LLMPP) cohort¹ of 233 patients with DLBCL (*t*-test, $P<0.001$).

To identify genes that were anti-expressed and co-expressed with *DTX1*, we computed Pearson's correlation between *DTX1* expression levels and other genes across 92 samples in the CGCI data set, which resulted in 471 inversely and 1832 positively correlated genes. Among the inversely correlated genes were, for example, *JAK2*, *CASP10*, *CD47*, $\beta 2M$ and *INFG1*, while *BCL6*, *PRKCA*, *RCOR1*, *CD22*, and *KMT2B*, were among the positively correlated ones. To identify biological pathways associated with the *DTX1* expression signature, we performed a pathway enrichment analysis for anti-expressed and co-expressed genes. The genes inversely correlated with *DTX1* were enriched, for example, for proteasomal degradation, IFN- γ signaling and JAK-STAT pathways (Online Supplementary Table S5), whereas positively correlated pathways included, for instance, chromatin remodeling, B-cell receptor (BCR) signaling and mitogen-activated protein kinase (MAPK) pathways (Online Supplementary Table S6). In addition, Notch signaling pathway effectors were found to correlate with *DTX1* expression.

DTX1 is an ubiquitin E3 ligase containing N-terminal Notch binding WWE domains, a proline-rich motif and a C-terminal really interesting new gene (RING) finger domain commonly found in ubiquitin E3 ligases.⁵⁻⁷ *DTX1* mediates Notch activation in *Drosophila*, but its exact role in Notch signaling in mammals has remained largely ambiguous. Interestingly, in the hematological context, *DTX1* can negatively regulate T-cell activation by targeting mitogen-activated protein kinase kinase kinase 1 (MEKK1 or MAP3K) and protein kinase C θ for degradation,^{8,9} and mediate degradation of hypoxia-inducible factor-1 α (HIF-1 α) in regulatory T cells (Tregs).¹⁰ However, in normal B cells and lymphomas, the function of *DTX1*

and the impact of mutations remain to be established. Somatic hypermutation has been predicted to aberrantly target the *DTX1* gene,¹¹ and *DTX1* mutations have previously been reported in follicular and splenic marginal zone lymphomas.^{12,13} Furthermore, *DTX1* mutations were also recently identified in a cohort consisting of Chinese patients with primary and relapsed DLBCL.¹⁴ Consistent with our findings, the mutations were almost exclusively localized to the WWE1 domain of the protein, and based on the Notch reporter gene assay in U2OS osteosarcoma cells, it was suggested that deleterious mutations impair the function of *DTX1* as a negative regulator of Notch. Our pathway enrichment analyses support the idea that there is a functional connection between *DTX1* and the Notch pathway. In addition, we identify other novel and potentially relevant associations. However, functional studies should be performed to establish the biological relevance of these findings.

In conclusion, we have used well characterized DLBCL patient cohorts to identify survival associated mutations in DLBCL, and validated the findings in a large and independent patient cohort. To our knowledge, this is the first report to examine the prognostic impact of *DTX1* aberrations in lymphomas. While further investigation of the molecular mechanisms driven by *DTX1* mutations remain to be established, our results indicate that these aberrations in the *DTX1* gene have a tumor-promoting role in DLBCL. Taken together, the results presented herein are promising and novel, and demonstrate that mutational status at diagnosis can provide prognostic value, independent of IPI, and improve risk stratification in DLBCL. (clinicaltrials.gov Identifier: 01502982)

Leo Meriranta,^{1,2} Annika Pasanen,^{1,2} Riku Louhimo,¹ Alejandra Cervera,¹ Amjad Alkodsí,¹ Matias Autio,^{1,2} Minna Taskinen,^{1,2} Ville Rantanen,¹ Marja-Liisa Karjalainen-Lindsberg,³ Harald Holte,⁴ Jan Delabie,^{5,6} Rainer Lehtonen,¹ Sampsa Hautaniemi¹ and Sirpa Leppä^{1,2}

¹Research Programs Unit, Genome-Scale Biology Program, Faculty of Medicine, University of Helsinki, Finland; ²Department of Oncology, Helsinki University Hospital Comprehensive Cancer Center, Finland; ³Department of Pathology, Helsinki University Hospital, Finland; ⁴Department of Oncology, Oslo University Hospital, Norway; ⁵Department of Pathology, University of Toronto, ON, Canada, and ⁶Department of Pathology, Oslo University Hospital, Norway

Acknowledgments: The authors would like to thank Anne Aarnio and Marika Tuukkanen for expert technical assistance, Marko Hyttiäinen and Kirsi Jäntti for valuable comments on the manuscript and Institute for Molecular Medicine, Finland (FIMM) and Biomedicum Genomics for sequencing services and the Cancer Genomics Characterization Initiative (CGCI) consortium for providing DLBCL data set (supported by NCI contract N01-CO-12400; <http://cgap.nci.nih.gov/cgi.html>). The study was supported by grants from the Academy of Finland (SH), Finnish Cancer Foundation (SL, SH), Juselius Foundation (SL, SH), University of Helsinki (SL), and Helsinki University Hospital (SL).

Correspondence: sirpa.leppa@helsinki.fi
doi:10.3324/haematol.2016.157495

Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at www.haematologica.org.

References

1. Lenz G, Wright G, Dave SS, et al. Stromal gene signatures in large-B-cell lymphomas. *N Engl J Med*. 2008;359(22):2313-2323.
2. Rosenwald A, Wright G, Chan WC, et al. The use of molecular profiling to predict survival after chemotherapy for diffuse large-B-cell lymphoma. *N Engl J Med*. 2002;346(25):1937-1947.
3. Pasqualucci L, Dalla-Favera R. The genetic landscape of diffuse large B-cell lymphoma. *Semin Hematol*. 2015;52(2):67-76.
4. Morin RD, Mungall K, Pleasance E, et al. Mutational and structural analysis of diffuse large B-cell lymphoma using whole-genome sequencing. *Blood*. 2013;122(7):1256-1265.
5. Matsuno K, Eastman D, Mitsiades T, et al. Human deltex is a conserved regulator of Notch signalling. *Nat Genet*. 1998;19(1):74-78.
6. Mukherjee A, Veraksa A, Bauer A, et al. Regulation of Notch signalling by non-visual beta-arrestin. *Nat Cell Biol*. 2005;7(12):1191-1201.
7. Takeyama K, Aguiar RC, Gu L, et al. The BAL-binding protein BBAP and related Deltex family members exhibit ubiquitin-protein isopeptide ligase activity. *J Biol Chem*. 2003;278(24):21930-21937.
8. Liu WH, Lai MZ. Deltex regulates T-cell activation by targeted degradation of active MEK1. *Mol Cell Biol*. 2005;25(4):1367-1378.
9. Hsu TS, Hsiao HW, Wu PJ, Liu WH, Lai MZ. Deltex1 promotes protein kinase C θ degradation and sustains Casitas B-lineage lymphoma expression. *J Immunol*. 2014;193(4):1672-1680.
10. Hsiao HW, Hsu TS, Liu WH, et al. Deltex1 antagonizes HIF-1 α and sustains the stability of regulatory T cells in vivo. *Nat Commun*. 2015;6:6353.
11. Khodabakhshi AH, Morin RD, Fejes AP, et al. Recurrent targets of aberrant somatic hypermutation in lymphoma. *Oncotarget*. 2012;3(11):1308-1319.
12. Rossi D, Trifonov V, Fangazio M, et al. The coding genome of splenic marginal zone lymphoma: activation of NOTCH2 and other pathways regulating marginal zone development. *J Exp Med*. 2012;209(9):1537-1551.
13. Green MR, Kihira S, Liu CL, et al. Mutations in early follicular lymphoma progenitors are associated with suppressed antigen presentation. *Proc Natl Acad Sci USA*. 2015;112(10):E1116-1125.
14. de Miranda NF, Georgiou K, Chen L, et al. Exome sequencing reveals novel mutation targets in diffuse large B-cell lymphomas derived from Chinese patients. *Blood*. 2014;124(16):2544-2553.