



May 18th, 6:30 PM - 7:00 PM

# Modeling cell proliferation in the transition from neurogenesis to gliogenesis

Christine Mennicke

*North Carolina State University, [cvmennic@ncsu.edu](mailto:cvmennic@ncsu.edu)*


Troy Ghashghaei

*North Carolina State University, [tghashg@ncsu.edu](mailto:tghashg@ncsu.edu)*

Mansoor A. Haider

*North Carolina State University, [m\\_haider@ncsu.edu](mailto:m_haider@ncsu.edu)*

Follow this and additional works at: <http://scholarscompass.vcu.edu/bamm>

 Part of the [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Physical Sciences and Mathematics Commons](#)

---

<http://scholarscompass.vcu.edu/bamm/2017/thursday/23>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact [libcompass@vcu.edu](mailto:libcompass@vcu.edu).

## Modeling cell proliferation in the transition from neurogenesis to gliogenesis

Christine Mennicke<sup>1</sup>, Troy Ghashghaei<sup>2</sup> and Mansoor A. Haider<sup>1</sup>

<sup>1</sup>Dept. of Mathematics, North Carolina State University, Raleigh NC, 27695

<sup>2</sup>Dept. of Molecular Biomedical Sciences, North Carolina State University, Raleigh NC, 27695

In the developing brain, radial glial progenitor cells follow patterns of proliferation and differentiation to produce neurons and glia. These progenitor cells first undergo neurogenesis, producing neurons in a fairly deterministic manner, and then a portion of the cell population switches to producing glia. We explore two different ways of modeling cell divisions, branching random walks and Galton-Watson processes, particularly focusing on their application to gliogenesis. A primary quantity of interest is the number of generations of cell division required to produce  $N$  cells from a single progenitor. We develop models that systematically determine the depth of all binary trees with  $N$  leaf nodes. This information is used to estimate the number of generations of cell division in data sets tracing cell lineages in the developing mouse brain. We examine the hypothesis that cell proliferation patterns are altered as development transitions from neurogenesis to gliogenesis.