

Lacunary series and stable distributions

István Berkes and Robert Tichy

Abstract By well known results of probability theory, any sequence of random variables with bounded second moments has a subsequence satisfying the central limit theorem and the law of the iterated logarithm in a randomized form. In this paper we give criteria for a sequence (X_n) of random variables to have a subsequence (X_{n_k}) whose weighted partial sums, suitably normalized, converge weakly to a symmetric stable distribution with parameter $0 < \alpha < 2$.

1 Introduction

It is known that sufficiently thin subsequences of general r.v. sequences behave like i.i.d. sequences. For example, Chatterji [9], [10] and Gaposhkin [15], [16] proved that if a sequence (X_n) of r.v.'s satisfies $\sup_n EX_n^2 < \infty$, then one can find a subsequence (X_{n_k}) and r.v.'s X and $Y \geq 0$ such that

$$\frac{1}{\sqrt{N}} \sum_{k \leq N} (X_{n_k} - X) \xrightarrow{d} N(0, Y) \quad (1)$$

and

$$\limsup_{N \rightarrow \infty} \frac{1}{\sqrt{2N \log \log N}} \sum_{k \leq N} (X_{n_k} - X) = Y^{1/2} \quad \text{a.s.}, \quad (2)$$

where $N(0, Y)$ denotes the distribution of the r.v. $Y^{1/2} \zeta$ where ζ is an $N(0, 1)$ r.v. independent of Y . Komlós [19] proved that under $\sup_n E|X_n| < \infty$ there exists a subsequence (X_{n_k}) and an integrable r.v. X such that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N X_{n_k} = X \quad \text{a.s.}$$

and Chatterji [8] showed that under $\sup_n E|X_n|^p < \infty$, $0 < p < 2$ the conclusion of the previous theorem can be changed to

$$\lim_{N \rightarrow \infty} \frac{1}{N^{1/p}} \sum_{k=1}^N (X_{n_k} - X) = 0 \quad \text{a.s.}$$

István Berkes

Institute of Statistics, Graz University of Technology, Kopernikusgasse 24, Graz, Austria. e-mail: berkes@tugraz.at. Research supported by FWF grants P24302-N18, W1230 and OTKA grant K106815.

Robert Tichy

Graz University of Technology, Institute of Mathematics A, Steyrergasse 30, 8010 Graz, Austria. e-mail: tichy@tugraz.at. Research supported by FWF grants P24302-N18 and W1230.

for some X with $E|X|^p < \infty$. Note the randomization in all these examples: the role of the mean and variance of the subsequence (X_{n_k}) is played by random variables X, Y . On the basis of these and several other examples, Chatterji [11] formulated the following heuristic principle:

Subsequence Principle. *Let T be a probability limit theorem valid for all sequences of i.i.d. random variables belonging to an integrability class L defined by the finiteness of a norm $\|\cdot\|_L$. Then if (X_n) is an arbitrary (dependent) sequence of random variables satisfying $\sup_n \|X_n\|_L < +\infty$ then there exists a subsequence (X_{n_k}) satisfying T in a mixed form.*

In a profound paper, Aldous [1] proved the validity of this principle for all limit theorems concerning the almost sure or distributional behavior of a sequence of functionals $f_k(X_1, X_2, \dots)$ of a sequence (X_n) of r.v.'s. Most "usual" limit theorems belong to this class; for precise formulations, discussion and examples we refer to [1]. On the other hand, the theory does not cover functionals f_k containing parameters (as in weighted limit theorems) or allows limit theorems to involve other type of uniformities. Such uniformities play an important role in analysis. For example, if from a sequence (X_n) of r.v.'s with finite p -th moments ($p \geq 1$) one can select a subsequence (X_{n_k}) such that

$$K^{-1} \left(\sum_{i=1}^N a_i^2 \right)^{1/2} \leq \left\| \sum_{i=1}^N a_i X_{n_i} \right\|_p \leq K \left(\sum_{i=1}^N a_i^2 \right)^{1/2}$$

for some constant $0 < K < \infty$, for every $N \geq 1$ and every $(a_1, \dots, a_N) \in \mathbb{R}^N$, then the subspace of L^p spanned by (X_n) contains a subspace isomorphic to Hilbert space. Such embedding arguments go back to the classical paper of Kadec and Pelczynski [18] and play an important role in Banach space theory, see e.g. Dacunha-Castelle and Krivine [12], Aldous [2]. In the theory of orthogonal series and in Banach space theory we frequently need subsequences (f_{n_k}) of a sequence (f_n) such that $\sum_{k=1}^{\infty} a_k f_{n_k}$ converges a.e. or in norm, after any permutation of its terms, for a class of coefficient sequences (a_k) . Here we need uniformity both over a class of coefficient sequences (a_k) and over all permutations of the terms of the series. A number of uniform limit theorems for subsequences have been proved by ad hoc arguments. Révész [22] showed that for any sequence (X_n) of r.v.'s satisfying $\sup_n EX_n^2 < \infty$ one can find a subsequence (X_{n_k}) and a r.v. X such that $\sum_{k=1}^{\infty} a_k (X_{n_k} - X)$ converges a.s. provided $\sum_{k=1}^{\infty} a_k^2 < \infty$. Under $\sup_n \|X_n\|_{\infty} < +\infty$, Gaposhkin [15] showed that there exists a subsequence (X_{n_k}) and r.v.'s X and $Y \geq 0$ such that for any real sequence (a_k) satisfying the uniform asymptotic negligibility condition

$$\max_{1 \leq k \leq N} |a_k| = o(A_N), \quad A_N = \left(\sum_{k=1}^N a_k^2 \right)^{1/2} \quad (3)$$

we have

$$\frac{1}{A_N} \sum_{k \leq N} a_k (X_{n_k} - X) \xrightarrow{d} N(0, Y) \quad (4)$$

and for any real sequence (a_k) satisfying the Kolmogorov condition

$$\max_{1 \leq k \leq N} |a_k| = o(A_N / (\log \log A_N)^{1/2}) \quad (5)$$

we have

$$\frac{1}{(2A_N \log \log A_N)^{1/2}} \sum_{k \leq N} a_k (X_{n_k} - X) = Y^{1/2} \quad \text{a.s.} \quad (6)$$

For a fixed coefficient sequence (a_k) the above results follow from Aldous' general theorems, but the subsequence (X_{n_k}) provided by the proofs depends on (a_k) and to find a subsequence working for all (a_k) simultaneously requires a uniformity which is, in general, not easy to establish and it can fail in important situations. (See Guerre and Raynaud [17] for a natural problem where uniformity is not valid.) In [1], Aldous used an equicontinuity argument to prove a permutation-invariant version of the theorem of Révész above, implying that every orthonormal system (f_n) contains a subsequence (f_{n_k}) which, using the standard

terminology, is an *unconditional convergence system*. This had been a long standing open problem in the theory of orthogonal series (see Uljanov [24], p. 48) and was first proved by Komlós [20]. In [3] we used the method of Aldous to prove extensions of the Kadec-Pelczynski theorem, as well as selection theorems for almost symmetric sequences. The purpose of the present paper is to use a similar technique to prove a uniform limit theorem of probabilistic importance, namely the analogue of Gaposhkin's uniform CLT (3)–(4) in the case when the limit distribution of the normed sum is a symmetric stable law with parameter $0 < \alpha < 2$. To formulate our result, we need some definitions. Using the terminology of [6], call the sequence (X_n) of r.v.'s *determining* if it has a limit distribution relative to any set A in the probability space with $P(A) > 0$, i.e. for any $A \subset \Omega$ with $P(A) > 0$ there exists a distribution function F_A such that

$$\lim_{n \rightarrow \infty} P(X_n < t \mid A) = F_A(t)$$

for all continuity points t of F_A . By an extension of the Helly-Bray theorem (see [6]), every tight sequence of r.v.'s contains a determining subsequence. Hence in studying the asymptotic behavior of thin subsequences of general tight sequences we can assume without loss of generality that our original sequence (X_n) is determining. By [6], Proposition 2.1, for any continuity point t of the limit distribution function F_Ω , the sequence $I\{X_n \leq t\}$ converges weakly in L^∞ to some r.v. G_t ; clearly $G_s \leq G_t$ a.s. for any $s \leq t$. (A sequence (ξ_n) of bounded r.v.'s is said to converge to a bounded r.v. ξ weakly in L^∞ if $E(\xi_n \eta) \rightarrow E(\xi \eta)$ for any integrable r.v. η . To avoid confusion, we will call ordinary weak convergence of probability measures *distributional convergence* and denote it by \xrightarrow{d} .) Using a standard procedure (see e.g. Révész [23], Lemma 6.1.4), by choosing a dense countable set D of continuity points of F_Ω , one can construct versions of G_t , $t \in D$ such that, for every fixed $\omega \in \Omega$, the function $G_t(\omega)$, $t \in D$ extends to a distribution function. Letting μ denote the corresponding measure, μ is called the *limit random measure* of (X_n) ; it was introduced by Aldous [1]; for properties and applications see [2], [3], [5], [6]. Clearly, μ can be considered as a measurable map from the underlying probability space (Ω, \mathcal{F}, P) to the space \mathcal{M} of probability measures on \mathbb{R} equipped with the Prohorov metric π . It is easily seen that for any A with $P(A) > 0$ and any continuity point t of F_A we have

$$F_A(t) = E_A(\mu(-\infty, t)), \tag{7}$$

where E_A denotes conditional expectation given A . Note that μ depends on the actual r.v.'s X_n , but the distribution of μ in (\mathcal{M}, π) depends solely on the distribution of the sequence (X_n) . The situation concerning the unweighted CLT for lacunary sequences can now be summarized by the following theorem.

Theorem 1. *Let (X_n) be a determining sequence of r.v.'s with limit random measure μ . Then there exists a subsequence (X_{n_k}) satisfying, together with all of its subsequences, the CLT (1) with suitable r.v.'s X and $Y \geq 0$ if and only if*

$$\int_{-\infty}^{\infty} x^2 d\mu(x) < \infty \quad \text{a.s.} \tag{8}$$

The sufficiency part of the theorem is contained in Aldous' general subsequence theorems in [1]; the necessity was proved in our recent paper [7]. Note that the condition for the CLT for lacunary subsequences of (X_n) is given in terms of the limit random measure of (X_n) and this condition is the exact analogue of the condition in the i.i.d. case, only the common distribution of the i.i.d. variables is replaced by the limit random measure. Note also that the existence of second moments of (X_n) (or the existence of any moments) is not necessary for the conclusion of Theorem 1.

In this paper we investigate the analogous question in case of a nonnormal stable limit distribution, i.e. the question under what conditions a sequence (X_n) of r.v.'s has a subsequence (X_{n_k}) whose weighted partial sums, suitably normalized, converge weakly to an α -stable distribution, $0 < \alpha < 2$. Let, for $c > 0$ and $0 < \alpha < 2$, $G_{\alpha,c}$ denote the distribution function with characteristic function $\exp(-c|t|^\alpha)$ and let $S = S(\alpha, c)$ denote the class of symmetric distributions on \mathbb{R} with characteristic function φ satisfying

$$\varphi(t) = 1 - c|t|^\alpha + o(|t|^\alpha) \quad \text{as } t \rightarrow 0. \tag{9}$$

Our main result is

Theorem 2. *Let $0 < \alpha < 2$, $c > 0$ and let (X_n) be a determining sequence of r.v.'s with limit random measure μ . Assume that $\mu \in S(\alpha, c)$ with probability 1. Then there exists a subsequence (X_{n_k}) such that for any real sequence (a_k) satisfying*

$$\max_{1 \leq k \leq N} |a_k| = o(A_N), \quad A_N = \left(\sum_{k=1}^N |a_k|^\alpha \right)^{1/\alpha} \quad (10)$$

we have

$$A_N^{-1} \sum_{k=1}^N a_k X_{n_k} \xrightarrow{d} G_{\alpha, c}.$$

Condition (9) holds provided the corresponding (symmetric) distribution function F satisfies

$$1 - F(x) = c_1 x^{-\alpha} + \beta(x) x^{-\alpha}, \quad x > 0$$

where $c_1 > 0$ is a suitable constant, $\beta(x)$ is non-increasing for $x \geq x_0$ and $\lim_{x \rightarrow \infty} \beta(x) = 0$. (See Berkes and Dehling [4], Lemma 3.2.) Apart from the monotonicity condition, this is equivalent to the fact that F is in the domain of normal attraction of a symmetric stable distribution. (See e.g. Feller [14], p. 581.) It is natural to ask if the conclusion of Theorem 2 remains valid (with a suitable centering factor) assuming only that $\mu \in S$ a.s. where S denotes the domain of normal attraction of a fixed stable distribution. From the theory in [1] it follows that the answer is affirmative in the unweighted case $a_k = 1$, but in the uniform weighted case the question remains open. Symmetry plays no essential role in the proof of Theorem 2; it is used only in Lemma 2 and at the cost of minor changes in the proof, (9) can be replaced by a condition covering nonsymmetric distributions as well. But since we do not know the optimal condition, we restricted our investigations to the case (9) where the technical details are the simplest and the idea of the proof becomes more transparent.

Given a sequence (X_n^*) of r.v.'s and a random measure μ defined on a probability space (Ω, \mathcal{F}, P) such that X_n^* are conditionally i.i.d. given μ with conditional distribution μ , the limit random measure of (X_n^*) is easily seen to be μ . Thus in the case $\mu \in S(\alpha, c)$ a.s., (X_n^*) provides a simple example for a sequence satisfying the conditions of Theorem 2. (Since (X_n^*) is exchangeable, in this case the conclusion of Theorem 2 holds for the whole sequence (X_n^*) without passing to any subsequence.) Theorem 2 shows that any deterministic sequence (X_n) with a limit random measure μ satisfying $\mu \in S(\alpha, c)$ a.s. has a subsequence (X_{n_k}) whose weighted partial sums behave, in a uniform sense, similarly to those of (X_n^*) .

2 Proof of Theorem 2

As the first step of the proof, we select a sequence $n_1 < n_2 < \dots$ of integers such that, after a suitable discretization of (X_n) , we have

$$P(X_{n_k} \in J | X_{n_1}, \dots, X_{n_{k-1}})(\omega) \longrightarrow \mu(\omega, J) \quad \text{a.s.} \quad (11)$$

for a large class of intervals J . This step follows exactly Aldous [1], see Proposition 11 of [1] for details. Let (Y_n) be a sequence of r.v.'s on (Ω, \mathcal{F}, P) such that, given \mathbf{X} and μ , the r.v.'s Y_1, Y_2, \dots are conditionally i.i.d. with distribution μ , i.e.,

$$P(Y_1 \in B_1, \dots, Y_k \in B_k | \mathbf{X}, \mu) = \prod_{i=1}^k P(Y_i \in B_i | \mathbf{X}, \mu) \quad \text{a.s.} \quad (12)$$

$$P(Y_j \in B | \mathbf{X}, \mu) = \mu(B) \quad \text{a.s.} \quad (13)$$

for any j, k and Borel sets B, B_1, \dots, B_k on the real line. Such a sequence (Y_n) always exists after redefining (X_n) and μ on a suitable, larger probability space; for example, one can define the triple $((X_n), \mu, (Y_n))$ on the product space $\mathbb{R}^\infty \times \mathcal{M} \times \mathbb{R}^\infty$ as done in [1], p. 72. This redefinition will not change the distribution of the sequence (X_n) and thus by Proposition 2.1 of [6] it remains determining. Since the random measure

μ depends on the variables X_n themselves and not only on the distribution of (X_n) , this redefinition will change μ , but not the joint distribution of (X_n) and μ on which our results depend. Using (11) and a martingale argument, in [1], Lemma 12 it is shown that

Lemma 1. *For every $\sigma(\mathbf{X})$ -measurable r.v. Z and any $j \geq 1$ we have*

$$(X_{n_k}, Z) \xrightarrow{d} (Y_j, Z) \quad \text{as } k \rightarrow \infty.$$

We now construct a further subsequence of (X_{n_k}) satisfying the conclusion of Theorem 2. By reindexing our variables, we can assume that Lemma 1 holds with $n_k = k$. For our construction we need some auxiliary considerations. For a (nonrandom) measure $\mu \in S(\alpha, c)$, the corresponding characteristic function φ satisfies

$$\varphi(t) = 1 - c|t|^\alpha + \beta(t)|t|^\alpha, \quad t \in \mathbb{R} \quad (14)$$

where β is a bounded continuous function on \mathbb{R} with $\beta(0) = 0$. Given $\mu_1, \mu_2 \in S(\alpha, c)$ with characteristic functions φ_1, φ_2 and corresponding functions β_1, β_2 in (14), define

$$\rho(\mu_1, \mu_2) = \sup_{0 \leq |t| \leq 1} |\beta_1(t) - \beta_2(t)| + \sum_{k=0}^{\infty} \frac{1}{2^k} \sup_{2^k \leq |t| \leq 2^{k+1}} |\beta_1(t) - \beta_2(t)|. \quad (15)$$

Clearly, ρ satisfies the triangle inequality and if $\rho(\mu_1, \mu_2) = 0$, then $\beta_1(t) = \beta_2(t)$ and consequently $\varphi_1(t) = \varphi_2(t)$ for all $t \in \mathbb{R}$ and thus $\mu_1 = \mu_2$. Hence, ρ is a metric on $S(\alpha, c)$. If $\mu, \mu_1, \mu_2, \dots \in S(\alpha, c)$ with corresponding characteristic functions $\varphi, \varphi_1, \varphi_2, \dots$ and functions $\beta, \beta_1, \beta_2, \dots$, then $\rho(\mu_n, \mu) \rightarrow 0$ implies that $\beta_n(t) \rightarrow \beta(t)$ and consequently $\varphi_n(t) \rightarrow \varphi(t)$ uniformly on compact intervals and thus $\mu_n \xrightarrow{d} \mu$. Conversely, if $\mu_n \xrightarrow{d} \mu$, then $\varphi_n(t) \rightarrow \varphi(t)$ uniformly on compact intervals and thus $\beta_n(t) \rightarrow \beta(t)$ uniformly on compact intervals not containing 0. Note that $\lim_{t \rightarrow 0} \beta_n(t) = 0$ for any fixed n by the definition of $S(\alpha, c)$; if this relation holds uniformly in n , then $\beta_n(t) \rightarrow \beta(t)$ will hold uniformly also on all compact intervals containing 0 and upon observing that (14) implies $|\beta(t)| \leq |t|^{-\alpha} |\varphi(t) - 1| + c \leq c + 2$ for $|t| \geq 1$ and thus the total contribution of the terms of the sum in (15) for $k \geq M$ is $\leq 4(c+2)2^{-M}$, it follows that $\rho(\mu_n, \mu) \rightarrow 0$. Thus if for a class $H \subset S(\alpha, c)$ we have $\lim_{t \rightarrow 0} \beta(t) = 0$ uniformly for all functions β corresponding to measures in H , then in H convergence of elements in Prohorov metric and in the metric ρ are equivalent.

Let now $\varphi(t) = \varphi(t, \omega)$ denote the characteristic function of the random measure $\mu = \mu(\omega)$. By the assumption $\mu \in S(\alpha, c)$ a.s. of Theorem 2, we have

$$\varphi(t, \omega) = 1 - c|t|^\alpha + \beta(t, \omega)|t|^\alpha, \quad t \in \mathbb{R}, \omega \in \Omega \quad (16)$$

where $\lim_{t \rightarrow 0} \beta(t, \omega) = 0$ a.s. Let $\xi_n(\omega) = \sup_{|t| \leq 1/n} |\beta(t, \omega)|$, then $\lim_{n \rightarrow \infty} \xi_n(\omega) = 0$ a.s. and thus by Egorov's theorem (see [13]) for any $\varepsilon > 0$ there exists a measurable set $A \subset \Omega$ with $P(A) \geq 1 - \varepsilon$ such that $\lim_{n \rightarrow \infty} \xi_n(\omega) = 0$ and consequently $\lim_{t \rightarrow 0} \beta(t, \omega) = 0$ uniformly on A . Considering A as a new probability space, we will show that there exists a subsequence (X_{n_k}) (depending on A) satisfying the conclusion of Theorem 2 together with all its subsequences. By a diagonal argument we can get then a subsequence (X_{n_k}) satisfying the conclusion of Theorem 2 on the original Ω . Thus without loss of generality we can assume in the sequel that the function $\beta(t, \omega)$ in (16) satisfies $\lim_{t \rightarrow 0} \beta(t, \omega) = 0$ uniformly in $\omega \in \Omega$ and thus by the remarks in the previous paragraph, in the support of the random measure μ the Prohorov metric and the metric ρ generate the same convergence.

Lemma 2. *Let $\mu_1, \mu_2 \in S(\alpha, c)$ satisfying (9), let Z_1, \dots, Z_n and Z_1^*, \dots, Z_n^* be i.i.d. sequences with respective distributions μ_1, μ_2 . Let $(a_1, \dots, a_n) \in \mathbb{R}^n$, $A_n = (\sum_{k=1}^n |a_k|^\alpha)^{1/\alpha}$, $\delta_n = \max_{1 \leq k \leq n} |a_k|/A_n$. Then for $|t| \delta_n \leq 1$ we have*

$$\left| E \exp \left(it A_n^{-1} \sum_{k=1}^n a_k Z_k \right) - E \exp \left(it A_n^{-1} \sum_{k=1}^n a_k Z_k^* \right) \right| \leq |t|^\alpha \rho(\mu_1, \mu_2) \quad (17)$$

where ρ is defined by (15).

Proof. Letting φ_1, φ_2 denote the characteristic function of the Z_k 's resp. Z_k^* 's and using (14), (10) and the inequality

$$\left| \prod_{k=1}^n x_k - \prod_{k=1}^n y_k \right| \leq \sum_{k=1}^n |x_k - y_k|,$$

valid for all $|x_k| \leq 1, |y_k| \leq 1$ we get that for $|t|\delta_n \leq 1$ the left hand side of (17) equals

$$\begin{aligned} & \left| \prod_{k=1}^n \varphi_1(ta_k/A_n) - \prod_{k=1}^n \varphi_2(ta_k/A_n) \right| \leq \sum_{k=1}^n |\varphi_1(ta_k/A_n) - \varphi_2(ta_k/A_n)| \\ & \leq \sum_{k=1}^n |\beta_1(ta_k/A_n) - \beta_2(ta_k/A_n)| |ta_k/A_n|^\alpha \leq \sup_{|x| \leq |t|\delta_n} |\beta_1(x) - \beta_2(x)| \sum_{k=1}^n |ta_k/A_n|^\alpha \\ & = |t|^\alpha \sup_{|x| \leq |t|\delta_n} |\beta_1(x) - \beta_2(x)| \leq |t|^\alpha \rho(\mu_1, \mu_2). \end{aligned}$$

Remark. The proof of Lemma 2 shows that for any $t \in \mathbb{R}$ the left hand side of (17) cannot exceed $|t|^\alpha \sup_{|x| \leq |t|\delta_n} |\beta_1(x) - \beta_2(x)|$, a fact that will be useful in the sequel.

Given probability measures ν_n, ν on the Borel sets of a separable metric space (S, d) we say, as usual, that $\nu_n \xrightarrow{d} \nu$ if

$$\int_S f(x) d\nu_n(x) \longrightarrow \int_S f(x) d\nu(x) \text{ as } n \rightarrow \infty \quad (18)$$

for every bounded, real valued continuous function f on S . (18) is clearly equivalent to

$$Ef(Z_n) \longrightarrow Ef(Z) \quad (19)$$

where Z_n, Z are r.v.'s valued in (S, d) (i.e. measurable maps from some probability space to (S, d)) with distribution ν_n, ν .

Lemma 3. (see [21]). Let (S, d) be a separable metric space and let ν, ν_1, ν_2, \dots be probability measures on the Borel sets of (S, d) such that $\nu_n \xrightarrow{d} \nu$. Let \mathcal{G} be a class of real valued functions on (S, d) such that

(a) \mathcal{G} is locally equicontinuous, i.e. for every $\varepsilon > 0$ and $x \in S$ there is a $\delta = \delta(\varepsilon, x) > 0$ such that $y \in S, d(x, y) \leq \delta$ imply $|f(x) - f(y)| \leq \varepsilon$ for every $f \in \mathcal{G}$.

(b) There exists a continuous function $g \geq 0$ on S such that $|f(x)| \leq g(x)$ for all $f \in \mathcal{G}$ and $x \in S$ and

$$\int_S g(x) d\nu_n(x) \longrightarrow \int_S g(x) d\nu(x) (< \infty) \text{ as } n \rightarrow \infty. \quad (20)$$

Then

$$\int_S f(x) d\nu_n(x) \longrightarrow \int_S f(x) d\nu(x) \text{ as } n \rightarrow \infty \quad (21)$$

uniformly in $f \in \mathcal{G}$.

Assume now that (X_n) satisfies the assumptions of Theorem 2, fix $t \in \mathbb{R}$ and for any $n \geq 1, (a_1, \dots, a_n) \in \mathbb{R}^n$ let

$$\psi(a_1, \dots, a_n) = E \exp \left(itA_n^{-1} \sum_{k=1}^n a_k Y_k \right), \quad (22)$$

where $A_n = (\sum_{k=1}^n |a_k|^\alpha)^{1/\alpha}$ and (Y_k) is the sequence of r.v.'s defined before Lemma 1. We show that for any $\varepsilon > 0$ there exists a sequence $n_1 < n_2 < \dots$ of integers such that

$$(1 - \varepsilon) \psi(a_1, \dots, a_k) \leq E \exp \left(itA_k^{-1} \sum_{i=1}^k a_i X_{n_i} \right) \leq (1 + \varepsilon) \psi(a_1, \dots, a_k) \quad (23)$$

for all $k \geq 1$ and all (a_k) satisfying (10); moreover, (23) remains valid for every further subsequence of (X_{n_k}) as well. To construct n_1 we set

$$\begin{aligned} Q(\mathbf{a}, n, \ell) &= \exp(itA_\ell^{-1}(a_1X_n + a_2Y_2 + \cdots + a_\ell Y_\ell)) \\ R(\mathbf{a}, \ell) &= \exp(itA_\ell^{-1}(a_1Y_1 + a_2Y_2 + \cdots + a_\ell Y_\ell)) \end{aligned}$$

for every $n \geq 1$, $\ell \geq 2$ and $\mathbf{a} = (a_1, \dots, a_\ell) \in \mathbb{R}^\ell$. We show that

$$E \left\{ \frac{Q(\mathbf{a}, n, \ell)}{\Psi(\mathbf{a})} \right\} \longrightarrow E \left\{ \frac{R(\mathbf{a}, \ell)}{\Psi(\mathbf{a})} \right\} \quad \text{as } n \rightarrow \infty \quad \text{uniformly in } \mathbf{a}, \ell. \quad (24)$$

(The right side of (24) equals 1.) To this end we recall that, given \mathbf{X} and μ , the r.v.'s Y_1, Y_2, \dots are conditionally i.i.d. with common conditional distribution μ and thus, given \mathbf{X}, μ and Y_1 , the r.v.'s Y_2, Y_3, \dots are conditionally i.i.d. with distribution μ . Thus

$$E(Q(\mathbf{a}, n, \ell) | \mathbf{X}, \mu) = g^{\mathbf{a}, \ell}(X_n, \mu) \quad (25)$$

and

$$E(R(\mathbf{a}, \ell) | \mathbf{X}, \mu, Y_1) = g^{\mathbf{a}, \ell}(Y_1, \mu), \quad (26)$$

where

$$g^{\mathbf{a}, \ell}(u, \mathbf{v}) = E \exp \left(itA_\ell^{-1} \left(a_1 u + \sum_{i=2}^{\ell} a_i \xi_i^{(\mathbf{v})} \right) \right) \quad (u \in \mathbb{R}^1, \mathbf{v} \in S)$$

and $(\xi_n^{(\mathbf{v})})$ is an i.i.d. sequence with distribution \mathbf{v} . Integrating (25) and (26), we get

$$E(Q(\mathbf{a}, n, \ell)) = E g^{\mathbf{a}, \ell}(X_n, \mu) \quad (27)$$

$$E(R(\mathbf{a}, \ell)) = E g^{\mathbf{a}, \ell}(Y_1, \mu) \quad (28)$$

and thus (24) is equivalent to

$$E \frac{g^{\mathbf{a}, \ell}(X_n, \mu)}{\Psi(\mathbf{a})} \longrightarrow E \frac{g^{\mathbf{a}, \ell}(Y_1, \mu)}{\Psi(\mathbf{a})} \quad \text{as } n \rightarrow \infty, \quad \text{uniformly in } \mathbf{a}, \ell. \quad (29)$$

We shall derive (29) from Lemmas 1–3. Recall that ρ is a metric on $S = S(\alpha, c)$; the remarks at the beginning of this section show that on the support of μ the metric ρ and the Prohorov metric π induce the same convergence and thus the same Borel σ -field; thus the limit random measure μ , which is a random variable taking values in (S, π) , can be also regarded as a random variable taking values in (S, ρ) . Also, μ is clearly $\sigma(\mathbf{X})$ measurable and thus $(X_n, \mu) \xrightarrow{d} (Y_1, \mu)$ by Lemma 1. (Recall that by reindexing, Lemma 1 can be assumed to hold for $n_k = k$.) Hence, (29) will follow from Lemma 3 (note the equivalence of (18) and (19)) if we show that the class of functions

$$\left\{ \frac{g^{\mathbf{a}, \ell}(t, \mathbf{v})}{\Psi(\mathbf{a})} \right\} \quad (30)$$

defined on the product metric space $(\mathbb{R} \times S, \lambda \times \rho)$ (λ denotes the ordinary distance on \mathbb{R}) satisfies conditions (a),(b) of Lemma 3. To see the validity of (a) let us note that by (12), (13), Y_n are conditionally i.i.d. with respect to μ with conditional distribution μ , moreover, we assumed without loss of generality that the characteristic function $\varphi(t, \omega)$ of $\mu(\omega)$ satisfies (16) with $\lim_{t \rightarrow 0} \beta(t, \omega) = 0$ uniformly in ω and thus applying Lemma 2 with $\varphi_1(t) = \varphi(t, \omega)$ and $\varphi_2(t) = \exp(-c|t|^\alpha)$ and using (10) and the remark after the proof of Lemma 2 it follows that there exists an integer n_0 and a positive constant c_0 such that $\Psi(\mathbf{a}) \geq c_0$ for $n \geq n_0$ and all (a_k) . Thus the validity of (a) follows from Lemma 2; the validity of (b) is immediate from $|g^{\mathbf{a}, \ell}(u, \mathbf{v})| \leq 1$. We thus proved relation (29) and thus also (24), whence it follows (note again that the right side of (24) equals 1) that

$$\psi(\mathbf{a})^{-1} E \exp(itA_\ell^{-1}(a_1X_n + a_2Y_2 + \cdots + a_\ell Y_\ell)) \longrightarrow 1 \quad (31)$$

as $n \rightarrow \infty$, uniformly in ℓ, \mathbf{a} . Hence given $\varepsilon > 0$, we can choose n_1 so large that

$$\begin{aligned} & |E \exp(itA_\ell^{-1}(a_1X_n + a_2Y_2 + \cdots + a_\ell Y_\ell)) - E \exp(itA_\ell^{-1}(a_1Y_1 + a_2Y_2 + \cdots + a_\ell Y_\ell))| \\ & \leq \frac{\varepsilon}{2} \psi(a_1, \dots, a_\ell) \end{aligned} \quad (32)$$

for every ℓ, \mathbf{a} and $n \geq n_1$. This completes the first induction step.

Assume now that n_1, \dots, n_{k-1} have already been chosen. Exactly in the same way as we proved (31), it follows that for $\ell > k$

$$\begin{aligned} & \psi(\mathbf{a})^{-1} E \exp(itA_\ell^{-1}(a_1X_{n_1} + \cdots + a_{k-1}X_{n_{k-1}} + a_kX_n + a_{k+1}Y_{k+1} + \cdots + a_\ell Y_\ell)) \\ & \longrightarrow \psi(\mathbf{a})^{-1} E \exp(itA_\ell^{-1}(a_1X_{n_1} + \cdots + a_{k-1}X_{n_{k-1}} + a_kY_k + \cdots + a_\ell Y_\ell)) \quad \text{as } n \rightarrow \infty \end{aligned}$$

uniformly in \mathbf{a} and ℓ . Hence we can choose $n_k > n_{k-1}$ so large that

$$\begin{aligned} & E \exp(itA_\ell^{-1}(a_1X_{n_1} + \cdots + a_{k-1}X_{n_{k-1}} + a_kX_n + a_{k+1}Y_{k+1} + \cdots + a_\ell Y_\ell)) \\ & \quad - E \exp(itA_\ell^{-1}(a_1X_{n_1} + \cdots + a_{k-1}X_{n_{k-1}} + a_kY_k + \cdots + a_\ell Y_\ell)) \\ & \leq \frac{\varepsilon}{2^k} \psi(a_1, \dots, a_\ell) \end{aligned} \quad (33)$$

for every $(a_1, \dots, a_\ell) \in \mathcal{R}^\ell$, $\ell > k$ and $n \geq n_k$. This completes the k -th induction step; the so constructed sequence (n_k) obviously satisfies

$$\begin{aligned} & E \exp(itA_\ell^{-1}(a_1X_{n_1} + \cdots + a_\ell X_{n_\ell})) - E \exp(itA_\ell^{-1}(a_1Y_1 + \cdots + a_\ell Y_\ell)) \\ & \leq \varepsilon \psi(a_1, \dots, a_\ell) \end{aligned}$$

for every $\ell \geq 1$ and $(a_1, \dots, a_\ell) \in \mathcal{R}^\ell$, i.e. (23) is valid. Since in the k -th induction step n_k was chosen in such a way that the corresponding inequalities (32) (for $k = 1$) and (33) (for $k > 1$) hold not only for $n = n_k$, but for all $n \geq n_k$ as well, relation (23) remains valid for any further subsequence of (X_{n_k}) .

We can now easily complete the proof of Theorem 2. Letting $\psi(a_1, \dots, a_n, t)$ denote the function defined by (22), the validity of (23) for (X_{n_k}) and its further subsequences and a diagonal argument yield a subsequence (X_{n_k}) such that for all rational t and all rational $\varepsilon > 0$ we have

$$(1 - \varepsilon) \psi(a_1, \dots, a_k, t) \leq E \exp\left(itA_k^{-1} \sum_{i=1}^k a_i X_{n_i}\right) \leq (1 + \varepsilon) \psi(a_1, \dots, a_k, t) \quad (34)$$

for $k \geq k_0(t, \varepsilon)$ and all (a_n) . Recall now that without loss of generality we assumed that the characteristic function $\varphi(t, \omega)$ of $\mu(\omega)$ satisfies (16) where $\lim_{t \rightarrow 0} \beta(t, \omega) = 0$ uniformly for $\omega \in \Omega$. Applying Lemma 2 with $\varphi_1(t) = \varphi(t, \omega)$, $\varphi_2(t) = \exp(-c|t|^\alpha)$, using the Remark after the proof of the lemma and integrating with respect to ω we get

$$|\varphi(a_1, \dots, a_k, t) - \exp(-c|t|^\alpha)| \leq |t|^\alpha \beta^*(|t| \delta_k) \quad (35)$$

for all $k \geq 1$, $t \in \mathbb{R}$ and all (a_k) , where $\beta^*(t)$ is a function satisfying $\lim_{t \rightarrow 0} \beta^*(t) = 0$ and $\delta_k = \max_{1 \leq j \leq k} |a_j|/A_k$. Since $\delta_k \rightarrow 0$ by (10), relations (34) and (35) imply

$$E \exp\left(itA_k^{-1} \sum_{i=1}^k a_i X_{n_i}\right) \longrightarrow \exp(-c|t|^\alpha) \quad \text{as } k \rightarrow \infty$$

for any rational t and any (a_k) satisfying (10), and consequently

$$A_k^{-1} \sum_{i=1}^k a_i X_{n_i} \xrightarrow{d} G_{\alpha, c}.$$

This completes the proof of Theorem 2.

Acknowledgements We would like to thank two anonymous referees for their comments on the paper leading to a substantial improvement of the presentation.

References

1. D. J. ALDOUS. Limit theorems for subsequences of arbitrarily-dependent sequences of random variables, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **40** (1977), 59–82.
2. D. J. ALDOUS. Subspaces of L^1 via random measures. *Trans. Amer. Math. Soc.* **267** (1981), 445–463.
3. I. BERKES. On almost symmetric sequences in L^p . *Acta Math. Hung.* **54** (1989) 269–278.
4. I. BERKES and H. DEHLING. Almost sure and weak invariance principles for random variables attracted by a stable law. *Prob. Theory Rel. Fields* **83** (1989), 331–353.
5. I. BERKES and E. PÉTER. Exchangeable r.v.'s and the subsequence principle. *Prob. Theory Rel. Fields* **73** (1986), 395–413.
6. I. BERKES and H. P. ROSENTHAL. Almost exchangeable sequences of random variables, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **70** (1985), 473–507.
7. I. BERKES and R. TICHY. On the central limit theorem for lacunary series. Preprint.
8. S. D. CHATTERJI. A general strong law. *Invent. Math.* **9** 1969/1970 235–245.
9. S. D. CHATTERJI. A principle of subsequences in probability theory: The central limit theorem. *Adv. Math.* **13** (1974), 31–54.
10. S. D. CHATTERJI. A subsequence principle in probability theory II. The law of the iterated logarithm. *Invent. Math.* **25** (1974), 241–251.
11. S. D. CHATTERJI. Un principe de sous-suites dans la théorie des probabilités. *Séminaire des probabilités VI, Strasbourg, Lecture Notes in Mathematics* 258, pp. 72–89. Springer, 1972.
12. D. DACUNHA-CASTELLE and J.L. KRIVINE. Sous-espaces de L^1 . *C. R. Acad. Sci. Paris Sr. A-B* 280 (1975), A645–A648.
13. D.F. EGOROV. Sur les suites de fonctions mesurables. *C.R. Acad. Sci. Paris* , **152** (1911) 244–246.
14. W. FELLER. An introduction to probability theory and its applications, Vol. II. Wiley, 1971.
15. V. F. GAPOSHKIN. Lacunary series and independent functions. *Russian Math. Surveys* **21** (1966), 3–82.
16. V. F. GAPOSHKIN. Convergence and limit theorems for subsequences of random variables. (Russian) *Teor. Veroyatnost. i Primenen.* **17** (1972), 401–423.
17. S. GUERRE and Y. RAYNAUD. On sequences with no almost symmetric subsequence. *Texas Functional Analysis Seminar 1985–1986 Longhorn Notes, Univ. of Texas* pp. 83–93. Austin, 1986.
18. M.I. KADEC and W. PEŁCZYŃSKI. Bases, lacunary sequences and complemented subspaces in the spaces L_p . *Studia Math.* **21** 1961/1962, 161–176.
19. J. KOMLÓS. A generalization of a problem of Steinhaus. *Acta Math. Acad. Sci. Hungar.* **18**, 217–229, (1967)
20. J. KOMLÓS. Every sequence converging to 0 weakly in L_2 contains an unconditional convergence sequence. *Ark. Mat.* **12** (1974), 41–49.
21. R. RANGA RAO. Relations between weak and uniform convergence of measures with applications. *Ann. Math. Stat.* **33** (1962), 659–680.
22. P. RÉVÉSZ, On a problem of Steinhaus. *Acta Math. Acad. Sci. Hung.* **16** (1965), 311–318.
23. P. RÉVÉSZ, The laws of large numbers. Academic Press, 1967.
24. P. ULJANOV. Solved and unsolved problems in the theory of trigonometric and orthogonal series. *Russian Math. Surveys* **19/1** (1964), 1–62.