

Hepatocellular Nuclear Factor 4 alpha (HNF-4 alpha) activation by saRNA rescues dyslipidemia and promotes favorable metabolic profile in a high fat diet (HFD) fed rat model.

K-W. Huang^{§1,2}, **V. Reebye**^{§3*}, K. Czysz³, S. Ciriello³, S. Dorman³, I. Reccia³, H.-S. Lai^{1,2}, L. Peng⁵, N. Kostomitsopoulos⁶, J. Nicholls³, R.S. Habib⁴, D.A. Tomalia⁷, P. Sætrom^{8,9}, E. Wilkes¹⁰, P. Cutillas¹⁰, J.J. Rossi^{11*}, N.A. Habib^{3*}.

¹Department of Surgery & Hepatitis Research Center, National Taiwan University Hospital ²Graduate Institute of Clinical Medicine, National Taiwan University College of Medicine, Zhongzheng Dist., Taipei City 10002, Taiwan (R.O.C.). ³Department of Surgery and Cancer, Imperial College London, London, W12 0NN (UK). ⁴MiNA Therapeutics Limited, London (UK). ⁵Centre Interdisciplinaire de Nanoscience de Marseille, 13288 Marseille (France). ⁶Biomedical Research Animal Facilities, Biomedical Research Foundation of the Academy of Athens, 11527 Athens (Greece). ⁷Department of Chemistry, University of Pennsylvania, Philadelphia, PA (USA). ⁸Department of Cancer Research and Molecular Medicine and ⁹Department of Computer and Information Science, Norwegian University of Science and Technology, NO-7489 Trondheim, (Norway). ¹⁰Cell Signalling and Proteomics Group, Centre for Haemato-Oncology, Barts Cancer Institute, Queen Mary University of London, London EC1M 6BQ (UK). ¹¹Division of Molecular Biology, Beckman Research Institute of City of Hope, Duarte, CA 91010 (USA).

[§] Joint primary authors.

Abstract

Background:

Nonalcoholic fatty liver disease (NAFLD) culminates in insulin resistance and metabolic syndrome. As yet there is no approved single agent that targets steatosis or its pathological progression to hepatitis. Hepatocyte-nuclear-factor 4-alpha (HNF-4 α) is at the centre of a complex transcriptional network where its disruption is directly linked to diabetes and steatosis. Re-activating HNF-4 α expression in NAFLD could therefore reverse pathology.

Aim:

Small activating RNAs (saRNAs), designed synthetically for upregulating HNF-4 α expression in hepatocytes, was administered intravenously (I.V) into a rat NAFLD model. These animals were sustained on a high fat diet (HFD) for 16 weeks to induce pathology. I.V. delivery (3x injections) of saRNAs (0.6mg/kg) beginning at week 16 was performed using 5-(G₅)-triethanolamine-core poly(amidoamine) dendrimers. Treatment continued for a further two weeks (1 x injection per week) whilst the animals were maintained on HFD.

Results:

saRNA treatment caused a significant increase in HNF4A transcript levels in the liver. This was also followed by a reduction in liver triglyceride; increased HDL/LDL ratio and decreased white adipose tissue/body weight ratio. Since HNF-4 α is a key transcriptional factor in hepatocytes we analysed saRNA transfected liver cells for global phosphoproteomic changes driven by HNF4A upregulation. We observed significant beneficial changes in proteins regulating in sphingolipid metabolism, fatty acid β -oxidation, ketogenesis, detoxification of reactive oxygen species and lipid transport

Conclusions:

We demonstrate that HNF-4 α (currently an undruggable target) can be specifically activated by saRNAs when delivered using nanoparticles with high tropism to the liver. HNF4A-saRNA treatment induced a favorable metabolic profile; thus potentially representing a single agent for the treatment of NAFLD and insulin resistance.

Keywords: Oligonucleotides; small activating RNA, nanoparticles, NAFLD, HNF4A, Fatty liver disease