



University of HUDDERSFIELD

University of Huddersfield Repository

Lai, L, Harris, Emma, West, R.M and Mackie, Sarah L.

What is the absolute risk of developing diabetes mellitus in patients with glucocorticoid-treated polymyalgia rheumatica and giant cell arteritis? a systematic review and meta-analysis

Original Citation

Lai, L, Harris, Emma, West, R.M and Mackie, Sarah L. (2017) What is the absolute risk of developing diabetes mellitus in patients with glucocorticoid-treated polymyalgia rheumatica and giant cell arteritis? a systematic review and meta-analysis. *Annals of the Rheumatic Diseases*, 76 (Supp 2). p. 613. ISSN 0003-4967

This version is available at <http://eprints.hud.ac.uk/id/eprint/32512/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

Medical or Research Professionals/Clinicians

Topic area: Clinical topics by disease

Topic: 18. Vasculitis

Submission N°: EULAR17-4942

WHAT IS THE ABSOLUTE RISK OF DEVELOPING DIABETES MELLITUS IN PATIENTS WITH GLUCOCORTICOID-TREATED POLYMYALGIA RHEUMATICA AND GIANT CELL ARTERITIS? A SYSTEMATIC REVIEW AND META-ANALYSIS

L. Lai¹, E. Harris^{1,2}, R. M. West³, S. L. Mackie^{1,4}

¹Leeds Institute for Rheumatic and Musculoskeletal Medicine, University of Leeds, ²School of Human and Health Sciences, University of Huddersfield, ³Leeds Institute of Health Sciences, University of Leeds, ⁴NIHR-Leeds Musculoskeletal Biomedical Research Unit, Leeds, United Kingdom

My abstract has been or will be presented at a scientific meeting during a 12 months period prior to EULAR 2017: No

Is the first author applying for a travel bursary and/or an award for undergraduate medical students?: Yes

Is the first author of this abstract an undergraduate medical student?: No

Please confirm that you will apply for the travel bursary on the EULAR website www.congress.eular.org: Yes


Background: Polymyalgia rheumatica (PMR) and giant cell arteritis (GCA) are treated with glucocorticoids (GCs) but long-term GC use is associated with diabetes mellitus (DM). The absolute incidence of this serious complication in this patient group remains unclear.

Objectives: To quantify the absolute risk of GC-induced DM in PMR and GCA in published literature.

Methods: We identified literature from inception to February 2016 reporting diabetes following exposure to oral GC in patients with PMR and/or GCA without pre-existing diabetes. A random-effects meta-analysis was performed to summarise the literature. Risk of bias was assessed using the Cochrane Collaboration tool.

Results: 21 eligible publications were identified. In studies of patients with GCA, mean cumulative GC dose was almost two times higher than in studies of PMR (8.9g vs 5.0g), with slightly longer treatment duration but much longer duration of follow-up (8.8years vs 4.4years). The incidence proportion (cumulative incidence) of patients who developed new-onset DM was 6% (95%CI: 3-9%) for PMR and 12% (95%CI: 8-17%) for GCA. Heterogeneity between studies was high ($I^2=78.2\%$), as there were differences in study designs, patient population, geographical locations and treatment strategies. Based on UK data on incidence rate of DM in the general population¹, the expected background incidence rate of DM over 4.4 years in PMR patients and 8.8 years in GCA patients (the duration of follow-up) would be 4.8% and 9.7%, respectively. Very little information on predictors of DM in PMR or GCA patients was found. The overall risk of bias was high for many of the observational studies, especially relating to definition and recording of outcome and prognostic variables.

Image/graph:



Conclusions: Physicians should screen patients treated for PMR/GCA for DM but it remains unclear what is the time-period of greatest risk and the influence of risk factors. Our meta-analysis produced plausible estimates of DM incidence in patients with PMR and GCA but there is insufficient published data to allow precise quantification of the DM risk or, crucially, which patients are at greatest risk.

References: ¹Sharma M et al. BMJ Open, 2016. 6(1): p.e010210

Disclosure of Interest: None declared