


CORRECTION

# Correction: Alternatively Activated (M2) Macrophage Phenotype Is Inducibile by Endothelin-1 in Cultured Human Macrophages

Stefano Soldano, Carmen Pizzorni, Sabrina Paolino, Amelia Chiara Trombetta, Paola Montagna, Renata Brizzolara, Barbara Ruaro, Alberto Sulli, Maurizio Cutolo

[Fig 1](#) has an incorrectly duplicated panel. The image for the IL-4 treated CD68 cells is an incorrect duplication of the CD68 ET-1 treated cells. Please see the correct [Fig 1](#) here.




**OPEN ACCESS**

**Citation:** Soldano S, Pizzorni C, Paolino S, Trombetta AC, Montagna P, Brizzolara R, et al. (2017) Correction: Alternatively Activated (M2) Macrophage Phenotype Is Inducibile by Endothelin-1 in Cultured Human Macrophages. PLoS ONE 12 (3): e0175238. <https://doi.org/10.1371/journal.pone.0175238>

**Published:** March 30, 2017

**Copyright:** © 2017 Soldano et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Fig 1. Evaluation of CD68, CD204 and CD206 expression in cultured THP-1-derived macrophages.** Immunocytochemistry of CD68 (marker of macrophage activation), CD204 and CD206 protein expression in cultured THP-1-derived macrophages (M0 macrophages) treated for 72 hours with ET-1 (100nM) and IL-4 (10ng/mL) alone, or pre-treated with ET<sub>A/B</sub>RA (bosentan, 10<sup>-5</sup>M) for 1 hour before being stimulated with ET-1. Cultured M0 macrophages maintained for 72 hours in RPMI at 5% of FBS were used as controls (M0-controls). Immunocytochemistry was performed on four

independent *in vitro* experiments. The detection of protein expression was performed evaluating the same number of cells by light microscopy (magnification 40X).

<https://doi.org/10.1371/journal.pone.0175238.g001>

## Reference

1. Soldano S, Pizzorni C, Paolino S, Trombetta AC, Montagna P, Brizzolara R, et al. (2016) Alternatively Activated (M2) Macrophage Phenotype Is Inducible by Endothelin-1 in Cultured Human Macrophages. PLoS ONE 11(11): e0166433. doi:[10.1371/journal.pone.0166433](https://doi.org/10.1371/journal.pone.0166433) PMID: [27846260](https://pubmed.ncbi.nlm.nih.gov/27846260/)