

FORUM

Dispersal of Egyptian Vultures *Neophron percnopterus*: the first case of long-distance relocation of an individual from France to Sicily

M. Di Vittorio^{a,b}, S. Henriquet^c, E. Kobierzycki^d, L. Luiselli^e, E.M. Hema^f, L. Murabito^a, G. Rannisi^a and P. López-López^g

^aGruppo Tutela Rapaci Sicily, Italy; ^bEcologia Applicata Italia srl, Rome, Italy; ^cLigue pour la Protection des Oiseaux Provence Alpes Côte d'Azur (LPO PACA), Hyères, France; ^dCoordination technique Plan National d'Actions France Vautour percnoptère, Bruges, Belgium; ^eNiger Delta Ecology and Biodiversity Conservation Unit, Department of Applied and Environmental Biology, Rivers State University of Science and Technology, Nkpolu, Port Harcourt, Nigeria; ^fUniversité Ouaga 1 Prof Joseph Ki -Zerbo / CUPD, Laboratoire de Biologie et Écologie Animales, Ouagadougou, Burkina Faso; ^gCavanilles Institute of Biodiversity and Evolutionary Biology, University of Valencia, Valencia, Spain

ABSTRACT

Knowledge of juvenile dispersal is important for understanding population dynamics and for effective conservation, particularly of geographically isolated raptor populations. Here, we report the first documented case of a long-distance movement of an Egyptian Vulture *Neophron percnopterus* from the French population to Sicily. This observation opens a new perspective for the conservation of the small and endangered Sicilian population of this species, providing evidence that persistence of the Italian population may be aided by new input from other countries.

ARTICLE HISTORY

Received 7 September 2016
Accepted 10 October 2016

The extent of juvenile dispersal and recruitment into a breeding population as sexual maturity is reached is a critical aspect of evolutionary and conservation biology (Greenwood 1980, Paradis *et al* 1998). Dispersal, a behaviour typical of several raptor species (Newton 1979, Pearce 2007), has important implications for population genetic and demographic processes (Elorriaga *et al* 2009), including the capacity for recolonization, (meta)population persistence, and breeding success (*eg* Forero *et al* 2002, Woodroffe 2003, Pearce 2007). Hence, knowledge of juvenile dispersal is important for understanding population dynamics and for implementing effective conservation (Walter 2000, Gosling 2003, Hobson *et al* 2004, Carrete *et al* 2007, Soutullo *et al* 2008), especially for threatened and long-distance migratory species such as the Egyptian Vulture *Neophron percnopterus* (Donazar 1993).

The Egyptian Vulture, a long-lived, cliff-nesting raptor, has a breeding range extending over Eurasia and Africa (Cramp & Simmons 1980). The Italian population is critically endangered according to IUCN criteria (Rondinini *et al* 2013) and breeds in the central and southern Apennines and in Sicily (Sarà & Di Vittorio 2003). This species has declined sharply throughout its range and is endangered worldwide (BirdLife International 2016). In Sicily, the population

has decreased since 1980 and there are currently only seven breeding pairs, the largest population in Italy (Andreotti & Leonardi 2009, Di Vittorio 2011, Di Vittorio *et al* 2016).

Anthropogenic factors such as poisoning, habitat degradation, reduction in food availability, and high mortality during migration (Oppel *et al* 2015) and in the African winter quarters, are possibly the main factors in the species' decline (Grande 2006, Grande *et al* 2008, Angelov *et al* 2013). Previous studies have underlined the link between philopatric behaviour and population dynamics (*eg* Grande 2006, Carrete *et al* 2007, Grande *et al* 2008), showing that natal dispersal distances are usually short, with dispersers recruiting in areas near their natal territories (Elorriaga *et al* 2009).

Movements of juveniles between France and Spain are well known (Donazar 1993, Elorriaga *et al* 2009), as demonstrated by the presence of French Egyptian Vultures in Spanish communal roosts (Donazar 2004), and the record of an individual banded in northern Spain and nesting in France (Elorriaga *et al* 2009). Moreover, the majority of the French population travels across the Iberian Peninsula annually during pre-breeding and post-breeding migration (Elorriaga *et al* 2009, García-Ripollés *et al* 2010), and reaches Africa through the Strait of Gibraltar, while others cross further east through the Levant (Ferguson-Lees &



Figure 1. The Egyptian Vulture observed in Sicily. The ring code is zoomed upper left (photograph by G. Rannisi).

Christie 2001, Meyburg *et al* 2004). The common migration route of French and Spanish populations could facilitate the exchange of individuals between these areas (Meyburg *et al* 2004).

Individuals from distant geographic areas can play an important role in the population dynamics of small

populations of raptors as a result of the ‘rescue effect’ (*ie* population recovery by immigration of new individuals), as recently evidenced for Bonelli’s Eagle in France (Lieury *et al* 2016). From comparisons between population viability models and survival rates, Grande (2006) found that the Egyptian Vulture population in the Ebro Valley, northeastern Spain, was declining slower than expected and suggested that immigration of adults from different populations might be responsible for population maintenance. In contrast, other small populations in southern Spain are declining at similar or even higher rates than those projected by population viability models, despite being located along the migratory route to Africa through the Strait of Gibraltar (Sanz-Aguilar *et al* 2015).

Egyptian Vultures from other countries have occasionally been reported in northern Italy: an individual from département du Gard, France, was observed in Udine, northeast Italy in July 2013 at a Griffon Vulture *Gyps fulvus* feeding station (F. Genero, pers comm) and a bird from département du Vaucluse was observed in Novi Ligure, northwest Italy, in 2015. Here, we report the first documented long-distance resighting of an Egyptian Vulture from the French



Figure 2. Natal site (Verdon Canyon, France: A) and the resighting site (Nebrodi Mountain, Sicily: B).

population. On 28 May 2016, we observed and photographed on the Nebrodi Mountain (northern Sicily) a fourth-calendar-year Egyptian Vulture marked with a white ring bearing the code 4M (Figure 1). This bird was observed together with another immature individual near a feeding station for Griffon Vultures. This bird was the first one ringed in the nest on 30 July 2013 in the Verdon Canyon, southeast France, where the Egyptian Vulture returned naturally on 1999 as a consequence of the reintroduction of the Griffon Vulture. To our knowledge, this is the first long-distance resighting of an Egyptian Vulture in Sicily (950 km; Figure 2). From a conservation perspective, exploratory movements of juveniles such as this, even if rare, may play an important role in the population dynamics of this endangered species. For example, long-distance dispersal could be underestimated (eg Koenig *et al* 1996, Woodroffe 2003, Saurola & Francis 2004, Grande *et al* 2008), as has been revealed by the results of several raptor-marking projects, together with wildlife telemetry and genetic studies (eg Arsenault *et al* 2005, Le Gouar *et al* 2006, Urios *et al* 2007, Cadahía *et al* 2009).

This observation is important because it introduces a new perspective for the conservation of the small and endangered Sicilian population of Egyptian Vultures, providing the first evidence that the Italian population could receive new input from other countries to facilitate its persistence. Moreover, the presence of this species near a feeding station highlights the importance of supplementary feeding interventions for the conservation of Egyptian Vultures (see eg Sarà *et al* 2009, López-López *et al* 2014, Arrondo *et al* 2015, Lieury *et al* 2015), which is particularly important at stopover sites during migration.

Acknowledgements

We thank Gruppo Tutela Rapaci Sicilia for field support during monitoring. This project was supported by the LIFE Project ConRaSi - LIFE14 NAT/IT/001017 CUP H86J15000 240006: Conservation of Raptors in Sicily. Thanks are due to Mario Lo Valvo and Fulvio Genero for their support to the project. Ana Sanz-Aguilar, an anonymous reviewer and Chris Redfern made valuable comments on earlier versions of the manuscript. Pascual López-López is supported by a 'Juan de la Cierva-incorporación' postdoctoral grant of the Spanish Ministry of Economy and Competitiveness (IJC1-2014-19190).

References

Andreotti, A. & Leonardi, G. (2009) Piano d'azione nazionale per il Capovaccaio (*Neophron percnopterus*). *Quaderni di Conservazione della Natura* 30, 1–115.

- Angelov, I., Hashim, I. & Oppel, S. (2013) Persistent electrocution mortality of Egyptian Vultures *Neophron percnopterus* over 28 years in East Africa. *Bird Conservation International* 23, 1–6.
- Arrondo, E., Cortés-Avizanda, A. & Donazar, J.A. (2015) Temporally unpredictable supplementary feeding may benefit endangered scavengers. *Ibis* 157, 648–651.
- Arsenault, D.P., Stacey, P.B. & Hooelzer, G.A. (2005) Mark-recapture and DNA fingerprinting data reveal high breeding-site fidelity, low natal philopatry, and low levels of genetic population differentiation in Flammulated Owls (*Otus flammeolus*). *Auk* 122, 329–337.
- BirdLife International (2016) *Species factsheet*: *Neophron percnopterus*. www.birdlife.org
- Cadahía, L., López-López, P., Urios, V. & Negro, J.J. (2009) Natal dispersal and recruitment of two Bonelli's Eagles *Aquila fasciata*: a four-year satellite tracking study. *Acta Ornithologica* 44, 193–198.
- Carrete, M., Grande, J.M., Tella, J.L., Sánchez-Zapata, J.A., Donazar, J.A., Díaz-Delgado, R. & Romo, A. (2007) Habitat, human-pressure, and social behavior: partialling out factors affecting large-scale territory extinction in an endangered vulture. *Biological Conservation* 136, 143–154.
- Cramp, S. & Simmons, K.E.L. (eds) (1980) *Handbook of the Birds of Europe, the Middle East and North Africa: the birds of the western Palearctic*. Volume 2: Hawks to Bustards. Oxford University Press, Oxford.
- Di Vittorio, M. (2011) *Raptors and biodiversity in Mediterranean pseudo-steppic habitat. Final report of post-doc research activity*. Department of Environmental Biology and Biodiversity, University of Palermo, Italy.
- Di Vittorio, M., Barbera, A., Di Trapani, E., Faraone, F.P., Ciaccio, A., Sciagura, N., D'Amico, D., Giacalone, G., Zafarana, M., Greci, S. & Sarto, A. (2016) Wintering of Egyptian Vultures (*Neophron percnopterus*) in Sicily: new data. *Arxius de Miscellània Zoològica* 14, 114–116.
- Donazar, J.A. (1993) *Los buitres ibéricos; biología y conservación*. J.M. Reyero, Madrid, Spain.
- Donazar, J.A. (2004) Alimoche común *Neophron percnopterus*. In *Libro Rojo de las aves de España* (eds Madroño, A., González, C. & Atienza, J.C.), pp 129–131. Dirección General para la Biodiversidad-SEO/BirdLife, Madrid, Spain.
- Elorriaga, J., Zuberogitia, I., Castillo, I., Azkona, A., Hidalgo, S., Astorkia, L., Ruiz-Moneo, F. & Iraeta, A. (2009) First documented case of long-distance dispersal in the Egyptian Vulture (*Neophron percnopterus*). *Journal of Raptor Research* 43, 142–145.
- Ferguson-Lees, J. & Christie, D.A. (2001) *Raptors of the World*. Christopher Helm., London.
- Forero, M.G., Donazar, J.A. & Hiraldo, F. (2002) Causes and fitness consequences of natal dispersal in a population of Black Kites. *Ecology* 83, 858–872.
- García-Ripollés, C., López-López, P. & Urios, V. (2010) First description of migration and wintering of adult Egyptian Vultures *Neophron percnopterus* tracked by GPS satellite telemetry. *Bird Study* 57, 261–265.
- Gosling, L.M. (2003) Adaptive behaviour and population viability. In *Animal Behavior and Wildlife Conservation* (eds Festa-Bianchet, M. & Apollonio, M.), pp 13–30. Island Press, Washington, DC.

- Grande, J.M. (2006)** *Factores limitantes antrópicos y naturales de poblaciones de aves carroñeras: el caso del Alimoche (Neophron percnopterus) en el Valle del Ebro*. PhD dissertation, Universidad de Sevilla, Spain.
- Grande, J.M., Serrano, D., Tavecchia, G., Carrete, M., Ceballos, O., Díaz-Delgado, R., Tella, J.L. & Donazar, J.A. (2008)** Survival in a long-lived territorial migrant: effects of life-history traits and ecological conditions in wintering and breeding areas. *Oikos* **118**, 580–590.
- Greenwood, P.J. (1980)** Mating systems, philopatry and dispersal in birds and mammals. *Animal Behaviour* **28**, 1140–1162.
- Hobson, K.A., Wassenaar, L.I. & Bayne, E. (2004)** Using isotopic variance to detect long-distance dispersal and philopatry in birds: an example with Ovenbirds and American Redstarts. *Condor* **106**, 732–743.
- Koenig, W.D., Van Vuren, D. & Hooge, P. (1996)** Detectability of dispersal distances in vertebrates. *Trends in Ecology and Evolution* **11**, 514–517.
- Le Gouar, P., Rigal, F., Boisselier-Dubayle, M.C., Samadi, S., Arthur, C., Choisy, J.P., Hatzofe, O., Henriquet, S., Lécuyer, P., Tessier, C., Susic, G. & Sarrazin, F. (2006)** Genetics of restored populations of Griffon Vultures in Europe and in France. In *Proceedings of the International Conference on Conservation and Management of Vulture Populations* (eds Houston, D.C. & Piper, S.), pp 116–126. Natural History Museum of Crete and WWF-Greece, Thessaloniki, Greece.
- Lieury, N., Gallardo, M., Ponchon, C., Besnard, A. & Millon, A. (2015)** Relative contribution of local demography and immigration in the recovery of a geographically-isolated population of the endangered Egyptian Vulture. *Biological Conservation* **191**, 349–356.
- Lieury, N., Besnard, A., Ponchon, C., Ravayrol, A. & Millon, A. (2016)** Geographically isolated but demographically connected: immigration supports efficient conservation actions in the recovery of a range-margin population of the Bonelli's Eagle in France. *Biological Conservation* **195**, 272–278.
- López-López, P., García-Ripollés, C. & Urios, V. (2014)** Food predictability determines space use of endangered vultures: implications for management of supplementary feeding. *Ecological Applications* **24**, 939–949.
- Meyburg, B.-U., Gallardo, M., Meyburg, C. & Dimitrova, E. (2004)** Migrations and sojourn in Africa of Egyptian Vultures (*Neophron percnopterus*) tracked by satellite. *Journal of Ornithology* **145**, 273–280.
- Newton, I. (1979)** *Population ecology of raptors*. T. & A.D. Poyser, Berkhamsted, UK.
- Oppel, S., Dobrev, V., Arkumarev, V., Saravia, V., Bounas, A., Kret, E., Velevski, M., Stoychev, S. & Nikolov, S.C. (2015)** High juvenile mortality during migration in a declining population of a long-distance migratory raptor. *Ibis* **157**, 545–557.
- Paradis, E., Baillie, S.R., Sutherland, W.J. & Gregory, R.D. (1998)** Patterns of natal and breeding dispersal in birds. *Journal of Animal Ecology* **67**, 518–536.
- Pearce, J.M. (2007)** Philopatry: a return to origins. *Auk* **124**, 1085–1087.
- Rondinini, C., Battistoni, A., Peronace, V. & Teofili, C. (eds) (2013)** *Lista Rossa IUCN dei Vertebrati Italiani*. Comitato Italiano IUCN e Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Rome.
- Sanz-Aguilar, A., Sánchez-Zapata, J.A., Carrete, M., Benítez, J.R., Ávila, E., Arenas, R., & Donazar, J.A. (2015)** Action on multiple fronts, illegal poisoning and wind farm planning, is required to reverse the decline of the Egyptian Vulture in southern Spain. *Biological Conservation* **187**, 10–18.
- Sarà, M. & Di Vittorio, M. (2003)** Factors influencing the distribution, abundance and nest-site selection of an endangered Egyptian Vulture (*Neophron percnopterus*) population in Sicily. *Animal Conservation* **6**, 317–328.
- Sarà, M., Greci, S. & Di Vittorio, M. (2009)** Status of Egyptian Vulture (*Neophron percnopterus*) in Sicily. *Journal of Raptor Research* **43**, 66–69.
- Saurola, P. & Francis, C.M. (2004)** Estimating population dynamics and dispersal distances of owls from nationally coordinated ringing data in Finland. *Animal Biodiversity and Conservation* **27**, 403–415.
- Soutullo, Á., López-López, P. & Urios, V. (2008)** Incorporating spatial structure and stochasticity in endangered Bonelli's Eagle's population models: implications for conservation and management. *Biological Conservation* **141**, 1013–1020.
- Urios, V., Soutullo, Á., López-López, P., Cadahía, L., Limiñana, R. & Ferrer, M. (2007)** The first case of successful breeding of a Golden Eagle *Aquila chrysaetos* tracked from birth by satellite telemetry. *Acta Ornithologica* **42**, 205–209.
- Walter, J.R. (2000)** Dispersal behavior: an ornithological frontier. *Condor* **102**, 479–481.
- Woodroffe, R. (2003)** Dispersal and conservation: a behavioral perspective on metapopulation persistence. In *Animal Behavior and Wildlife Conservation* (eds Festa-Bianchet, M. & Apollonio, M.), pp 33–48. Island Press, Washington, DC.